

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Modelling drivers of mangrove propagule dispersal and restoration of abandoned shrimp farms

**D. Di Nitto¹, P. L. A. Erfemeijer^{2,5}, J. K. L. van Beek², F. Dahdouh-Guebas^{1,3},
L. Higazi¹, K. Quisthoudt¹, L. P. Jayatissa⁴, and N. Koedam¹**

¹Biocomplexity Research Focus c/o Laboratory of Plant Biology and Nature Management, Mangrove Management Group, Vrije Universiteit Brussel – VUB, Pleinlaan 2, 1050 Brussels, Belgium

²DELTA RES, P.O. Box 177, 2600 MH Delft, The Netherlands

³Laboratoire d'Écologie des Systèmes et Gestion des Ressources, Département de Biologie des Organismes, Faculté des Sciences, Université Libre de Bruxelles – ULB, CP 169, Avenue F. D. Roosevelt 50, 1050 Bruxelles, Belgium

⁴Department of Botany, University of Ruhuna, Matara, Sri Lanka

⁵Sinclair Knight Merz (SKM), P.O. Box H615, Perth WA 6008, Australia

Received: 21 December 2012 – Accepted: 7 January 2013 – Published: 28 January 2013

Correspondence to: D. Di Nitto (diana.dinitto@gmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Propagule dispersal of four mangrove species *Rhizophora mucronata*, *R. apiculata*, *Ceriops tagal* and *Avicennia officinalis* in the Pambala-Chilaw Lagoon Complex (Sri Lanka) was studied by combining a hydrodynamic model with species-specific knowledge on propagule dispersal behaviour. Propagule transport was simulated using a finite-volume advection-diffusion model to investigate the effect of dispersal vectors (tidal flow, freshwater discharge and wind), trapping agents (retention by vegetation) and seed characteristics (buoyancy) on propagule dispersal patterns. Sensitivity analysis showed that smaller propagules, like the oval-shaped propagules of *Avicennia officinalis*, dispersed over larger distances and were most sensitive to changing values of retention by mangrove vegetation compared to larger, torpedo-shaped propagules of *Rhizophora* spp. and *C. tagal*. Directional propagule dispersal in this semi-enclosed lagoon with a small tidal range was strongly concentrated towards the edges of the lagoon and channels. Short distance dispersal appeared to be the main dispersal strategy for all four studied species, with most of the propagules being retained within the vegetation. Only a small proportion (max. 5%) of propagules left the lagoon through a channel connecting the lagoon with the open sea. Wind significantly influenced dispersal distance and direction once propagules entered the lagoon or adjacent channels. Implications of these findings for mangrove restoration were tested by simulating partial removal in the model of dikes around abandoned shrimp ponds to restore tidal hydrology and facilitate natural recolonisation by mangroves. The specific location of dike removal, (with respect to the vicinity of mangroves and independently suitable hydrodynamic flows), was found to significantly affect the resultant quantities and species of inflowing of propagules and hence the potential effectiveness of natural regeneration. These results demonstrate the value of propagule dispersal modelling in guiding hydrological restoration efforts that aim to facilitate natural mangrove regeneration.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

World aquaculture production continues to grow rapidly. Mainly in Asia and Latin America, shrimp farming has emerged as a major source of employment and income for many people (Naylor et al., 2000; Rönnbäck et al., 2002). However, when unregulated, these expanding economical activities can have major negative environmental and social impacts (Patil and Krishan, 1997; Rönnbäck, 2001; Rönnbäck et al., 2002). Over the years, many aquaculture ponds have been abandoned as it appeared that insufficient attention was paid to appropriate site selection, sustainable farm design and effective regulatory and institutional support (FAO, 2002; Lewis III et al., 2002). The conversion of mangrove ecosystems into brackish aquaculture ponds has occurred in many countries which all had significant mangrove resources (Dahdouh-Guebas et al., 2002; Foell et al., 1999; Lewis III et al., 2002). Paradoxically, sustainability and productivity of fish and shrimp aquaculture is often strongly dependent on the provision of mangrove goods (e.g. fry and broodstock) as well as services (e.g. erosion control and water quality maintenance) (Beveridge et al., 1997; Kautsky et al., 2000; Rönnbäck, 1999; Rönnbäck et al., 2003).

To date there has been little coordinated research effort into pond restoration or rehabilitation (Stevenson et al., 1999). Basically there are three options: (1) rehabilitation of the pond sites into sustainable shrimp production, (2) rehabilitation into an alternative, sustainable use or (3) restoration of the environmental conditions within these ponds and the surrounding area to re-establish a productive wetland ecosystem (Bosire et al., 2008; Lewis III, 2005; Stevenson and Burbridge, 1997; Stevenson et al., 1999). Mangroves in disused pond areas can be restored since it has been reported that they can self-repair or successfully undergo secondary succession over periods of 15–30 yr (Stevenson et al., 1999). Regarding pond restoration, it is important to first restore the normal tidal flooding regime that is typically disrupted by existing farm infrastructure. Secondly, the availability of propagules or seedlings from adjacent mangrove stands is crucial for successful natural regeneration within these ponds. Only when natural

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



recovery is not initiated after alleviating the latter potential disturbances, should actual planting of propagules, collected seeds or cultivated seedlings be envisaged (Bosire et al., 2008; Lewis III, 2005; Stevenson et al., 1999). Understanding of mangrove autoecology, and more specifically the patterns of their reproduction, propagule distribution and establishment, is indispensable to pinpoint which particular stresses are inhibiting natural mangrove regeneration and to design and implement a successful mangrove restoration project (Bosire et al., 2008; Lewis III, 2005; Lewis III and Marshall, 1998; Stevenson et al., 1999). Nonetheless, mangrove restoration remains a complex matter. In the absence of studies quantifying the environmental conditions of disused ponds, there has been some debate on their re-use potential. The excavation and construction of shrimp ponds leads to soil oxidation, transforming the “potential acid sulphate soils” of a natural mangrove area into actual acid sulphate soils (Lewis III et al., 2003; Stevenson and Burbridge, 1997; Stevenson et al., 1999).

Although soil recovery is possible to some extent, it is not known to what degree moderately acid sulphate soils limit mangrove recolonization. Ecological restoration of mangroves, including the monitoring of mangrove hydrology and propagule availability, received little attention until recent years (Lewis III, 1999). Implementing an ecological engineering project to restore mangroves is however not as simple as just breaching some dikes.

In Sri Lanka, poor water quality and poor site selection are amongst some of the main causes of large scale aquaculture production failure (Jayasinghe, 1995). According to estimates by the Ministry of Fisheries of the North-western Province, more than 1200 ha of mangroves have been totally destroyed for the purpose of shrimp farming (De Silva and Jacobson, 1995). Although natural mangrove regeneration has occurred within some abandoned aquaculture ponds in the Pambala-Chilaw Lagoon Complex, most ponds remain uncolonised even after several years of abandonment (Quisthoudt, 2007).

In this study we focus on mangrove propagule dispersal within its ecological context and more specifically in view of shrimp pond restoration projects. How can

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



species-specific propagule dispersal help understand a wider framework of mangrove vegetation structure dynamics, especially when considering long distance dispersal vs. short distance dispersal? As the first aim of this study, a numerical model was constructed to simulate and study the effect of dispersal vectors (tidal flow, freshwater discharge, wind), trapping agents (retention by vegetation) and seed characteristics (buoyancy) on propagule distribution in the Pambala-Chilaw Lagoon Complex (Sri Lanka). The second aim of this study focused on the applicability of this model in view of mangrove restoration in abandoned shrimp farm areas. By modelling propagule dispersal, we examined to what extent the removal of certain parts of outer pond dikes could ensure sufficient propagule inflow from adjacent mangroves and consequently allow for natural regeneration. Propagule dispersal of 4 mangrove species *Rhizophora mucronata* Lamk., *Rhizophora apiculata* BL., *Ceriops tagal* (Perr.) C.B. Robinson and *Avicennia officinalis* L. was simulated through a combination of hydrodynamic modelling and species-specific dispersal modelling based on field data. This pioneer study is the first modelling exercise to date that applies such a combined bio-physical modelling set-up to simulate mangrove propagule dispersal. As many pioneer studies are not holistic, we acknowledge the reductionistic edge of this study regarding settlement processes and the effect of wave action on propagule dispersal. Yet, we think it is timely to explore possible propagule behaviour based on a substantial amount of field data against the backdrop of often repeated but little tested views on mangrove dispersal ecology.

2 Materials and methods

For the present study, a model was developed to simulate the transport by hydrodynamic flows of hydrochorous mangrove propagules with certain characteristics (shape, buoyancy) from the moment they are detached from their parental tree to their subsequent dispersal throughout the Pambala-Chilaw Lagoon Complex in Sri Lanka. The general methodology was derived from experiences gained in the two modeling studies

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



on the dispersal of eelgrass seeds (2008) and fish larvae (2009). This combined biophysical model set-up was performed by means of the Delft3D-modelling suite which contains several interacting modules to simulate flows, waves, sediment transport, water quality, morphological developments and ecology (Lesser et al., 2004; Roelvink and Van Banning, 1994). As this approach is new within the discipline of mangrove ecology, we provide an elaborated overview of the methodology below (Fig. 2).

2.1 Study area

The Pambala-Chilaw Lagoon Complex (Fig. 1) is situated along the west coast of Sri Lanka, near the small town “Chilaw” (07°35′48″ N, 079°47′25″ E) and within the island’s intermediate climate zone (Mueller-Dombois, 1968). The study area is about 17 km long and 4.5 km wide and is surrounded by mangrove forests and shrimp farms, many of which are currently abandoned. Flooding of these mangrove areas mainly occurs due to the heavy rainfalls during wet seasons; the micro-tidal regime limits the tidal range to $\gg 1$ m. Chilaw Lagoon is an intermittently closed tidal lagoon with a connection to open ocean through a narrow and long channel (Dutch Canal), which is joined at the most northern and southern ends of the lagoon. These entrances are temporarily open or closed depending on local sandbar formation and movement, which varies seasonally and between years (Baranasuriya, 2001). The northern entrance may be closed incidentally (usually during dry summers), while the southern entrance (Thoduwawa) is closed throughout most of the year. However, due to local economic activities, both entrances are periodically dredged to allow boat traffic and to avoid floods upstream.

There are no rivers discharging directly into Chilaw Lagoon (Fig. 1). Most freshwater influx stems from the Karabalan Oya catchment (and possibly Deduru Oya) and rainwater runoff channels discharging at regular intervals. The Karabalan Oya and the Deduru Oya have catchment areas of 596 km² and 2.647 km² and annual mean discharge in the order of 8 m³ s⁻¹ and 36 m³ s⁻¹ respectively (UNEP/GPA, 2003).

The mangrove forests within this area are typical fringe or riverine forests (Lugo and Snedaker, 1974) with an irregular distribution along a complex of creeks (Fig. 1). There

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are 16 mangrove species present in the lagoon (Jayatissa et al., 2002), of which *Rhizophora mucronata*, *Rhizophora apiculata*, *Ceriops tagal* and *Avicennia officinalis* are the most dominant in the study area. Shrimp farming has encroached into this area, expanding to 25 ha by 1998 in the Pambala region, mainly at the expense of mangrove forests (approx. 13 ha) and coconut plantations (approx. 11 ha) (Dahdouh-Guebas et al., 2002). The location of mangrove areas and shrimp farms (active and non-active) around the lagoon is shown in Fig. 1. Weather patterns in Sri Lanka, in particular the seasonal rainfall variations, are determined by a monsoon regime and coincide with a major change in the wind field together with the southward and northward shift of the Intertropical Convergence Zone (Domroes and Ranatunge, 1992). The Northeast Monsoon (December to early March) is weaker and shorter-lived than the Southwest Monsoon (mid-May to early October), which is reflected in the wind patterns respectively orientated from north to northeast and from west to south (Swan, 1983).

2.2 Field measurements

2.2.1 Input field data for the hydrodynamic model

Water level and salinity measurements were taken at several stations within the lagoon from 1 April 1998 until 29 June 1998. Time series of water levels and salinity variations were recorded respectively by means of Micro Tide pressure gauges from Coastal Leasing, USA with an absolute accuracy of ± 1.5 cm and an Aanderaa Environmental Monitoring System (EMS) of which the accuracy amounts to ± 0.1 ‰. The locations of the instruments (Fig. 1, white dots) were selected in order to specify model boundary conditions and to collect data for model calibration and validation at control points. Data were recorded at 10-min intervals. Additionally, in January–February 2007, daily salinity measurements were recorded at several points along the Pambala-Chilaw Lagoon Complex (Fig. 1, red dots).

Bathymetry was measured in the field in January–February 2007. Depth values were recorded along line transects within the lagoon, channels and creeks by means of a

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



hand-held echo sounder (deeper areas, > 0.6 m) and by means of aluminium tower ruler divided into 0.5 cm intervals (shallower areas). These measurements were corrected for tides by subtracting the corresponding heights of the tide from the daily measurements.

Data concerning precipitation, wind speed and direction were obtained from the Meteorology Department in Colombo (Sri Lanka). Daily precipitation data (mm) was available for the station in Puttalam for the period 1997–2006, while 3-hourly data regarding wind speed and direction were available from the Puttalam station for the period 2002–2007. Unfortunately, discharge measurements were not available, except (as stated above) for annual measurements of the total annual freshwater influx derived from the Irrigation Department in Sri Lanka (UNEP/GPA, 2003).

2.2.2 Input field data for the propagule dispersal model

Experiments on propagule buoyancy were conducted in situ during January–February 2007. Propagules of *Rhizophora mucronata*, *R. apiculata*, *Ceriops tagal* and *Avicennia officinalis* were collected in the field by either shaking the branches from a mangrove tree or by picking mature propagules from the tree based on the colour of the cotyledons and the “ease” by which they detach. These propagules were placed in water basins with different salinity values (35, 30, 25, 20, 10, 5 and 0‰) and tested on their buoyancy during a time span of two weeks with a total of 14 sampling times. “Buoyancy” is here defined as “floating in the water, regardless of orientation and position within the water column”, whereas “buoyancy period” is the period of time that this happens. In a microtidal mangrove area, large areas can remain unflooded for a substantial amount of time. This means that propagules can fall from their parental tree and lay on the mangrove floor to dry. The effect of drying (for 1, 2 and 3 weeks) on propagule buoyancy of *Rhizophora mucronata*, *R. apiculata* and *Ceriops tagal* was also investigated (Higazi, 2008). In addition, every three days and for a period of three weeks, successive propagule counts were conducted within 38 permanent quadrats of 4 m² in the mangrove forest near Pambala. These quadrats

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were chosen based on the environmental factors “slope”, “top soil texture” and “root complex”. For each quadrat, the number of propagules on the forest floor was recorded and the percentage coverage of the adult trees visually estimated (Higazi, 2008). These values will be used as input values for the number of propagules released at different locations within the advection-diffusion model (see Sect. 2.4.).

2.3 Hydrodynamic model

The hydrodynamic model was constructed by means of Delft3D-FLOW. Three-dimensional unsteady flow and transport phenomena resulting from tidal and meteorological forcing were simulated by solving well-established shallow-water hydrodynamic equations (Lesser et al., 2004; Stelling, 1983). The model equations, formulated in orthogonal curvilinear coordinates, were discretised onto a staggered Arakawa-C grid and time-integrated by means of an ADI (Alternating Direction Implicit) numerical scheme in horizontal directions and by Crank-Nicolson along the vertical. The latter was discretised by terrain following coordinates through σ -transformation (Leendertse, 1987; Stelling, 1983). This code was extended on the one hand with transport of salt and heat content and on the other hand with the $k-\varepsilon$ model (Launder and Spalding, 1982) for vertical exchange of horizontal momentum and matter or heat, possibly subjected to density stratification. Along the open (sea) boundaries, constituents from tidal harmonics of water level patterns were imposed. The solution of this modelling process was mass conserving at every grid cell and time step (2 min) and coupled offline to the advection-diffusion model Delft3D-WAQ. For the computation of the surface roughness, a Manning roughness coefficient, n , of $0.024 \text{ s m}^{1/3}$ was used as an input value for the calculation of the Chèzy friction coefficient, which is depth-dependent and therefore calculated each time step.

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.1 Model grid resolution and bathymetry

By means of Delft3D-RGFGRID, a model grid was developed consisting in total of 21 111 computational elements and covering the whole study area. The horizontal dimension covered grid cells with a resolution in the order of 35 m. The vertical dimension was represented by subdividing the water column into 5 layers, each representing 20 % of the water depth, following a sigma-coordinated approach to ensure sufficient vertical resolution in the near-coastal zone (Stelling and Van Kester, 1994). Run time of the hydrodynamic model for a two months – long simulation required approximately 5 h.

The 5-layered coupled communication output files, generated every hour, were subsequently aggregated vertically to 1 layer and then used as input files for the advection-diffusion model to simulate the biological transport modelling of mangrove propagules (see below). Since the propagules spend most of the dispersal phase floating on the water surface, due to their buoyance characteristics, and rapidly sink to the bottom at the end of their flotation period, aggregation to a 2-D model was considered acceptable for the purpose of this study. Test runs revealed that differences in dispersal patterns between vertically aggregated and multi-layered Delft3D-WAQ runs were negligible. Horizontal aggregation of grid resolution was not applied in any of the model runs.

For the generation and interpolation of the bathymetry, Delft3D-Quickin was used. A shapefile containing the horizontal (X, Y) and vertical (Z) coordinates of the measured depth points was generated in ArcGIS 8.3 (ESRI) and subsequently imported in Delft3D-Quickin as a sample file. All sample points were interpolated by triangulation, a method which is best suited for data sets with a resolution that is about equal to or smaller than the grid resolution. The sample points were first organised into a Delaunay network (Raper, 1990), after which grid values were interpolated.

2.3.2 Model forcing

The hydrodynamic model was forced using temporally varying meteorological data comprising of a horizontal wind velocity and direction component archived every 3 h

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from 2002–2007 at a station in Puttalam, situated north of the study area. For the two seasons, April–May 1998 (wet season) and July–August 2007 (dry season), modelled in Delft3D-FLOW, time series of wind velocity and direction were derived by averaging over the 5 consecutive years.

One discharge point was assigned at the junction of the Karabalan Oya with the southern part of the Dutch Canal, where salinity was assumed to be constant (0 ppt). Discharge rates were derived from scaling the total annual discharge according to averaged monthly precipitation values.

2.3.3 Calibration and validation of hydrodynamics

Calibration of the hydrodynamics was accomplished based on available water levels within the northern part of the Dutch Canal (Fig. 1) for the period 1 April 1998 until 29 June 1998. Several adjustments were made to the boundary conditions in order to obtain an optimal similarity between modelled and observed water level amplitudes and phases for a large number of tidal constituents, more specifically by altering the phase difference between the north and south boundary within the tidal constituents. Model validation was done by using salinity patterns recorded within the Chilaw Lagoon (Fig. 1) from 15 May 1998 until 29 June 1998. Goodness-of-fit between model results and field measurements were tested by means of two approaches: (1) r^2 (square of the correlation coefficient), and (2) bias n (normalised bias) calculated according to Jolliff et al. (2009) as follows:

$$\text{Bias } n = \text{Bias} / (\sigma d) \quad (1)$$

with Bias = Mean (model) – Mean (data) and σd = standard deviation of the data. The r^2 (square of Pearson product-moment correlation coefficient) between the salinity measured in the field vs. the salinity generated by the model was 0.85 meaning that the range and general trend of the data sets were good (Fig. 3). Bias n (normalized bias) was –0.51 which indicates that the model average deviated considerably from the field measurements. This was most likely due to the fact that accurate freshwater discharge

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data were not available for Chilaw Lagoon. However, the salinity values showed correspondence between model and field measurements throughout the simulated period, with highest and lowest values occurring in the same periods. As we do not intend to generate conclusions on absolute quantitative values of propagule dispersal, we consider the hydrodynamic model as acceptable for the purpose of the sensitivity analysis and relative interpretation of the propagule dispersal scenario modelling results.

2.4 Advection-diffusion model

Delft3D-WAQ, the water quality module of Delft3D, calculates the transport of substances by numerically solving the advection-diffusion reaction equation on a predefined computational grid and for a wide range of model substances (Postma, 1988). The Delft3D-WAQ model considers computational elements (or segments) as volumes that are linked to each other and where transport between segments are derived from dedicated models (e.g. simulated in Delft3D-FLOW). Internally, Delft3D-WAQ multiplies fluxes with concentrations to obtain masses for each time step across internal and external boundaries and includes a large selection of numerical solution schemes.

In order to simulate the propagule transport of different mangrove species, we used a generic advection-diffusion model that was originally developed to simulate the dispersal of eelgrass seeds via floating generative shoots (Erftemeijer et al., 2008). We have opted for the finite-volume method offered by Delft3D-WAQ, rather than a particle tracking method. While the latter method offers the possibility to follow particles as individuals and assign them specific properties, mangrove propagules are passive dispersal units assumed to react similarly. Since the process parameters mainly depend on their shape and buoyancy properties (orientation and position in the water column), this approach is sufficient for the purpose of this study, i.e. differentiate between species and to model the dispersal of different cohorts of propagules per species, each with their own buoyancy properties. When properly used, both finite-volume methods and particle tracking model approaches should provide comparable results (Zhang and Chen, 2007).

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4.1 Definition of processes and parameters

When simulating propagule dispersal, species-specific knowledge on buoyancy duration, longevity, period required for establishment and obligated dispersal (time taken for viable, floating propagules to develop lateral roots) are very important. Propagule dispersal characteristics of four different mangrove species (*Rhizophora mucronata*, *R. apiculata*, *Ceriops tagal* and *Avicennia officinalis*) were obtained from published data and additional experiments by our group (Table 1). Buoyancy characteristics can change with varying salinities, and could influence propagule dispersal patterns in different seasonal periods with dissimilar discharge rates.

Retention and release of propagules within mangrove vegetation is another equally important parameter with respect to hydrochorous dispersal. Retention means that propagules can be retained within the vegetation for a certain period after which they can be released again to an adjacent vegetation patch. The retention and release values implemented within this model were experimentally established and found to be different between species; smaller, oval-shaped propagules (like *A. officinalis*) having smaller retention values and higher release values than larger, torpedo-shaped propagules (like *R. mucronata*). Retention and release values were also dependent on the vegetation structure, varying from open to dense. Given the lack of published data on this complicated matter, different schemes of retention and release were applied varying from minimum and intermediate to maximum values. Finally, the effect of wind speed and direction on propagule dispersal by water currents was considered by applying a wind drag function (3 % of wind speed sensu Tsanis (1989) on surface currents in the model.

Specific discharge points were assigned to release the propagule “concentrations” at 10 different places within the mangrove areas where their parental trees were known to be present. The amount of propagules released during one run by means of Delft3D-WAQ (Table 2) was based on the successive density of available propagules counted in the field within the permanent quadrats dominated by the concerned species

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(> 50% of adult tree coverage). For each species, the average number of propagules at the beginning of the counts was listed as well as the average increase of propagules after a time span of three weeks.

2.4.2 Sensitivity analysis and seasonal variability

5 A sensitivity analysis was done to determine how “sensitive” the output of the model is to changes in parameter input values of the model, i.e. by varying the buoyancy period and the values of retention and release. Initial input values of propagule concentrations at the moment of release (T_0) were kept constant for all species.

10 In addition, seasonal variability was analysed by modelling and comparing propagule dispersal within two seasons, the wet season and the dry season respectively from April to May and July to August. This was done for all scenarios (see below). We acknowledge the seasonal variability of propagule production (Table 1) (Kumara, 2001) and the fact that the assigned release points represent a realistic situation which is however more extensive if all spatiotemporal release points would be considered. Therefore, these results were interpreted only for a relative comparison between species.

2.4.3 Scenarios

20 The main objective of this study was to investigate the effect of dispersal vectors (tidal flow, freshwater discharge, wind), trapping agents (retention by vegetation) and seed characteristics (buoyancy) on propagule dispersal. In addition, we simulated the effect of dike removal on mangrove regeneration in abandoned shrimp farm areas.

2.4.4 Scenario 1: What is the effect of species-specific buoyancy characteristics on propagule dispersal?

The obligated dispersal period, which is the time that propagules remain viable and start developing lateral roots while floating, was modelled for the four different

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mangrove species, (i.e. *Rhizophora mucronata*, *R. apiculata*, *Ceriops tagal* and *Avicennia officinalis*) based on published data on their buoyancy properties.

2.4.5 Scenario 2: What is the effect of drying of propagules on their dispersal?

The dispersal of fresh propagules vs. for 1 week, 2 weeks and 3 weeks dried propagules was modelled with the intention of examining the effect of drying-induced buoyancy changes on the dispersal extent of these propagules. The values regarding the buoyancy factor were based on the findings of the buoyancy field experiments, available for the species *Rhizophora mucronata*, *R. apiculata* and *Ceriops tagal* (Higazi, 2008).

2.4.6 Scenario 3: What is the effect of wind on propagule dispersal?

The effect of sporadic high wind velocities on maximum propagule dispersal extents was investigated for the species *Rhizophora mucronata*, *R. apiculata*, *Ceriops tagal* and *Avicennia officinalis*. Especially in large, open water bodies, like Chilaw Lagoon, high wind velocities can generate small waves in the upper layer of the water column which could strongly influence the dispersal of propagules. Within this scenario the effect of additional wind was included by implementing the maximum values per month derived from the 3-hourly data available from the Meteorological Department in Colombo, Sri Lanka. The effect of waves was not modelled.

2.4.7 Scenario 4: What is the effect of dike removal (rehabilitation of shrimp pond areas) on propagule dispersal?

Removal of parts of outer dikes of abandoned shrimp farms was simulated to assess to what extent this could assure the inflow of propagules from adjacent mangroves and facilitate natural regeneration. Observation points were added in the two main shrimp farm areas A (SA1-SA7) and B (SB1-SB5) (Fig. 1). In approach 1, part (60 m) of an outer dike was removed at the north-eastern side of shrimp farm A and the northern side of shrimp farm B, while in approach 2 part (60 m) of the outer dike at

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the south-eastern side of shrimp farm A and the north-western side of shrimp farm B was removed. The locations of dike removal were chosen based on a preliminary assessment of potentially favourable hydrodynamic flows towards these abandoned shrimp farms. Sediment transport and related geomorphologic processes after dike removal were not considered.

2.4.8 Analysis of model output

Model output of the propagule dispersal modelling for the different scenarios was post-processed in two ways:

Distribution patterns: graphic contour plots of the temporal and spatial distribution of propagules (densities per m^2).

Time series plots: propagule densities (N/m^2 , N = number) at selected observation points, accumulated over time during the modelled time series.

3 Results

3.1 Sensitivity analysis and seasonal variability

Model results (wet season) for the four species indicated that smaller propagules, like the oval-shaped propagules of *Avicennia officinalis*, were the most sensitive to changing values of retention and release (Fig. 4a). The minimum retention scheme (in combination with high release values) signified a wide dispersal extent of *A. officinalis* along both the northern and the southern parts of the lagoon. The maximum retention scheme (in combination with low release values) limited the dispersal of this species to the southernmost part of the lagoon. These results indicate that propagules of *A. officinalis* released from the southern part of the lagoon might also be important for replenishing sites along the northern part if retention is low. Irrespective of the retention scheme, part of the propagules released from adult trees situated on the lagoon banks within the northern mangrove area, were lost from the lagoon system through

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the northern part of the Dutch Canal. A small portion of these propagules (max. 5 %) had the ability to disperse along the northern mangrove area under the minimum retention scheme, while a maximum retention scheme prevented distribution of *Avicennia* into this part of the lagoon. Larger, torpedo-shaped propagules represented by the genus *Rhizophora* appeared to be less sensitive to changes in values of retention and release. All retention schemes (minimum, intermediate and maximum) restricted the dispersal extent of *R. mucronata* to the southernmost part of the lagoon (Fig. 4b). The latter was true for a varying buoyancy period of 1 week, 2 weeks and 4 weeks (results not shown).

Results for the dry season confirmed that *A. officinalis* remained the most sensitive species with respect to changes in retention and its effect on the propagule dispersal extent, while sensitivity to varying buoyancy periods was less significant for all species. While some dispersal of *A. officinalis*, *R. mucronata* and *R. apiculata* propagules occurred during the dry season, there was no significant spatial dispersal pattern for *Cerriops tagal* propagules in the model (data not shown). During dry seasons, flooding of the inner mangrove areas is rare, but genera that also occur along the lagoon and riverbanks, such as *Avicennia* and *Rhizophora*, can still disperse along water bodies without being retained within the inner mangrove vegetation. Inversely, species restricted to the northern mangrove areas and occupying a more landward position, such as *C. tagal*, are rarely inundated and therefore have no opportunity for any propagule dispersal during the dry season. *Rhizophora apiculata* and *R. mucronata* displayed dispersal patterns similar to *A. officinalis*, although less distinct in the dry season.

3.2 Scenario 1: Effect of species-specific buoyancy characteristics on propagule dispersal

The results of scenario 1 show that *Avicennia officinalis* had a more extensive propagule dispersal range in the dry season than in the wet season (Fig. 5). The minimum retention scheme of the dry season resulted in dispersal of its propagules along all parts of the lagoon as well as along all connecting channels. Differences in dispersal

patterns of *Rhizophora apiculata* and *R. mucronata* between the wet season and the dry season were less pronounced in comparison to the dispersal patterns of *A. officinalis*. *Ceriops tagal* had no propagule dispersal during the dry season yet an extensive dispersal during the wet season under conditions of minimal retention.

3.3 Scenario 2: Effect of drying of propagules on their dispersal

Figure 6 shows the model output distribution plots for *Rhizophora mucronata* when comparing the propagule dispersal extent of fresh propagules vs. propagules dried for 3 weeks. Field experiments revealed that (1) buoyancy differences mainly occurred when propagules were naturally dried for 3 weeks and (2) drying only affected *Rhizophora* species. Model results for the wet season indicated that the drying of propagules for 3 weeks of *Rhizophora mucronata* within the lagoon resulted in a more concentrated extent of dispersal near the release points (Fig. 6), while still allowing for some propagule distribution of *R. mucronata* throughout the southern part of Dutch Canal, Pol Ela and Marambettiya Ela. Similar results were obtained for *R. apiculata* (data not shown). The main difference in dispersal between fresh propagules and 3 weeks dried propagules (both species) was that a portion of the dried propagules released within the mangrove area north of the lagoon had not left the system along the northern part of the Dutch Canal. During the dry season, propagule distribution was also concentrated near the release points (results not shown).

3.4 Scenario 3: Effect of wind on propagule dispersal

“Additional wind” (high velocities) had a significant effect on the dispersal of propagules (Fig. 7 for *Rhizophora mucronata*). Irrespective of the retention scheme, distribution patterns clearly indicated that with high wind velocities, propagules of *Avicennia officinalis*, *Rhizophora apiculata* and *R. mucronata* tend to disperse more to the southern part of the study area along the channel Marambettiya Ela. With respect to seasonal variation, distribution patterns were similar, however dispersal of propagules along the

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Marambettiya Ela was less pronounced during the dry season. Only for *Cerriops tagal* differences between propagule dispersal patterns with or without the “additional” wind were negligible (results not shown).

3.5 Scenario 4: Effect of dike removal for rehabilitation of shrimp pond areas

5 A noteworthy difference was observed between the two approaches of partial outer dike removal (Fig. 8a and b). When considering propagule inflow from adjacent mangrove areas, approach 1 (north-east/north dike removal) was the most effective, as propagule concentrations within shrimp farm area A and B were much higher than those generated by approach 2 (south-east/north-west dike removal). Minimum, intermediate and maximum retention all showed comparable distribution patterns with the same order of magnitude (results not shown). Additionally, a seasonal comparison indicated that propagule inflow in shrimp farm area A and B was most efficient during the wet season when water levels rose to guarantee flooding of these areas (Fig. 8a).
10 Figure 8c shows the different concentrations of each species during the wet season when parts of the outer dikes were removed according to approach 1. As *Cerriops tagal* only occurs within the northern mangrove area, concentrations of this species were highest in shrimp farm area A, an area where *Rhizophora apiculata* and *R. mucronata* are also equally present. The most abundant species entering shrimp farm area B were *Rhizophora apiculata* and *R. mucronata*, yet the presence of *Avicennia officinalis*
15 was not insignificant, considering the relatively low initial concentrations of propagules released in the model. The extent of the propagule inflow within shrimp farm area A reached observation point SA1, though with a lower magnitude than the inflow towards observation point SB3 in shrimp farm area B (Fig. 1). Propagule dispersal towards observation points SA6 and SA7 was obstructed due to the presence of dikes of two shrimp farms that are still active. The effect of high wind velocities played a significant additional role in the dispersal of propagules during the wet season towards the designated shrimp farm areas (results not shown). Concerning shrimp farm area A, the effect of wind was variable during the time span of propagule dispersal for all species,
25

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mangrove propagules, in this study defined as the movement of propagules leaving the system, has been acknowledged by many authors (Clarke, 1993; Sauer, 1988; Sen-
gupta et al., 2005; Stieglitz and Ridd, 2001). As an example, Clarke (1993) discovered
that during one single flood tide propagules of *Avicennia marina* could disperse as far
as 500 m from their release point along tidal creeks that enter Jervis Bay (Australia).
Within the Pambala-Chilaw Lagoon complex, it appeared that hydrodynamics allows for
only a small part of the propagules (max. 5% for each species) to leave the system via
the northern part of the Dutch Canal. Few propagules of *R. apiculata* and *R. mucronata*
were indeed found along the northern sea mouth, strangled within a pile of waste along
the sides of the channel (personal observation). However, Short Distance Dispersal
(SDD) appeared to be the main dispersal strategy for all four concerned species in
this modelling exercise. The majority of propagules remained within the lagoon and
its adjacent channels, and often near the parental trees. Field experiments support
these findings as modal or average propagule movement of *Rhizophora* and even *Avi-
cennia* was found to be limited and concentrated near the parental trees, especially
within mature forest stands (Clarke and Myerscough, 1993; De Ryck, 2009; McGuin-
ness, 1997; Sousa et al., 2007). Water-buoyant propagules may set out to colonize and
establish new stands but like in most plant species, they will rather strand in the vicin-
ity of the parent trees to replenish existing stands (Duke et al., 1998; Harper, 1977;
Levin et al., 2003; Sousa et al., 2007) with a higher chance of suitable environmen-
tal conditions. Stranding and self-planting are known dispersal strategies of the family
Rhizophoraceae (Van Speybroeck, 1992). Stranding of these propagules does not per
se imply long distance dispersal, as dispersion can occur in the vicinity of the parental
mangrove trees. Self-planting on the other hand entails that a propagule falls from the
parental tree with the possibility to self-plant underneath. Nevertheless, questions con-
cerning the ecological advantage of long distance vs. short distance dispersal arise.
As the sensitivity analysis showed, propagules of a typical pioneer species *Avicennia
officinalis*, could reach areas further afield if their obligated dispersal period (ODP),
more specifically the time taken for viable propagules to start developing lateral roots,

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

would exceed 2/3 weeks. The ODP of *R. apiculata* and *R. mucronata* provides chances for colonization of new areas, yet LDD of these species is inhibited by retention within the vegetation. Furthermore, the effect of drying of propagules, especially in areas with a small tidal range, concentrates propagule distribution near their release points and can further inhibit LDD. Longevity, the period required for establishment and the period of obligated dispersal are therefore vital factors to determine the ability of propagules to survive dispersal both locally and across large expanses of ocean (Drexler, 2001). A comparison between *Rhizophora* spp. species by Drexler (2001) pointed out that *Rhizophora mucronata* propagules are better equipped for LDD, yet have lower rate of survival concerning establishment than propagules of *R. apiculata*, which in their turn have a shorter longevity.

Few studies on the hydrochory of propagules have addressed the effect of wind on dispersal patterns. Results of the present study indicate that, irrespective of the retention scheme, wind can have a significant influence on their dispersal distance and direction once propagules enter the lagoon or adjacent channels. The effect of wind on propagule dispersal was generalized for all concerned species in the present study. However, differences in size, weight and shape may further alter wind-induced dispersal patterns of the different species. The effect of wave action on propagule dispersal, which was not included in this study, also deserves attention in future studies.

Although this pioneer modelling exercise showed promising results with respect to propagule dispersal processes, it is still in its early stages. We therefore emphasize the importance of additional field experiments to quantify the dispersal distances and directions of each species in different environmental settings (see Sousa et al., 2007), as well as propagule retention by vegetation (see Chang et al., 2008).

4.2 Implications of propagule dispersal for shrimp pond rehabilitation

Tidal flooding regime and propagule availability are key issues when restoring mangroves in abandoned shrimp pond areas (Lewis III, 2005). Our results indicate that, irrespective to the retention schemes, a computer-based ecological engineering project

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



can provide valuable information on the most suitable locations of propagule inflow (approach 1) through simulating the removal of parts of outer pond dikes. Inflow of propagules of different species clearly depends on the location of dike removal, suitable hydrodynamic flows and on the presence of these particular species in adjacent mangrove stands. The latter was shown in this study for *Ceriops tagal*, which appeared more likely to colonize abandoned shrimp farm areas in the northern part of the Chilaw lagoon. Given the extent of this species present distribution within the lagoon and its inner mangrove character, propagule dispersal to abandoned shrimp farm areas in southern parts is limited. Wind velocity played an additional role influencing the dispersal of propagules towards the designated shrimp farm areas.

Favourable dispersal patterns alone do not guarantee successful establishment and persistence of these species within the disused shrimp farms. Prior to seedling establishment, processes such as dispersal towards the concerning shrimp farm areas and propagule predation are likely to influence initial patterns of distribution and abundance (Cannicci et al., 2008; Dahdouh-Guebas et al., 1997; Dahdouh-Guebas et al., 1998; McGuinness, 1994; McKee, 1995; Osborne and Smith, 1990). Once arrived in the abandoned shrimp farms, other factors like physico-chemical characteristics of the sediment (Delgado et al., 2001), predation (Cannicci et al., 2008), acid sulphate soils (Sammut and Hanafi, 2000), interspecific competition and frequency of inundation (Kitaya et al., 2002) may further affect the success of establishment, early growth and survival of seedlings and ultimately determine the success of natural regeneration.

Several researchers have recently focussed on the potential role of mangroves as purifiers of effluents and sediment derived from shrimp aquaculture ponds (Costanzo et al., 2004; Jackson et al., 2003; Shimoda et al., 2005), but research results concerning seedling growth and interspecific competition within abandoned ponds are scarce. Rajendran and Katherisan (1996) studied the effect of effluent from a shrimp pond on growth in terms of shoot dry weight of 5 mangrove species (*Avicennia marina* (Forsk.) Vierh., *A. officinalis*, *Ceriops decandra* (Griff.) Ding Hou, *Rhizophora mucronata* and *R. apiculata*). Raw effluents had a negative effect on the shoot dry weight

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the species *R. apiculata* and *C. decandra* while effluents diluted by 70% improved the shoot biomass production of all mangrove seedlings. General studies concerning the effects of soil moisture, salinity and sediment accretion on propagule establishment could give more insights on the potential for natural regeneration within abandoned shrimp farms at Pambala-Chilaw Lagoon. Desiccated soils within these shrimp farms may prove to be unsuitable for propagule establishment and in addition, possible sediment accretion during flooding may cause further stresses. Survival of planted propagules of *Ceriops tagal* was found to be correlated with soil moisture and salinity and was lower in cleared areas than in small light gaps within a north Australian mangrove (McGuinness, 1997). Mortality of *R. apiculata* seedlings was found to be closely related to soil hardness (Komiyama et al., 1998). These findings suggest that the indirect effects of light on soil conditions may be more critical than its direct effects on the plants themselves. *Rhizophora apiculata* seedlings appear to be inefficient colonizers of coastal areas exposed to sudden events of high (> 4 cm) sediment accretion (Terrados et al., 1997; Thampanya et al., 2002). Furthermore, competition with non-mangrove species could pose an additional limiting factor for mangrove regeneration in the abandoned shrimp farms, unless they function as a trap for propagule recruitment.

Despite these constraints, we have in situ observed some degree of natural regeneration within a few abandoned ponds of shrimp farm area B in favour of *Rhizophora* spp. Case studies elsewhere (southern Thailand) also indicate that the potential exists for converting abandoned shrimp ponds areas back to mangroves within a period of about 5 to 10 yr provided that there is sufficient recruitment of viable propagules and hydrological conditions are restored (Lewis III et al., 2002).

While seed ecology is a well developed field, the understanding of processes driving propagule dispersal has not been advancing. On the contrary, it remained stuck in speculation even in spite of its importance towards socially and ecologically induced pond abandonment. Studies on mangrove ecology indicate that a wide variety of factors, including propagule buoyancy, tolerance to salinity, desiccation, disturbance, stochastic events, competition and predation, may affect the distribution and

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



abundance of species. Regarding mangrove establishment, there are however few studies that examine the relative importance of these different factors for individual species, but also for the interaction between these species in different environmental settings. Pond restoration projects therefore require insight on both propagule dispersal towards and establishment within “stressful” areas, or knowledge on means to proactively diminish these stresses in order to stimulate natural regeneration.

Acknowledgements. We thank the people of the Small Fishers Federation Lanka, especially Chamara Jayasooriya for his assistance in the field. Many thanks are also due to E. M. S. Wijeratne from the National Aquatic Resources Research and Development Agency in Colombo (Sri Lanka) for providing the water level measurements and to Firmijn Zijl from Deltares for his guidance during the set-up of the hydrodynamic model. This research was funded by the Flemish Interuniversity Council (VLIR) the Fonds David and Alice Van Buuren and D. D. has a VLIR PhD Scholarship. This work was in part presented at the International Interdisciplinary Conference “Hydropredict 2008” (20–24 September, Czech Republic, Prague) and in part at the Meeting on Mangrove ecology, functioning and Management – MMM3 (2–6 July 2012, Sri Lanka, Galle).

References

- Baranasuriya, P. W.: Hydrographic investigations for the design of an anchorage in a complex lagoon estuary, A Spatial Odyssey: 42nd Australian Surveyors Congress, 2001.
- Beveridge, M. C. M., Phillips, M. J., and Macintosh, D. J.: Aquaculture and the environment: the supply of and demand for environmental goods and services by Asian aquaculture and the implications for sustainability, *Aquac. Res.*, 28, 797–807, 1997.
- Bosire, J. O., Dahdouh-Guebas, F., Walton, M., Crona, B. I., Lewis, R. R., Field, C., Kairo, J. G., and Koedam, N.: Functionality of restored mangroves: A review, *Aquat. Bot.*, 89, 251–259, 2008.
- Cannicci, S., Burrows, D., Fratini, S., Smith, T. J., Offenberg, J., and Dahdouh-Guebas, F.: Faunal impact on vegetation structure and ecosystem function in mangrove forests: A review, *Aquat. Bot.*, 89, 186–200, 2008.

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Chang, E. R., Veeneklaas, R. M., Buitenwerf, R., Bakker, J. P., and Bouma, T. J.: To move or not to move: determinants of seed retention in a tidal marsh, *Funct. Ecol.*, 22, 720–727, 2008.
- Clarke, P. J.: Dispersal of grey mangrove (*Avicennia marina*) propagules in Southeastern Australia, *Aquat. Bot.*, 45, 195–204, 1993.
- 5 Clarke, P. J. and Myerscough, P. J.: The intertidal distribution of the grey mangrove (*Avicennia marina*) in southeastern Australia: the effects of physical conditions, interspecific competition, and predation on propagule establishment and survival, *Aust. J. Ecol.*, 18, 307–315, 1993.
- Clarke, P. J., Kerrigan, R. A., and Westphal, C. J.: Dispersal potential and early growth in 14 tropical mangroves: do early life history traits correlate with patterns of adult distribution?, *J. Ecol.*, 89, 648–659, 2001.
- 10 Costanzo, S. D., O'Donohue, M. J., and Dennison, W. C.: Assessing the influence and distribution of shrimp pond effluent in a tidal mangrove creek in north-east Australia, *Mar. Pollut. Bull.*, 48, 514–525, 2004.
- Dahdouh-Guebas, F., Verneirt, M., Tack, J. F., and Koedam, N.: Food preferences of *Neosar-matium meinerti* de Man (Decapoda: Sesarinae) and its possible effect on the regeneration of mangroves, *Hydrobiologia*, 347, 83–89, 1997.
- 15 Dahdouh-Guebas, F., Verneirt, M., Tack, J. F., Van Speybroeck, D., and Koedam, N.: Propagule predators in Kenyan mangroves and their possible effect on regeneration, *Mar. Freshwater Res.*, 49, 345–350, 1998.
- 20 Dahdouh-Guebas, F., Verheyden, A., De Genst, W., Hettiarachchi, S., and Koedam, N.: Four decade vegetation dynamics in Sri Lankan mangroves as detected from sequential aerial photography: A case study in Galle, *Bull. Mar. Sci.*, 67, 741–759, 2000.
- Dahdouh-Guebas, F., Zetterström, T., Rönnbäck, P., Troell, M., Wickramasinghe, A., and Koedam, N.: Recent changes in land-use in the Pambala-Chilaw Lagoon complex (Sri Lanka) investigated using remote sensing and GIS: conservation of mangroves vs. development of shrimp farming, *Environment, Development and Sustainability*, 4, 185–200, 2002.
- 25 De Ryck, D.: Moving and settling: Experiments on the dispersal and establishment of hydrochorous propagules, Master Thesis, Vrije Universiteit Brussel, Brussels, 2009.
- De Silva, R. I. and Jacobson, S.: Mundel Lake (Sri Lanka): An important habitat for migratory shorebirds, Published by the Wildlife and Nature Protection Society of Sri Lanka, 1995.
- 30 Delgado, P., Hensel, P. F., Jimenez, J. A., and Day, J. W.: The importance of propagule establishment and physical factors in mangrove distributional patterns in a Costa Rican estuary, *Aquat. Bot.*, 71, 157–178, 2001.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Domroes, M. and Ranatunge, E.: The orthogonal structure of Monsoon rainfall variation over Sri Lanka, *Theor. Appl. Climatol.*, 46, 109–114, 1992.
- Drexler, J. Z.: Maximum longevities of *Rhizophora apiculata* and *R. mucronata* propagules, *Pac. Sci.*, 55, 17–22, 2001.
- 5 Duke, N. C., Ball, M. C., and Ellison, J. C.: Factors influencing biodiversity and distributional gradients in mangroves, *Global Ecol. Biogeogr. Lett.*, 7, 27–47, 1998.
- Erftemeijer, P. L. A., van Beek, J. K. L., Ochieng, C. A., Jager, Z., and Los, H. J.: Eelgrass seed dispersal via floating generative shoots in the Dutch Wadden Sea: a model approach, *Mar. Ecol. Prog. Ser.*, 358, 115–124, 2008.
- 10 Erftemeijer, P. L. A., van Beek, J. K. L., Bolle, L. J., Dickey-Collas, M., and Los, H. F. J.: Variability in transport of fish eggs and larvae, I, Modelling the effects of coastal reclamation, *Mar. Ecol. Prog. Ser.*, 390, 167–181, 2009.
- FAO: Review of the state of world aquaculture, FAO Fisheries Circular No. 886, Rev. 2, Rome, FAO, 2003.
- 15 Foell, J., Harrison, E., and Stirrat, R. L.: Participatory approaches to natural resource management – the case of coastal zone management in the Puttalam District. Summary findings of DFID-funded research “participatory mechanisms for sustainable development of coastal ecosystems”, Project R6977, School of African and Asian studies, Falmer, Brighton, UK, University of Sussex, 1999.
- 20 Harper, J. L.: *Population Biology of Plants*, Academic Press, London, 1977.
- Higazi, L.: De verspreiding en vestiging van mangrove-propagulen in het micro-getijdengebied Pambala-Chilaw Lagune (Sri Lanka) – een experimentele aanpak, MSc. thesis, Vrije Universiteit Brussel, Brussels, 2008.
- Honnay, O., Verhaeghe, W., and Hermy, M.: Plant community assembly along dendritic networks of small forest streams, *Ecology and Freshwater Fish*, 82, 1691–1702, 2001.
- 25 Jackson, C. J., Preston, N., Burford, M. A., and Thompson, P. J.: Managing the development of sustainable shrimp farming in Australia: the role of sedimentation ponds in treatment of farm discharge water, *Aquaculture*, 226, 23–34, 2003.
- Jansson, R., Zinko, U., Merritt, D. M., and Nilsson, C.: Hydrochory increases riparian plant species richness: a comparison between a free-flowing and a regulated river, *J. Ecol.*, 93, 1094–1103, 2005.
- 30 Jayasinghe, J. M. P. K.: Sri Lanka, FAO/NACA 1995, Regional Study and Workshop on the Environmental Assessment and Management of Aquaculture Development (TCPIRAS12253),

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bangkok, Thailand: NACA Environment and Aquaculture Development Series No. I, Network of Aquaculture Centres in Asia-Pacific, 357–376, 1995.

Jayatissa, L. P., Dahdouh-Guebas, F., and Koedam, N.: A review of the floral composition and distribution of mangroves in Sri Lanka, *Bot. J. Linn. Soc.*, 138, 29–43, 2002.

5 Jolliff, J. K., Kindle, J. C., Shulman, I., Penta, B., Friedrichs, M. A. M., Helber, R., and Arnone, R. A.: Summary diagrams for coupled hydrodynamic-ecosystem model skill assessment, *J. Mar. Sy.*, 76, 64–82, 2009.

Kautsky, N., Ronnback, P., Tedengren, M., and Troell, M.: Ecosystem perspectives on management of disease in shrimp pond farming, *Aquaculture*, 191, 145–161, 2000.

10 Kitaya, Y., Jintana, V., Piriyaoytha, S., Jaijing, D., Yabuki, K., Izutani, S., Nishimiya, A., and Iwasaki, M.: Early growth of seven mangrove species planted at different elevations in a Thai estuary, *Trees-Struct. Funct.*, 16, 150–154, 2002.

Komiyama, A., Tanapermpool, P., Havanond, S., Maknual, C., Patanaponpaiboon, P., Sumida, A., Ohnishi, T., and Kato, S.: Mortality and growth of cut pieces of viviparous mangrove (*Rhizophora apiculata* and *R. mucronata*) seedlings in the field condition, *Forest Ecol. Manag.*, 112, 227–231, 1998.

Kumara, G. W. N. T.: A study of some biological aspects pertaining to conservation and sustainable use of Rhizophoraceae mangroves in Sri Lanka with particular emphasis on phenology, Msc. Thesis, University of Ruhuna, Sri Lanka, 2001.

20 Launder, B. E. and Spalding, D. B.: *Mathematical Models of Turbulence*, New York, Academic Press, 1982.

Leendertse, J. J.: A three-dimensional alternating direction implicit model with iterative fourth order dissipative non-linear advection terms WD-3333-NETH, Rijkswaterstaat, 1987.

Lesser, G. R., Roelvink, J. A., Van Kester, J. A. T. M., and Stelling, G. S.: Development and validation of a three-dimensional morphological model, *Coast. Eng.*, 51, 883–915, 2004.

25 Levin, S. A., Muller-Landau, H. C., Nathan, R., and Chave, J.: The ecology and evolution of seed dispersal: a theoretical perspective, *Annu. Rev. Ecol., Evol. Syst.*, 34, 575–604, 2003.

Lewis III, R. R.: Key concepts in successful ecological restoration of mangrove forests, 12–32, *Proceedings of the TCE-Workshop No. II, Coastal Environmental Improvement in Mangrove/Wetland Ecosystems*, 18–23 August 1998, Ranong, Thailand, 1999, Danish, S. E.: Asian Collaboration in Tropical Coastal Ecosystems (TCE) Research and Training, NACA, P.O. Box 1040, Bangkok, Thailand 10903, Ecological engineering for successful management and restoration of mangrove forests, *Ecol. Eng.*, 24, 403–418, 2005.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lewis III, R. R. and Marshall, M. J.: Principles of successful restoration of shrimp aquaculture ponds back to mangrove forests, 327 in World Aquaculture Society Book of Abstracts, Aquaculture '98, Las Vegas, Nevada, Abstract, 1998.
- Lewis III, R. R., Erfteemeijer, P. L. A., Sayaka, A., and Kethkaew, P.: Mangrove rehabilitation after shrimp aquaculture: A case study in progress at the Don Sak National Forest Reserve, Surat, Thani, Southern Thailand, Case Study 13, Annexes to the Thematic Review on Coastal Wetland Habitats and Shrimp in: Aquaculture, edited by: Macintosh, D. J., Phillips, M. J., Lewis III, R. R., and Clough, B., Case Studies 7-13, Report prepared under the World Bank, NACA, WWF and FAO Consortium Program on Shrimp Farming and the Environment, 108–128, 2002.
- Lewis III, R. R., Phillips, M. J., Clough, B., Macintosh, D. J.: Thematic Review on Coastal Wetland Habitats and Shrimp Aquaculture, Report prepared under the World Bank, NACA, WWF and FAO Consortium Program on Shrimp Farming and the Environment, Published by the Consortium, Report , 2003.
- Lugo, A. E., Snedaker, S. C.: The ecology of mangroves, *Annu. Rev. Ecol. Syst.*, 5, 39–64, 1974.
- McGuinness, K. A.: The climbing behaviour of *Cerithidea anticipata* (Mollusca: Gastropoda): the roles of physical and biological factors, *Aust. J. Ecol.*, 19, 283–289, 1994.
- McGuinness, K. A.: Dispersal, establishment and survival of *Cerriops tagal* propagules in a north Australian mangrove forest, *Oecologia*, 109, 80–87, 1997.
- McKee, K. L.: Seedling recruitment patterns in a Belizean mangrove forest – Effects of establishment ability and physicochemical Factors, *Oecologia* 101, 448–460, 1995.
- Mueller-Dombois, D.: Ecogeographic analysis of a climate map of Ceylon with particular reference to vegetation, Including climate diagram map of Ceylon at 1:506 880, *The Ceylon Forester*, 8, 39–58, 1968.
- Naylor, R. L., Goldburg, R. J., Primavera, J. H., Kautsky, N., Beveridge, M. C. M., Clay, J., Folke, C., Lubchenco, J., Mooney, H., and Troell, M.: Effect of aquaculture on world fish supplies, *Nature*, 405, 1017–1024, 2000.
- Nilsson, C., Andersson, E., Merritt, D. M., and Johansson, M. E.: Differences in riparian flora between riverbanks and river lakeshores explained by dispersal traits, *Ecology and Freshwater Fish*, 83, 2878–2887, 2002.
- Osborne, K. and Smith, T. J. I.: Differential predation on mangrove propagules in open and closed canopy forest habitats, *Vegetatio*, 89, 1–6, 1990.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Patil, P. G. and Krishan, M.: The Kandaleru shrimp farming industry and its impact on the rural economy, *Agricultural Economics Research Review*, 10, 293–308, 1997.
- Postma, L.: DELWAQ Users manual, Version 3.0: WL |Delft Hydraulics, 1988.
- Quisthoudt, K.: Mangrove en garnalenkweek in de Lagune van Chilaw (Sri Lanka): Toestand en perspectieven verlaten garnaalkwekerijen Vrije Universiteit Brussel, Brussels, Belgium, 2007.
- Rabinowitz, D.: Dispersal properties of mangrove propagules, *Biotropica*, 10, 47–57, 1978.
- Rajendran, N. and Kathiresan, K.: Effect of effluent from a shrimp pond on shoot biomass of mangrove seedlings, *Aquac. Res.*, 27, 745–757, 1996.
- Raper, J.: *Three-dimensional Applications in Geographical Information Systems*, London, UK, Taylor & Francis, 1990.
- Roelvink, J. A. and Van Banning, G. K. F. M.: Design and development of DELFT3D and application to coastal morphodynamics, *Proc. Hydroinformatics*, Verwey, Minns, 1994.
- Rönnbäck, P.: The ecological basis for economic value of seafood production supported by mangrove ecosystems, *Ecological Economics* 29, 235–252, 2001, Shrimp aquaculture, State of the art, Swedish International Development Agency (SIDA), Stockholm & Swedish EIA Centre, Swedish University of Agricultural Sciences (SLU), Uppsala, Sweden. Report, 1999.
- Rönnbäck, P., Bryceson, I., and Kautsky, N.: Coastal Aquaculture development in eastern Africa and the Western Indian Ocean: Prospects and problems for food security and local economies, *Ambio*, 31, 537–542, 2002.
- Rönnbäck, P., Troell, M., Zetterstrom, T., and Babu, D. E.: Mangrove dependence and socio-economic concerns in shrimp hatcheries of Andhra Pradesh, India, *Environ. Conserv.*, 30, 344–352, 2003.
- Sammut, J. and Hanafi, A.: Remediation and Management of Shrimp Ponds Excavated in Acid Sulphate Soils, Section in Report to World Bank, FAO, NACA and WWF, Bangkok, Thailand, 2000.
- Sauer, J. D.: *Plant migration: The dynamics of geographic patterning in seed plant species*, Berkeley: University of California Press, 1988.
- Sengupta, R., Middleton, B., Yan, C., Zuro, M., and Hartman, H.: Landscape characteristics of *Rhizophora mangle* forests and propagule deposition in coastal environments of Florida (USA), *Landscape Ecol.*, 20, 63–72, 2005.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Shimoda, T., Srithong, C., and Aryuthaka, C.: Attempt at purification of effluent and sediment in shrimp aquaculture ponds using mangrove trees, *Jarq-Jpn. Agr. Res. Q.*, 39, 139–145, 2005.
- 5 Sousa, W. P., Kennedy, P. G., Mitchell, B. J., and Ordonez, B. M.: Supply-side ecology in mangroves: Do propagule dispersal and seedling establishment explain forest structure?, *Ecol. Monogr.*, 77, 53–76, 2007.
- Stelling, G. S.: On the construction of computational methods for shallow water flow problems, PhD thesis, Delft University of Technology, Delft, The Netherlands, 1983.
- 10 Stelling, G. S. and Van Kester, T. J. A.: On the approximation of horizontal gradients in sigma coordinates for bathymetry with steep bottom slopes, *Int. J. Numer. Meth. Fl.*, 18, 915–935, 1994.
- Stevenson, N. J. and Burbridge, P. R.: Abandoned Shrimp Ponds: Options for Mangrove Rehabilitation, *International Newsletter of Coastal Management, Special Edition 1*, 1997.
- 15 Stevenson, N. J., Lewis, R. R., and Burbridge, P. R.: Disused Shrimp Ponds and Mangrove Rehabilitation, in: *An International Perspective on Wetland Rehabilitation*, edited by: Streever, W., 277–297, 1999.
- Stieglitz, T. and Ridd, P. V.: Trapping of mangrove propagules due to density-driven secondary circulation in the Normanby River estuary, NE Australia, *Mar. Ecol. Prog. Ser.*, 211, 131–142, 2001.
- 20 Swan, B.: *An Introduction to the Coastal Geomorphology of Sri Lanka*, Colombo, UNESCO, 1983.
- Terrados, J., Thampanya, U., Srichai, N., Kheowongsri, P., Geertz-Hansen, O., Boromtharath, S., Panapitukkul, N., and Duarte, C. M.: The effect of increased sediment accretion on the survival and growth of *Rhizophora apiculata* seedlings, *Estuar. Coast. Shelf Sci.*, 45, 697–701, 1997.
- 25 Thampanya, U., Vermaat, J. E., and Terrados, J.: The effect of increasing sediment accretion on the seedlings of three common Thai mangrove species, *Aquat. Bot.*, 74, 315–325, 2002.
- Tsanis, I. K.: Simulation of wind-induced water currents, *J. Hydraul. Eng.-ASCE*, 115, 1113–1134, 1989.
- 30 UNEP/GPA: *Study on Better Practices for Shrimp Farming in Chilaw and Puttlam districts of Sri Lanka*, Study prepared by Small Fishers Federation (Nirodhawardane HDLU, Thilak CR, Rajapaksha RMBU, P. Munasinghe), The Hague, The Netherlands, 2003.

Van Speybroeck, D.: Regeneration strategy of mangroves along the Kenya coast: a first approach, *Hydrobiologia*, 247, 243–251, 1992.

Zhang, Z. and Chen, Q.: Comparison of the eulerian and lagrangian methods for predicting particle transport in enclosed spaces, *Atmos. Environ.*, 41, 5236–5248, 2007.

BGD

10, 1267–1312, 2013

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 1. Propagule dispersal characteristics of the species *Ceriops tagal*, *Rhizophora mucronata*, *R. apiculata* and *Avicennia officinalis* (e.g. Clarke, 1993; Clarke et al., 2001; Drexler, 2001; Rabinowitz, 1978). S = salt water, F = fresh water, na = data not available.

Species	<i>Ceriops tagal</i>	<i>Rhizophora mucronata</i>	<i>Rhizophora apiculata</i>	<i>Avicennia officinalis</i>
Family	Rhizophoraceae	Rhizophoraceae	Rhizophoraceae	Acanthaceae
Vivipary	Viviparous	Viviparous	Viviparous	Cryptoviviparous
Propagule type	Seedling	Seedling	Seedling	Seedling
Shape	Torpedo-shaped	Torpedo-shaped	Torpedo-shaped	Oval-shaped
Dispersal part	Hypocotyl	Hypocotyl	Hypocotyl	Cotyledon
Mean fresh mass of propagules (g) (SE)	11.70 (2.40)	48.40 (14.51)	20.60 (5.10)	1.10 (0.11)
Mean length of propagules (cm)	30.00 (3.50)	39.90 (6.62)	24.20 (2.80)	1.83 (0.05)
Predominant buoyancy pattern in saltwater	Floater	Floater	Floater	Floater
Predominant buoyancy pattern in freshwater	Floater	Floater	Floater	Floater (after initially sinking)
Obligated dispersal	na	±40 days	±40 days	7–21 days
Time until root initiation (days), averaged over all salinity treatments	Starting from ±14 days	15–40 days	15–40 days	7–10 days
% of propagules with roots after 23 days (SE)	8.3 (3.3)	na	na	100 (0.00)
Longevity (days)	na	150	89	110 days S Longer F
Flowering (Sri Lanka, Negombo)	April–June, September–December	February–March, June–August	April–June	March–July
Fruiting (falling of propagules) (Sri Lanka, Negombo)	June–August, December	July–August	June–October	April–September

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Table 2. Input values of propagule “concentrations” in Delft3D-WAQ based on the successive propagule counts. Given the nature of the advection-diffusion model, the input requires the numbers of propagules to be converted into concentrations (g m^{-3}).

Genera	Rhizophora	Ceriops	Avicennia
Average number of propagules present at the start of the propagule counts	54	75	12
Concentration of propagules as initial input value in WAQ (g m^{-3} per s, time step = 15 min)	0.0600	0.0833	0.0133
Average increase of the number of propagules during the propagule counts	60	61	14
Concentration of propagule increase as input value in WAQ (g m^{-3} per s, time step = 15 min)	0.0667	0.0678	0.0156

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. (A) Representation of Sri Lanka (Dahdouh-Guebas et al., 2000), (B) an IKONOS image (2002) showing the Pambala-Chilaw Lagoon Complex., (C) Indication of the water level and salinity measurements points (both in 1998 and 2007), the observation points in Delft3D-WAQ within shrimp farm area A and B, the active and non-active shrimp farms, the dense and open mangrove vegetation and the channels. Locations of outer dike removal (approach 1 and 2) are also indicated. EMS = Environmental Monitoring System. DC = Dutch Canal, PE = Pol Ela, ME = Marambettiya Ela, BE = Bate Ela.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.




Fig. 2. Overview of the applied methodology.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



SENSITIVITY ANALYSIS

Different schemes of retention, buoyancy period of 1 week
Avicennia officinalis




Fig. 4a. Results of the sensitivity analysis: distribution patterns for the species *Avicennia officinalis* when varying the retention schemes from minimum, intermediate to maximum retention values. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

SENSITIVITY ANALYSIS

Different schemes of retention, buoyancy period of 1 week
Rhizophora mucronata




Fig. 4b. Results of the sensitivity analysis: distribution patterns for the species *Rhizophora mucronata* when varying the retention schemes from minimum, intermediate to maximum retention values. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Scenario 1: 'What is the effect of species-specific buoyancy characteristics on propagule dispersal?'

A) *Avicennia officinalis*




Fig. 5a. Results of scenario 1 (effect of species-specific buoyancy) showing the distribution plots of *Avicennia officinalis*. WS = wet season, DS = dry season, ODP = obligated dispersal period. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Scenario 1: 'What is the effect of species-specific buoyancy characteristics on propagule dispersal?'

B) *Ceriops tagal*




Fig. 5b. Results of scenario 1 (effect of species-specific buoyancy) showing the distribution plots of *Ceriops tagal*. WS = wet season, DS = dry season, ODP = obligated dispersal period. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Scenario 1: 'What is the effect of species-specific buoyancy characteristics on propagule dispersal?'

C) *Rhizophora apiculata*




Fig. 5c. Results of scenario 1 (effect of species-specific buoyancy) showing the distribution plots of *Rhizophora apiculata*. WS = wet season, DS = dry season, ODP = obligated dispersal period. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Scenario 2: 'What is the effect of drying of propagules on their dispersal?'

Buoyancy period of 2 weeks Intermediate retention scheme

WET SEASON

A) *Rhizophora mucronata*




Fig. 6a. Results of scenario 2 (effect of drying) showing the distribution plots of *Rhizophora mucronata* in the wet season. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Scenario 2: 'What is the effect of drying of propagules on their dispersal?'

Buoyancy period of 2 weeks Intermediate retention scheme

DRY SEASON

B) *Rhizophora apiculata*




Fig. 6b. Results of scenario 2 (effect of drying) showing the distribution plots of *Rhizophora apiculata* in the dry season. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Results of scenario 3 (the effect of “additional wind”) showing the distribution plots of propagules of *Avicennia officinalis* and *Rhizophora mucronata*. Scale is applicable to all distribution patterns. Red, yellow and light blue colours indicate a high, medium and low concentration of propagules respectively.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modelling drivers of mangrove propagule dispersal

D. Di Nitto et al.

Scenario 4: 'What is the effect of dike removal (shrimp pond rehabilitation) on propagule dispersal?'




Fig. 8. Results of scenario 4 (the effect of dike removal) showing the time-series plots of *Avicennia officinalis*, *Ceriops tagal* and *Rhizophora apiculata* and *Rhizophora mucronata*. A comparison was made between **(A)** seasons (wet vs. dry season), **(B)** approaches of outer dike removal (In approach 1 60 m of an outer dike was removed at the north-eastern side of shrimp farm A and the northern side of shrimp farm B, while in approach 2 part 60 m of the outer dike at the south-eastern side of shrimp farm A and the north-western side of shrimp farm B was removed) and **(C)** the different species. ObsP Ax = observation point in shrimp farm area A, ObsP Bx = observation point in shrimp farm area B. X-axes = buoyancy period (days), Y-axes = propagule density m^{-2} .