

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Comparing soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau¹, M. J. B. Swallow¹, C. E. Prescott², S. J. Grayston², and S.-W. Oh³

¹Department of Renewable Resources, University of Alberta, 442 Earth Sciences Building, Edmonton, AB T6G 2E3, Canada

²Department of Forest and Conservation Sciences, Faculty of Forestry, University of British Columbia, Vancouver, BC V6T 1Z4, Canada

³Department of Chemistry, Mokpo National University, Muan, Chonnam 534-729, Republic of Korea

Received: 27 December 2012 – Accepted: 21 April 2013 – Published: 30 April 2013

Correspondence to: S. A. Quideau (sylvie.quideau@ualberta.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Emulating the variability that exists in the natural landscape prior to disturbance should be a goal of soil reconstruction and land reclamation efforts following resource extraction. Long-term ecosystem sustainability within reclaimed landscapes can only be achieved with the re-establishment of biogeochemical processes between reconstructed soils and plants. In this study, we assessed key soil biogeochemical attributes (nutrient availability, organic matter composition, and microbial communities) in reconstructed, novel, anthropogenic ecosystems covering different reclamation treatments following open-cast mining for oil extraction. We compared the attributes to those present in a range of natural soils representative of mature boreal forest ecosystems in the same area of northern Alberta. Soil nutrient availability was determined in situ with resin probes, organic matter composition was described with ^{13}C nuclear magnetic resonance spectroscopy and soil microbial community structure was characterized using phospholipid fatty acid analysis. Significant differences among natural ecosystems were apparent in nutrient availability and seemed more related to the dominant tree cover than to soil type. When analyzed together, all natural forests differed significantly from the novel ecosystems, in particular with respect to soil organic matter composition. However, there was some overlap between the reconstructed soils and some of the natural ecosystems in nutrient availability and microbial communities, but not in organic matter characteristics. Hence, our results illustrate the importance of considering the range of natural landscape variability, and including several soil biogeochemical attributes when comparing novel, anthropogenic ecosystems to the mature ecosystems that constitute ecological targets.

1 Introduction

Processes of soil formation do not vary randomly across natural landscapes. Instead, their spatial pattern is determined by the combined actions of defined soil-forming

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of biogeochemical fluxes (Prescott et al., 2000a). Factors influencing forest floor quality include the overstory and understory vegetation growing on site, stand age, and mineral parent material (Lamarche et al., 2004). Boreal and tundra ecosystems have tightly regulated internal biogeochemical cycling between plants and soils (Bashkin, 2003), and their soil organic matter composition, microbial communities, and plant communities are interlinked (Ohtonen and Väre, 1998; Hannam et al., 2004, 2006; Eskelinen et al., 2009). The equilibrium between soil chemical and biochemical properties that exists in undisturbed soils may be disrupted following disturbance (Chaer et al., 2009). Prior research in the Athabasca oil sands region demonstrated that organic matter composition, nutrient availability, and microbial communities in these reclaimed soils differed from mature upland forest soils (Turcotte et al., 2009; Rowland et al., 2009; Dimitriu et al., 2010). These past studies looked at each of these variables separately, and did not specifically examine the inherent variability present in the natural landscape.

The objective of this study was to determine how soil biogeochemical functioning of novel ecosystems reconstructed following oil sands mining compared to the boreal forests that constitute targets for restoration. We addressed this objective by concurrently assessing several key soil attributes that we used as surrogates of ecosystem biogeochemical functioning, namely: soil nutrient availability, organic matter composition, and microbial communities. We examined how variable these attributes were in natural ecosystems, and investigated potential links amongst them. One of the ecological goals of soil reclamation is to emulate the variability that exists in the natural landscape prior to disturbance. Hence, we were interested in testing whether the soils that had developed on any of the reclaimed sites were similar in any of the relevant measurable attributes to natural soils across the natural range of variability that exists in this region.

2 Materials and methods

2.1 Study sites

Climate in the Athabasca oil sands region presents sharp seasonal contrasts. Winters are long and cold, with a mean monthly air temperature for January of -19°C (Beckingham and Archibald, 1996). Summers are short, but warm, with a mean monthly air temperature for July of 17°C , an estimated frost-free period (0°C) of 60 to 70 days, and a plant growing season averaging between 90 and 100 days (Crown and Twardy, 1970). Mean annual precipitation is 478 mm, and falls predominantly as rain (342 mm) during the summer months. Till, fluvial and lacustrine plains were deposited in the Fort McMurray Region from the advance and retreat of glacial ice and subsequent postglacial sorting following the last Wisconsin glaciation from 70 000 to 10 000 yr ago (Crown and Twardy, 1970). These surficial deposits constitute the geological parent materials in which soils from the experimental area have developed. Organic soils are found in depressional and level landscape positions where surface waters accumulate, and generally contain the decomposed remains of sphagnum and feather mosses, Labrador tea and black spruce (*Picea mariana* (Mill) BSP). Dystric brunisolic soils (Soil Classification Working Group, 1998) are well drained and have developed in acidic, coarse-textured, glaciofluvial and eolian materials. Gray Luvisols are well to moderately well drained and have developed from till and glaciolacustrine sediments (Crown and Twardy, 1970).

A total of 41 sites were chosen from the network of soil and vegetation long-term monitoring plots established in the Athabasca oil sands region in 2000 (OSVR, 1998). All sites are located within an 86 km radius around Ft. McMurray, Alberta, Canada ($56^{\circ}43' \text{N}$ $111^{\circ}21' \text{W}$). Fifteen undisturbed sites were selected to span the range of natural variability of forested ecosystems within the region (Table 1). According to the Northern Alberta Ecosite classification system (Beckingham and Archibald, 1996), sites increase in nutrient and water content from ecosite a1 to ecosite d3 (Table 1). Ecosites a and b may be water deficient for periods of the year, occur on brunisolic soils and exhibit either a canopy dominated by jack pine (*Pinus banksiana* Lamb.)

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

for the a1 ecosites, or stands mixed with aspen (*Populus tremuloides* Michx.) and conifers (ecosites b1 and b3). Ecosites d occur on luvisolic soils and are defined by their mesic moisture regime and medium nutrient regime (Beckingham and Archibald, 1996). Changes in canopy composition along a natural successional trajectory entail a shift from pure aspen stands (d1), to mixed stands of aspen and white spruce (*Picea glauca* (Moench) for the d2 ecosites, and finally to stands of pure white spruce (d3).

Land reclamation practices following open-cast mining involve placement of several layers of various soil materials, geological deposits, and products of the mining activity in order to reconstruct soils. For this study, a total of 26 sites were selected to capture the five main reclamation practices that have been used in the Oil Sands region (Table 1). Overburden (OB), which is the geological substrate (> 3 m depth) removed to access the mineable oil sands, may encompass lean oil sands (< 10 % oil), as well as Cretaceous and Pleistocene deposits. Tailings sand (TS), the residue remaining once bitumen extraction is complete, is primarily composed of a medium to fine sand (> 90 % sand). Salvaged mineral soils and surficial geological deposits (< 3 m depth) are referred to as “secondary” materials (2°). Organic amendments applied to cover these mineral substrates are referred to as “peat mineral mixes” (PM) as they consist of a mixture of peat materials and their underlying mineral substrates salvaged from low land areas. Finally, the reclamation practice of “direct placement” (DP) refers to the use of surficial peat and mineral materials (< 1 m) salvaged from adjacent areas and placed directly on the reclaimed sites; i.e., with no stockpiling. Following soil reconstruction, sites are typically sown with barley (*Hordeum vulgare* L.) during the first year to control erosion, and are subsequently planted to nursery-grown white spruce, aspen, and jack pine.

2.2 Field sampling and laboratory analysis

At each of the 41 study sites (15 natural and 26 reclaimed), one composite representative sample was obtained in July 2005 by randomly collecting ten mineral soil samples (0–10 cm) within 1–2 m from the perimeter edge of 10 m by 40 m plots that had been

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

previously established (OSVR, 1998). All samples were transported in an ice-box to the laboratory, and sieved (< 4 mm) to exclude the fresh litter materials corresponding to the L horizon (Soil Classification Working Group, 1998). Samples destined to organic matter characterization were oven-dried at 60°C , while samples for microbial characterization were kept frozen at -86°C until they were freeze-dried in preparation for analysis.

In-situ nutrient bioavailability was estimated by incubating plant root simulator (PRSTM) probes (Western Ag Innovations Inc., Saskatoon, SK), which include resin membranes to capture nutrient anions and cations from the soil solution. Four pairs of cation and anion probes were incubated randomly at 5 to 10 cm depth within each site. Following incubation, the probes were washed with deionized water, and returned to Western Ag Innovations Inc. for elution with 0.5M HCl and nutrient concentration analysis, including colorimetric ammonium (NH_4^+), nitrate (NO_3) and phosphate (PO_4^{3-}) analysis on a segmented flow Autoanalyzer III (Bran and Lubbe, Inc., Buffalo, NY), and P, K, S, Ca, Mg, Al, Fe, Mn, Cu, Zn, B, and Pb analysis by ICP spectroscopy (PerkinElmer Optima 3000-DV, PerkinElmer Inc., Shelton, CT). The probe data reported here (mmol per 10 cm^{-2} of resin area per day) correspond to a total burial period of 95 days (Rowland et al., 2009).

For the characterization of soil microbial communities using phospholipid fatty acid (PLFA) analysis, polar lipids were extracted from freeze-dried samples using a modified Bligh and Dyer protocol (Hannam et al., 2007; Dimitriu et al., 2010). Extracts were purified on pre-packed silicic acid columns (Agilent Technologies, Wilmington, DE, USA) to separate polar lipids from neutral lipids and glycolipids, and subsequently subjected to mild alkaline methanolysis. The resulting fatty-acid methyl esters were separated using an Agilent 6890 Series capillary gas chromatograph (Agilent Technologies, Wilmington, DE, USA) equipped with a 25 m Ultra 2 (5 %-phenyl)-methylpolysiloxane column using hydrogen as the carrier gas. Individual peaks were identified and quantified (nmol g^{-1}) using the MIDI peak identification software (MIDI Inc., Newark, DE, USA). The standardized $X : Y @ Z$ nomenclature for fatty acids was used to identify PLFAs, where X is

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the number of carbon atoms, Y is the number of double bonds, and Z is the position of the first double bond from the aliphatic end (ω) of the molecule. Prefixes “ i ” and “ a ” indicate branching at the second and third carbon atom, respectively, from the ω end; the suffix “ c ” corresponds to a c transfiguration; and 2OH or 3OH indicates an hydroxyl group at the second or third carbon from the carboxyl end of the molecule, respectively.

Samples for organic matter characterization were isolated using a combination of density separation and sieving to yield > 53 mm low-density materials (Turcotte et al., 2009). The chemical composition of these materials was analyzed using ramped-cross-polarisation (RAMP-CP) ^{13}C NMR spectroscopy on a Bruker Avance 400 ($B = 9.4\text{T}$, $\nu_L(^{13}\text{C}) = 100.6$ MHz) spectrometer, using a 13-kHz spinning frequency, a 1H 90° pulse width of 4.0 ms, a 1 ms contact time and a 5 s pulse delay. Four to eight thousand scans were acquired for each sample. The ^{13}C chemical shifts were referenced relative to TMS ($\delta_{\text{iso}} = 0.0$ ppm), and the Bruker’s WIN-NMR package was used to estimate the relative integrated areas (% of total spectral area) of the following five regions between 0 and 192 ppm: 0–45 ppm region-, attributed to alkyl carbons; 45–112 ppm, attributed to O-alkyl carbons; 112–140 ppm, attributed to aromatic carbons; 140–165 ppm, attributed to phenolic carbons; and 140–192 ppm, attributed to carbonyl carbons.

2.3 Statistical analysis

Due to the multivariate nature of the dataset, we used non-metric multi-dimensional scaling (NMS) ordinations to identify potential patterns in the soil bioavailable nutrients, organic matter composition, and microbial community composition among natural sites and reclamation treatments. This ordination technique, which presents the advantage of not requiring normal distribution, organizes complex datasets in a reduced dimensional space (typically two dimensions) as to reveal similarities or dissimilarities in the original dataset structure. All analyses were conducted using PCORD software (Version 5, MjM Software Design, Gleneden Beach, OR, USA). In terms of nutrient bioavailability, a total of 14 ions was used in the ordination, while five spectral NMR areas were included for organic matter characterization. Analysis of the microbial

communities used all PLFAs with < 20 carbons, which corresponded to a total of 80 PLFAs. Prior to the NMS analysis, all PLFAs were proportionalized on a mol% basis and were arcsine square-root transformed.

Data groupings were tested for significant differences in the NMS analysis using a multi-response permutation procedure (MRPP). Analyses were completed using the Sorensen distance, and Bonferroni corrections were used to control family error rate for among grouping comparisons. In addition, indicator species analyses were performed using the data groupings shown to be different by MRPP. This statistical method generates an indicator value based on the abundance and frequency of a particular variable (individual nutrient, NMR spectral area, or PLFA) in a given data grouping. The statistical significance of the indicator value was tested against a randomized Monte Carlo test.

3 Results

3.1 Biogeochemical attributes in natural boreal forest soils

For the natural forest soils (Fig. 1), three distinct ecosite groups were discriminated (Fig. 1). All ordinations attained a very low stress (< 10 in all cases) after 40–70 iterations. Solutions for the nutrients and organic matter variables were two-dimensional and the two axes explained > 95% of the data variance. Analysis of the microbial communities led to a three-dimensional solution, where the two axes presented on Fig. 1 accounted for 78% of the variance. Analysis of the PRS-probe data (nutrients) showed a clear separation of three significantly different groups: group 1, which included the a1 and b1 ecosites; group 2, corresponding to the d1 ecosites; and group 3, which included the b3, d2, and d3 ecosites. Preliminary analyses had indicated that the individual ecosite types contained in either group 1 or group 3 were not significantly different from one another.

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Separation of these three groups was weaker but still significant for the PLFA data (microbial communities) as evidenced by their larger p value than the nutrients (Table 2). Similarly, the T value obtained from the MRPP analysis, which is an index of the separation among groups, and the A value, which indicates within group homogeneity, were smaller for the microbial communities (-4.5 and 0.15 , respectively) than for the nutrients (-7.3 and 0.51). The individual groups 1, 2, and 3 which, for the PRS data (nutrients), formed three distinct clusters clearly separated on the NMDS graph (Fig. 1), yielded fairly large T values (≥ 4) in all pair-wise comparisons (Table 2). The T values were smaller, and the p values larger, for the PLFA data set, particularly comparing groups 1 and 3. This weaker pair-wise separation is also noticeable on Fig. 1 there is some overlap among cluster points for the different groups derived from the PLFA ordination. For organic matter composition (^{13}C NMR), separation among the three groups was much weaker than for either the nutrients or microbial communities, as illustrated with by their larger p value (0.07) and smaller T and A values (Table 2).

The differences among the natural sites in nutrients and microbial communities were linked to their respective tree covers (pine, aspen, and/or spruce) rather than to the associated soil types. Indeed, the b1 and b3 ecosites, which both occurred on brunisolic soils belonged to two different groups, while group 3 encompassed both Brunisols and Luvisols (Table 1). Group 1 was characterized by a cover of pine trees, either in pure stands (a1 ecosites) or in mixed stands (b1 ecosites); group 2 corresponded to pure aspen stands (d1 ecosites), and group 3 was characterized by the presence of white spruce, either in pure stands (d3 ecosites) or in mixed stands (b3 and d2 ecosites).

Cations (Ca, Mg, K, and Na) were the only nutrients that were found to be significantly different among the three ecosite groupings (i.e., having a p value ≤ 0.05 and an indicator value > 30) in the indicator species analysis (Table 3). The d1 ecosites, which support pure aspen stands, were different from the other two groups in that they had greater available Ca and Mg. The indicator species analysis also detected seven significant PLFAs, with six of the seven indicative of the d1 ecosites. No significant indicators of ecosite types were found in the NMR data.

3.2 Comparison of novel and natural ecosystem attributes

The novel ecosystems failed to show any differentiation among the different reclamation treatments (Fig. 2). As opposed to the natural ecosites, which separated according to tree cover, the novel ecosystems also failed to show any separation by vegetation (Table 2). As illustrated by higher p values (0.25–0.53), separation by vegetation was even weaker than by reclamation treatment, with associated p values ranging from 0.09 to 0.15 for the three biogeochemical criteria (soil nutrients, organic matter, and microbial communities). There was no correlative relationship between the age of the novel ecosystems (Table 1) and the biogeochemical variables (data not shown).

As the novel ecosystems did not differ from one another, we compare them as a group (i.e.; group 4) to the three distinct natural ecosite groups (Fig. 2). Separation between the natural and novel sites was most marked for the NMR data, as seen from the largest T and A values, and the smallest p value (Table 2). In all cases, the separation of natural and novel sites was very strong, and a low stress (< 9) was obtained after 48–70 iterations for all attributes. As was the case for the natural sites only (Fig. 1), solutions for the nutrients and organic matter variables as presented on Fig. 2 were two-dimensional and $> 95\%$ of the data variance was explained by the two axes. In the case of the microbial communities, the solution was three-dimensional, and the two axes presented on Fig. 2 accounted for 55% of the data variance.

The novel ecosystems (group 4) strongly differed from all natural site groups in organic matter composition, as demonstrated by extremely small p values for all pair-wise comparisons with group 4 (Table 2). The pattern for both the microbial communities and nutrients was different in that group 2 (d1 ecosites) was much closer to the reclaimed sites than either group 1 or group 3. In the case of the nutrients, group 2 was enclosed within the larger point cluster composed of group 4 sites (Fig. 2), and the corresponding p value exceeded 0.05 (Table 2); the same relationship, albeit weaker with a marginally significant p value (0.02) between groups 2 and 4, was observed for

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the microbial communities (Fig. 2 and Table 2). For both variables, the reconstructed soils were furthest away from group 1 (a1 and b1 ecosites).

Indicator species analysis identified a few additional nutrients and PLFAs that were representative of the novel ecosystems (Tables 3 and 4). The reconstructed soils were characterized by high sulphur and nitrate concentrations, and the presence of two monounsaturated PLFAs (16:1 ω 11*c* and 17:1 ω 9*c*).

4 Discussion

4.1 Environmental controls on natural variability

For the natural soils, vegetation cover was the main factor influencing separation among the different ecological units. The influence of tree species on key biogeochemical processes has been previously documented in western Canadian forests, including influences on nutrient availability (Prescott et al., 2000c; Prescott and Vesterdal, 2005; Jerabkova et al., 2006), microbial communities (Leckie et al., 2004; Grayston and Prescott, 2005; Hannam et al., 2006), and organic matter characteristics (Hannam et al., 2004). In our study, the importance of the coniferous component in mixed stands was obvious, as the presence of either spruce or pine trees determined how boreal forest soils grouped in terms of nutrient availability. The unique influence of white spruce on forest floor microbial communities has been previously reported in mixed-wood forests from northwestern Alberta, which are comparable to the forests studied here (Hannam et al., 2006; Swallow et al., 2009).

The relative importance of biotic versus abiotic determinants of soil biogeochemical processes is difficult to tease apart. Biotic factors such as forest vegetation type can be closely linked to factors such as soil organic matter composition (Quideau et al., 2001; Merilä et al., 2010), microbial community composition (Prescott and Grayston, 2013) and nutrient availability (Binkley and Giardina, 1998). Other studies have shown that abiotic conditions including climate and geology can have a stronger effect than

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

forest canopy composition on forest floor nutrient concentrations and soil humus type (Lamarche et al., 2004; Ponge et al., 2011). The issue of scale may explain these apparent discrepancies. On a regional scale, climate conditions were the controlling factor influencing soil microbial communities (Brockett et al., 2012) and humus forms (Ponge et al., 2011), while at landscape and local scales, differences in soil biogeochemical processes were more directly derived from differences in tree species (Hannam et al., 2004; Grayston and Prescott, 2005). In our study, base cations (Ca, Mg, K) were the only nutrients identified as indicative of specific ecosite groupings (Table 3). Although cation availability in the long-term is controlled by mineral weathering from the soil parent material, plant uptake and biocycling mechanisms can influence availability especially in the forest floor and upper mineral soil (Quideau et al., 1996). In our study, pure aspen stands were associated with higher Ca and Mg concentrations. This is consistent with other studies reporting higher base cation concentrations under broadleaved tree species including aspen (Prescott et al., 2004; Hobbie et al., 2006).

In the present study, differences among mature forest soils were quite clear for nutrient availability, and were partially mirrored by differences in microbial communities (Fig. 1 and Table 2). Below-ground communities and soil biodiversity are often assumed to be directly correlated to above-ground biotic factors, including vegetation composition (Wardle et al., 2004; Wall et al., 2010). Further, the regulation of carbon sources in boreal forests by heterotrophic soil microorganisms is considered to be closely associated with organic matter quality (Merilä et al., 2010), with microbial activity typically being limited by the availability of carbon rather than other nutrients (Ekblad and Nordgren, 2002). However, in our study, the patterns of organic matter composition were much weaker than patterns observed for nutrient availability and microbial communities. It may be that solid-state NMR, while providing an overall fingerprint of organic matter composition is not a good proxy for carbon availability. It may also be that organic matter characteristics as probed by NMR, are not as readily quantifiable as nutrient availability and PLFA data, and so do not show strong relationships with other factors. In all cases, it was evident that soil nutrient availability and microbial

communities were linked to above-ground vegetation. It seems that the soils underneath a pure aspen canopy (group 2) were different from the rest in that high Ca and Mg concentrations and different microbial communities were associated with this group (Tables 3 and 4). While none of the identified PLFAs were specific biomarkers of bacterial groups or fungi, three PLFAs were straight-chain hydroxy acids, which are common components of Gram-negative lipopolysaccharides (Ratledge and Wilkinson, 1988).

4.2 Novel vs. natural ecosystems

The novel ecosystems differed from the natural boreal forests in the three biogeochemical criteria measured, although there was overlap for some of these sites in terms of nutrient availability. Soil organic matter composition, while displaying a relatively weak separation among natural ecosites, differed the most between the novel and the natural ecosystems. Microbial communities in the reconstructed soils differed significantly from all natural ecosite groupings, albeit less so than did organic matter. Reconstructed soils were characterized by the presence of two unsaturated PLFAs, one of which (16 : 1@11) has been reported to increase in concentration following litter amendment (Esperschütz et al., 2011). Characterization of organic matter and microbial communities, as well as the interrelationship between them has often been used as simple measurements of reclamation success (e.g.; Insam and Dosch, 1988; Banning et al., 2008; Hahn and Quideau, 2012). The chemical and physical characteristics of soil organic matter have been proposed as sensitive indicators related to changing environmental conditions (Leifeld and Kögel-Knabner, 2005; Turcotte et al., 2009), while the reestablishment of soil microbial communities similar to undisturbed environments has been recognized as a crucial target in mine restoration (Mummey et al., 2002; Machulla et al., 2005; Harris, 2009).

Although the reconstructed soils examined do not fully resemble the natural forest soils, there is some evidence that they may, with time and soil development, come to better emulate natural soils. Based on plant community composition and soil nutrient availability, Rowland et al. (2009) concluded that some of the reclamation treatments

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

appear to develop into functioning forest ecosystems similar to those that occur naturally in this area after about 25 yr. Studies of chronosequences of reconstructed soils under similar tree canopies detected a measurable evolution of organic matter composition with time (Sorenson et al., 2011). However, temporal changes in the soil microbial communities of the same novel ecosystems have been harder to establish (Dimitriu et al., 2010; Hahn and Quideau, 2012), suggesting that, at least in this boreal environment, soil microbial response may be lagging behind changes in organic matter composition and nutrient availability.

Novel ecosystems were characterized by higher N and S concentrations than the natural sites (Table 3). This is consistent with the higher nitrate concentrations previously reported for oil sands reconstructed soils compared to the natural forest soils (McMillan et al., 2007; Rowland et al., 2009), and is likely due to the close proximity of the novel ecosystems to industrial emissions and greater atmospheric N deposition (Bytnerowicz et al., 2010; Hemsley, 2012). In all cases, the novel ecosystems (group 4) and the d1 ecosites (group 2) significantly differed in organic matter but had comparable nutrient availability (Fig. 2 and Table 2). This implies a disconnection between organic matter composition and nutrient availability in these soils. Disruption of the equilibrium between soil organic matter and biological activity may provide an index to quantify the amplitude of soil degradation (Chaer et al., 2009). This elegant paradigm is worth pursuing in future studies of novel ecosystems.

5 Implications for reconstructed boreal ecosystems

Biogeochemical processes affecting soils occur at vastly different time scales (Duchaufour, 1960; Targulian and Krasilnikov, 2007). Effects of long-term pedogenesis on boreal forest soils may be more limited than in other forest biomes as they are relatively young soils with low mean annual temperatures that limit the rate of biological and pedogenic processes. For instance, in most Canadian soils, the extent of mineral weathering and clay alteration is minimal (Komada, 1979), since the prevalent pedogenic processes

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

occurring in boreal soils; i.e., podzolization and lessivage, typically require several millennia to result in diagnostic soil horizons (e.g.; Parsons and Herriman, 1976; Barrett and Scaetzi, 1992). In boreal forest ecosystems, biogeochemical cycling between trees and soils is shallower than in other settings, and may be more closely associated to the forest floor than to the underlying mineral soils (Prescott et al., 2000b; Bashkin, 2003). It is therefore important that efficient organic matter-associated processes are re-established in novel boreal ecosystems, perhaps even more so than in other more temperate forest biomes. Results from this study, which showed some divergence between nutrient availability, microbial communities, and organic matter composition illustrate the importance of including several biogeochemical criteria when assessing novel boreal ecosystems.

According to the Society of Ecological Restoration (SER, 2004), a restored ecosystem should exhibit similar diversity and community structure as reference (target) sites (Ruiz-Jaen and Aide, 2005). This concept was originally developed with above-ground communities in mind but should be extended to include belowground communities (Harris, 2009; Wall et al., 2010). In our study, reconstructed soils differed from natural soils in all three measured biogeochemical attributes (nutrient availability, organic matter composition, and microbial communities), and so cannot be considered to be restored ecosystems (according to the SER criteria) at this stage of development. While there is evidence that the reclaimed soils will share many attributes of natural soils within a few decades, it is unlikely that they will fully emulate natural soil horization, which develops over thousands of years. As pointed out by Chadzon (2008), novel forests emerging on human modified landscapes cannot be expected to match the composition and structure of the original forest cover. In a similar fashion, reconstructed soils cannot be expected to fully emulate natural soils. Reconstructed soils in this region may therefore represent a case of novel ecosystems with novel species and novel abiotic conditions as defined by Hobbs et al. (2009). It may be more effective and realistic to determine if the reconstructed soils are able to maintain comparable ecosystem services and overall functioning. Answering this question requires that

we better understand the links between structure and function in these reconstructed, novel ecosystems.

Acknowledgements. Direct financial support was provided by the Natural Sciences and Engineering Research Council (NSERC) of Canada and the Canadian Oil Sands Network for Research and Development. We would like to acknowledge Isabelle Turcotte and Sara Rowland, who conducted the field work and the necessary soil sampling.

References

- Banning, N. C., Grant, C. D., Jones, D. L., and Murphy, D. V.: Recovery of soil organic matter, organic matter turnover, and nitrogen cycling in a post-mining forest rehabilitation chronosequence, *Soil Biol. Biochem.*, 40, 2021–2031, 2008.
- Barrett, L. R. and Schaetzl, R. J.: An examination of podzolization near Lake Michigan using chronofunctions, *Can. J. Soil Sci.*, 72, 527–541, 1992.
- Bashkin, V. N.: *Modern Biogeochemistry*, Kluwer Academic Press, Dordrecht, the Netherlands, 561 pp., 2003.
- Beckingham, J. D. and Archibald, J. H.: Field guide to ecosites of northern Alberta, Special Report 5, UBC Press, Vancouver, BC, Canada, 1996.
- Binkley, D. and Giardina, C.: Why do tree species affect soils? The warp and woof of tree-soil interactions, *Biogeochemistry*, 42, 89–106, 1998.
- Bradshaw, A. D.: Ecological principles and land reclamation practice, *Landscape Plan.*, 11, 35–48, 1984.
- Bradshaw, A. D.: Restoration of mined lands – using natural processes, *Ecol. Eng.*, 8, 255–269, 1997.
- Brockett, B. F. T., Prescott, C. E., and Grayston, S. J.: Soil moisture is the major factor influencing microbial community structure and enzyme activities across seven biogeoclimatic zones in western Canada, *Soil Biol. Biochem.*, 44, 9–20, 2012.
- Bytnerowicz, A., Fraczek, W., Schilling, S., and Alexander, D.: Spatial and temporal distribution of ambient nitric acid and ammonia in the Athabasca Oil Sands Region, Alberta, *J. Limnol.*, 69, 11–21, 2010.

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Chaer, G., Myrold, D., and Bottomley, P. J.: A soil quality index based on the equilibrium between soil organic matter and biochemical properties of undisturbed coniferous forest soils of the Pacific Northwest, *Soil Biol. Biochem.*, 41, 822–830, 2009.

Chapin, F. S., Torn, M. S., and Tateno, M.: Principles of ecosystem sustainability, *Am. Nat.*, 148, 1016–1037, 1996.

Chazdon, R. L.: Beyond deforestation: restoring forests and ecosystem services on degraded lands, *Science*, 320, 1458–1460, 2008.

Crown, P. H. and Twardy, A. G.: Soils of the Fort McMurray region, Alberta, and their relation to agricultural and urban development, Contribution M70-2, Alberta Institute of Pedology, University of Alberta, 52 pp., 1970.

Dimitriu, P. A., Prescott, C. E., Quideau, S. A., and Grayston, S. J.: Impact of reclamation of surface-mined boreal forest soils on microbial community composition and function, *Soil Biol. Biochem.*, 42, 2289–2297, 2010.

Duchaufour, P.: Précis de pédologie, Masson, Paris, France, 482 pp., 1960.

Ekblad, A. and Nordgren, A.: Is growth of soil microorganisms in boreal forests limited by carbon or nitrogen availability?, *Plant Soil*, 242, 115–122, 2002.

Eskelinen, A., Stark, S., and Männistö, M.: Links between community composition, soil organic matter quality and microbial communities in contrasting tundra habitats, *Oecologia*, 161, 113–123, 2009.

Esperschütz, J., Pérez-de-Mora, A., Schreiner, K., Welzl, G., Buegger, F., Zeyer, J., Hagedorn, F., Munch, J. C., and Schlöter, M.: Microbial food web dynamics along a soil chronosequence of a glacier forefield, *Biogeosciences*, 8, 3283–3294, doi:10.5194/bg-8-3283-2011, 2011.

Frouz, J., Pizl, V., Cienciala, E., and Kalcik, J.: Carbon storage in post-mining forest soil, the role of tree biomass and soil bioturbation, *Biogeochemistry*, 94, 111–121, 2009.

Grayston, S. J. and Prescott, C. E.: Microbial communities in forest floors under four tree species in coastal British Columbia, *Soil Biol. Biochem.*, 37, 1157–1167, 2005.

Hahn, A. and Quideau, S. A.: Long-term effects of organic amendments on the recovery of plant and soil microbial communities following disturbance in the Canadian boreal forest, *Plant Soil*, 363, 331–344, doi:10.1007/s11104-012-1306-4, 2012.

Hannam, K. D., Quideau, S. A., Oh, S.-W., Kishchuk, B. E., and Wasylishen, R. E.: Forest floor composition in aspen- and spruce-dominated stands of the boreal mixedwood forest, *Soil Sci. Soc. Am. J.*, 68, 1735–1743, 2004.

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Hannam, K. D., Quideau, S. A., and Kishchuk, B. E.: Forest floor microbial communities in relation to stand composition and timber harvesting in northern Alberta, *Soil Biol. Biochem.*, 38, 2565–2575, 2006.
- Harris, J. A.: Soil microbial communities and restoration ecology: facilitators or followers, *Science*, 325, 573–574, 2009.
- Hemsley, T.: Ecological response of atmospheric nitrogen deposition on reconstructed soils in the Athabasca Oil Sands region, M.Sc. Thesis, Department of Renewable Resources, University of Alberta, Edmonton, 128 pp., 2012.
- Hobbie, S. E., Reich, P. B., Oleksyn, J., Ogdahl, M., Zytkowski, R., Hale, C., and Karolewski, P.: Tree species effects on decomposition and forest floor dynamics in a common garden, *Ecology*, 87, 2288–2297, 2006.
- Hobbs, R. J., Higgs, E., and Harris, J. A.: Novel ecosystems: implications for conservation and restoration, *Trends Ecol. Evol.*, 24, 599–605, 2009.
- Ibáñez, J. J. and Bockheim, J. G. (Eds.): *Pedodiversity*, Taylor and Francis, CRC Press, Boca Raton, FL, USA, 2013.
- Insam, H. and Domnsch, K. H.: Relationship between soil organic carbon and microbial biomass on chronosequences of reclamation sites, *Microb. Ecol.*, 15, 177–188, 1988.
- Jenny, H.: *Factors of Soil Formation*, Dover Publications Inc., New York, 281 pp., 1941.
- Jerabkova, L., Prescott, C. E., and Kishchuk, B. E.: Nitrogen availability in soil and forest floors of contrasting types of boreal mixedwood forests, *Can. J. Forest Res.*, 36, 112–122, 2006.
- Kardol, P. and Wadle, D. A.: How understanding aboveground-belowground linkages can assist restoration ecology, *Trends Ecol. Evol.*, 25, 670–679, 2010.
- Komada, H.: Clay minerals in Canadian soils: their origin, distribution, and alteration, *Can. J. Soil Sci.*, 59, 37–58, 1979.
- Lamarche, J., Bradley, R., Paré, D., Légaré, S., and Bergeron, Y.: Soil parent material may control forest floor properties more than stand type or stand age in mixedwood boreal forests, *Ecoscience*, 11, 228–237, 2004.
- Leckie, S. E., Prescott, C. E., and Grayston, S. J.: Forest floor microbial community response to tree species and fertilization of regenerating coniferous forests, *Can. J. Forest Res.*, 34, 1426–1435, 2004.
- Leifeld, J. and Kögel-Knabner, I.: Soil organic matter fractions as early indicators for carbon stock changes under different land-use?, *Geoderma*, 124, 143–155, 2005.

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Machulla, G., Bruns, M. A., and Scow, K. M.: Microbial properties of mine spoil materials in the initial stages of soil development, *Soil Sci. Soc. Am. J.*, 69, 1069–1077, 2005.
- McBratney, A. R.: On variation, uncertainty and informatics in environmental soil management, *Aust. J. Soil Sci.*, 30, 913–935, 1992.
- 5 McMillan, R., Quideau, S. A., MacKenzie, M. D., and Birjukova, O. V.: Nitrogen mineralization and microbial biomass in reclaimed boreal forest soils, *J. Environ. Qual.*, 36, 1470–1478, 2007.
- Merilä, P., Malmivaara-Lämsä, M., Spetz, P., Stark, S., Vierikko, K., Derome, J., and Fritze, H.: Soil organic matter quality as a link between microbial community structure and vegetation composition along a successional gradient in a boreal forest, *Appl. Soil Ecol.*, 46, 259–267, 10 2010.
- Mummey, D. L., Stahl, P. D., and Buyer, J. S.: Microbial biomarkers as an indicator of ecosystem recovery following surface mine reclamation, *Appl. Soil Ecol.*, 21, 251–259, 2002.
- Ohtonen, R. and Väre, H.: Vegetation composition determines microbial activities in a boreal forest soil, *Microb. Ecol.*, 36, 328–335, 1998.
- 15 Oil Sands Vegetation Reclamation Committee: Guidelines for reclamation to forest vegetation in the Alberta Oil Sands Region. #ESD/LM/99-1, Alberta Environmental Protection – Environmental Service, Edmonton, 1998.
- Ponge, J.-F., Jabiol, B., and Gégout, J.-C.: Geology and climate conditions affect more humus forms than forest canopies at large scale in temperate forests, *Geoderma*, 162, 187–195, 20 2011.
- Prescott, C. E. and Grayston, S. J.: Tree species influence on microbial communities in litter and soil: current knowledge and research needs, *Forest Ecol. Manag.*, in press, 2013.
- Prescott, C. E. and Vesterdal, L.: Effects of British Columbia tree species on forest floor chemistry, in: *Trees and Soil Interactions, Implications to Global Climate Change*, NATO Science Series, edited by: Menyailo, O. and Binkley, D., Kluwer Academic, Dordrecht, the Netherlands, 2005.
- 25 Prescott, C. E., Maynard, D. G., and Laiho, R.: Humus in northern forests: friend or foe?, *Forest Ecol. Manag.*, 133, 23–36, 2000a.
- 30 Prescott, C. E., Vesterdal, L., Pratt, J., Venner, K. H., de Montigny, L. M., and Trofymow, J. A.: Nutrient concentrations and nitrogen mineralization in forest floors of single species conifer plantations in coastal British Columbia, *Can. J. Forest Res.*, 30, 1341–1352, 2000b.

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Prescott, C. E., Zabek, L. M., Staley, C. L., and Kabzems, R.: Decomposition of broadleaf and needle litter in forests of British Columbia: influences of litter type, forest type and litter mixtures, *Can. J. Forest Res.*, 30, 1742–1750, 2000c.

Prescott, C. E., Vesterdal, L., Simard, S. W., and Preston, C. M.: Influence of initial chemistry on decomposition of foliar litter in contrasting forest types in British Columbia, *Can. J. Forest Res.*, 34, 1714–1729, 2004.

Quideau, S. A., Chadwick, O. A., Graham, R. C., and Wood, H. B.: Base cation biogeochemistry and weathering under oak and pine: a controlled long-term experiment, *Biogeochemistry*, 35, 377–398, 1996.

Quideau, S. A., Chadwick, O. A., Benesi, A., Graham, R. C., and Anderson, M. A.: A direct link between forest vegetation type and soil organic matter composition, *Geoderma*, 104, 41–60, 2001.

Ratledge, C. and Wilkinson, S. G. (Eds.): *Microbial Lipids*, vol. 1, Academic Press, London, UK, 1988.

Rowland, S. M., Prescott, C. E., Grayston, S. J., Quideau, S. A., and Bradfield, G. E.: Recreating a functional forest soil in reclaimed oilsands in northern Alberta: an approach for measuring success in ecological restoration, *J. Environ. Qual.*, 38, 1580–1590, 2009.

Ruiz-Jaen, M. C. and Aide, T. M.: Restoration success: how is it being measured?, *Rest. Ecol.*, 13, 569–577, 2005.

Society for Ecological Restoration (SER) International Science and Policy Working Group: The SER international primer on ecological restoration, Society for Ecological Restoration International, Tucson, Arizona, available at: <http://www.ser.org>, last access: September 2012, 2004.

Soil Classification Working Group: The Canadian System of Soil Classification, Agriculture and Agri-Food Canada Publication 1646 (Revised), 1998.

Sorenson, P. T., Quideau, S. A., MacKenzie, M. D., Landhäusser, S. M., and Oh, S. W.: Forest floor development and biochemical properties in reconstructed boreal forest soils, *Appl. Soil Ecol.*, 49, 139–147, 2011.

Swallow, M., Quideau, S. A., MacKenzie, M. D., and Kishchuk, B. E.: Microbial community structure and function: the effect of silvicultural burning and topographic variability in northern Alberta, *Soil Biol. Biochem.*, 41, 770–777, 2009.

Targulian, V. O. and Krasilnikov, P. V.: Soil system and pedogenic processes: self-organization, time scales, and environmental significance, *Catena*, 71, 373–381, 2007.

BGD

10, 7521–7548, 2013

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Turcotte, I., Quideau, S. A., and Oh, S.-W.: Organic matter quality in reclaimed boreal forest soils following oilsands mining, *Org. Geochem.*, 40, 510–519, 2009.

Vetterlein, D. and Hüttnl, R. F.: Can applied organic matter fulfil similar functions as soil organic matter? Risk-benefit analysis for organic matter application as a potential strategy for rehabilitation of disturbed ecosystems, *Plant Soil*, 213, 1–10, 1999.

Wall, D. H., Bardgett, R. D., and Kelly, E. F.: Biodiversity in the dark, *Nat. Geosci.*, 3, 297–298, 2010.

Wardle, D. A., Bardgett, R. D., Klironomos, J. N., Setälä, H., van der Putten, W. H., and Wall, D. H.: Ecological linkages between aboveground and belowground biota, *Science*, 304, 1629–1633, 2004.

Table 1. Site characteristics.

Ecosite or reclamation treatment	Site	Location (° N, ° W)	Soil	Vegetation	Age (yr)
a1	10	57.0735, -111.5944	Dystric Brunisol	pine	48
a1	27	57.5051, -111.4370	Dystric Brunisol	pine	82
b1	29	57.1020, -111.6402	Dystric Brunisol	aspen and pine	48
b1	62	57.5012, -111.5208	Dystric Brunisol	aspen and pine	67
b1	63	57.4980, -111.5229	Dystric Brunisol	aspen and pine	63
b3	2	57.0053, -111.4517	Dystric Brunisol	white spruce and aspen	101
b3	49	57.1030, -111.6367	Dystric Brunisol	white spruce and aspen	52
d1	4	56.9534, -111.7230	Gray Luvisol	aspen	75
d1	8	57.2604, -111.4775	Gray Luvisol	aspen	57
d1	61	56.4390, -111.1902	Gray Luvisol	aspen	58
d2	19	56.4139, -111.1942	Gray Luvisol	white spruce and aspen	73
d2	50	56.6448, -111.0946	Gray Luvisol	white spruce and aspen	79
d2	57	56.3924, -111.4405	Gray Luvisol	white spruce and aspen	75
d3	20	57.3234, -111.2254	Gray Luvisol	white spruce	163
d3	21	57.2852, -111.2743	Gray Luvisol	white spruce	87
A	36	57.0648, -111.6618	PM/2°/TS	white spruce	6
A	37	57.1005, -111.6455	PM/2°/TS	white spruce	20
A	40	57.0591, -111.6682	PM/2°/TS	white spruce and aspen	11
A	87	57.0844, -111.6094	PM/2°/TS	white spruce and aspen	3
A	88	57.0915, -111.6299	PM/2°/TS	white spruce and aspen	12
A	89	57.0906, -111.6794	PM/2°/TS	white spruce and aspen	1
B	3	57.0663, -111.6637	DP/TS	pine	8
B	38	57.0470, -111.6563	DP/TS	aspen and pine	14
B	39	57.1021, -111.6610	DP/TS	white spruce and aspen	4
B	42	57.0748, -111.6665	DP/TS	aspen and pine	5
B	46	56.9991, -111.7318	DP/TS	white spruce and aspen	6
E	7	56.9968, -111.5913	PM/2°/OB	white spruce	7
E	14	57.0009, -111.5744	PM/2°/OB	aspen	6
E	43	56.9909, -111.5638	PM/2°/OB	white spruce and aspen	18
E	86	56.9988, -111.6151	PM/2°/OB	aspen	1
H	1	56.9776, -111.4640	PM/TS	pine	9
H	16	56.9872, -111.5300	PM/TS	white spruce and pine	5
H	17	56.9819, -111.5015	PM/TS	pine	9
H	24	56.9986, -111.4608	PM/TS	white spruce	30
H	30	56.9817, -111.5179	PM/TS	pine	29
I	25	57.0236, -111.4998	PM/OB	white spruce and pine	17
I	28	56.9909, -111.5373	PM/OB	white spruce	30
I	32	56.9987, -111.5482	PM/OB	white spruce and aspen	18
I	34	56.9881, -111.5406	PM/OB	white spruce	21
I	75	56.9120, -111.4192	PM/OB	white spruce and aspen	1

PM: mixture of peat and mineral substrate; 2°: secondary (subsoil) material; TS: Tailing sands; DP: direct placement; OB: overburden.

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[⏪](#) [⏩](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Table 2. Multi-response permutation procedure results for soil nutrient supply (PRS), organic matter composition (NMR), and microbial community structure (PLFA).

	Nutrients			Organic matter			Microbial communities		
	<i>T</i>	<i>A</i>	<i>p</i>	<i>T</i>	<i>A</i>	<i>p</i>	<i>T</i>	<i>A</i>	<i>p</i>
Natural ecosites (NMS groups)									
1 vs. 2 vs. 3	-7.3	0.51	2×10^{-5}	-1.6	0.09	0.07	-4.5	0.15	3×10^{-4}
1 vs. 2	-4.0	0.58	1×10^{-3}	-0.6	0.06	0.23	-3.3	0.20	9×10^{-3}
1 vs. 3	-5.7	0.36	5×10^{-3}	-1.6	0.08	0.07	-2.6	0.09	0.02
2 vs. 3	-5.1	0.42	2×10^{-3}	-1.0	0.06	0.15	-3.0	0.08	6×10^{-3}
Novel ecosystems									
Among treatments	-1.2	0.05	0.13	-1.4	0.06	0.09	-1.0	0.03	0.15
Among vegetation types	-0.6	0.03	0.25	-0.2	0.01	0.53	-0.3	< 0.01	0.47
All ecosystems									
Novel (group 4) vs. all natural	-17.6	0.25	2×10^{-8}	-22.3	0.36	$< 10^{-8}$	-15.8	0.12	$< 10^{-8}$
1 vs. 4	-14.8	0.38	2×10^{-7}	-14.2	0.28	3×10^{-7}	-1.8	0.14	3×10^{-7}
2 vs. 4	-1.8	0.04	0.06	-10.0	0.19	7×10^{-6}	-2.4	0.06	0.02
3 vs. 4	-13.8	0.32	8×10^{-7}	-14.8	0.23	2×10^{-8}	-3.7	0.25	6×10^{-3}

Group 1: a1 and b1 ecosites; group 2: d1 ecosites; group 3: b3, d2, and d3 ecosites; and group 4: all novel ecosystems.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Table 3. Nutrient indicator species analysis associated with site groups. Each value represents the mean indicator with standard deviation in parenthesis, and the highest indicator value is in bold.

Nutrient	Indicator Value				Monte Carlo $p < 0.05$	Indicator Value				Monte Carlo $p < 0.05$	
	Mean	1	2	3		Mean	1	2	3		4
NO ₃						33.5 (4.2)	18	20	21	42	0.048
Ca	39.9 (3.6)	17	50	33	0.009	28.2 (1.5)	14	31	21	34	< 0.001
Mg	41.3 (4.4)	17	57	26	< 0.001	27.7 (1.4)	16	36	22	27	< 0.001
K	42.0 (4.7)	19	22	59	< 0.001	31.0 (3.0)	21	24	40	15	0.010
S						30.9 (2.9)	11	26	20	42	0.002
Na	40.7 (4.0)	58	9	34	< 0.001	32.0 (3.5)	42	16	30	12	0.011

Group 1: a1 and b1 ecosites; group 2: d1 ecosites; group 3: b3, d2, and d3 ecosites; and group 4: all novel ecosystems.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.

Table 4. PLFA indicator species analysis associated with site groups. Each value represents the mean indicator with standard deviation in parenthesis, and the highest indicator value is in bold.

PLFA	Indicator Value			Monte Carlo $p < 0.05$	Indicator Value					Monte Carlo $p < 0.05$	
	Mean	1	2		3	Mean	1	2	3		4
<i>a</i> 15:0	32.0 (12.6)	0	69	13	0.010	27.3 (8.13)	0	44	8	24	0.040
15:1 ω 6 c	27.2 (13.7)	0	87	2	0.004	19.9 (10.5)	0	76	2	2	0.002
15:1 ω 8 c	37.6 (11.2)	80		12	< 0.001	21.4 (10.8)	78		11	0	< 0.001
16:1 ω 11 c						29.5 (4.34)	2	21	8	40	0.003
<i>i</i> 15:0 3OH	32.0 (12.4)	0	70	13	0.005	24.6 (9.77)	0	52	9	9	0.020
<i>i</i> 17:1 ω 9 c						21.9 (10.7)	0			44	0.030
16:0 2OH	29.8 (13.4)	0	71	8	0.020	19.0 (10.1)	0	66	8	1	0.003
<i>i</i> 17:0 3OH	27.3 (13.9)	0	89	2	0.001	20.3 (10.4)	0	75	1	3	< 0.001
19:1 ω 9(11) c	27.2 (13.8)	0	86	2	0.003	21.1 (10.7)	0	68	2	5	0.003

Group 1: a1 and b1 ecosites; group 2: d1 ecosites; group 3: b3, d2, and d3 ecosites; and group 4: all novel ecosystems.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.




Fig. 1. NMDS ordination biplots for soil nutrient supply (PRS), organic matter composition (NMR), and microbial community structure (PLFA) for the natural sites. Each point on the ordination corresponds to one site. Description of soil type and dominant vegetation for each ecosite are given in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Soil biogeochemical processes in novel and natural boreal forest ecosystems

S. A. Quideau et al.




Fig. 2. NMDS ordination biplots for soil nutrient supply (PRS), organic matter composition (NMR), and microbial community structure (PLFA) for the natural and the novel ecosystems. Each point on the ordination corresponds to one site. Description of soil type and dominant vegetation for each ecosite and reclamation treatment are given in Table 1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion