

Modeling
micro-topographic
controls

F. Cresto Aleina et al.

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Modeling micro-topographic controls on boreal peatland hydrology and methane fluxes

F. Cresto Aleina^{1,2}, B. R. K. Runkle^{3,4}, T. Kleinen¹, L. Kutzbach³, J. Schneider⁵,
and V. Brovkin¹

¹Max Planck Institute for Meteorology, Hamburg, Germany

²Max Planck Institute for Biogeochemistry, Jena, Germany

³Institute of Soil Science, Center for Earth System Research and Sustainability. Universität Hamburg, Hamburg, Germany

⁴Department of Biological and Agricultural Engineering, University of Arkansas, Fayetteville, AR, USA

⁵Institute for Environmental Sciences, University of Koblenz-Landau, Landau in der Pfalz, Germany

Received: 05 June 2015 – Accepted: 18 June 2015 – Published: 08 July 2015

Correspondence to: F. Cresto Aleina (fabio.cresto-aleina@mpimet.mpg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Small-scale surface heterogeneities can influence land–atmosphere fluxes and therefore carbon, water and energy budgets on larger scale. This effect is of particular relevance for high-latitude ecosystems, because of the great amount of carbon stored in their soils. We introduce a novel micro-topographic model, the Hummock–Hollow (HH) model, which explicitly represents small-scale surface elevation changes. By computing the water table at the small scale, and by coupling the model with a process-based model for soil methane processes, we are able to model effects of micro-topography on hydrology and methane emissions in a typical boreal peatland. In order to assess the effect of micro-topography on water balance and methane emissions of the peatland we compare two versions of the model, one with a representation of micro-topography and a classical single-bucket model version, and show that the temporal variability in the model version with micro-topography performs better if compared with local data. Accounting for micro-topography almost triples the cumulative methane flux over the simulated time-slice. We found that the single-bucket model underestimates methane emissions because of its poor performance in representing hydrological dynamics. The HH model with micro-topography captures the spatial dynamics of water and methane fluxes, being able to identify the hotspots for methane emissions. The model also identifies a critical scale (0.01 km^2) which marks the minimal resolution for the explicit representation of micro-topography in larger-scale models.

1 Introduction

Peatlands cover only about 3% of the global land surface (Wieder et al., 2009), but they play a fundamental role in the global carbon cycle (Blodau, 2002; Limpens et al., 2008). In boreal latitudes peatlands and wetlands are one of the major natural sources of methane to the atmosphere (e.g., Bousquet et al., 2006). During the Holocene peatlands have functioned as a sink of atmospheric carbon (Smith et al., 2004), and Yu

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(2012); Yu et al. (2011) recently estimated the amount of carbon stored in northern peatlands of about 547 (473–621) Pg, significantly larger than previous estimates of 270–370 Pg (e.g., Turunen et al., 2002). Recent efforts have tried to reproduce peatland and wetland extent and carbon accumulation in various Dynamic Global Vegetation Models (DGVMs), (i.e., Schuldt et al., 2013; Kleinen et al., 2012; Wania et al., 2009a, b). The WETland and Wetland CH₄ Inter-comparison of Models Project (WETCHIMP) (Melton et al., 2013; Wania et al., 2013) revealed the large variability among the different models in wetland extent and in the parameterization of hydrological and biogeochemical processes such as methane emissions. It is also clear that all of these models lack the explicit representation of fine-scale heterogeneities and sub-grid processes.

We propose a novel method that takes into account sub-grid scale processes, and directly assess their impact on peatland ecosystems, from greenhouse gas emissions, to the water budget. Previous studies have suggested that micro-topography in peatlands (micro-relief with a typical spatial scale of 1 m × 1 m) may influence GHG emissions (Baird et al., 2009b), but the extent of the micro-topography influence in land-atmosphere fluxes is yet to be determined. Small-scale processes can have significant effects in the peatland carbon cycle (Baird et al., 2009a), and local surface models (e.g., Nungesser, 2003; Bohn et al., 2013) also highlighted how local surface heterogeneities matter for the water balance in northern peatlands. Acharya et al. (2015) recently linked the self-organization occurring in patterned peatlands to local small-scale interaction among micro-topography units. Cresto Aleina et al. (2013) instead showed that the importance of small-scale surface heterogeneities in estimating water table change in permafrost-generated soil patterns. Observations in northern peatlands also showed the position of the water table has a fundamental control on greenhouse gas emissions, since it changes the depth of the oxic zone, i.e., the region where methane gas diffusing from below can be oxidized and therefore released as CO₂ instead (Couwenberg and Fritz, 2012). Because of the high global warming potential of CH₄ relative to CO₂ a robust estimation of methane emissions is essential to evaluate the climate impact of natural wetlands and peatlands (Kirschke et al., 2013). To compute a consistent green-

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


house balance over the region, one should consider the small-scale properties and how the water table and the soil surface heterogeneously change within the environment (Bellisario et al., 1999; Camill and Clark, 1998; Law et al., 2002). Process-based models recently suggested that hydrological heterogeneities at the landscape scale between different wetland types (i.e., between fens and bogs) control different water table responses under a changing climate forcing (Gong et al., 2012). This phenomenon can potentially influence the carbon fluxes from peatlands at regional scales (Gong et al., 2013). Van der Ploeg et al. (2012) showed how micro-topography exerts dominant control in hydrological processes in wetlands. On the other hand, there has not yet been a quantification of the micro-topography effects on methane emissions, nor is there a proper way to represent these effects on larger-scale models. The general issue of scale interactions in the climate–biosphere system is therefore of particular interest in northern peatlands, where large emissions of greenhouse gases are influenced by the small-scale surface heterogeneities.

Here, we developed a mechanistic model operating at the landscape scale for a typical boreal peatland, in order to assess the importance of surface micro-topography for the water balance and the methane fluxes. These small-scale surface heterogeneities are typical in peatlands, and consist of elevated and relatively drier zones, called *hummocks*, and lower and relatively wetter zones, called *hollows*. We calibrated this landscape-scale model (Hummock–Hollow model, HH model hereafter) with data from an elevated bog in the Ust-Pojeg mire complex, in the Komi Republic, Russia. A number of recent studies have analyzed this site's peat characteristics and depth (Pluchon et al., 2014), and provide measures of fluxes of water vapor (Runkle et al., 2012), carbon dioxide (Schneider et al., 2012), and methane (Gažovič et al., 2010; Wolf, 2009), as well as the energy and water balance (Runkle et al., 2014) and spatial distribution of dissolved organic carbon (DOC) (Avagyan et al., 2014b) and biogeochemical elements (Avagyan et al., 2014a).

In this paper we present the new model for peatland micro-topography and the measurements we used for tuning and evaluating model performances. We then proceed to

analyze the influence of micro-topography on water balance and methane processes comparing the output from the model including a representation of micro-topography with the output of a version of the model without any small-scale information.

2 Methods

5 We developed the Hummock–Hollow (HH) model to evaluate peatland micro-topographic controls on land–atmosphere fluxes. We represented the surface sub-grid scale heterogeneities with a square lattice. The model works at a landscape-scale of 1 km × 1 km, and each model grid-cell represents a micro-topographic feature, namely a hummock or a hollow, with dimensions of 1 m × 1 m. We compute for each grid cell
10 both water table balance and methane emissions. We explicitly represent the micro-topography characterizing the peatland surface, and therefore we parameterize the heterogeneous hydrological properties of the peatland soil (i.e., surface and the sub-surface water fluxes). Due to such a fine representation of the micro-scale, we can upscale emissions and water balance at the landscape-scale by averaging the local quantities over the whole domain. To investigate the micro-topographic effect, we created a second version of the model (hereafter *Single Bucket*), in which we represent the whole peatland in a single bucket with parameters averaged over the whole 1 km × 1 km domain.

2.1 The HH model

20 We tuned the model with micro-topographic data from a peatland of Northwest Russia, the Ust-Pojeg mire in the Komi Republic (61°56' N, 50°13' E, 119 m a.s.l.). Many studies have focused on this study site as a typical boreal peatland, as the mire complex displays different kinds of peatland types, from an ombrogenous bog, to a minerogenous fen, to a transitional zone with the surrounding forest. In order to simulate small-scale
25 surface heterogeneities, we use field data to initialize the surface elevation in the model.

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We focus our modeling framework on the bog, in order to exclude the influence of the subsurface water input that characterizes the fen ecosystem.

We initialize the soil surface elevation with elevation data collected through surveying with a theodolite. The circumference and points on the height of each hummock and within representative hollows in a 40 m × 60 m grid were surveyed and a surface between them interpolated. In order to consistently represent the surface elevation, we inferred the statistical distribution of the field data. We fitted the histogram of elevational data (x) with a generalized 3-parameter gamma distribution multiplied by the maximum number of counts in the histogram:

$$f(x) = N_{\max} \frac{pb^q x^{q-1} e^{-(bx)^p}}{\Gamma(q/p)}, \quad b > 0, \quad q > 0, \quad p > 0. \quad (1)$$

Values of parameters for Eq. (1) are reported in the following section. We randomly pick a value from the statistical distribution, and we assign it to each grid cell. We assume that with this procedure we statistically capture the peatland micro-topography. If the grid cell at the position i, j has an elevation $H_{i,j}$ (height in Fig. 1) above a surface level $H_0 = 20$ cm then we assume that such a grid cell is a *Hummock*, otherwise it is a *Hollow*.

Along with the micro-topography, we initialize each grid-cell with two other properties which affect methane production and hydrology respectively: peat depth $pd(i, j)$ and absolute elevation $sl(i, j)$. We model the peat profile (peat depth) according to in situ measurements, and we use this information in the methane emission model as a surrogate for the amount of carbon available for decomposition and GHG emissions. The equations used for pd initialization can be found in the Appendix. The second term $sl(i, j)$ parameterizes the elevation of each grid cell in respect the lowest part of the bog. Following observations, we assume the slope of the bog to have a uniform linear dependence on y , which represents the distance in meters from the origin. We therefore assign to each cell an absolute elevation according to the formula:

$$sl(i, j) = HS(1 - y(i, j)/m_y) \quad (2)$$

Modeling
micro-topographic
controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where $m_y = 1000$ m is the longitudinal dimension of the landscape-box representing our bog and $HS = 3$ m is the maximal height of the bog. The slope regulates water flow over the peatland (see Sect. 2.2.1).

2.2 Water table dynamics

5 We compute the water table position at each time step. We start the simulations at the end of April and we end it at the end of October, when the cold season starts. After the snowmelt (not simulated) most of the landscape becomes inundated. For each cell at position i, j we compute the water balance following the equation:

$$\frac{dW_{i,j}}{dt} = \frac{P - ET + R_{i,j}}{s_{i,j}} \quad (3)$$

10 where $W_{i,j}$ is the water table level in the grid cell at the position (i, j) relative to the surface level, P is the precipitation input, ET is the evapotranspiration, $R_{i,j}$ is the lateral runoff, $s_{i,j}$ is the drainable porosity, and t is time, and the time step for the simulation is $\delta t = 1$ day. We randomly initialize water table level at the beginning of the simulations using values measured by Schneider et al. (2012) typical for hummocks and hollows:

$$15 \quad W_{i,j}^1 \in \begin{cases} [-150 \quad -50] & \text{mm if Hummock} \\ [-10 \quad 100] & \text{mm if Hollow} \end{cases} \quad (4)$$

Where $W_{i,j}^1$ is the water table level at the first time step. In the initialization of the *Single Bucket* model version we use: $W_{SB}^1 \in [-50 \quad 100]$ mm, where W_{SB}^1 is the uniform water table level for the whole peatland at the first time step.

20 Values of P and ET are uniform over the whole landscape, i.e., we do not apply any downscaling (see Sect. 2.5).

Term $s_{i,j}$ is the drainable porosity, and it varies spatially both horizontally and vertically. If water table is above the surface level, $s_{i,j} = 1$, whereas if the water table is

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



below the surface, we assume it to be heterogeneous within the environment, and we choose two different values for hummocks and hollows. Values of $s_{i,j}$ are described in Table 1. By dynamically representing the water table at the micro-topographic level, we are able to distinguish the subsurface water flow in hummocks and hollows. We expect the spatial pattern of water table depth to change over the season, as the water flow, along with the evapotranspiration, removes water from the system making it progressively drier.

2.2.1 Lateral runoff

The lateral runoff term R is the computed water flux among the different grid cells, and it allows the exchange of water through the soil. We define it as $R_{i,j}^{\text{in}} - R_{i,j}^{\text{out}}$, where $R_{i,j}^{\text{in/out}}$ is the sum of Darcy's and Manning's flows D and M :

$$R_{i,j}^{\text{in/out}} = D_{i,j}^{\text{in/out}} + M_{i,j}^{\text{in/out}} \quad (5)$$

These terms represent the subsurface water flow and the overland water flow respectively and are computed in mm day^{-1} . We parameterize the subsurface water flow as:

$$D_{i,j}^{\text{in/out}} = d_{i,j} \frac{\Delta W_{i,j}}{m_{i,j}^{x/y}} \quad (6)$$

The term $m_{i,j}^{x/y}$ is the distance between the centers of cells i and j , and $\Delta W_{i,j}$ is the difference between the water table in the two adjacent grid cells. With respect to the classical expression for Darcy's Law, we then consider the term $d_{i,j}$ to represent the hydraulic conductivity, and its dimensions are length per time. It is spatially dependent, since we assume a different value for hummocks and hollows, as reported in Table 1. Parameters are chosen in the order of magnitude of the ones measured by Clymo (2004). Manning's formula describes instead the velocity of an overland flow driven by

gravity:

$$M_{i,j}^{\text{in/out}} = \begin{cases} \left(\frac{k}{Mc_{i,j}}\right) \text{Rad}_{i,j}^{\frac{2}{3}} \Delta S_{i,j}^{\frac{1}{2}} & \text{if } W_{i,j} > 0 \\ 0 & \text{otherwise} \end{cases} \quad (7)$$

The overland water flow is also dependent on the soil heterogeneity because we assume a difference in the hummock-hollow surface roughness due to the increased vascular plant cover in hummocks. In the equation, this term is represented by the dimensionless number $Mc_{i,j}$, as displayed in Table 1. In Eq. (7), $\Delta S_{i,j}^{\frac{1}{2}}$ represents the difference in slope between the cells i and j and k is a conversion factor of $1 \text{ m}^{1/3} \text{ s}^{-1}$. The term $\text{Rad}_{i,j}$ is the hydraulic radius, defined as $\text{Rad}_{i,j} = A_{i,j}/p_{i,j}$, where $A_{i,j}$ is the cross sectional area of flow, in our model:

$$A_{i,j} = m_{i,j}^{x/y} (W_{i,j} - H_{i,j}) \quad (8)$$

and $p_{i,j}$ is the wetted perimeter, or the perimeter of the cross sectional area $A_{i,j}$ at contact with water, in our model:

$$p_{i,j} = 2(W_{i,j} - H_{i,j})m_{i,j}^{x/y} \quad (9)$$

where $W_{i,j} - H_{i,j}$ is the elevation of water table with respect to the surface and $m_{i,j}^{x/y}$ is the lateral extent of the grid cell.

Manning's flow occurs only if the water table is above the surface level, whereas Darcy's flow is continuous. This hydrological representation is one of the main differences between this new approach and the classical bucket model, and an important driver of the model's seasonal hydrological dynamics. The heterogenous surface and the interactions among the different grid cells represent at a fine spatial scale the interactions among hummocks and hollows in typical peatlands. Values for parameters are displayed in Table 1.

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Coupling with a process-based model for CH₄ emissions

We propose an explicit parameterization of methane fluxes, by coupling the micro-topographic water balance model to a process-based model for methane emissions, in order to more consistently quantify the effect of surface heterogeneities on GHG fluxes. The model developed by Walter and Heimann (2000) is a quite general model for methane emissions, and can be applied to peatlands in different environments. In particular, it is the same model which is built into some DGVMs (i.e., Kleinen et al., 2012). We tune the Walter and Heimann (2000) model parameters to perform in a typical peatland at the latitude of the Ust-Pojeg mire complex. We couple the methane model at each grid-cell, and we compute methane fluxes for each hummock and each hollow. We average over the whole landscape in order to upscale the local fluxes at the landscape-scale. The process-based model for methane emissions provides an output of methane fluxes $F_{\text{CH}_4}^{i,j}$ as a function of water table, available carbon as a function of peat depth, net primary productivity (NPP), and soil temperature (T):

$$F_{\text{CH}_4}^{i,j}(t) = f\left(W_{i,j}(t), z_{i,j}^{\text{soil}}, \text{NPP}(t), T(t)\right) \quad (10)$$

Where $W_{i,j}$ is the water table depth with respect to the surface computed at each position i, j , $z_{i,j}^{\text{soil}}$ is the soil depth, and NPP and T are at the daily time scale. The soil depth takes into account that each grid cell has a different peat depth. We sum the peat depth to the height of the acrothelm (the part of peat containing living plants, which we assume to be $H_{i,j}$). This quantity $z_{i,j}^{\text{soil}}$ is a proxy for the amount of carbon available by adding to the peat depth the micro-topography height:

$$z_{i,j}^{\text{soil}} = \text{pd}_{i,j} + H_{i,j} \quad (11)$$

Values of parameters used in the model are listed in the Appendix.

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Sensitivity analysis

In order to test the robustness of our results, we change the key parameters of the model. We call hereafter *HH Standard configuration* the version of the model with parameter values described above.

5 In particular, we test our model by varying the grid size. In the *HH Standard configuration* the number of grid cells $N = 10^6$, and the grid cells have a length size of $m_{x/y} = 1$ m. We compare the output of the model with micro-topography with a classical Single Bucket model, i.e. a model where $N = 1$ and $m_{x/y} = 10^3$ m. To test the scale dependency of the model, we perform simulations that gradually increase the number of grid cells (i.e., decreasing the grid cell size), and we compute the cumulative methane emissions over the whole season, from the end of April to the end of October. We aim to find a critical scale at which the model performance does not change qualitatively.

15 Another parameter to be tested is the drainable porosity $s_{i,j}$, which plays a key role in the water balance, according to Eq. (3). By changing this value, the water table responds differently. In particular, for small values of the parameter, the water table variations within the soil are amplified. We also explore the importance of the difference in drainable porosity among hummock and hollows. We also studied the dependence of the hydrological properties of the model by varying to the parameters in Table 1, which mainly influence the velocity of the surface and subsurface flow among the grid cells. We also ran the HH model forced by half the NPP values in the *HH Standard configuration*, and then we forced with NPP values which are twice the original ones in order to assess the goodness of our assumption on NPP values (see following sub-section).

25 2.5 Forcing data

The HH model is forced with prescribed precipitation and evapotranspiration (Eq. 3). We use precipitation from Schneider et al. (2012), and evapotranspiration computed by

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



them passed the Kolmogorov–Smirnov test. We then assume that $f(x)$ fits our data well enough to be used for our purposes as parameterization of the distribution for micro-topographic elevation, and we proceed to initialize the micro-topographic model by assigning at each grid cell a value randomly picked from $f(x)$. The best fit parameters in Eq. (1) are: ($b = 5.8$, $q = 8.9$, $p = 1.5$). It is worthy to notice that for this peatland the distribution of surface elevation is not bimodal, as shown in previous studies (Eppinga et al., 2008). In fact, the two peaks in the histogram are too close to be resolved by the fitting distribution (Fig. 2).

3.2 Hydrology

Because of the random initialization we performed ensemble simulations with 30 ensemble members and we compared two versions of the model. Each ensemble member differs for initialized water table and, in the *Microtopography* version of the HH model, for micro-topography configuration. The ensemble members differ very little with each other in the *Microtopography* version, whereas the span of water table positions computed by the *Single Bucket* version is much larger.

The difference in water table dynamics between the two model versions is significant (Fig. 3). The shaded areas represent the standard deviations from the mean over the 30 ensemble members. In the *Microtopography* version, this shaded area is so small to be invisible in the figure ($\approx \pm 3$ mm). This is due to the fact that the only random parameters are the micro-topographic height and the water table position at the beginning of the simulation. Since in the *Microtopography* version the model has a large number of cells ($N = 10^6$) all realizations are very similar to each other. Because of the initialization in both versions, the water table at the beginning of the simulations lies above the average surface level and floods the peatland. In the model version *Single Bucket*, the water flows out of the peatland much faster than in the *Microtopography* version. As a result, due to the strong surface (at the immediate beginning of the season) and subsurface flow, the water table drops quickly below 400 mm from the average surface level, thus increasing the oxic zone depth as the simulation proceeds. In the *Microtopography*

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



version, instead, the sub-grid scale interactions delay the runoff. The more ragged surface of the hummocky peatland prevents the strong surface flow from draining the peatland in the first month of simulation. As a consequence, the peatland soil remains wetter for longer time, enhancing anoxic conditions where the water table lays near the surface level. Therefore, the fast decrease in the water table position at the beginning of the season in the *Single Bucket* version disappears. The average behavior of the water table dynamics is similar to site observations that show a much smoother water table decrease, as in Runkle et al. (2014); Schneider et al. (2012). The simulated water table dynamics also capture the late increase in average water table visible in the measurements due to abrupt large precipitation inputs.

With the HH model in the *Microtopography* version we are able to study the differences in water table position between hummocks and hollows, in order to compare the performance of the model against field data. In Fig. 4 we represent the distribution of simulated water table position relative to the surface level in box plots for hummocks (orange boxes), hollows (red boxes). To evaluate the effects of our hydrological model, we also plot the water table position computed by a realization of the HH model in the *Single Bucket* version (blue line). We compare model output against in situ measurements from the bog region (black crosses) for both micro-types. For completeness, we also compare the model performances against measurements from other regions of the Ust-Pojeg mire complex, i.e. the fen region (black circles in the figure). We chose these specific dates of Fig. 4 because of the relative abundance of chamber measurements of methane for both hummocks and hollows, which we will describe in the following sub-sections. The model performs generally well against the data, with most of the measurements among the whiskers of the box plots. The model seems to slightly overestimate the hummock water table towards the mid of August and to underestimate the hollow water table at the beginning of September, but the measurements still fall among the whiskers of the box. The model without micro-topography, instead, consistently underestimates water table position. In particular, the computed water table is lower than the one of the deepest measurement for all analyzed dates.

Modeling micro-topographic controls

F. Cresto Aleina et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In our simulation, the presence of outliers in the negative tail of the distribution shows that the data are skewed to the left. The skewness of both hummock and hollow distributions is negative for most of the simulation, becoming weakly positive for hummocks only at the beginning of the simulation, when water table is well above the surface level.

5 Afterwards, the skewness of the hummock distribution reaches larger negative values than in the hollow distribution, as the hummocks are generally drier than the hollows. Accordingly, the excess kurtosis of the distributions in all cases is positive, since the tails of the histograms are fatter than they would be if data were normally distributed. In particular, the left tail becomes fatter as more and more grid cells become dry.

10 3.3 Methane emissions

By coupling the HH model with a process-based model for methane emissions, we are able to simulate the dynamics of methane over the warm season in the Ust-Pojeg mire complex. The HH model enables us also to distinguish among emissions from different micro-topographic units, in order to better compare the performances of the model against chamber measurements.

15 We compute methane emissions for each cell of the squared lattice. Due to the heterogeneous patterns of soil properties and water table position that we analyzed in the previous section, the emissions of methane are not uniform over the landscape. We study the impact of such heterogeneous emission pattern by distinguishing among hummock and hollow distributions of methane emissions. We compare the box plots of methane emissions at different time steps with chamber measurements taken in the bog part of the Ust-Pojeg mire complex (Wolf, 2009). We compare model output against in situ measurements from the bog region (black crosses) for both micro-types (Fig. 5). For completeness, we also compare the model performances against measurements from other regions of the Ust-Pojeg mire complex, i.e. the fen region (black circles in the figure). As expected, methane emissions differ between hummocks (orange boxes) and hollows (red boxes), being the latter ones generally wetter and thus displaying a shallower oxic zone. We also compare the averaged output of the model in the *Single*

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bucket version (blue lines in the figure) against the distributions of methane emissions from the *Microtopography* version.

The large increase in methane emissions at the end of July and at the beginning of August, as we can also see in the averaged fluxes (Fig. 6), is due to the combination of a higher temperature forcing and the presence of shallow oxic one in most of the model grid cells in the *Microtopography* version. By the end of August, instead, despite the comparable water table position, temperatures are much lower. This phenomenon causes a significant decrease in methane emissions, only partially mirrored by chamber measurements. The HH model coupled with the Walter and Heimann (2000) methane emission model captures the general trend and the magnitude of methane emissions, but towards the end of the season it seems to fail in representing large methane emissions, in particular from a hummock at the end of August and from two hollows in September, whereas in the rest of the season most of the measurements fall between, or near to, the whiskers of the boxes. Because of the generally good agreement of simulated water table depth with measurements, we exclude a bias coming from the hydrological model. The only other potential bias comes from the methane emission model, which seems to be overly sensitive to temperature variations.

Despite such differences in fluxes, in general the measurements fall in the range of the simulated fluxes by the *Microtopography* version. The agreement is particularly good for hollows towards the center of the season, when methane emissions are higher.

The *Single Bucket* version represents an average over the whole region, but it nevertheless produces outputs which are not outside the range of the measurements. The inability of this version to represent a distribution of the large spread in methane emissions and in particular of hollows as methane emission hotspots, though, leads to an overall underestimation of the averaged fluxes (Fig. 6). The spatially-explicit representation of the methane flux distribution, instead, is essential for the *Microtopography* version to better capture the magnitude of the measured fluxes in average.

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sion because such resolution is approximatively the dimension of the micro-topographic features in the field. We computed then the cumulative CH_4 emissions for each simulation to test the dependence of emissions on grid cell size (Fig. 7). The cumulative emissions increase almost linearly for an increasing N , if $N \leq 10^2$. By increasing further the number of cells, the cumulative emissions stabilize after slightly decreasing for $10^2 < N \leq 3 \times 10^2$, but they do not largely change. Such different behavior depends on the different representation of water table dynamics, which delays the drying as the surface becomes more diverse in micro-topography, i.e., if the number of grid cells increases. In particular, in the first three months of simulation the water is retained within the system as the number of hydrological interactions between the different grid cells increases, and the water table lays more and more in proximity of the average soil surface. This change explains the increase in emissions as $N = 10^2$. By $N > 10^2$, instead, the water table behavior asymptotically approaches water table dynamics for $N = 10^6$, i.e. for model resolution of 1 m, hence changes in cumulative emissions are no longer significant. In our model, therefore, we can identify a critical scale for micro-topographic controls on hydrology and methane emissions at about 0.01 km^2 .

Among other parameters, the drainable porosity $s_{i,j}$ in Eq. (3) has a direct impact on water table position and therefore indirectly on methane emissions. We chose the amount of water the peat soil can retain in the *Standard configuration* based on Kolka et al. (2011). This parameter can vary greatly in different peatlands, and therefore the values in our *Standard configuration* are an assumption we need to test. In order to assess the model sensitivity to this parameter we make drainable porosity uniform over the whole region, i.e., we do not distinguish between hummocks and hollows. We changed $s_{i,j}$ uniformly at steps of 0.1 from 0.2 to 1.0 to test how water table changes its position. The average position is generally lower with respect to the dynamics of Fig. 3, but the differences are not qualitatively significant, nor can we see any large effect in methane emissions. This is due to the methane emission model by Walter and Heimann (2000) being very sensitive to temperature variations, which in this test did not vary from the forcing of the *Standard configuration*. For very low values of $s_{i,j}$,

water table variation are larger and in general we observe both a deeper water table and lower methane emissions than in the *Standard configuration*, but the model still produces higher emissions if a representation of micro-topography is included.

The results proved to be robust also by changing the other parameters in Table 1 across a wide range of values. In particular, also for this analysis we eliminated the differences in hydrological parameters between hummocks and hollows, and the results were qualitatively the same as in the *Standard configuration*.

We also tested the influence of NPP forcing on model performance. In particular, we changed NPP values for 0.1 to 5 times the values in the *Standard configuration*. The differences that we saw in model output (see Appendix) were small or negligible and did not qualitatively alter the results. This finding confirms the robustness of our results, and the assumption that the bias introduced by not considering NPP for mosses, but only for C3 grasses was neglectable for the purposes of this study.

4 Summary and conclusions

We developed the Hummock–Hollow (HH) model, a new model representing the hydrology and the properties of the micro-topographic features typical of a boreal peatland, working at the resolution of 1 m^2 . This novel model presents a physical representation of the peatland micro-topography with the help of in situ measurements in the Ust-Pojeg mire complex in the Komi Republic, Russia. After inferring the statistical distribution of micro-topography data, we used this result to randomly assign elevation values at the grid cells. The explicit representation of the micro-topography allows the HH model to distinguish water table and methane fluxes among hummocks and hollows, thus identifying the role of the diverse micro-topographic features in water table and methane flux dynamics. To assess the effects of the micro-topographic controls on these two observables, we created a model version which averages all quantities (*Single Bucket*), thus not distinguishing among hummocks and hollows, and which reproduces how a global or regional model would represent the landscape. We compared

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the output of the two model versions with in situ measurements of water table depth and methane fluxes.

Overall, the model version with micro-topographic representation performs better in comparison with hydrological data, as the water table position simulated by the *Single Bucket* version of the model is constantly deeper than the measurements. The *Single Bucket* version simulates a drier peatland, because the strong runoff washes away the water at the beginning of the simulation. The flow is instead diminished in the model version with a representation of micro-topography, since the more rugged, hummocky surface delays the water discharge. This phenomenon allows for wetter conditions, leading to a general good agreement with field data in the *Microtopography* version working at 1 m² resolution. The HH model therefore correctly captures not only the averaged water table dynamic, but also the heterogeneity in water table depth distribution among hummocks and hollows, as micro-topography slows down the water runoff.

By changing the water table dynamics, the micro-topography affects methane emissions. The water table position in respect to the surface level changes regulates the depth of the oxic zone, i.e. the region where methane can be oxidized and therefore the methane emissions are drastically reduced (as seen experimentally, e.g., in Couwenberg and Fritz, 2012). In our simulations, the *Single Bucket* version generally underestimates the averaged methane fluxes, because of the overly deep water table towards the central months of the simulations. The spatially explicit version of the HH model, instead, is able to produce an output of methane emission distributions, which as expected identifies the hollow grid cells as hotspots for methane emissions.

We progressively increased the scale of the model, i.e., reducing the number of grid cells, and we identified a critical scale at which the model results do not change for an increased resolution. This critical scale for the grid cells is 0.01 km² in the HH model, which is still too small for the investigated processes to be explicitly included in a global or regional model. Therefore we argue that further developments are needed towards a new parameterization of peatland surface which takes into account the upscaled effects of micro-topography. The identification of a critical scale for the representation

BGD

12, 10195–10232, 2015

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of micro-topography on a global scale requires the application of the HH model to other peatlands, with a more structured and patterned micro-topography, which is the object of future investigations.

This last result limits the applicability of the model to landscape-scale studies, because of the computational non-feasibility of including our findings directly into a global model. We tested the HH model only for one particular peatland, and even though we believe the peatland system we studied to represent a rather typical system in boreal peatlands, further studies are needed to consistently assess the global relevance of our results.

Further potential developments of the HH model include coupling the model to other process-based models for greenhouse gas emissions, CO₂ in particular, such as the one developed by Wania et al. (2010). Such a coupling would lead to the estimation of the micro-topography controls on the total carbon emissions of a typical boreal peatland. Along this line of the development, the implementation of a peat accumulation module (e.g., the one presented by Kleinen et al., 2012) could potentially assess the micro-topographic controls not only on hydrology and carbon emissions, but also on the long term carbon cycle. Further model developments involve an explicit representation of the energy balance, in order to eliminate some of the biases introduced by the forcing timeseries, and to fully represent the dynamics of other processes we now ignore, i.e., micro-topography controls on evapotranspiration, heat fluxes, and potential feedbacks among the different components of the model. The coupling with a model for moss dynamics (e.g., Porada et al., 2013) will make the model able to investigate dynamic and micro-topography dependent insulation properties of the soil.

The good agreement between the HH model and data shows that an explicit representation of micro-topography is fundamental in predicting landscape-scale hydrology and, as a secondary effect of surface heterogeneities, methane emissions. This study highlights the need of effective upscaling procedure in order to parameterize the effects of local surface heterogeneities across scales, ranging from the micro-scale (1 m) to the GCM scale (30–500 km).

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The article processing charges for this open-access publication were covered by the Max Planck Society.

References

- 5 Acharya, S., Kaplan, D. A., Casey, S., Cohen, M. J., and Jawitz, J. W.: Coupled local facilitation and global hydrologic inhibition drive landscape geometry in a patterned peatland, *Hydrol. Earth Syst. Sci.*, 19, 2133–2144, doi:10.5194/hess-19-2133-2015, 2015. 10197
- Avagyan, A., Runkle, B., Hartmann, J., and Kutzbach, L.: Spatial variations in pore-water biogeochemistry greatly exceed temporal changes during baseflow conditions in a boreal river valley mire complex, Northwest Russia, *Wetlands*, 34, 1171–1182, doi:10.1007/s13157-014-0576-4, 2014a. 10198
- 10 Avagyan, A., Runkle, B. R., and Kutzbach, L.: Application of high-resolution spectral absorbance measurements to determine dissolved organic carbon concentration in remote areas, *J. Hydrol.*, 517, 435–446, doi:10.1016/j.jhydrol.2014.05.060, 2014b. 10198
- 15 Baird, A. J., Belyea, L. R., and Morris, P. J.: Upscaling of peatland-atmosphere fluxes of methane: small-scale heterogeneity in process rates and the pitfalls of the “Bucket-and-Slab” models, in: *Carbon Cycling in northern peatlands*, edited by: Baird, A. J., Belyea, L. R., Comas, X., Reeve, A. S., and Slater, L. D., vol. 184, American Geophysical Union, 37–53, available at: <http://cat.inist.fr/?aModele=afficheN&cpsidt=23940034>, 2009a. 10197
- 20 Baird, A. J., Belyea, L. R., and Morris, P. J.: Carbon cycling in northern peatlands, in: *Geophysical Monograph Series*, edited by: Baird, A. J., Belyea, L. R., Comas, X., and Reeve, A. S., vol. 184, American Geophysical Union, Washington, DC, doi:10.1029/GM184, 2009b. 10197
- Bellisario, L. M., Bubier, J. L., Moore, T. R., and Chanton, J. P.: Controls on CH₄ emissions from a northern peatland, *Global Biogeochem. Cy.*, 13, 81–91, doi:10.1029/1998GB900021, 1999. 10198
- 25 Blodau, C.: Carbon cycling in peatlands – a review of processes and controls, *Environ. Rev.*, 10, 111–134, doi:10.1139/A02-004, 2002. 10196
- Bohn, T. J., Podest, E., Schroeder, R., Pinto, N., McDonald, K. C., Glagolev, M., Filippov, I., Maksyutov, S., Heimann, M., Chen, X., and Lettenmaier, D. P.: Modeling the large-scale effects of surface moisture heterogeneity on wetland carbon fluxes in the West Siberian Lowland, *Biogeosciences*, 10, 6559–6576, doi:10.5194/bg-10-6559-2013, 2013. 10197
- 30

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bousquet, P., Ciais, P., Miller, J., Dlugokencky, E., Hauglustaine, D., Prigent, C., Van der Werf, G., Peylin, P., Brunke, E.-G., Carouge, C., Langenfelds, R. L., Lathière, J., Papa, F., Ramonet, M., Schmidt, M., Steele, L. P., Tyler, S. C., and White, J.: Contribution of anthropogenic and natural sources to atmospheric methane variability, *Nature*, 443, 439–443, 2006. 10196

Camill, P. and Clark, J. S.: Climate change disequilibrium of boreal permafrost peatlands caused by local processes, *Am. Nat.*, 151, 207–222, available at: <http://www.jstor.org/stable/10.1086/286112>, 1998. 10198

Clymo, R. S.: Hydraulic conductivity of peat at Ellergower Moss, Scotland, *Hydrol. Process.*, 18, 261–274, doi:10.1002/hyp.1374, 2004. 10202

Couwenberg, J. and Fritz, C.: Towards developing IPCC methane “emission factors” for peatlands (organic soils), *Mires and Peat*, 100, 1–17, 2012. 10197, 10215

Cresto Aleina, F., Brovkin, V., Muster, S., Boike, J., Kutzbach, L., Sachs, T., and Zuyev, S.: A stochastic model for the polygonal tundra based on Poisson–Voronoi diagrams, *Earth Syst. Dynam.*, 4, 187–198, doi:10.5194/esd-4-187-2013, 2013. 10197

Eppinga, M., Rietkerk, M., Borren, W., Lapshina, E., Bleuten, W., and Wassen, M.: Regular surface patterning of peatlands: confronting theory with field data, *Ecosystems*, 11, 520–536, doi:10.1007/s10021-008-9138-z, 2008. 10208

Gažovič, M., Kutzbach, L., Schreiber, P., Wille, C., and Wilmking, M.: Diurnal dynamics of CH₄ from a boreal peatland during snowmelt, *Tellus B*, 62, 133–139, doi:10.1111/j.1600-0889.2010.00455.x, 2010. 10198

Gong, J., Wang, K., Kellomäki, S., Zhang, C., Martikainen, P. J., and Shurpali, N.: Modeling water table changes in boreal peatlands of Finland under changing climate conditions, *Ecol. Model.*, 244, 65–78, doi:10.1016/j.ecolmodel.2012.06.031, 2012. 10198

Gong, J., Kellomäki, S., Wang, K., Zhang, C., Shurpali, N., and Martikainen, P. J.: Modeling CO₂ and CH₄ flux changes in pristine peatlands of Finland under changing climate conditions, *Ecol. Model.*, 263, 64–80, doi:10.1016/j.ecolmodel.2013.04.018, 2013. 10198

Kirschke, S., Bousquet, P., Ciais, P., Saunio, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J.-F., Langenfelds, R. L., Le Quere, C., Naik, V., O’Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P.,

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013. 10197

Kleinen, T., Brovkin, V., and Schuldt, R. J.: A dynamic model of wetland extent and peat accumulation: results for the Holocene, *Biogeosciences*, 9, 235–248, doi:10.5194/bg-9-235-2012, 2012. 10197, 10204, 10216

Kolka, R. K., Sebestyen, S. D., Verry, E. S., and Brooks, K. N.: *Peatland Biogeochemistry and Watershed Hydrology at the Marcell Experimental Forest*, CRC Press, Taylor and Francis Group, 2011. 10213

Law, B., Falge, E., Gu, L., Baldocchi, D., Bakwin, P., Berbigier, P., Davis, K., Dolman, A., Falk, M., Fuentes, J., Goldstein, A., Granier, A., Grelle, A., Hollinger, D., Janssens, I., Jarvis, P., Jensen, N., Katul, G., Mahli, Y., Matteucci, G., Meyers, T., Monson, R., Munger, W., Oechel, W., Olson, R., Pilegaard, K., Paw U, K. T., Thorgeirsson, H., Valentini, R., Verma, S., Vesala, T., Wilson, K., and Wofsy, S.: Environmental controls over carbon dioxide and water vapor exchange of terrestrial vegetation, *Agr. Forest Meteorol.*, 113, 97–120, doi:10.1016/S0168-1923(02)00104-1, {FLUXNET} 2000 Synthesis, 2002. 10198

Limpens, J., Berendse, F., Blodau, C., Canadell, J. G., Freeman, C., Holden, J., Roulet, N., Rydin, H., and Schaepman-Strub, G.: Peatlands and the carbon cycle: from local processes to global implications – a synthesis, *Biogeosciences*, 5, 1475–1491, doi:10.5194/bg-5-1475-2008, 2008. 10196

Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, doi:10.5194/bg-10-753-2013, 2013. 10197

Nungesser, M. K.: Modelling microtopography in boreal peatlands: hummocks and hollows, *Ecol. Model.*, 165, 175–207, doi:10.1016/S0304-3800(03)00067-X, 2003. 10197

Pluchon, N., Hugelius, G., Kuusinen, N., Kuhry, P., Pluchon, N., Hugelius, G., and Kuusinen, N.: The Holocene storage in two boreal peatlands of Northeast European Russia, *The Holocene*, 24, 1126–1136, doi:10.1177/0959683614523803, 2014. 10198

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Porada, P., Weber, B., Elbert, W., Pöschl, U., and Kleidon, A.: Estimating global carbon uptake by lichens and bryophytes with a process-based model, *Biogeosciences*, 10, 6989–7033, doi:10.5194/bg-10-6989-2013, 2013. 10206, 10216
- Runkle, B., Wille, C., Gažovič, M., Wilmking, M., and Kutzbach, L.: The surface energy balance and its drivers in a boreal peatland fen of northwestern Russia, *J. Hydrol.*, 511, 359–373, doi:10.1016/j.jhydrol.2014.01.056, 2014. 10198, 10206, 10209
- Runkle, B. R. K., Wille, C., Gažovič, M., and Kutzbach, L.: Attenuation correction procedures for water vapour fluxes from closed-path eddy-covariance systems, *Bound.-Lay. Meteorol.*, 142, 401–423, doi:10.1007/s10546-011-9689-y, 2012. 10198
- Schneider, J., Kutzbach, L., and Wilmking, M.: Carbon dioxide exchange fluxes of a boreal peatland over a complete growing season, Komi Republic, NW Russia, *Biogeochemistry*, 111, 485–513, doi:10.1007/s10533-011-9684-x, 2012. 10198, 10201, 10205, 10209
- Schuldt, R. J., Brovkin, V., Kleinen, T., and Winderlich, J.: Modelling Holocene carbon accumulation and methane emissions of boreal wetlands – an Earth system model approach, *Biogeosciences*, 10, 1659–1674, doi:10.5194/bg-10-1659-2013, 2013. 10197
- Smith, L. C., MacDonald, G. M., Velichko, A. A., Beilman, D. W., Borisova, O. K., Frey, K. E., Kremenetski, K. V., and Sheng, Y.: Siberian peatlands a net carbon sink and global methane source since the early holocene, *Science*, 303, 353–356, doi:10.1126/science.1090553, 2004. 10196
- Turunen, J., Tomppo, E., Tolonen, K., and Reinikainen, A.: Estimating carbon accumulation rates of undrained mires in Finland–application to boreal and subarctic regions, *The Holocene*, 12, 69–80, doi:10.1191/0959683602hl522rp, 2002. 10197
- Van der Ploeg, M. J., Appels, W. M., Cirkel, D. G., Oosterwoud, M. R., Witte, J.-P., and van der Zee, S.: Microtopography as a driving mechanism for ecohydrological processes in shallow groundwater systems, *Vadose Zone J.*, 11, 52–62, available at: <https://www.soils.org/publications/vzj/abstracts/11/3/>, 2012. 10198
- Walter, B. P. and Heimann, M.: A process-based, climate-sensitive model to derive methane emissions from natural wetlands: application to five wetland sites, sensitivity to model parameters, and climate, *Global Biogeochem. Cy.*, 14, 745–765, 2000. 10204, 10206, 10211, 10213, 10218
- Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model: 1. Evaluation and sensitivity of physical land surface processes, *Global Biogeochem. Cy.*, 23, 1–19, doi:10.1029/2008GB003412, 2009a. 10197

Modeling micro-topographic controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model: 2. Evaluation and sensitivity of vegetation and carbon cycle processes, *Global Biogeochem. Cy.*, 23, 1–15, doi:10.1029/2008GB003413, 2009b. 10197

Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, *Geosci. Model Dev.*, 3, 565–584, doi:10.5194/gmd-3-565-2010, 2010. 10216

Wania, R., Melton, J. R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Chen, G., Eliseev, A. V., Hopcroft, P. O., Riley, W. J., Subin, Z. M., Tian, H., van Bodegom, P. M., Kleinen, T., Yu, Z. C., Singarayer, J. S., Zürcher, S., Lettenmaier, D. P., Beerling, D. J., Denisov, S. N., Prigent, C., Papa, F., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: methodology of a model inter-comparison project (WETCHIMP), *Geosci. Model Dev.*, 6, 617–641, doi:10.5194/gmd-6-617-2013, 2013. 10197

Wieder, R. K., Turetsky, M. R., and Vile, M. A.: Peat as an archive of atmospheric, climatic and environmental conditions, in: *The Wetlands Handbook*, edited by: Maltby, E. and Barker, T., 1 edn., Blackwell Publishing, 96–112, 2009. 10196

Wolf, U.: Above- and belowground methane dynamics of a peatland ecosystem of varying vegetation composition during summer in the Komi Republic, Russia, 2009. 10198, 10210

Yu, Z., Beilman, D. W., Froking, S., MacDonald, G. M., Roulet, N. T., Camill, P., and Charman, D. J.: Peatlands and their role in the global carbon cycle, *EOS T. Am. Geophys. Un.*, 92, 97–98, doi:10.1029/2011EO120001, 2011. 10197

Yu, Z. C.: Northern peatland carbon stocks and dynamics: a review, *Biogeosciences*, 9, 4071–4085, doi:10.5194/bg-9-4071-2012, 2012. 10196




Figure 1. Schematics of the HH model. The model represents a 1 km × 1 km peatland, and works at a 1 m resolution. It is therefore able to resolve the micro-topographical features such as hummocks and hollows. Each grid cell has an elevation which is randomly assigned from the distribution of elevation data collected in the Ust-Pojeg mire complex in the Komi Republic, Russia. For each grid cell we simulate a dynamical water table, which changes with precipitation (P in the figure), evapotranspiration (ET), and lateral runoff (R). In the *Single Bucket* model version all quantities are averaged, in order to assess the effect of the representation of micro-topography if compared to a mean field approximation.

Modeling
micro-topographic
controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. Comparison between the Ust-Pojeg mire topographic data collected in a field survey and a generalized 3-parameters gamma distribution multiplied by maximum number of counts in the histogram. The good visual agreement is confirmed by the positive results from a Kolmogorov–Smirnov test at 95% confidence ($P \geq 0.5$).

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Ensemble simulations of water table dynamics. The *Microtopography* version of the HH model (black line) is compared to the *Single Bucket* version (red line). The water table in the *Microtopography* version is averaged over the whole model region. The micro-topography affects water table position by delaying the runoff, because of the complex interactions among the grid cells, as resolved in a finer scale model. The shaded areas represent the standard deviation over 30 simulations (but their small size renders them invisible in the microtopography case).

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)






Figure 4. Box plots of simulated water table depth in hummocks (orange boxes) and hollows (red boxes) against measurements from the bog (black crosses) and from the rest of the peatland (black circles). Grey crosses are outliers in the simulation. The blue line represents the output of the model without representation of micro-topography. The box plot parameters regulating the length of the whiskers are the default values for covering 99.3% of the data if the data are normally distributed.

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 5. Box plots of simulated methane emissions in hummocks (orange boxes) and hollows (red boxes) against measurements from the bog (black crosses) and from the rest of the peatland (black circles). Grey crosses are outliers in the simulation. The blue line represents the output of the model without representation of micro-topography. The box plot parameters regulating the length of the whiskers are the default values for covering 99.3% of the data if the data are normally distributed.

Modeling
micro-topographic
controls

F. Cresto Aleina et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 7. Dependence of the cumulative methane fluxes computed from the end of April to the end of October on the grid size of the micro-topographic model. The x axis shows the squared root of the number of cells N , as $\sqrt{N} = m_x = m_y$, where m_x and m_y are the resolution horizontally and vertically, respectively. The x axis is on a logarithmic scale. Cumulative methane emissions greatly increase with increasing N if $\sqrt{N} \leq 10$, i.e., decreasing cell size from 1 to 0.01 km² in a 1 km² domain. After this threshold, cumulative methane emissions stabilize as the number of grid cells increases. This phenomenon is mirrored by the average position of the water table dynamics (not shown), which by increasing N asymptotically approaches the water table dynamics for $N = 10^6$.

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 8. Dependence of the cumulative methane fluxes computed from the end of April to the end of October on NPP forcing. The x axis shows the ratio between the simulation NPP (NPP_{Sim}) and the NPP of the *Standard configuration* (NPP_{St}). The vertical scale is the same as in Fig. 7, to show that differences among the simulations are small. Black crosses show cumulative emissions for the model in the *Microtopography* configuration, and red crosses cumulative emissions for the model in the *Single Bucket* configuration.

**Modeling
micro-topographic
controls**

F. Cresto Aleina et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

