

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

This discussion paper is/has been under review for the journal Biogeosciences (BG). Please refer to the corresponding final paper in BG if available.

Carbon export and transfer to depth across the Southern Ocean Great Calcite Belt

S. Z. Rosengard^{1,2}, P. J. Lam^{1,3}, W. M. Balch⁴, M. E. Auro¹, S. Pike¹, D. Drapeau⁴, and B. Bowler⁴

¹Woods Hole Oceanographic Institution, Woods Hole, MA, USA

²Massachusetts Institute of Technology, Cambridge, MA, USA

³University of California, Santa Cruz, CA, USA

⁴Bigelow Laboratory for Ocean Sciences, Boothbay Harbor, ME, USA

Received: 29 December 2014 – Accepted: 6 January 2015 – Published: 10 February 2015

Correspondence to: S. Z. Rosengard (srosengard@whoi.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Sequestration of carbon by the marine biological pump depends on the processes that alter, remineralize and preserve particulate organic carbon (POC) during transit to the deep ocean. Here, we present data collected from the Great Calcite Belt, a calcite-rich band across the Southern Ocean surface, to compare the transformation of POC in the euphotic and mesopelagic zones of the water column. The ^{234}Th -derived export fluxes and size-fractionated concentrations of POC, particulate inorganic carbon (PIC), and biogenic silica (BSi) were measured from the upper 1000 m of 27 stations across the Atlantic and Indian sectors of the Great Calcite Belt. POC export out of the euphotic zone was correlated with BSi export. PIC export was not, but did correlate positively with POC flux transfer efficiency. Moreover, regions of high BSi concentrations, which corresponded to regions with proportionally larger particles, exhibited higher attenuation of $> 51 \mu\text{m}$ POC concentrations in the mesopelagic zone. The interplay among POC size partitioning, mineral composition and POC attenuation suggests a more fundamental driver of POC transfer through both depth regimes in the Great Calcite Belt. In particular, we argue that diatom-dominated communities produce large and labile POC aggregates, which generate high export fluxes but also drive more remineralization in the mesopelagic zone. We observe the opposite in communities with smaller calcifying phytoplankton, such as coccolithophores. We hypothesize that these differences are influenced by inherent differences in the lability of POC exported by different phytoplankton communities.

1 Introduction

The biological pump sequesters atmospheric carbon dioxide (CO_2) in the ocean (Volk and Hoffert, 1985) by way of phytoplankton-driven CO_2 fixation, followed by the sinking of this fixed particulate organic carbon (POC) as aggregates and fecal pellets down the water column (Riley et al., 2012). The quantity per unit area and time of POC exiting

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the base of the euphotic zone defines the export flux, while export efficiency represents the fraction of bulk primary production comprising this flux (Buesseler, 1998). In the mesopelagic zone ~ 1000 m below, export flux attenuates due to remineralization mediated by zooplankton grazing and bacteria (Buesseler and Boyd, 2009; Giering et al., 2014; Martin et al., 1987). The flux of this processed organic carbon leaving the mesopelagic zone, only $\leq 10\%$ of export flux, directly scales with the quantity of atmospheric CO_2 sequestered by the marine biological pump over hundreds to thousands of years.

On average, only $\sim 1\%$ of the organic matter produced by phytoplankton in the surface reaches the deep sea (Martin et al., 1987). However, export and sequestration flux vary widely by region, as do export efficiencies and attenuation of export flux (Buesseler and Boyd, 2009; Buesseler et al., 2007; Henson et al., 2011, 2012b; Martin et al., 1987; Thomalla et al., 2008). Such variations may drive observed differences in the weight percent of organic carbon deposited at the sediment surface (Hedges and Oades, 1997), suggesting that the overall strength of the biological pump as a carbon sink is not globally uniform. These geographical differences have spurred decades of research into how mechanisms in the shallower ocean – the euphotic and mesopelagic zones – alter sinking particulate organic matter during vertical transit.

As an example, Armstrong et al. (2002), Klaas and Archer (2002) and Francois et al. (2002) posited that mineral associations with sinking organic carbon could explain these variations. Their ballast hypothesis model suggested that minerals enhanced the biological pump (1) by increasing the density, and consequently, the sinking speed of particulate organic matter and (2) by inhibiting organic carbon remineralization down the water column. Expediting vertical transit decreases the time for remineralization to act on sinking particulate organic matter, increasing its chances of remaining intact on its way down. The authors observed that calcite flux in the bathypelagic zone (> 1000 m) explains roughly half of the variation in the magnitude of POC flux reaching the deep sea (Klaas and Archer, 2002), and may also account for some of the observed geographical variation in POC flux attenuation with depth (Francois et al., 2002).

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In its conception and infancy, the ballast hypothesis was based upon observed correlations between mineral and organic carbon fluxes in the deep (> 1000 m) sea. Yet, evidence for the ballast mechanism in the euphotic and mesopelagic zones remains equivocal, as deeper correlations are scarcely matched by shallower ocean observations (Le Moigne et al., 2012). Several surface regions do not exhibit ballast correlations between mineral flux and POC flux (e.g., Thomalla et al., 2008; Henson et al., 2012b). In the Atlantic and Southern Oceans, Le Moigne et al. (2012) found a significant fraction of POC export flux to remain unassociated with minerals altogether. Moreover, tank incubations simulating POC and mineral suspensions yield conflicting results: some have observed mineral associations to increase aggregate sinking rates (Engel et al., 2009), while others find no such effect (Passow and De la Rocha, 2006). De La Rocha et al. (2008) even suggest that sticky polymers from POC might ballast sinking minerals, rather than vice-versa.

The scarcity of evidence supporting a shallow ocean ballast mechanism suggests that the transit of particulate organic carbon in the surface, mesopelagic and deeper ocean is mechanistically de-coupled (Lam et al., 2011; Lomas et al., 2010). Indeed, the debate surrounding the ballast hypothesis arises from a deeper issue of whether the mechanisms that influence carbon export from the euphotic zone are the same as those that control its remineralization in the mesopelagic zone, and/or its transfer beyond the mesopelagic zone into the deep sea.

The following report compares the export of organic carbon from the euphotic zone with its transfer through the mesopelagic zone across the region of the Great Calcite Belt (Balch et al., 2011a, 2014; Fig. 1). Spanning across the Southern Ocean, particularly between the Subtropical and Polar Fronts, the Great Calcite Belt defines a highly reflective band observed from space during each austral spring and summer. Its high reflectivity is caused by calcite-rich surface waters produced by coccolithophore blooms in the Southern Ocean. In this zone, coccolithophores are more abundant than in regions north and south of the Belt. South of the Polar Front, coccolithophore

abundances decline dramatically as dissolved silica concentrations increase and diatoms flourish (Balch et al., 2011a).

Spanning a large range in surface mineral concentrations, primary productivity, and phytoplankton community composition (Balch et al., 2011a), the Great Calcite Belt provides an excellent opportunity to assess the processes controlling organic carbon export, export efficiency, and attenuation of POC concentration ([POC]) with depth. Here, we report estimates of ^{234}Th -derived POC fluxes and [POC] through both the euphotic and mesopelagic zones within the Atlantic and Indian sectors of the Great Calcite Belt. We focus on the upper 1000 m of the Great Calcite Belt because the attenuation of POC flux and concentration is most dramatic within this depth interval (Martin et al., 1987; Lam et al., 2011). As the following discussion illuminates, this study additionally weighs the ballast hypothesis against other mechanisms hypothesized to control the transfer of organic carbon through the water column, and ultimately into the deep sea, where carbon residence time modulates atmospheric $p\text{CO}_2$ and climate over hundreds to thousands of years (Kwon et al., 2009).

2 Methods

2.1 Field site

Samples from the Great Calcite Belt were collected during two research cruises, GB1 and GB2, which transited the Atlantic and Indian sectors of the Great Calcite Belt during the austral summer of 2011 and 2012, respectively (Fig. 1), concurrent with the putative coccolithophore bloom (Balch et al., 2011a). In 2011, for cruise GB1 (MV1101), the R/V *Melville* crossed the Atlantic sector from Punta Arenas, Chile to Cape Town, South Africa, sampling between 39 and 59° S. One year later, for cruise GB2 (RR1202), the R/V *Revelle* crossed the Indian sector from Durban, South Africa to Perth, Australia, sampling between 37 and 60° S (Table 1). Both cruise tracks crossed the Subtropical, Subantarctic and Polar fronts, which are approximately located at 40, 45 and 52° S

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(e.g., Belkin and Gordon, 1996; Sokolov and Rintoul, 2009), respectively, defining observed shifts in temperature and nutrient characteristics of the surface ocean.

Each day during GB1 and GB2, 30-L Niskin samples were collected pre-dawn for measuring primary production. A Biospherical Instruments (San Diego, CA) sensor was mounted on the CTD/rosette and referenced to a deck sensor mounted on the ship's superstructure to measure Photosynthetically Available Radiation (PAR) during the casts. Water was then sampled at fixed light depths relative to surface irradiance: 36.5, 21.1, 11.7, 3.55, 1.93 and 0.28 %. The light depths were calculated two ways: (a) between 10:00 and 14:00 LT (during daylight hours), percentages of surface irradiance were derived directly from the downcast PAR profile immediately preceding bottle firing, or (b) at all other times, the light levels were back-calculated from the previously-determined relationship between beam transmittance and diffuse attenuation of PAR (Balch et al., 2011b). From these casts, primary production rates were measured using the ^{14}C microdiffusion technique (Paasche and Brubak, 1994) with modifications by Balch et al. (2000) (see also Fabry and Balch, 2010).

2.2 Size-fractionated particle collection

We report measurements of total and particulate ^{234}Th activity and size-fractionated particle composition from 27 stations (Fig. 1; Table 1).

Size-fractionated particles were collected at eight depths in the upper 1000 m of fourteen stations from GB1 and thirteen stations from GB2, using modified battery operated in situ pumps (McLane WTS-LV). The modified pumps directed seawater through two flow paths (Lam et al., 2014), each of which passed through a "mini-MULVFS" filter holder designed to retain large particles (Bishop et al., 2012). Seawater first passed through 51 μm polyester pre-filters in both filter holders for collection of large ($> 51 \mu\text{m}$) size-fraction particles, and then through paired 0.8 μm polyethersulfone (SuporTM) filters in one flow path and paired 1 μm quartz fiber (WhatmanTM QMA) filters in the other flow path, both of which collected small ($< 51 \mu\text{m}$) size-fraction particles. Immediately after collection, half to all of the $> 51 \mu\text{m}$ size-fraction particles from one

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flow path were rinsed off of the polyester pre-filters and onto 25 mm 1 μm Sterlitech silver filters using 0.2 μm -filtered seawater, and dried at 50 $^{\circ}\text{C}$ for subsequent analysis of particulate ^{234}Th , particulate organic carbon (POC), and particulate inorganic carbon (PIC, or calcium carbonate). Subsamples of QMA filters were likewise dried at 50 $^{\circ}\text{C}$ for ^{234}Th and POC analysis in the < 51 μm size-fraction. Finally, the polyester pre-filters from the other flow path and Supor filters were dried in a laminar flow hood at room temperature.

In the euphotic zone, where most POC is produced, these operationally defined size fractions allude primarily to the structure of phytoplankton communities producing POC (e.g., large diatoms would be found in > 51 μm size-fraction particles). In the mesopelagic zone, which extends from the base of the euphotic zone to 1000 m in depth, > 51 μm POC is predominantly comprised of phytoplankton and bacterial biomass that has been repackaged into aggregates and fecal pellets. The > 51 μm particles collected at station GB1-85 illustrate these different size-fraction interpretations by depth. Shallower particles collected at 25 and 73 m, the base of the euphotic zone, are mainly comprised of intact phytoplankton cells (Fig. 2a and b). By contrast, deeper particles collected at 173 m exhibit the features of particulate aggregates and fecal pellets (Fig. 2c).

2.3 Particle composition

Bulk concentrations of POC, PIC, biogenic silica (BSi), and particulate ^{234}Th activity were measured in both < 51 and > 51 μm fractions of particles collected at each station and depth. Particulate ^{234}Th activities in all sub-fractions of > 51 μm (25 mm silver filters) and < 51 μm (25 mm QMA filters) samples were measured using low level Risø beta counters immediately on the ship and in the lab at least six ^{234}Th half-lives post-collection for background activity.

After counting for ^{234}Th background activity, ~ 25 % of the silver filter (~ 115 L equivalent) was fumed with concentrated hydrochloric acid to remove inorganic carbon, before measuring > 51 μm [POC] using an elemental CHN analyzer. A similar protocol

was followed to measure $< 51 \mu\text{m}$ [POC] from one 12 mm-diameter subsample of each QMA filter, representing $\sim 1\%$ of the entire sample ($\sim 5\text{ L}$ equivalent). Vertical profiles of > 51 and $< 51 \mu\text{m}$ [POC] below the euphotic zone were fitted to a power-law function to describe the attenuation of [POC] with depth, based on a function first applied to POC flux by Martin et al. (1987) and then analogously to POC concentration by Lam and Bishop (2007),

$$[\text{POC}]_z = [\text{POC}]_0 \left(\frac{z}{z_{\text{PAR}}} \right)^{-b} \quad (1)$$

where, at most stations, z_{PAR} represents the depth of 0.3% photosynthetically active radiation (see Sect. 2.4). The exponent b represents the attenuation coefficient, with higher attenuation coefficients (more negative exponents) for profiles with greater attenuation of $> 51 \mu\text{m}$ [POC] with depth. We focus our discussion on the attenuation of $> 51 \mu\text{m}$ [POC], because we assume that they contribute disproportionately to sinking fluxes compared to the $< 51 \mu\text{m}$ size fraction (McCave, 1975; Lam and Bishop, 2007; Lam et al., 2011). Figure 3 and Table 2 show all significant ($p < 0.05$) power law fits for $> 51 \mu\text{m}$ [POC] profiles.

PIC in the samples was assumed to be biomineral calcium carbonate (CaCO_3), and was derived from particulate calcium (Ca) corrected for salt Ca using a seawater 0.0382 Ca : Na (g : g) ratio (Lam and Bishop, 2007; Pilson, 2012). In the in situ pump samples, salt-derived Ca typically accounted for $\sim 60\%$ of total Ca. The $> 51 \mu\text{m}$ PIC size-fraction concentrations were measured mainly in subsamples of remaining pre-filter material and occasionally in sub-fractions of the silver filters, if the former were unavailable. The $< 51 \mu\text{m}$ size fraction [PIC] was measured in three 12 mm circular QMA subsamples, representing $\sim 15\text{ L}$ or $\sim 3\%$ of the sample. Subsamples were leached in 0.6 N ultrapure Sea-StarTM Baseline hydrochloric acid (HCl) at 60°C for 12–16 h. The leachate was subsequently filtered through a $0.4 \mu\text{m}$ polycarbonate membrane filter, diluted to 0.12 N HCl, and spiked with 1 ppb of Indium as an internal standard. The spiked leachate solution was then analyzed for Ca, Na and P using

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

an Element 2 sector-field inductively-coupled plasma mass spectrometer (ICP-MS) in medium and high resolution. Counts per second were converted to concentration using external mixed element standard curves.

For measuring $> 51 \mu\text{m}$ and $< 51 \mu\text{m}$ [BSi], prefilter or Supor subsamples, respectively, were leached in 0.2N sodium hydroxide at 85°C for one hour, and analyzed by standard spectrophotometric detection of the blue silico-molybdate complex in each leachate within 24 h of the leach (Strickland and Parsons, 1968; Brzezinski and Nelson, 1989). Absorbance through each sample was converted to concentration using an external Si standard curve.

2.4 ^{234}Th -derived flux estimates

Particle fluxes were estimated at each station by measuring the water-column disequilibrium between ^{234}Th and ^{238}U in the upper 350 m of the water-column (Savoye et al., 2006). ^{234}Th is the radioactive daughter of ^{238}U with a short enough half-life (24.1 days) relative to ^{238}U such that it is assumed to be in secular equilibrium with its parent isotope in the absence of particle scavenging (i.e., ^{234}Th activity = ^{238}U activity). Disequilibria between the two isotope activities in the water column are attributed to the scavenging of ^{234}Th by sinking particles (Savoye et al., 2006). Integrating the deficit in ^{234}Th relative to ^{238}U provides a measure of particle flux down the water column (Buesseler et al., 2006). Because of the short half-life of ^{234}Th , deviation from secular equilibrium exists only in regions of high particle flux. Thus, ^{234}Th -based flux estimates are most frequently applied in the euphotic zone of the ocean where particle export is maximal.

^{234}Th - ^{238}U deficits were determined by measuring total water-column activities of both isotopes. ^{238}U activity ($A_{\text{U-238}}$) profiles were calculated from salinity by the following relationship (Owens et al., 2011):

$$A_{\text{U-238}} \left(\frac{\text{dpm}}{\text{L}} \right) = (0.0786 \cdot \text{Salinity}) - 0.315 \quad (2)$$

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Total water-column ^{234}Th activity ($A_{\text{Th-234}}$) profiles were determined from 4 L seawater samples collected by CTD casts down to 300–350 m at each station (Pike et al., 2005). Shortly after collection, each 4 L seawater sample was acidified to pH 2 using concentrated nitric acid (HNO_3), spiked with 1 g of ^{230}Th of a known activity (50.06 dpm g^{-1}) as a yield monitor, equilibrated for 8 h, and finally brought up to pH 8.5 using ammonium hydroxide (NH_4OH) (van der Loeff et al., 2006). Manganese chloride (MnCl_2) and potassium permanganate (KMnO_4) were added to the neutralized seawater to form a manganese oxide (MnO_2) precipitate, which efficiently scavenges both natural ^{234}Th and added ^{230}Th . After 12 h, the precipitate was filtered onto a quartz fiber filter, dried at 50 °C, and then mounted beneath a sheet of Mylar and aluminum foil. ^{234}Th activity in the precipitate was measured on board by low level Risø beta counters and post-cruise after at least six ^{234}Th half-lives for background activity. The ^{230}Th spike was recovered by fully dissolving the MnO_2 precipitate, adding a 1 g spike of ^{229}Th of a known activity (69.74 dpm g^{-1}), and measuring $^{229}\text{Th} : ^{230}\text{Th}$ ratios on an Element 2 sector-field ICP-MS in low resolution. Recovery of ^{230}Th spike was derived from this ratio, and used to correct for inefficiencies in the scavenging of total seawater ^{234}Th by MnO_2 precipitation.

To calibrate beta counting efficiency for each cruise, total deep water (i.e., below 2000 m) ^{234}Th activities were compared to total deep water ^{238}U activities, as measured in 4–5 replicate samples from 2 to 3 deep water CTD casts during each cruise (at 5000 m during GB1, and at 2500 m during GB2). Beta counting efficiencies were adjusted such that ^{234}Th and ^{238}U activities were equal in these deep measurements, as secular equilibrium would be expected at such depths. We only report upper water-column activities (< 350 m) after correcting for experimental efficiencies in both the seawater collection process and beta detector counting. Uncertainties in the total ^{234}Th activity profiles averaged 4.5% and were propagated from errors associated with counting statistics, recoveries, and beta-counting efficiency.

To calculate ^{234}Th export flux, ^{234}Th activity deficits were integrated down to the base of the euphotic zone (z_{PAR}) (Buesseler et al., 2008; Thomalla et al., 2008):

$$^{234}\text{Th Flux} \left(\frac{\text{dpm}}{\text{m}^2 \text{d}} \right) = \int_0^{z_{\text{PAR}}} (A_{\text{U-238}} - A_{\text{Th-234}}) dz \quad (3)$$

At most stations, the export depth, z_{PAR} , was chosen to be the depth where light levels were 0.3% of surface-level PAR. The exception was station GB2-27, which did not include a PAR measurement profile. For this station, the z_{PAR} value of 105 m was defined as the depth where the transmissometry-based particle concentration decreased. These export depths were compared to one additional metric describing particle concentration in seawater: the depths where ^{234}Th and ^{238}U activities re-established secular equilibrium, or $z_{\text{Th/U}}$. We explore the sensitivity of ^{234}Th flux estimates to choice of z_{PAR} in Sects. 3 and 4.1.

^{234}Th flux estimates were converted to POC, PIC and BSi fluxes by multiplication with ratios of $> 51 \mu\text{m}$ POC, PIC, and BSi concentrations to particulate ^{234}Th activity in samples at z_{PAR} (Thomalla et al., 2008; Sanders et al., 2010):

$$\text{POC Flux} \left(\frac{\mu\text{mol}}{\text{m}^2 \text{d}} \right) = [\text{POC}] : A_{\text{Th-234}} \cdot ^{234}\text{Th Flux} \left(\frac{\text{dpm}}{\text{m}^2 \text{d}} \right) \quad (4)$$

$$\text{PIC Flux} \left(\frac{\mu\text{mol}}{\text{m}^2 \text{d}} \right) = [\text{PIC}] : A_{\text{Th-234}} \cdot ^{234}\text{Th Flux} \left(\frac{\text{dpm}}{\text{m}^2 \text{d}} \right) \quad (5)$$

$$\text{Si Flux} \left(\frac{\mu\text{mol}}{\text{m}^2 \text{d}} \right) = [\text{BSi}] : A_{\text{Th-234}} \cdot ^{234}\text{Th Flux} \left(\frac{\text{dpm}}{\text{m}^2 \text{d}} \right) \quad (6)$$

2.5 Interpolation of data

In all cases where ^{234}Th activity, > 51 and $< 51 \mu\text{m}$ [POC] and mineral concentrations, and $> 51 \mu\text{m}$ particulate ^{234}Th measurements were unavailable at z_{PAR} , linear

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



interpolations between the sampling depths above and below z_{PAR} were used to estimate a value at the export depth (Table 1). The > 51 and $< 51 \mu\text{m}$ size-fraction POC concentrations were interpolated by the power law attenuation function when fits were significant ($p < 0.05$), or linearly when these power-law fits were not significant or inconsistent with the broader shape of the [POC] profile at that particular station. In general, corresponding POC : ^{234}Th , BSi : ^{234}Th , and PIC : ^{234}Th ratios are quotients of these interpolated values except as noted in Tables 2 and 3.

3 Results

^{234}Th activity profiles were measured over the upper 300–350 m at the 27 stations of cruises GB1 and GB2 (Fig. 4; Table S1 in the Supplement). Each activity profile is associated with two metrics that have been used in previous studies to define the export depth (see Sect. 2.4): the base of the euphotic zone (z_{PAR}), which we define at 0.3 % surface photosynthetically available radiation (PAR) (e.g., Buesseler and Boyd, 2009), and $z_{Th/U}$, where ^{234}Th and ^{238}U activities re-establish secular equilibrium (Table 1). In most stations, profiles exhibited ^{234}Th activity deficits over a range from surface to 75–170 m in depth, below which ^{234}Th activity generally returned to secular equilibrium with ^{238}U activity, within error. The notable exceptions were profiles at stations GB1-6, and GB1-16, which did not return to secular equilibrium by 170 m in depth. Considering that stations GB1-6 and GB1-16 are closest to shore, their sustained ^{234}Th deficits may have been influenced by lateral advection of particles from the continental shelf. At these stations, $z_{Th/U}$ depths were approximated by the depth below which ^{234}Th activities remain constant with depth. For example, at station GB1-6, $z_{Th/U} = 130$ m because below this depth ^{234}Th activities remained relatively constant.

In the Atlantic sector, sampled in January–February 2011, all observed z_{PAR} depths were significantly shallower than $z_{Th/U}$ depths (Student's t test $p < 0.05$); on average, z_{PAR} was 66 ± 44 % shallower than $z_{Th/U}$. By contrast, in the Indian sector, sampled

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



roughly a year later in February–March 2012, z_{PAR} was not significantly different from $z_{\text{Th/U}}$ ($p > 0.05$), and the average relative difference was $-6 \pm 29\%$. In general, when water-column ^{234}Th activity is at steady-state, the euphotic zone should correspond to the region of ^{234}Th deficit relative to ^{238}U (Buesseler et al., 2008; Buesseler and Boyd, 2009), i.e., z_{PAR} should equal $z_{\text{Th/U}}$.

Using integrated activity deficits, export fluxes of ^{234}Th , POC, PIC, and BSi at z_{PAR} were estimated at the 27 sites (Figs. 5 and 6; Table 3). Overall mean ^{234}Th fluxes at z_{PAR} were $1413 \pm 432 \text{ dpm m}^{-2} \text{ d}^{-1}$ (mean ± 1 SD), and ranged from 717 to $2437 \text{ dpm m}^{-2} \text{ d}^{-1}$ at stations GB2-112 and GB1-6, respectively. Mean derived POC fluxes at z_{PAR} were $4.5 \pm 3.9 \text{ mmol m}^{-2} \text{ d}^{-1}$, ranging from 0.97 to $20 \text{ mmol m}^{-2} \text{ d}^{-1}$ at stations GB2-112 and GB1-85, respectively. Mean PIC fluxes were $1.2 \pm 1.7 \text{ mmol m}^{-2} \text{ d}^{-1}$, and ranged from 0.067 to $6.2 \text{ mmol m}^{-2} \text{ d}^{-1}$ at stations GB2-73 and GB1-59, respectively. Finally, mean BSi fluxes at z_{PAR} were $3.8 \pm 5.8 \text{ mmol m}^{-2} \text{ d}^{-1}$, ranging from 0.17 to $28 \text{ mmol m}^{-2} \text{ d}^{-1}$ at stations GB2-46 and GB1-85, respectively. Higher POC export stations frequently corresponded with higher BSi export stations (e.g., station GB1-85), but less so with higher PIC export stations.

The highest and lowest measured biomineral (PIC and BSi) fluxes at z_{PAR} were in GB1 and GB2, respectively, but mean values were not significantly different between ocean basins because of high variability within each basin (Fig. 6). However, mean POC fluxes at z_{PAR} were significantly higher in GB1 (mean ± 1 SD = $6.0 \pm 4.9 \text{ mmol m}^{-2} \text{ d}^{-1}$) than in GB2 ($3.0 \pm 1.7 \text{ mmol m}^{-2} \text{ d}^{-1}$) (Student's t test $p > 0.05$). Because POC: ^{234}Th values did not differ between GB1 and GB2 ($p < 0.05$), we attribute this inter-basin difference in POC fluxes primarily to significantly higher ^{234}Th fluxes in GB1 ($1574 \pm 463 \text{ dpm m}^{-2} \text{ d}^{-1}$) relative to fluxes in GB2 ($1240 \pm 330 \text{ dpm m}^{-2} \text{ d}^{-1}$).

Further, there were significant latitudinal differences among export fluxes and particulate composition ratios in three temperature/nutrient regimes across both sectors (Fig. 1; Table 4): (1) north of 45°S , the approximate location of the Subantarctic

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

front, where temperatures exceeded $\sim 10^\circ\text{C}$; (2) south of 52°S , the approximate location of the Polar Front (e.g., Belkin and Gordon, 1996; Sokolov and Rintoul, 2009), where temperatures remained below $\sim 5^\circ\text{C}$; and (3) between 45°S and 52°S , where temperatures ranged from $\sim 5\text{--}10^\circ\text{C}$. The $> 51\ \mu\text{m}$ size-fraction POC: ^{234}Th values at z_{PAR} were significantly lower in the most equatorward zone north of 45°S , where average ratios were $1.9 \pm 0.9\ \mu\text{mol dpm}^{-1}$. The highest average ratios, south of 52°S , were $5.4 \pm 3.0\ \mu\text{mol dpm}^{-1}$, illustrating the wide variation in POC: ^{234}Th ratios with ecosystem type (Buesseler et al., 2006; Jacquet et al., 2011). Likewise, zonally averaged POC export fluxes in the most equatorward zone ($2.7 \pm 2.3\ \text{mmol m}^{-2}\ \text{d}^{-1}$) were significantly lower than average fluxes in the most poleward zone ($8.0 \pm 6.3\ \text{mmol m}^{-2}\ \text{d}^{-1}$). BSi: ^{234}Th values were significantly different in all three zones, with highest average ratios south of 52°S ($7.1 \pm 4.1\ \mu\text{mol dpm}^{-1}$) and smallest ratios north of 45°S ($0.3 \pm 0.1\ \mu\text{mol dpm}^{-1}$). Similarly, average BSi export fluxes were also significantly different from each other in all three zones, with the greatest average values south of 52°S ($10 \pm 8.7\ \text{mmol m}^{-2}\ \text{d}^{-1}$), and lowest values north of 45°S ($0.35 \pm 0.16\ \text{mmol m}^{-2}\ \text{d}^{-1}$). Finally, PIC: ^{234}Th ratios, which averaged $0.72 \pm 0.85\ \mu\text{mol dpm}^{-1}$ across all zones, and PIC export fluxes were not significantly different from each other in any zone defined by these latitudinal bands.

These fluxes are sensitive to the choice of export depth (z_{PAR} or $z_{\text{Th/U}}$), not only because the export depth determines the magnitude of ^{234}Th flux by influencing the integrated ^{234}Th deficit, but also because the export depth determines which POC: ^{234}Th ratio best describes particles sinking from the chosen depth. Across stations, the depth metrics z_{PAR} and $z_{\text{Th/U}}$ differed from each other to varying extents (Fig. 4; Table 1). As exemplified by stations GB1-92, GB1-16, and GB2-100, POC fluxes changed significantly between z_{PAR} and $z_{\text{Th/U}}$ (Fig. 5b and c; Table S2). At station GB1-92, where z_{PAR} was 40 m shallower than $z_{\text{Th/U}}$, POC flux decreased from $8.0\ \text{mmol m}^{-2}\ \text{d}^{-1}$ at z_{PAR} to $5.1\ \text{mmol m}^{-2}\ \text{d}^{-1}$ at $z_{\text{Th/U}}$. In contrast, at station GB1-16, where z_{PAR} was 80 m shallower than $z_{\text{Th/U}}$, POC fluxes increased from 5.9 to

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

6.6 mmol m⁻² d⁻¹. At station GB2-100, one of few stations where z_{PAR} was deeper than $z_{\text{Th/U}}$, POC fluxes decreased from 3.3 to 1.5 mmol m⁻² d⁻¹ going deeper. At this station, the POC : ²³⁴Th ratio at $z_{\text{Th/U}}$ was 102 % greater than ratios at z_{PAR} , while ²³⁴Th fluxes at $z_{\text{Th/U}}$ were 6 % greater than fluxes at z_{PAR} , demonstrating that changes in particle composition disproportionately contributed to the observed difference in POC export at z_{PAR} and $z_{\text{Th/U}}$. By contrast, at station GB1-16, the relative change in ²³⁴Th fluxes from z_{PAR} to $z_{\text{Th/U}}$ (+29 %) contributed more to the increase in POC flux with depth than the relative change in POC : ²³⁴Th ratio (-13 %). Finally, for station GB1-92, the relative change in ²³⁴Th flux with depth (-19 %) was similar to the relative change in POC : ²³⁴Th with depth (-21 %), demonstrating that the export flux estimate was equally sensitive to changes in both parameters.

4 Discussion

The following discusses these flux measurements in the context of other Southern Ocean observations, and hypotheses surrounding the transformation of sinking organic carbon within the euphotic and mesopelagic zones of the water column.

4.1 Choice of export depth

The two possible depths we use to calculate export flux, z_{PAR} and $z_{\text{Th/U}}$, are significantly different in the Atlantic sector, which influences the magnitude of flux estimated (see Sect. 3). We offer here two possible and not mutually exclusive explanations for why $z_{\text{Th/U}}$ depths were on average deeper than z_{PAR} depths at GB1 stations.

One hypothesis is that the ²³⁴Th-²³⁸U profiles used to calculate export fluxes may not have been at steady-state during the time of sampling on the GB1 cruise. Non-steady state conditions in the ²³⁸U-²³⁴Th system do occur during phytoplankton blooms, particularly during their decline and ascent (Savoye et al., 2006; Buesseler et al., 2009).

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For example, a recent and rapid increase in the near-surface particle concentration could decrease the depth of light penetration faster than the ^{238}U - ^{234}Th system can adjust, leading to a Z_{PAR} measured on station that is shallower than the $Z_{\text{Th/U}}$, which reflects conditions prior to the rapid increase. Since the GB1 cruise in the Atlantic sector took place a month earlier in the growing season (January–February 2011) than the GB2 cruise in the Indian sector (February–March 2012), the two sectors may have been sampled at different stages of the seasonal bloom, contributing to differences in agreement between Z_{PAR} and $Z_{\text{Th/U}}$. Satellite chlorophyll time-series, if well-resolved, can shed light on how dynamic primary production was around the time of sampling at each station of GB1 and GB2, whether rapid (i.e., within three weeks) changes in particle production and sinking fluxes from a bloom could have decoupled ^{234}Th - ^{238}U deficits from light profiles into the surface ocean of the Great Calcite Belt. Eight-day composites of chlorophyll imagery from December 2010 to February 2011 were required to overcome spatial patchiness in the data due to clouds, and indicate that the changes leading up to sampling during GB1 were not consistent across all stations where $Z_{\text{PAR}} < Z_{\text{Th/U}}$. At several stations, productivity declined towards the sampling date; at others, productivity did not change or increased towards the sampling date. Moreover, out of the three stations where $Z_{\text{PAR}} = Z_{\text{Th/U}}$, only one experienced relatively constant primary productivity in the month preceding sampling. In GB2, where the differences between Z_{PAR} and $Z_{\text{Th/U}}$ were not significant, primary productivity tended to be constant preceding more sampling stations. Nonetheless, as in GB1, several locations still experienced increasing or decreasing productivity in the weeks before sampling, despite having a similar Z_{PAR} and $Z_{\text{Th/U}}$.

The inability of the chlorophyll series to unequivocally resolve the differences between Z_{PAR} and $Z_{\text{Th/U}}$ points to other possible mechanisms underlying the discrepancy. One other mechanism, which does not necessarily preclude non-steady state in the ^{234}Th system, is sinking particle production below the euphotic zone Z_{PAR} (Trull et al., 2008). Physical aggregation and fecal pellet production by zooplankton grazing in the region below Z_{PAR} (i.e., the upper mesopelagic zone) can increase

the speed and total abundance of sinking of particles by transforming phytoplankton biomass exiting the euphotic zone, thereby contributing to sustained ^{234}Th deficits below z_{PAR} (Steinberg et al., 2008; Wilson et al., 2008; Abramson et al., 2010). Why this occurs only in GB1 and not GB2 is not known.

For example, the $\sim 70\text{ m}$ difference in z_{PAR} and $z_{\text{Th/U}}$ at a station like GB1-85 (Table 1) may be attributed to additional production or repackaging of sinking particles in the upper mesopelagic zone, causing ^{234}Th deficits to persist beyond the euphotic zone of primary productivity, and a deeper $z_{\text{Th/U}}$. Images of $> 51\ \mu\text{m}$ particles from this station highlight the changing nature of $> 51\ \mu\text{m}$ particles with depth (Fig. 2), from primarily large phytoplankton in the euphotic zone to predominantly fecal pellets in the mesopelagic zone. The difference in POC fluxes measured at both depths may arise from the evolution of these particles during vertical transit, from predominantly intact and relative buoyant diatoms at z_{PAR} to degraded, sinking fecal pellets produced between z_{PAR} and $z_{\text{Th/U}}$.

Going forward, it is most important to keep in mind how the choice of export depth impacts flux estimates. For this study, all export fluxes are defined by z_{PAR} so that they can be compared with integrated primary production measurements (Buesseler and Boyd, 2009). Non-steady-state effects of ^{234}Th profiles on export fluxes will not be considered further because we do not have Lagrangian observations at multiple time points necessary to detect such effects (Buesseler et al., 2003; Resplandy et al., 2012).

4.2 Comparison of export fluxes to previous studies

The ^{234}Th fluxes we report are generally within range of measurements from other Southern Ocean studies (e.g., Le Moigne et al., 2013; Shimmield et al., 1995; Rutgers Van Der Loeff et al., 1997, 2011; Buesseler, 1998, 2001, 2003; Cochran et al., 2000; Friedrich and van der Loeff, 2002; Coppola et al., 2005; Morris et al., 2007; Thomalla et al., 2008; Savoye et al., 2008; Rodriguez y Baena et al., 2008; Jacquet et al., 2011; Zhou et al., 2012; Planchon et al., 2013). By contrast, the POC fluxes we report are on average three times lower due to lower POC: ^{234}Th

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



biomineral fluxes estimated by Henson et al. (2012b), respectively, who used a steady-state model of nutrient uptake against nutrient export (Sarmiento et al., 2002, 2004). The Henson et al. method used annual climatologies of nutrient concentration profiles for their estimates, whereas the ^{234}Th -derived export method used here integrates over several weeks in the growing season. This difference in timescales of integration likely accounts for the smaller biomineral fluxes in Henson et al. (2012b).

4.3 Export efficiency

We found no significant relationship observed between integrated primary productivity and POC flux at z_{PAR} , highlighting the variable export efficiency across GB1 and GB2. Export efficiencies, or “Ez-ratios” (Buesseler and Boyd, 2009), were calculated as the ratio of POC flux at z_{PAR} to total integrated primary production in the euphotic zone (Fig. 7b; Table 3). Mean export efficiencies were 0.26 ± 0.19 , and ranged from 0.04 to 0.77 at stations GB1-16 and GB2-63, respectively. The lack of association between primary productivity and POC export flux confirms previously observed decoupling between the factors that drive export and those that modulate primary productivity (Buesseler et al., 2001; Coppola et al., 2005; Maiti et al., 2010).

4.4 Vertical attenuation of POC flux and concentration

At most stations, both POC flux and $> 51 \mu\text{m}$ [POC] decline with depth below z_{PAR} as a result of remineralization. In the following, we use two metrics to describe POC transfer in the mesopelagic zone: (1) the attenuation of $> 51 \mu\text{m}$ [POC] in the mesopelagic zone, expressed as the attenuation coefficients extracted from power-law fits of mesopelagic $> 51 \mu\text{m}$ [POC] (exponent from Eq. 1) and (2) the POC flux transfer efficiency (T_{100}), defined as the fraction of ^{234}Th -based POC flux that survives remineralization and is transferred 100 m below z_{PAR} (Buesseler and Boyd, 2009). The first metric describes the disappearance of POC concentration, and applies to the

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



entire mesopelagic zone; the second metric describes the survival of POC flux, and applies to the upper mesopelagic zone.

The mean T_{100} was 0.71 ± 0.38 , ranging from 0.20 to 1.8 at stations GB2-119 and GB1-25, respectively (Fig. 7c; Table 2), generally falling within the spread of values observed globally as well as specifically in the Southern Ocean (Buesseler and Boyd, 2009). At stations GB1-6, GB1-16, GB1-25, GB1-59, and GB2-106, T_{100} values are greater than 1.0 and reflect an increase in POC flux with depth between Z_{PAR} and 100 m below Z_{PAR} (Fig. 5b and d). At GB1-6, GB1-16 and GB1-59, the ^{234}Th - ^{238}U disequilibrium extends relatively deep (> 200 m) into the water column, thus leading to continually increasing ^{234}Th flux with depth, suggesting that either renewed particle production at depth or lateral advection of particles away from these coastal stations could sustain the ^{234}Th deficit below Z_{PAR} . Moreover, because Z_{PAR} depths are significantly shallower than $Z_{Th/U}$ in most GB1 stations, including GB1-6, GB1-16 and GB1-59, the transfer efficiency calculation at these stations in GB1 captures an increase in ^{234}Th flux between Z_{PAR} and 100 m below Z_{PAR} . Thus, for the following discussion, it is important to view transfer efficiency values with the caveat that GB1 and GB2 stations display different ^{234}Th - ^{238}U disequilibria profiles with respect to Z_{PAR} and $Z_{Th/U}$, and this difference impacts all calculations that use a ^{234}Th flux component.

At the two other stations for which $T_{100} > 1$, GB1-25 and GB2-106, the increases in POC flux below Z_{PAR} arise primarily from increasing POC : ^{234}Th ratios rather than increasing ^{234}Th flux with depth. The increase in these ratios results from a faster decrease in particulate ^{234}Th activity compared to changes in $> 51 \mu\text{m}$ [POC] with depth. This is unexpected and all other stations where $> 51 \mu\text{m}$ [POC] decreases more quickly than particulate ^{234}Th activity due to organic carbon remineralization. We suspect a sampling issue and thus exclude the T_{100} transfer efficiencies from these two stations from further consideration.

The general decline in POC flux with depth at most stations is mirrored by a decrease in $> 51 \mu\text{m}$ [POC], both of which are a result of remineralization. Attenuation coefficients from power-law fits of mesopelagic $> 51 \mu\text{m}$ [POC] at 22 stations describe

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shown) arise from differences in interpolation methods for POC : ^{234}Th , BSi : ^{234}Th , and PIC : ^{234}Th ratios at z_{PAR} (Table 3).

Pearson correlation tests between shallow POC export and the two biomineral fluxes revealed a significantly positive correlation between POC and BSi fluxes ($p \ll 0.001$, $r^2 = 0.77$). By contrast, there was no significant relationship between shallow POC and PIC fluxes ($p = 0.24$, $r^2 = 0.06$). Both BSi and POC export fluxes tend to increase poleward from the region north of the Subtropical/Subantarctic fronts to the inter-frontal zone to the region south of the Polar front (Figs. 5b, 6a and b). Station GB1-85, which sits just south of the Polar Front ($\sim 52^\circ \text{S}$), is a high BSi and POC flux outlier. When removed, the BSi flux vs. POC flux correlation remains significant, though weaker ($r^2 = 0.43$), suggesting that although this correlation is strongly influenced by station GB1-85, the shallow BSi ballast association still remains valid for the rest of the dataset.

We also compared POC export fluxes to both PIC and BSi export fluxes simultaneously by multiple linear regression:

$$\text{POC Flux} = (m_{\text{BSi}} \cdot \text{BSi Flux}) + (m_{\text{PIC}} \cdot \text{PIC Flux}) + \text{constant} \quad (7)$$

The multiple linear regression only explains an additional 5% of the variance in POC flux at z_{PAR} ($r^2 = 0.82$, $p \ll 0.001$), affirming that BSi flux explains most of the variation in POC export fluxes at z_{PAR} across the Atlantic and Indian sectors of the Great Calcite Belt region.

The carrying capacities of BSi and PIC for POC, as defined by the slopes m_{BSi} and m_{PIC} in the multiple linear regression, are 0.60 and 0.50, on a per-mole basis, and 0.23 and 0.13, on a per-weight basis, respectively, assuming $12 \times 2.199 \text{ g mol}^{-1} \text{ POC} \times 2.199$, $67.3 \text{ g SiO}_2 \cdot 0.4 \text{ H}_2\text{O mol}^{-1} \text{ BSi}$ and $100.1 \text{ g CaCO}_3 \text{ mol}^{-1} \text{ PIC}$ (Klaas and Archer, 2002). The unassociated POC flux, as defined by the constant in Eq. (7), is $1.7 \text{ mmol POC m}^{-2} \text{ d}^{-1}$, or $44 \text{ mg POC m}^{-2} \text{ d}^{-1}$. These carrying capacities for POC are 2–10 times higher than global biomineral carrying capacities of deeper ($> 2000 \text{ m}$) flux data ($m_{\text{BSi}} = 0.025\text{--}0.026$, $m_{\text{PIC}} = 0.070\text{--}0.074$; Klaas and Archer, 2002), reflecting how POC remineralization with depth

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

consistently reduces apparent mineral carrying capacities between the base of the euphotic zone and the deep sea.

These upper ocean carrying capacities, especially m_{PIC} , are considerably different than corresponding per-weight carrying capacities reported in the *Crozex* study in the Indian sector of the Southern Ocean ($m_{\text{BSi}} = 0.16$, $m_{\text{PIC}} = -0.11$, constant = $105 \text{ mg POC m}^{-2} \text{ d}^{-1}$) (Le Moigne et al., 2012). But, as the *Crozex* study was carried out several months earlier in the growing season than our sampling of the same area within the Great Calcite Belt, seasonal changes in the phytoplankton communities and their associated food webs could account for the differences in upper ocean carrying capacities. The Le Moigne et al. (2012) study also highlighted that variable ecosystem composition contributed to regional variations in upper ocean carrying capacities (Le Moigne et al., 2014), echoing a contemporaneous study that showed that even the deep ($> 1500 \text{ m}$) flux carrying capacities have statistically significant spatial variability (Wilson et al., 2012).

It is worth noting that Le Moigne et al. (2012) included lithogenic minerals in their multiple linear regressions. We did not measure lithogenic minerals on GB1 and GB2, as we assumed lithogenic fluxes to be small in the Southern Ocean due to low terrestrial dust inputs (e, g., Honjo et al., 2000). While omitting this lithogenic component from the multiple linear regression could potentially impact derived m_{BSi} and m_{PIC} values, lithogenic material is nonetheless unlikely to be an important carrier of POC flux because of its low flux in the Southern Ocean. Indeed, regional studies have found that the lithogenic carrying capacity (Wilson et al., 2012) and the lithogenic-associated POC fluxes (Le Moigne et al., 2012) are very low in the Southern Ocean.

4.6 Mineral-POC flux correlations in the mesopelagic zone

To directly test whether minerals facilitate POC transfer through the upper mesopelagic zone of the water column as well, we compared flux transfer efficiencies 100 m below the base of the euphotic zone (T_{100}) with BSi and PIC fluxes at Z_{PAR} (Fig. 9c and d). If the mineral ballast model were to apply to the upper mesopelagic zone, one would

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



expect greater transfer efficiencies (i.e., lower attenuation of POC flux) in regions of higher mineral export. The data highlight several key differences between the role of minerals in the euphotic and upper mesopelagic zones. For one, the correlation between PIC flux and T_{100} , excluding values at GB1-25 and GB2-106, is significantly positive ($p < 0.001$, $r^2 = 0.39$). The relationship remains even when assessing data from each cruise individually (for GB1, $p = 0.047$, $r^2 = 0.34$; for GB2, $p = 0.009$, $r^2 = 0.52$), lending further support to a potential role for PIC in POC transfer through the upper mesopelagic zone.

Further, there was no significant correlation, with or without GB1-25 and GB2-106 T_{100} values, between BSi export fluxes in GB2 and T_{100} . However, higher particulate biogenic silica concentrations ($> 51 \mu\text{m}$ [BSi]) at z_{PAR} did correspond with greater attenuation of $> 51 \mu\text{m}$ [POC] below z_{PAR} ($p = 0.004$, $r^2 = 0.35$; Fig. 10a), suggesting that in contrast to its role in the euphotic zone, BSi is associated with greater degradation in the mesopelagic zone of the water column.

4.7 Other controls on POC transfer

The correlation between the attenuation of $> 51 \mu\text{m}$ [POC] and the size fractionation of POC ($\% > 51 \mu\text{m}$ [POC]) at z_{PAR} is even stronger than with $> 51 \mu\text{m}$ [BSi] ($p \ll 0.001$, $r^2 = 0.63$; Fig. 10b). GB1-85 appears to be an outlier for both relationships in Fig. 10, but especially for the relationship between $> 51 \mu\text{m}$ [POC] attenuation and $> 51 \mu\text{m}$ [BSi] (Fig. 10a). The correlation remains significant when the high [BSi] value from station GB1-85 is removed. Notably, the power law fit at GB1-85 is not very good in the upper mesopelagic; fitting $> 51 \mu\text{m}$ [POC] between z_{PAR} and 500 m yields a better fit (higher r^2 ; see Fig. 3) with a higher attenuation coefficient of 2.35 (compared to 1.7 for the entire mesopelagic zone). This modified upper mesopelagic attenuation at GB1-85 improves the overall correlations between the attenuation coefficient and both $> 51 \mu\text{m}$ [BSi] ($p \ll 0.001$, $r^2 = 0.60$) and $\% > 51 \mu\text{m}$ [POC] ($p \ll 0.001$; $r^2 = 0.78$),

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



further strengthening the argument that $> 51 \mu\text{m}$ [BSi] and $\% > 51 \mu\text{m}$ [POC] at z_{PAR} are important factors in POC transfer in the upper mesopelagic zone.

The relationships between the attenuation of $> 51 \mu\text{m}$ [POC] and $> 51 \mu\text{m}$ [BSi] and particle size fractionation may arise from a more fundamental feature shared by both BSi-rich and large-particle stations of the Great Calcite Belt: diatom-dominated phytoplankton communities. Indeed, we also observe a strong correlation between $> 51 \mu\text{m}$ [BSi] and $\% > 51 \mu\text{m}$ [POC] at z_{PAR} ($p \ll 0.001$, $r^2 = 0.65$; not shown). This is a consistent feature across diatom-rich populations, which produce large, BSi-rich organic aggregates that sink rapidly out of the euphotic zone (Michaels and Silver, 1988; Buesseler, 1998; Thomalla et al., 2006). Thus, characteristics describing ecosystem structure may underlie the correlation between BSi export and POC export in the Great Calcite Belt (Francois et al., 2002; Thomalla et al., 2008; Henson et al., 2012a, b).

However, ecosystem composition does not directly explain why larger particles exported into the mesopelagic zone are remineralized more vigorously hundreds of meters below (Fig. 10b). It is paradoxical that the same large particles that sink quickly out of the euphotic zone would then remineralize faster, as well. This association between attenuation coefficient and particle size suggests that these particles sink more slowly than expected in the mesopelagic zone given their size (for example, as a result of high porosity and low excess density), and/or that they are subject to faster remineralization compared to regions with more POC in the small size fraction. Francois et al. (2002) noted a negative relationship between bathypelagic transfer efficiency and opal flux, and attributed this to increased lability in large diatom aggregates. Though we do not observe any negative correlation between upper mesopelagic transfer efficiency (T_{100}) and BSi fluxes at z_{PAR} , we suggest that potentially higher degradability of POC produced by diatom-dominated communities may similarly explain the relationship between particle size and $> 51 \mu\text{m}$ [POC] attenuation in the upper mesopelagic zone.

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The view of POC quality as a driving factor behind POC transfer argues for a deterministic role of euphotic zone community structure in POC transfer below the euphotic zone. It supports the conventional perspective that diatom-dominated communities are strong exporters of large, sinking POC particles out of the euphotic zone (Buesseler, 1998; Guidi et al., 2009), but also adds to the growing view that these communities have poor transfer efficiency and high attenuation through the mesopelagic zone (Francois et al., 2002; Guidi et al., 2009; Henson et al., 2012b, a).

For instance, station GB1-85, with over half of [POC] in the $> 51 \mu\text{m}$ size class fraction in the euphotic zone (Fig. 6c; Table 3), has a low $> 51 \mu\text{m}$ [PIC]:[BSi] ratio of 0.035 at z_{PAR} (indicated in log-scale in Fig. 10a and b), which indicates a community dominated by diatoms producing large BSi-rich aggregates (Fig. 2). Station GB1-85 exhibits a high export efficiency (Ez-ratio = 0.38, within the upper quartile of the data set), and the highest $> 51 \mu\text{m}$ [POC] and export fluxes at z_{PAR} (Figs. 5b, 7b, and 8a; Table 3). Notably, $> 51 \mu\text{m}$ [POC] values in the lower mesopelagic zone are also the highest at GB1-85, despite attenuating greatly below z_{PAR} (attenuation coefficient = 1.7) (Figs. 3, 8b and c; Table 2). But, because of high attenuation, proportionally less organic carbon transfers to the deep sea at GB1-85. The same diatom-rich communities that vigorously export POC also have a short length scale of remineralization, and ultimately may not sequester as much organic carbon in the deep ocean or draw down as much atmospheric CO_2 (Kwon et al., 2009) as would be expected considering the magnitude of export alone.

In contrast to the model diatom community, station GB1-25 is BSi-deplete, with a $> 51 \mu\text{m}$ [PIC]:[BSi] ratio of 4.4 at z_{PAR} (indicated in log-scale in Fig. 10a), indicating the dominance of coccolithophores over diatoms. With proportionally less POC in the $> 51 \mu\text{m}$ size-fraction (only 3.2%) (Figs. 6c and 10b; Table 3), $> 51 \mu\text{m}$ [POC] at GB1-25 attenuates little through the mesopelagic zone (attenuation coefficient = 0.4, the lowest of the data set) such that a third of the $> 51 \mu\text{m}$ [POC] at z_{PAR} remains at 1000 m, compared to only 1.4% at station GB1-85 (Fig. 3). At GB1-25, export efficiency is very low (Ez-ratio = 0.04), suggesting that the particles exiting the euphotic zone here have

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

been recycled vigorously in the euphotic zone prior to export, which may explain their low $> 51 \mu\text{m}$ [POC] and high proportion in the $< 51 \mu\text{m}$ size-fraction at Z_{PAR} . In the mesopelagic zone, these particles are not very reactive and thus remineralize very little, perhaps sequestering a higher proportion of the CO_2 fixed in the euphotic zone.

Several other stations with proportionally more small particles and weaker $> 51 \mu\text{m}$ [POC] attenuation in the mesopelagic zone exhibit higher $> 51 \mu\text{m}$ [PIC] than $> 51 \mu\text{m}$ [BSi] at Z_{PAR} (labeled in the lower left quadrant of Fig. 10b), suggesting that export regimes dominated by coccolithophores over diatoms consistently transfer less reactive POC to the mesopelagic zone. Artificial roller tank experiments have demonstrated that coccolithophore cultures can produce smaller, more compact aggregates than diatom cultures, partly because of smaller cell sizes (Iversen and Ploug, 2010). However, smaller size does not necessarily mean slower sinking velocities (e.g., McDonnell and Buesseler, 2010). Iversen and Ploug (2010) showed that the higher excess density of these small aggregates generated faster sinking speeds than similarly sized pure diatom aggregates. Another roller tank study that compared aggregate formation by calcifying vs. non-calcifying coccolithophores observed that aggregates formed from calcifying coccolithophores were smaller but faster sinking (Engel et al., 2009). In regions like the Great Calcite Belt, dense coccolithophore populations may similarly export small, highly degraded and compact particles out of the euphotic zone. As a result, the remineralization length scale of calcareous assemblages is likely to be longer than regions dominated by diatom assemblages. These communities would efficiently transfer POC towards the base of the mesopelagic zone, even if the magnitude of exported POC is not as high as in diatom-dominated regions (Thomalla et al., 2008; Guidi et al., 2009; Henson et al., 2012b). This may explain why higher PIC export fluxes are associated with higher transfer efficiencies but not higher POC flux at Z_{PAR} (Fig. 9), and also why the ballast association between PIC and POC fluxes appears only at greater depths (Francois et al., 2002; Klaas and Archer, 2002).

Attenuation coefficients for $> 51 \mu\text{m}$ [POC] across diatom-dominated regions exhibit a great spread ($\text{SD} = 0.47$), ranging from 0.47 to 1.88, such that not all stations

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dominated by diatoms (i.e., $> 51 \mu\text{m [PIC] : [BSi]} < 1$ at z_{PAR}) have proportionally larger particles or higher b values (e.g., stations GB1-70, GB1-77 and GB2-87; Fig. 10b). In contrast, attenuation coefficients across coccolithophore-dominated regions (i.e., $> 51 \mu\text{m [PIC] : [BSi]} \geq 1$ at z_{PAR}) exhibit a lower SD (0.31) and a smaller range, 0.35–1.12. The greater variance in attenuation across BSi-rich regions may result from sampling the diatom populations at different seasons of the bloom cycle (Lam et al., 2011), and implies that there may be less seasonality in POC transfer to depth in coccolithophore-dominated regions. Indeed, massive diatom export events with high transfer efficiency through the mesopelagic zone have been observed (Martin et al., 2011; Smetacek et al., 2012), so there are clearly conditions that can lead to efficient mesopelagic POC transfer from diatom blooms.

It is worth noting that $> 51 \mu\text{m [PIC] : [Si]}$ ratios did increase with depth at most stations of the Great Calcite Belt, as might be expected because BSi is undersaturated in seawater. The possibility that BSi dissolves faster than PIC in particles sinking through the mesopelagic zone would complicate the connections we draw between diatom-dominated communities in the euphotic zone and the attenuation of $> 51 \mu\text{m [POC]}$. But, there are no associations between the magnitude of $[PIC] : [BSi]$ increase and $> 51 \mu\text{m [BSi]}$ at z_{PAR} , $> 51 \mu\text{m [PIC]}$ at z_{PAR} or $> 51 \mu\text{m [POC]}$ attenuation with depth, suggesting that the issue of differential dissolution should not significantly impact our earlier interpretations. In the future, directly evaluating the degradability of sinking POC using organic characterization techniques (e.g., ramped pyrolysis or biomarker isolation) (e.g., Wakeham et al., 2002; Rosenheim et al., 2008, 2013; Rosenheim and Galy, 2012) would greatly improve our ability to track the transformation of POC produced by different ecosystem assemblages across the Great Calcite Belt.

5 Conclusion

In summary, we argue here that phytoplankton assemblages play a fundamental role (Francois et al., 2002; Thomalla et al., 2008; Henson et al., 2012a, b) in determining the

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fate of POC export through the Great Calcite Belt region, the effect of which sometimes, but not always, appears as a mineral ballast mechanism in the euphotic zone (Lam et al., 2011; Henson et al., 2012a; Lima et al., 2013). Though shallow BSi export fluxes were strongly correlated with POC export fluxes, they are also associated with diatom communities that produce larger particles that attenuate more quickly through the mesopelagic zone, such that proportionally less POC reaches the lower mesopelagic zone, and proportionally more is returned to the water column as remineralized carbon (dissolved inorganic and organic carbon).

The Supplement related to this article is available online at
doi:10.5194/bgd-12-2843-2015-supplement.

Author contributions. S. Z. Rosengard, the primary author, participated in the GB2 field work, sample analysis in lab and writing. P. J. Lam contributed to field work during GB2, and participated in both data interpretation and editing the manuscript. W. M. Balch, supplied the primary productivity and light profile data included here, and provided valuable feedback during writing. M. E. Auro and S. Pike participated in field work and sample analysis during and after GB1. D. Drapeau and B. Bowler contributed to field work during both GB1 and GB2, as well as sample analysis, particularly with respect to primary productivity and light profile data.

Acknowledgements. Many thanks to Ken Buesseler for discussions and lending us field equipment; Dan Ohnemus, Angela Warner, Michael Brown, Rebecca Fowler, and Marina Van der Eb for help at sea; and Laura Lubelczyk, Paul Henderson, and Scott Birdwhistell for analytical work/assistance. This work was funded by NSF OCE-0960880 to P. J. Lam, and NSF OCE-0961660, NASA NNX11A072G and NNX11AL93G to W. M. Balch.

References

Abramson, L., Lee, C., Liu, Z. F., Wakeham, S. G., and Szlosek, J.: Exchange between suspended and sinking particles in the northwest Mediterranean as inferred from the organic

composition of in situ pump and sediment trap samples, *Limnol. Oceanogr.*, 55, 725–739, 2010.

Armstrong, R. A., Lee, C., Hedges, J. I., Honjo, S., and Wakeham, S. G.: A new, mechanistic model for organic carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals, *Deep-Sea Res. Pt. II*, 49, 219–236, 2002.

Balch, W. M., Drapeau, D., and Fritz, J.: Monsoonal forcing of calcification in the Arabian Sea, *Deep-Sea Res. Pt. II*, 47, 1301–1337, 2000.

Balch, W. M., Drapeau, D. T., Bowler, B. C., Lyczkowski, E., Booth, E. S., and Alley, D.: The contribution of coccolithophores to the optical and inorganic carbon budgets during the Southern Ocean Gas Exchange Experiment: new evidence in support of the Great Calcite Belt hypothesis, *J. Geophys. Res.*, 116, C00F06, doi:10.1029/2011JC006941, 2011a.

Balch, W. M., Poulton, A. J., Drapeau, D. T., Bowler, B. C., Windecker, L. A., and Booth, E. S.: Zonal and meridional patterns of phytoplankton biomass and carbon fixation in the Equatorial Pacific Ocean, between 110° W and 140° W, *Limnol. Oceanogr.*, 59, 1715–1732, 2011b.

Balch, W. M., Drapeau, D. T., Bowler, B. C., Lyczkowski, E. R., Lubelczyk, L. C., Painter, S. C., and Poulton, A. J.: Surface biological, chemical, and optical properties of the Patagonian Shelf coccolithophore bloom, the brightest waters of the Great Calcite Belt, *Limnol. Oceanogr.*, 59, 1715–1732, 2014.

Belkin, I. M. and Gordon, A. L.: Southern Ocean fronts from the Greenwich meridian to Tasmania, *J. Geophys. Res.-Oceans*, 101, 3675–3696, 1996.

Bishop, J. K. B., Lam, P. J., and Wood, T. J.: Getting good particles: accurate sampling of particles by large volume in-situ filtration, *Limnol. Oceanogr.-Meth.*, 10, 681–710, 2012.

Brzezinski, M. A. and Nelson, D. M.: Seasonal changes in the silicon cycle within a Gulf Stream warm-core ring, *Deep-Sea Res.*, 36, 1009–1030, 1989.

Buesseler, K., Lamborg, C. H., Cai, P., Escoube, R., Johnson, R., Pike, S., Masque, P., McGillicuddy, D., and Verdeny, E.: Particle fluxes associated with mesoscale eddies in the Sargasso Sea, *Deep-Sea Res. Pt. II*, 55, 1426–1444, 2008.

Buesseler, K. O.: The decoupling of production and particulate export in the surface ocean, *Global Biogeochem. Cy.*, 12, 297–310, 1998.

Buesseler, K. O. and Boyd, P.: Shedding light on processes that control particle export and flux attenuation in the twilight zone of the open ocean, *Limnol. Oceanogr.*, 54, 1210–1232, 2009.

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Buesseler, K. O., Ball, L., Andrews, J., Cochran, J. K., Hirschberg, D. J., Bacon, M. P., Flerer, A., and Brzezinski, M.: Upper ocean export of particulate organic carbon and biogenic silica in the Southern Ocean along 170° W, *Deep-Sea Res. Pt. II*, 48, 4275–4297, 2001.

Buesseler, K. O., Barber, R. T., Dickson, M. L., Hiscock, M. R., Moore, J. K., and Sambrotto, R.: The effect of marginal ice-edge dynamics on production and export in the Southern Ocean along 170° W, *Deep-Sea Res. Pt. II*, 50, 579–603, 2003.

Buesseler, K. O., Benitez-Nelson, C. R., Moran, S. B., Burd, A., Charette, M., Cochran, J. K., Coppola, L., Fisher, N. S., Fowler, S. W., and Gardner, W. D.: An assessment of particulate organic carbon to thorium-234 ratios in the ocean and their impact on the application of ²³⁴Th as a POC flux proxy, *Mar. Chem.*, 100, 213–233, doi:10.1016/j.marchem.2005.10.013, 2006.

Buesseler, K. O., Lamborg, C. H., Boyd, P. W., Lam, P. J., Trull, T. W., Bidigare, R. R., Bishop, J. K. B., Casciotti, K. L., Dehairs, F., Elskens, M., Honda, M., Karl, D. M., Siegel, D. A., Silver, M. W., Steinberg, D. K., Valdes, J., Van Mooy, B., and Wilson, S.: Revisiting carbon flux through the ocean's twilight zone, *Science*, 316, 567–570, 2007.

Buesseler, K. O., Pike, S., Maiti, K., Lamborg, C. H., Siegel, D. A., and Trull, T. W.: Thorium-234 as a tracer of spatial, temporal and vertical variability in particle flux in the North Pacific, *Deep-Sea Res. Pt. I*, 56, 1143–1167, 2009.

Cochran, J. K., Buesseler, K. O., Bacon, M. P., Wang, H. W., Hirschberg, D. J., Ball, L., Andrews, J., Crossin, G., and Flerer, A.: Short-lived thorium isotopes (Th-234, Th-228) as indicators of POC export and particle cycling in the Ross Sea, Southern Ocean, *Deep-Sea Res. Pt. II*, 47, 3451–3490, 2000.

Coppola, L., Roy-Barman, M., Mulrow, S., Povinec, P., and Jeandel, C.: Low particulate organic carbon export in the frontal zone of the Southern Ocean (Indian sector) revealed by Th-234, *Deep-Sea Res. Pt. I*, 52, 51–68, 2005.

De La Rocha, C. L., Nowald, N., and Passow, U.: Interactions between diatom aggregates, minerals, particulate organic carbon, and dissolved organic matter: further implications for the ballast hypothesis, *Global Biogeochem. Cy.*, 22, GB4005, doi:10.1029/2007GB003156, 2008.

Engel, A., Szlosek, J., Abramson, L., Liu, Z., and Lee, C.: Investigating the effect of ballasting by CaCO₃ in *Emiliania huxleyi*: I. Formation, settling velocities and physical properties of aggregates, *Deep-Sea Res. Pt. II*, 56, 1396–1407, 2009.

Fabry, V. J. and Balch, W. M.: Direct measurements of calcification rates in planktonic organisms, in: *Guide to Best Practices in Ocean Acidification Research and Data Reporting*,

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



edited by: Riebeseil, U., Fabry, V. J., Hansson, L., and Gattuso, J.-P., European Project on Ocean Acidification (EPOCA), Bremerhaven, Germany, 185–196, 2010.

Francois, R., Honjo, S., Krishfield, R., and Manganini, S.: Factors controlling the flux of organic carbon to the bathypelagic zone of the ocean, *Global Biogeochem. Cy.*, 16, 1087, doi:10.1029/2001GB001722, 2002.

Friedrich, J. and van der Loeff, M. M. R.: A two-tracer (Po-210-Th-234) approach to distinguish organic carbon and biogenic silica export flux in the Antarctic Circumpolar Current, *Deep-Sea Res. Pt. I*, 49, 101–120, 2002.

Giering, S. L. C., Sanders, R., Lampitt, R. S., Anderson, T. R., Tamburini, C., Boutrif, M., Zubkov, M. V., Marsay, C. M., Henson, S. A., Saw, K., Cook, K., and Mayor, D. J.: Reconciliation of the carbon budget in the ocean's twilight zone, *Nature*, 507, 480–483, 2014.

Glover, D. M., Jenkins, W. J., and Doney, S. C.: *Modeling Methods for Marine Science*, Cambridge University Press, New York, 2011.

Guidi, L., Stemmann, L., Jackson, G. A., Ibanez, F., Claustre, H., Legendre, L., Picheral, M., and Gorsky, G.: Effects of phytoplankton community on production, size and export of large aggregates: a world-ocean analysis, *Limnol. Oceanogr.*, 54, 1951–1963, 2009.

Hedges, J. I. and Oades, J. M.: Comparative organic geochemistries of soils and marine sediments, *Org. Geochem.*, 27, 319–361, 1997.

Henson, S., Lampitt, R., and Johns, D.: Variability in phytoplankton community structure in response to the North Atlantic Oscillation and implications for organic carbon flux, *Limnol. Oceanogr.*, 57, 1591–1601, doi:10.4319/lo.2012.57.6.1591, 2012a.

Henson, S. A., Sanders, R., Madsen, E., Morris, P. J., Le Moigne, F., and Quartly, G. D.: A reduced estimate of the strength of the ocean's biological carbon pump, *Geophys. Res. Lett.*, 38, L04606, doi:10.1029/2011GL046735, 2011.

Henson, S. A., Sanders, R., and Madsen, E.: Global patterns in efficiency of particulate organic carbon export and transfer to the deep ocean, *Global Biogeochem. Cy.*, 26, GB1028, doi:10.1029/2011GB004099, 2012b.

Honjo, S., Francois, R., Manganini, S., Dymond, J., and Collier, R.: Particle fluxes to the interior of the Southern Ocean in the Western Pacific sector along 170° W, *Deep-Sea Res. Pt. II*, 47, 3521–3548, 2000.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Iversen, M. H. and Ploug, H.: Ballast minerals and the sinking carbon flux in the ocean: carbon-specific respiration rates and sinking velocity of marine snow aggregates, *Biogeosciences*, 7, 2613–2624, doi:10.5194/bg-7-2613-2010, 2010.

Jacquet, S. H. M., Lam, P. J., Trull, T., and Dehairs, F.: Carbon export production in the subantarctic zone and polar front zone south of Tasmania, *Deep-Sea Res. Pt. II*, 58, 2277–2292, 2011.

Klaas, C. and Archer, D. E.: Association of sinking organic matter with various types of mineral ballast in the deep sea: implications for the rain ratio, *Global Biogeochem. Cy.*, 16, 1116–1129, 2002.

Kwon, E. Y., Primeau, F., and Sarmiento, J. L.: The impact of remineralization depth on the air–sea carbon balance, *Nat. Geosci.*, 2, 630–635, 2009.

Lam, P. J. and Bishop, J. K. B.: High biomass, low export regimes in the Southern Ocean, *Deep-Sea Res. Pt. II*, 54, 601–638, 2007.

Lam, P. J., Doney, S. C., and Bishop, J. K. B.: The dynamic ocean biological pump: insights from a global compilation of particulate organic carbon, CaCO₃, and opal concentration profiles from the mesopelagic, *Global Biogeochem. Cy.*, 25, GB3009, doi:10.1029/2010GB003868, 2011.

Lam, P. J., Ohnemus, D. C., and Auro, M. E.: Size-fractionated major particle composition and concentrations from the US GEOTRACES north Atlantic zonal transect, *Deep-Sea Res. Pt. II*, doi:10.1016/j.dsr2.2014.11.020, online first, 2014.

Le Moigne, F. A. C., Sanders, R. J., Villa-Alfageme, M., Martin, A. P., Pabortsava, K., Planquette, H., Morris, P. J., and Thomalla, S. J.: On the proportion of ballast versus non-ballast associated carbon export in the surface ocean, *Geophys. Res. Lett.*, 39, L15610, doi:10.1029/2012GL052980, 2012.

Le Moigne, F. A. C., Henson, S. A., Sanders, R. J., and Madsen, E.: Global database of surface ocean particulate organic carbon export fluxes diagnosed from the ²³⁴Th technique, *Earth Syst. Sci. Data*, 5, 295–304, doi:10.5194/essd-5-295-2013, 2013.

Le Moigne, F. A. C., Pabortsava, K., Marcinko, C. L. J., Martin, P., and Sanders, R. J.: Where is mineral ballast important for surface export of particulate organic carbon in the ocean?, *Geophys. Res. Lett.*, 41, doi:10.1002/2014GL061678, online first, 2014.

Lima, I. D., Lam, P. J., and Doney, S. C.: Dynamics of particulate organic carbon flux in a global ocean model, *Biogeosciences*, 11, 1177–1198, doi:10.5194/bg-11-1177-2014, 2014.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lomas, M. W., Steinberg, D. K., Dickey, T., Carlson, C. A., Nelson, N. B., Condon, R. H., and Bates, N. R.: Increased ocean carbon export in the Sargasso Sea linked to climate variability is countered by its enhanced mesopelagic attenuation, *Biogeosciences*, 7, 57–70, doi:10.5194/bg-7-57-2010, 2010.
- 5 Maiti, K., Benitez-Nelson, C. R., and Buesseler, K. O.: Insights into particle formation and remineralization using the short-lived radionuclide, Thorium-234, *Geophys. Res. Lett.*, 37, L15608, doi:10.1029/2010GL044063, 2010.
- Martin, J. H., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: Vertex – carbon cycling in the northeast Pacific, *Deep-Sea Res.*, 34, 267–285, 1987.
- 10 Martin, P., Lampitt, R. S., Jane Perry, M., Sanders, R., Lee, C., and D’Asaro, E.: Export and mesopelagic particle flux during a north Atlantic spring diatom bloom, *Deep-Sea Res. Pt. I*, 58, 338–349, 2011.
- McCave, I. N.: Vertical flux of particles in the ocean, *Deep Sea Research*, 22, 491–502, 1975.
- McDonnell, A. M. P. and Buesseler, K. O.: Variability in the average sinking velocities of marine particles, *Limnol. Oceanogr.*, 55, 2085–2096, 2010.
- 15 Michaels, A. F. and Silver, M. W.: Primary production, sinking fluxes and the microbial food web, *Deep-Sea Res.*, 35, 473–490, 1988.
- Morris, P. J., Sanders, R., Turnewitsch, R., and Thomalla, S.: Th-234-derived particulate organic carbon export from an island-induced phytoplankton bloom in the Southern Ocean, *Deep-Sea Res. Pt. II*, 54, 2208–2232, 2007.
- 20 Owens, S., Buesseler, K., and Sims, K.: Re-evaluating the ^{238}U -salinity relationship in seawater: implications for the ^{238}U – ^{234}Th disequilibrium method, *Mar. Chem.*, 127, 31–39, 2011.
- Paasche, E. and Brubak, S.: Enhanced calcification in the coccolithophorid *Emiliana huxleyi* (Haptophyceae) under phosphorus limitation, *Phycologia*, 33, 324–330, 1994.
- 25 Passow, U. and De la Rocha, C. L.: Accumulation of mineral ballast on organic aggregates, *Global Biogeochem. Cy.*, 20, GB1013, doi:10.1029/2005GB002579, 2006.
- Pike, S. M., Buesseler, K. O., Andrews, J., and Savoye, N.: Quantification of Th-234 recovery in small volume sea water samples by inductively coupled plasma-mass spectrometry, *J. Radioanal. Nucl. Ch.*, 263, 355–360, 2005.
- 30 Pilson, M. E. Q.: *An Introduction to the Chemistry of the Sea*, Cambridge University Press, New York, 2012.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Sanders, R., Morris, P. J., Poulton, A. J., Stinchcombe, M. C., Charalampopoulou, A., Lucas, M. I., and Thomalla, S. J.: Does a ballast effect occur in the surface ocean?, *Geophys. Res. Lett.*, 37, L08602, doi:10.1029/2010GL042574, 2010.
- Santschi, P., Murray, J. W., Baskaran, M., Benitez-Nelson, C. R., Guo, L., Hung, C.-C., Lamborg, C., Moran, S. B., Passow, U., and Roy-Barman, M.: Thorium speciation in seawater, *Mar. Chem.*, 100, 250–268, 2006.
- Sarmiento, J. L., Dunne, J., Gnanadesikan, A., Key, R. M., Matsumoto, K., and Slater, R.: A new estimate of the CaCO₃ to organic carbon export ratio, *Global Biogeochem. Cy.*, 16, 1107, doi:10.1029/2002GB001919, 2002.
- Sarmiento, J. L., Gruber, N., Brzezinski, M. A., and Dunne, J. P.: High-latitude controls of thermocline nutrients and low latitude biological productivity, *Nature*, 427, 56–60, 2004.
- Savoie, N., Benitez-Nelson, C., Burd, A. B., Cochran, J. K., Charette, M., Buesseler, K. O., Jackson, G. A., Roy-Barman, M., Schmidt, S., and Elskens, M.: Th-234 sorption and export models in the water column: a review, *Mar. Chem.*, 100, 234–249, 2006.
- Savoie, N., Trull, T. W., Jacquet, S. H. M., Navez, J., and Dehairs, F.: ²³⁴Th-based export fluxes during a natural iron fertilization experiment in the Southern Ocean (KEOPS), *Deep-Sea Res. Pt. II*, 55, 841–855, 2008.
- Shimmield, G. B., Ritchie, G. D., and Fileman, T. W.: The impact of marginal ice-zone processes on the distribution of Pb-210, Po-210 and Th-234 and implications for new production in the Bellingshausen Sea, Antarctica, *Deep-Sea Res. Pt. II*, 42, 1313–1335, 1995.
- Smetacek, V., Klaas, C., Strass, V. H., Assmy, P., Montresor, M., Cisewski, B., Savoie, N., Webb, A., d’Odivio, F., Arrieta, J. M., Bathmann, U., Bellerby, R., Berg, G. M., Croot, P., Gonzalez, S., Henjes, J., Herndl, G. J., Hoffmann, L. J., Leach, H., Losch, M., Mills, M. M., Neill, C., Peeken, I., Röttgers, R., Sachs, O., Sauter, E., Schmidt, M. M., Schwarz, J., Terbrüggen, A., and Wolf-Gladrow, D.: Deep carbon export from a Southern Ocean iron-fertilized diatom bloom, *Nature*, 487, 313–319, 2012.
- Sokolov, S. and Rintoul, S. R.: Circumpolar structure and distribution of the Antarctic Circumpolar Current fronts: 1. Mean circumpolar paths, *J. Geophys. Res.*, 114, C11018, doi:10.1029/2008JC005108, 2009.
- Steinberg, D. K., Van Mooy, B. A. S., Buesseler, K. O., Boyd, P. W., Kobari, T., and Karl, D. M.: Bacterial vs. zooplankton control of sinking particle flux in the ocean’s twilight zone, *Limnol. Oceanogr.*, 53, 1327–1338, 2008.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Strickland, J. D. and Parsons, T. R.: A Practical Handbook of Seawater Analysis, Fisheries Research Board of Canada, Ottawa, 1968.
- Thomalla, S., Turnewitsch, R., Lucas, M., and Poulton, A.: Particulate organic carbon export from the north and south Atlantic gyres: the $^{234}\text{Th}/^{238}\text{U}$ disequilibrium approach, *Deep-Sea Res. Pt. II*, 53, 1629–1648, 2006.
- Thomalla, S. J., Poulton, A. J., Sanders, R., Turnewitsch, R., Holligan, P. M., and Lucas, M. I.: Variable export fluxes and efficiencies for calcite, opal, and organic carbon in the Atlantic Ocean: a ballast effect in action?, *Global Biogeochem. Cy.*, 22, GB1010, doi:10.1029/2007GB002982, 2008.
- Trull, T. W., Bray, S. G., Buesseler, K. O., Lamborg, C. H., Manganini, S., Moy, C., and Valdes, J.: In situ measurement of mesopelagic particle sinking rates and the control of carbon transfer to the ocean interior during the Vertical Flux in the Global Ocean (VERTIGO) voyages in the North Pacific, *Deep-Sea Res. Pt. II*, 55, 1684–1695, 2008.
- Volk, T. and Hoffert, M. I.: Ocean carbon pumps: analysis of relative strengths and efficiencies in ocean-driven atmospheric CO_2 changes, *Geophysical Monographs*, 32, 99–110, 1985.
- Wakeham, S. G., Peterson, M. L., Hedges, J. I., and Lee, C.: Lipid biomarker fluxes in the Arabian Sea, with a comparison to the equatorial Pacific Ocean, *Deep-Sea Res. Pt. II*, 49, 2265–2301, 2002.
- Wilson, J. D., Barker, S., and Ridgwell, A.: Assessment of the spatial variability in particulate organic matter and mineral sinking fluxes in the ocean interior: implications for the ballast hypothesis, *Global Biogeochem. Cy.*, 26, GB4011, doi:10.1029/2012GB004398, 2012.
- Wilson, S. E., Steinberg, D. K., and Buesseler, K. O.: Changes in fecal pellet characteristics with depth as indicators of zooplankton repackaging of particles in the mesopelagic zone of the subtropical and subarctic North Pacific Ocean, *Deep-Sea Res. Pt. II*, 55, 1636–1647, 2008.
- Zhou, K., Nodder, S. D., Dai, M., and Hall, J. A.: Insignificant enhancement of export flux in the highly productive subtropical front, east of New Zealand: a high resolution study of particle export fluxes based on ^{234}Th : ^{238}U disequilibria, *Biogeosciences*, 9, 973–992, doi:10.5194/bg-9-973-2012, 2012.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. POC fluxes, concentrations, and attenuation of $> 51 \mu\text{m}$ [POC] in the mesopelagic zone. Attenuation coefficient is the exponent from significant power law fits to $> 51 \mu\text{m}$ [POC]. $Z_{\text{PAR}} + 100 \text{ m}$ is 100 m below Z_{PAR} , as defined in the Table 1 caption. Transfer efficiency is POC flux at $Z_{\text{PAR}} + 100 \text{ m}$ divided by POC flux at Z_{PAR} . Deep $> 51 \mu\text{m}$ [POC] was measured at 1000 and 800 m for GB1 and GB2, respectively. POC flux errors are propagated from ^{234}Th flux, and POC : ^{234}Th errors.

Cruise	Station	Depth	$> 51 \mu\text{m}$ [POC] Attenuation Coefficient	^{234}Th Flux at $Z_{\text{PAR}} + 100 \text{ m}$	POC : Th at $Z_{\text{PAR}} + 100 \text{ m}$	POC Flux at $Z_{\text{PAR}} + 100 \text{ m}$	Transfer Efficiency	$> 51 \mu\text{m}$ [POC] ($\geq 800 \text{ m}$)
–	–	m	unitless	$\text{dpm m}^{-2} \text{d}^{-1}$	$\mu\text{mol dpm}^{-1}$	$\text{mmol m}^{-2} \text{d}^{-1}$	unitless	μM
GB1	6	179	0.8	$3319 \pm 128^{\text{c}}$	1.7	5.7 ± 0.31	1.00	0.030
GB1	16	162	1.1	$2567 \pm 116^{\text{c}}$	2.4	6.1 ± 0.30	1.04	No data
GB1	25	162	0.4	$1,074 \pm 125$	2.5	2.7 ± 0.37	1.76	0.013
GB1	32	169	0.9	1581 ± 186	1.3	2.0 ± 0.25	0.86	0.006
GB1	38	221	No fit	911 ± 206	1.6	1.5 ± 0.35	0.70	0.026
GB1	46	163	1.0	1937 ± 146	1.6	3.1 ± 0.27	0.4	0.009
GB1	59	160	0.6	$2582 \pm 126^{\text{c}}$	3.7	9.5 ± 0.56	1.29	0.014
GB1	70	200	0.6	1414 ± 248	3.5	5.0 ± 0.90	0.90	0.024
GB1	77	198	0.5	1903 ± 162	2.1	4.0 ± 0.41	0.44	0.012
GB1	85	173	1.7^{a}	$2,076 \pm 207$	3.9	8.1 ± 0.83	0.41	0.035
GB1	92	159	1.1	1339 ± 170	3.7	4.9 ± 0.64	0.61	0.019
GB1	101	181	0.8	1774 ± 135	1.7	3.0 ± 0.24	0.83	0.019
GB1	109	176	1.0	1719 ± 97	1.1	1.9 ± 0.13	0.87	0.006
GB1	117	162	1.1	1258 ± 86	1.2	1.5 ± 0.13	0.87	0.005
GB2	5	178	0.5	$1402 \pm 3706^{\text{c}}$	1.1	1.5 ± 6.1	0.5	No data
GB2	27	205	No fit	$2,063 \pm 205$	1.2	2.5 ± 0.29	0.71	No data
GB2	36	190	1.5	$1,077 \pm 194$	0.9	0.93 ± 0.18	0.48	0.011
GB2	43	208	1.9	$1,247 \pm 200$	2.2	2.7 ± 0.45	0.54	0.005
GB2	53	181	No fit	$1,013 \pm 220$	2.0	2.0 ± 0.45	0.49	No data
GB2	63	209	1.8	$1,292 \pm 262$	1.7	2.1 ± 0.45	0.31	0.014
GB2	73	193	1.5	807 ± 189	1.9	1.6 ± 0.37	0.48	0.008
GB2	87	207	0.7	$1,213 \pm 196$	1.6	1.9 ± 0.34	0.60	0.013
GB2	93	213	2.3^{b}	469 ± 249	1.6	0.77 ± 0.42	0.53	0.001
GB2	100	213	0.8	$1,132 \pm 190$	0.7	0.80 ± 0.15	0.52	0.014
GB2	106	202	0.9	$1,405 \pm 186$	1.3	1.8 ± 0.25	1.63	0.017
GB2	112	176	1.3	270 ± 186	0.9	0.23 ± 0.21	0.24	0.007
GB2	119	192	No fit	756 ± 218	0.8	0.57 ± 0.17	0.20	0.013

^a Attenuation coefficient is 2.35 when only fitting $> 51 \mu\text{m}$ [POC] measurements at depths $< 500 \text{ m}$ (Fig. 3).

^b Outlier approximated by Chauvenet's Theorem (Glover et al., 2011).

^c Values were estimated by linear interpolation of values at upper and lower depths around $Z_{\text{PAR}} + 100 \text{ m}$.

"no data": no measurements at these depths.

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. POC, biomineral and ^{234}Th concentrations and fluxes at z_{PAR} . Ez-ratio is ^{234}Th -derived POC flux at z_{PAR} divided by integrated primary productivity. The % > 51 μm [POC] metric is the fraction of total [POC] in the > 51 μm size fraction. POC and biomineral flux errors are propagated from ^{234}Th flux, and POC : ^{234}Th errors.

Cruise	Station	z_{PAR} m	^{234}Th Flux dpm m $^{-2}$ d $^{-1}$	> 51 μm [POC]	> 51 μm [Bsi]	> 51 μm [PIC]	> 51 μm Th activity
				μM			dpm L $^{-1}$
GB1	6	79	2437 \pm 100	0.23 ^b	0.03 ^a	0.124 ^a	0.07 ^a
GB1	16	62	1933 \pm 71	0.38	0.08	0.390	0.12
GB1	25	62	862 \pm 46 ^a	0.04	0.005 ^a	0.015 ^a	0.02
GB1	32	69	1304 \pm 116	0.07	0.01	0.027	0.04
GB1	38	121	809 \pm 126	0.04	0.003	0.017	0.01
GB1	46	63	2123 \pm 69	0.23	0.005	0.059	0.06
GB1	59	60	1844 \pm 102	0.09	0.10	0.072	0.02
GB1	70	100	1280 \pm 94	0.11	0.06 ^a	0.001 ^a	0.02
GB1	77	98	1485 \pm 105	0.03	0.03	0.002	0.01
GB1	85	73	1858 \pm 94	2.50	3.44	0.124	0.23
GB1	92	59	1639 \pm 77	0.40	0.46	0.020	0.08
GB1	101	81	1763 \pm 82	0.19	0.05	0.013	0.09
GB1	109	76	1524 \pm 76	0.19 ^a	0.05 ^a	0.027 ^a	0.14 ^a
GB1	117	62	1177 \pm 50	0.21	0.02	0.032	0.15
GB2	5	78	1889 \pm 5207	0.08 ^b	0.01 ^a	0.048 ^a	0.05 ^a
GB2	27	105	1869 \pm 160	0.08 ^a	0.10 ^a	0.060 ^a	0.04 ^a
GB2	36	90	988 \pm 89	0.43	0.28	0.074	0.22
GB2	43	108	1221 \pm 153	0.74 ^a	0.62 ^a	0.041 ^a	0.18 ^a
GB2	53	81	1058 \pm 100 ^a	0.54 ^a	0.80 ^a	0.081 ^a	0.14 ^a
GB2	63	109	1229 \pm 138	0.71 ^a	1.04 ^a	0.028 ^a	0.13 ^a
GB2	73	93	977 \pm 108	0.21 ^b	1.13 ^a	0.014 ^a	0.20 ^a
GB2	87	107	1299 \pm 115	0.06 ^b	0.30 ^a	0.041 ^a	0.06 ^a
GB2	93	113	1142 \pm 137	0.07 ^a	0.01 ^a	0.023 ^a	0.05 ^a
GB2	100	113	1112 \pm 130	0.08	0.02	0.006	0.06
GB2	106	102	1394 \pm 82 ^a	0.09 ^b	0.04 ^a	0.024 ^a	0.12 ^a
GB2	112	76	717 \pm 97	0.22 ^b	0.17 ^a	0.087 ^a	0.36 ^a
GB2	119	92	1223 \pm 124	0.51 ^a	0.12 ^a	0.048 ^a	0.22 ^a

^a Values at z_{PAR} estimated by linear interpolation of values at upper and lower depths around z_{PAR} .

^b > 51 μm [POC] values interpolated by significant power-law fits (Fig. 3).

"no data": not enough depths were sampled and analyzed to interpolate at z_{PAR} .

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Table 3. Continued.

Cruise	Station	POC:Th μmol dpm ⁻¹	POC Flux mmol m ⁻² d ⁻¹	Bsi:Th μmol dpm ⁻¹	BSi Flux mmol m ⁻² d ⁻¹	PIC:Th μmol dpm ⁻¹	PIC Flux mmol m ⁻² d ⁻¹	Primary Productivity mmol m ⁻² d ⁻¹	Ez-ratio unitless	% > 51 μm [POC] %
GB1	6	2.3 ^a	5.7 ± 0.26	0.4	0.9 ± 0.04	1.8	4.3 ± 0.20	42	0.14	8.8
GB1	16	3.0	5.9 ± 0.68	0.6	1.2 ± 0.14	3.1	6.1 ± 0.70	165	0.04	17.7
GB1	25	1.8	1.6 ± 0.11	0.2	0.2 ± 0.02	0.7	0.6 ± 0.04	35	0.04	3.2
GB1	32	1.8	2.3 ± 0.21	0.3	0.3 ± 0.03	0.7	0.9 ± 0.08	11	0.21	3.9
GB1	38	2.7	2.2 ± 0.35	0.2	0.2 ± 0.03	1.2	0.9 ± 0.15	21	0.10	8.4
GB1	46	4.1	8.8 ± 0.38	0.1	0.2 ± 0.02	1.1	2.2 ± 0.10	13	0.67	5.3
GB1	59	4.0	7.3 ± 0.52	4.6	8.6 ± 0.53	3.4	6.2 ± 0.39	26	0.28	5.3
GB1	70	4.3	5.5 ± 0.44	3.5 ^a	4.5 ± 0.35	0.1 ^a	0.1 ± 0.09	10	0.53	10.6
GB1	77	6.0	9.0 ± 1.3	5.6	8.3 ± 0.98	0.4	0.7 ± 0.23	57	0.16	3.6
GB1	85	10.8	20 ± 1.1	14.9	28 ± 1.5	0.5	1.0 ± 0.05	53	0.38	52.0
GB1	92	4.9	8.0 ± 0.40	5.6	9.3 ± 0.46	0.2	0.4 ± 0.02	26	0.31	11.3
GB1	101	2.0	3.6 ± 0.18	0.5	0.9 ± 0.04	0.1	0.2 ± 0.01	22	0.17	12.5
GB1	109	1.4	2.1 ± 0.11	0.4	0.6 ± 0.03	0.2	0.3 ± 0.02	14	0.16	21.0
GB1	117	1.4	1.7 ± 0.07	0.1	0.2 ± 0.01	0.2	0.3 ± 0.01	18	0.09	6.6
GB2	5	1.6	3.0 ± 8.8	0.2	0.4 ± 1.2	1.0	1.9 ± 5.2	8.2	0.37	7.6
GB2	27	1.9	3.5 ± 0.31	2.2	4.0 ± 0.34	1.3	2.5 ± 0.22	8.0	0.44	6.7
GB2	36	2.0	2.0 ± 0.18	1.3	1.3 ± 0.12	0.3	0.3 ± 0.03	12	0.16	15.6
GB2	43	4.1	5.0 ± 0.63	3.4	4.2 ± 0.52	0.2	0.3 ± 0.04	12	0.43	37.8
GB2	53	3.9	4.1 ± 0.39	5.7	6.1 ± 0.58	0.6	0.6 ± 0.06	16	0.25	22.5
GB2	63	5.6 ^a	6.9 ± 0.78	8.1	9.9 ± 1.1	0.2	0.3 ± 0.03	9.0	0.77	33.2
GB2	73	3.3 ^a	3.2 ± 0.35	5.6	5.4 ± 0.60	0.1	0.1 ± 0.01	8.8	0.36	17.6
GB2	87	2.5 ^a	3.2 ± 0.39	4.7	6.1 ± 0.54	0.6	0.8 ± 0.14	11	0.29	3.4
GB2	93	1.3	1.5 ± 0.25	0.2	0.3 ± 0.06	0.4	0.5 ± 0.14	12	0.12	4.3
GB2	100	1.4	1.5 ± 0.19	0.3	0.3 ± 0.04	0.1	0.1 ± 0.02	14	0.11	12.8
GB2	106	0.8	1.1 ± 0.86	0.3 ^a	0.4 ± 0.02	0.2	0.3 ± 0.02	22	0.05	12.2
GB2	112	1.4 ^a	1.0 ± 0.13	0.5	0.3 ± 0.05	0.2	0.2 ± 0.02	no data	no data	13.3
GB2	119	2.3	2.8 ± 0.29	0.5	0.7 ± 0.07	0.2	0.3 ± 0.03	17	0.17	21.5

^a Values at z_{PAR} estimated by linear interpolation of values at upper and lower depths around z_{PAR} .

^b > 51 μm [POC] values interpolated by significant power-law fits (Fig. 3).

"no data": not enough depths were sampled and analyzed to interpolate at z_{PAR} .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Table 4. Mean \pm SDs of ^{234}Th fluxes, POC : ^{234}Th , BSi : ^{234}Th , PIC : ^{234}Th , POC fluxes, and biomineral fluxes at z_{PAR} , divided by three latitude zones. 45°S marks the approximate latitude of the Subantarctic front, while 52°S marks the approximate latitude of the Polar front (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009).

Lat. zone $^\circ\text{S}$	^{234}Th Flux at z_{PAR} $\text{dpm m}^{-2} \text{d}^{-1}$	POC : Th at z_{PAR} $\mu\text{mol dpm}^{-1}$	POC Flux at z_{PAR} $\text{mmol m}^{-2} \text{d}^{-1}$	BSi : Th at z_{PAR} $\mu\text{mol dpm}^{-1}$	BSi Flux at z_{PAR} $\text{mmol m}^{-2} \text{d}^{-1}$	PIC : Th at z_{PAR} $\mu\text{mol dpm}^{-1}$	PIC Flux at z_{PAR} $\text{mmol m}^{-2} \text{d}^{-1}$	# stn –
36–45	1.3 ± 0.44	1.9 ± 0.9	2.7 ± 2.3	0.3 ± 0.1	0.33 ± 0.17	0.5 ± 0.4	0.73 ± 0.76	10
45–52	1.5 ± 0.50	2.8 ± 1.2	4.4 ± 2.2	2.3 ± 2.2	3.4 ± 3.3	1.1 ± 1.2	2.0 ± 2.4	11
52–> 60	1.4 ± 0.30	5.4 ± 3.0	8.0 ± 6.3	7.1 ± 4.1	10 ± 8.7	0.3 ± 0.2	0.49 ± 0.4	6

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.




Figure 1. Cruise tracks across the Atlantic (cruise GB1) and Indian (cruise GB2) sectors of the Great Calcite Belt showing sea surface temperature along the two transects. Station numbers where only ^{234}Th and size-fractionated particles were sampled are indicated by crosses. The two horizontal dashed lines at 45 and 52° S represent the approximate locations of the Subantarctic and Polar fronts, respectively (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Station GB1-85




Figure 2. Digital images of $> 51 \mu\text{m}$ filters from station GB1-85 (refer to Fig. 1 for station location). $> 51 \mu\text{m}$ particles are from **(a)** 25 m in the euphotic zone, **(b)** 73 m, which corresponds to Z_{PAR} , as defined in Table 1, and **(c)** at 173 m, below both metrics of export depth, Z_{PAR} and $Z_{\text{Th/U}}$ (Table 1). $> 51 \mu\text{m}$ particles appear as intact cells in the euphotic zone **(a, b)** and as fecal pellets below Z_{PAR} **(c)**.

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







Figure 3. Significant power law fits of $> 51 \mu\text{m}$ [POC] below z_{PAR} , according to Eq. (1). Only the 22 significant fits are shown as lines. Three stations are highlighted to show the range in $> 51 \mu\text{m}$ [POC] attenuation across GB1 and GB2 profiles (symbols represent measurements): GB1-85 had the highest POC concentration through the water column and an attenuation coefficient of 1.7; GB1-25 had the lowest attenuation coefficient (0.4); GB2-43 had the highest attenuation coefficient (1.9) (Table 2). Fitting GB1-85 $> 51 \mu\text{m}$ [POC] measurements between z_{PAR} and 500 m yields a higher attenuation coefficient of 2.35. Refer to Fig. 1 for station locations.

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





Carbon export and transfer to depth across the Great Calcite Belt


S. Z. Rosengard et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.



Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 4. Total ^{234}Th and ^{238}U activity profiles measured at 14 stations of GB1 and 13 stations of GB2 (note different x axis for station GB2-5) (Table S1). Error bars for ^{234}Th activity are propagated errors. ^{238}U is calculated from salinity. All ^{234}Th activity profiles exhibit a deficit relative to ^{238}U activity at the surface, and mostly return to equilibrium with ^{234}U within error at depth of $z_{\text{Th/U}}$ (Table 1). Refer to Fig. 1 for station locations.

BGD

12, 2843–2896, 2015

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.




Figure 5. Distribution of ^{234}Th flux and ^{234}Th -derived POC flux at 27 stations along GB1 and GB2. **(a)** ^{234}Th fluxes at z_{PAR} range from 717 to 2437 $\text{dpm m}^{-2} \text{d}^{-1}$ at stations GB2-112 and GB1-6, respectively. **(b)** POC fluxes at z_{PAR} range from 0.97 to 20 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB2-112 and GB1-85, respectively. **(c)** POC fluxes at $z_{\text{Th/U}}$ range from 0.57 to 12 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB2-112 and GB1-85, respectively (Table S2). **(d)** POC fluxes at 100 m below z_{PAR} range from 0.23 to 9.5 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB2-112 and GB1-59, respectively. A few station numbers discussed in the text are indicated. Red outlines distinguish stations where fluxes are greater at the specified depth than at z_{PAR} . The two horizontal dashed lines at 45 and 52° S represent the approximate locations of the Subantarctic and Polar fronts, respectively (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009). Refer to Fig. 1 for other station locations. z_{PAR} and $z_{\text{Th/U}}$ are defined as in Table 1.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.




Figure 6. Distribution of BSi flux, PIC flux, and % > 51 μm [POC], the percent of total [POC] in the > 51 μm size class, at z_{PAR} (Table 1) along GB1 and GB2. **(a)** BSi fluxes range from 0.17 to 28 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB1-46 and GB1-85, respectively. **(b)** PIC fluxes range from 0.067 to 6.2 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB2-73 and GB2-59, respectively. **(c)** The proportion of [POC] in the > 51 μm size-fraction at z_{PAR} ranges from 3.3 to 52 % at stations GB1-25 and GB1-85, respectively. A few station numbers discussed in the text are indicated. The two horizontal dashed lines at 45 and 52° S represent the approximate locations of the Subantarctic and Polar fronts, respectively (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009). Refer to Fig. 1 for other station locations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.




Figure 7. Distribution of primary productivity, export efficiency, and transfer efficiency along GB1 and GB2. **(a)** Primary productivity integrated through the euphotic zone ranges from 8.0 to 165 $\text{mmol m}^{-2} \text{d}^{-1}$ at stations GB2-27 and GB1-16, respectively. **(b)** Export efficiency (Ez-ratio) at z_{PAR} (Table 1), which is the ratio of ^{234}Th -derived POC flux at z_{PAR} to primary productivity integrated to z_{PAR} , ranges from 0.04 to 0.77 at stations GB1-16 and GB2-63, respectively. **(c)** Transfer efficiency at z_{PAR} , which is the ratio of POC flux 100 m below z_{PAR} to POC flux at z_{PAR} , ranges from 0.20 to 1.8 at stations GB1-119 and GB1-25, respectively. A few station numbers discussed in the text are indicated. The two horizontal dashed lines at 45 and 52° S represent the approximate locations of the Subantarctic and Polar fronts, respectively (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009). Refer to Fig. 1 for other station locations.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Carbon export and transfer to depth across the Great Calcite Belt

S. Z. Rosengard et al.




Figure 8. Distribution and vertical attenuation coefficient of $> 51 \mu\text{m}$ [POC]. **(a)** $> 51 \mu\text{m}$ POC concentrations at z_{PAR} (Table 1) range from 0.03 to $2.5 \mu\text{M}$ at stations GB1-77 and GB1-85, respectively. **(b)** $> 51 \mu\text{m}$ [POC] at the deepest pump depth in the lower mesopelagic zone (800 – 1000 m). Concentrations range from 0.001 to $0.035 \mu\text{M}$ at stations GB2-93 and GB1-85, respectively. **(c)** Attenuation coefficient from significant power-law fits of 22 $> 51 \mu\text{m}$ [POC] profiles, excluding GB2-93 (see Sect. 4.4). A few station numbers discussed in the text are indicated. The two horizontal dashed lines at 45 and 52° S represent the approximate locations of the Subantarctic and Polar fronts, respectively (Belkin and Gordon, 1996; Sokolov and Rintoul, 2009). Refer to Fig. 1 for other station locations.




Figure 9. ^{234}Th -derived POC flux as a function of (a) PIC flux and (b) BSi flux at z_{PAR} . POC flux transfer efficiency between z_{PAR} and $z_{\text{PAR}} + 100\text{ m}$ (T_{100} , defined in Sect. 4.4) as a function of (c) PIC flux and (d) BSi flux at z_{PAR} . Significant linear relationships are plotted as a solid blue line. T_{100} values at GB1-25 and GB2-106 were excluded from all correlations (Sect. 4.4). Colorbar indicates longitude of stations – GB1 and GB2 stations are in cool and warm colors, respectively. Refer to Fig. 1 for more specific station locations.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	






Figure 10. Attenuation coefficient as a function of **(a)** $> 51 \mu\text{m}$ [BSi] at z_{PAR} and **(b)** the proportion of [POC] in the $> 51 \mu\text{m}$ size-fraction at z_{PAR} . The open circle indicates where GB1-85 would plot with a higher attenuation coefficient of 2.35, derived from fitting $> 51 \mu\text{m}$ [POC] at depths between z_{PAR} and 500 m. Significant linear relationships using the lower and higher attenuation coefficient values for GB1-85 are shown as solid and dashed lines, respectively; p and r^2 values are provided for the solid lines. The colorbar is the natural logarithm of the ratio of $> 51 \mu\text{m}$ PIC : BSi at z_{PAR} . We interpret all warm colors > 0 to indicate coccolithophore-dominated stations, and all cool values < 0 to indicate diatom-dominated stations. A few station numbers discussed in the text are indicated. Refer to Fig. 1 for station locations.