

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

# Particle fluxes in the deep Eastern Mediterranean basins: the role of ocean vertical velocities

L. Patara<sup>1</sup>, N. Pinardi<sup>2</sup>, C. Corselli<sup>3</sup>, E. Malinverno<sup>3</sup>, M. Tonani<sup>4</sup>, R. Santoleri<sup>5</sup>,  
and S. Masina<sup>1,4</sup>

<sup>1</sup>Centro Euro-Mediterraneo per i Cambiamenti Climatici (CMCC), Viale Aldo Moro 44, 40 127 Bologna, Italy

<sup>2</sup>Laboratorio SINCEM, Università di Bologna, Via S. Alberto 163, 48 100 Ravenna, Italy

<sup>3</sup>Dipartimento di Scienze Geologiche e Geotecnologie, Università Milano-Bicocca, Piazza della Scienza 4, 20 126 Milano, Italy

<sup>4</sup>Istituto Nazionale di Geofisica e Vulcanologia (INGV), Viale Aldo Moro 44, 40 127 Bologna, Italy

<sup>5</sup>Istituto di Scienze dell'Atmosfera e del Clima–C. N. R., Via del Fosso del Cavaliere 100, 00 133 Rome, Italy

Received: 25 April 2008 – Accepted: 28 May 2008 – Published: 12 August 2008

Correspondence to: L. Patara (patara@bo.ingv.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

This paper analyzes the relationship between deep sedimentary fluxes and ocean current vertical velocities in an offshore area of the Ionian Sea, the deepest basin of the Eastern Mediterranean Sea. Sediment trap data are collected at 500 m and 2800 m depth in two successive moorings covering the period September 1999–May 2001. A tight coupling is observed between the upper and deep traps and the estimated particle sinking rates are higher than  $200 \text{ m day}^{-1}$ . The current vertical velocity field is computed from a high resolution Ocean General Circulation Model simulation and from the wind stress curl. Current vertical velocities are larger and more variable than Ekman vertical velocities, yet the general patterns are alike. Current vertical velocities are generally smaller than  $1 \text{ m day}^{-1}$ : we therefore exclude a direct effect of downward velocities in determining high sedimentation rates. However, we find that upward velocities in the subsurface layers of the water column are positively correlated with deep particle fluxes. We thus hypothesize that upwelling would produce an increase in upper ocean nutrient levels – thus stimulating primary production and grazing – a few weeks before an enhanced vertical flux is found in the sediment traps. By analyzing the delayed effects of ocean vertical velocities on deep particle fluxes we envisage a spectrum of particle sinking speeds ranging from about  $100 \text{ m day}^{-1}$  to more than  $200 \text{ m day}^{-1}$ . High particle sedimentation rates may be attained by means of rapidly sinking fecal pellets produced by gelatinous macro-zooplankton. Other sedimentation mechanisms, such as dust deposition, are also considered in explaining large pulses of deep particle fluxes. The fast sinking rates estimated in this study might be an evidence of the efficiency of the biological pump in sequestering organic carbon from the surface layers of the deep Eastern Mediterranean basins.

**BGD**

5, 3123–3156, 2008

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



# 1 Introduction

The Eastern Mediterranean Sea is characterized by an anti-estuarine basin scale circulation through the Sicily Channel, with the entrance of surface nutrient-depleted waters and the exiting of nutrient-enriched intermediate waters (Malanotte-Rizzoli et al., 1997; Pinardi and Masetti, 2000). Superimposed to this large scale circulation pattern, a seasonal wind-forced sub-basin scale circulation is evident (Pinardi and Navarra, 1993; Molcard et al., 2002), with the dominant north-westerly wind component giving a characteristic separation between the southern regions, dominated by anticyclonic motion, and the northern regions, dominated by cyclonic motion (Pinardi et al., 2005). The Eastern Mediterranean general circulation is moreover importantly affected by interannual scale variability mainly in response to surface fluxes (Korres et al., 2000; Demirov and Pinardi, 2002).

The open areas of the Ionian Sea, which is the deepest basin of the Eastern Mediterranean Sea, are considered to be highly oligotrophic: surface chlorophyll concentrations are generally lower than  $0.5 \text{ mg m}^{-3}$  and a deep chlorophyll maximum is found at around 80–100 m depth (Boldrin et al., 2002; Malinverno et al., 2003; Ignatiades, 2005). The nutricline in the Eastern Mediterranean Sea is normally well below the euphotic layer. Klein et al. (1999) measure in the year 1987 a 300–400 m deep nutricline in the southeastern Ionian Sea, while Ediger and Yilmaz (1996) measure in the Levantine Basin a 600 m deep nutricline in correspondence of anticyclonic structures. In the Ionian Sea, oligotrophic features increase according to a north-south and west-east gradient (Bricaud et al., 2002; Casotti et al., 2003), in response to different water mass characteristics and possibly circulation patterns. In this oligotrophic region coccolithophores are an important constituent of the phytoplankton assemblage larger than  $3 \mu\text{m}$  (Boldrin et al., 2002; Malinverno et al., 2003) and their calcitic remains give a major contribution to the calcium carbonate export to the deep layers (Ziveri et al., 2000).

Sinking mechanisms and vertical particle fluxes in open ocean areas were studied in

**BGD**

5, 3123–3156, 2008

---

## Particle fluxes and ocean vertical velocities

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Particle fluxes and  
ocean vertical  
velocities**L. Patara et al.

---

the past by means of laboratory experiments, in situ observations and sediment traps (Honjo et al., 1984; Alldredge and Gotshalk, 1988; Asper et al., 1992; Stemmann et al., 2002; Lee et al., 2004). Aggregation inside zooplankton fecal pellets is considered to be one of the main mechanisms for organic matter sedimentation in the deep ocean (Honjo et al., 1976; Fowler et al., 1987; Knappertsbusch and Brummer, 1995). Other sinking agents include the formation of “marine snow” particles (Lampitt et al., 1993; Turner, 2002) possibly enhanced by the presence of transparent exopolymer particles (Engel, 2004). However the processes affecting particle flux magnitude and sedimentation rates in the ocean interior are not yet fully understood.

Vertical velocities of the general circulation might play a relevant role in affecting particle fluxes in two opposite ways: on one hand they could directly increase sedimentation rates through downwelling velocities; on the other hand they could produce, via upwelling velocities, an indirect effect through an enhancement of nutrient levels and primary productivity in the upper ocean.

Current vertical velocities in the upper ocean are directly forced by the wind field through the wind stress curl. Thus Ekman vertical velocities, generated by the wind stress curl, were used in previous studies as a proxy of the ocean currents vertical velocities and a relation with biological patterns was found (Andersen and Prieur, 2000; Agostini and Bakun, 2002). However, biogeochemical processes are directly influenced by smaller scale circulation features (Lévy et al., 2001) which might not be correctly represented by the large scale Ekman vertical velocities. These small scale features, called mesoscales, are pervasive in the Mediterranean Sea and in particular in the Ionian Sea (Robinson et al., 1987; Nittis et al., 1993).

Our goal is to understand the effect of ocean currents vertical velocities on particle fluxes observed in the ocean interior. Particles are sampled during the period 1999–2001 by means of sediment traps moored at 500 m and 2800 m depth in two offshore sites located in the southeastern Ionian Sea, i.e. the Urania and Bannock stations. We analyze total particle flux, including all sinking particles, and coccolithophore flux in terms of the calcitic plates – the coccoliths – constituting the coccolithophores external

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

shell.

Current vertical velocities are difficult to directly measure because of their extremely low values, which are commonly less than  $1 \text{ m day}^{-1}$ . We therefore analyze vertical velocities simulated by a high resolution Ocean General Circulation Model (OGCM) for the years 1999–2001. The OGCM vertical velocities are compared with the wind induced Ekman vertical velocities in order to verify the consistency between the two fields. The OGCM vertical velocity component is then correlated with the particle fluxes from the sediment traps.

The paper is organized as follows: Sect. 2 gives some details regarding the particle flux observations and the computation of the wind-induced and current vertical velocity fields. Section 3 investigates the main results of this work, concentrating on the role of ocean vertical velocities on the observed particle fluxes. In Sect. 4 we discuss the major findings and in Sect. 5 we give a few concluding remarks.

## 2 Methods

### 2.1 Particle flux data from sediment traps

Sediment trap particle flux data analyzed in this work were collected in the offshore stations Urania and Bannock located in the southeastern Ionian Sea (Fig. 1). The sediment trap time series at the Urania station covers the period 15 September 1999–9 May 2001 in two successive moorings, the first one from 15 September 1999 to 2 June 2000 (hereafter referred to as Mooring 1), the second one from 30 May 2000 to 9 May 2001 (hereafter referred to as Mooring 2). The sampling frequency varies between 10–11 days for Mooring 1 and 14–15 days for Mooring 2.

Soon after recovery, the samples were splitted on board in 8 aliquots using a pneumatic splitter. One aliquote, intended for the analysis of biogenic components, was subsequently splitted into additional sample fractions by means of a rotary splitter. Of these fractions one was filtered on a pre-weighted cellulose acetate filter, dried and

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



then weighted. The measure of total particle flux in units of  $\text{mg m}^{-2} \text{day}^{-1}$  was then calculated by knowing the sample fraction weight, the sediment trap aperture area, and the sampling frequency. Another fraction, designed for the total coccolith flux computation, was first centrifuged in order to disassemble particle aggregates, then filtered on a cellulose acetate filter and finally used for coccolith counting. The measure of total coccolith flux in units of  $\text{nC m}^{-2} \text{day}^{-1}$  thus represents the sum of coccoliths sinking alone and of those deriving from disaggregated coccospheres. Additional technical details regarding the sampling procedures may be found in Malinverno et al. (in preparation).

Flux data at the Urania station are available at both 500 m and 2800 m depth for Mooring 1 and at 500 m depth for Mooring 2. At the Bannock station only total particle flux at 2800 m depth is available (see Table 1).

A few technical problems were encountered during the sediment trap deployment and recovery and during the processing of some samples. At the Urania site, a few samples from the Mooring 1 time series collected at 2800 m depth dissolved in the laboratory after being weighted and before the coccolith count. The total particle flux time series is therefore complete, whereas a few samples are missing from the coccolith flux time series. Moreover during the sediment trap recovery of Mooring 1, the last collecting bottle remained open during the ascent through the water column. The last samples of the two time series – 500 m depth and 2800 m depth – may therefore give an overestimated value of the true particle flux. At the Bannock site, the sediment trap rotation mechanism stopped after the 16th temporal interval of Mooring 1, i.e. after 6 March 2000. Particles sinking during the remaining part of the time series (7 March–2 June 2000) were then collected in one single vial during a period of 88 days.

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 Vertical velocities calculation

The model used in this study is derived from the OPA code (Madec et al., 1998), it has a horizontal resolution of  $1/16^\circ \times 1/16^\circ$  and the vertical coordinate is a full cell z-level step-wise discretization with 72 unevenly spaced levels. Vertical resolution is enhanced in the upper layers and gradually decreases with depth (Tonani et al., 2008).

The current vertical velocity field is computed as a diagnostic variable according to the continuity equation for an incompressible fluid:

$$w(z) = w(-H) - \int_{-H}^z \left( \frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} \right) dz' \quad (1)$$

where  $u$  and  $v$  are the horizontal velocity components,  $w$  is the vertical velocity component and  $-H$  is the bottom depth. The kinematic bottom boundary condition for vertical velocity can be expressed as:

$$w(-H) = -\mathbf{U}_h \cdot \nabla_h H, \quad (2)$$

where  $\mathbf{U}_h = (u, v)$  and  $\nabla_h = (\partial/\partial x, \partial/\partial y)$ . Because of the bottom topography discretization used,  $\nabla_h H$  is equal to zero, thus vertical velocity at the ocean bottom will also be zero. The disadvantage of the z-level step-wise vertical discretization is that topographic gradients less than the grid aspect ratio are not well resolved unless very high vertical resolution is used. This may lead in some cases to small scale vertical velocities with artificially high amplitudes. This behavior was observed by Pacanowski and Gnanadesikan (1997) who compare a full cell and a partial cell discretization, which gives a better representation of topographic gradients. In the latter case the effect of Eq. (2) is mitigated and the current vertical velocity field appears to be smoother and less affected by small-scale noise.

We compare the vertical velocities computed with Eq. (1) with the Ekman vertical velocities, which are calculated as a function of the curl of the wind stress  $\boldsymbol{\tau} = (\tau_x, \tau_y)$ ,

**BGD**

5, 3123–3156, 2008

### Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



according to:

$$w_E(x, y, \delta_E) = \frac{1}{\rho_0 f} \left( \mathbf{k} \cdot \nabla_h \times \boldsymbol{\tau} + \frac{\beta \tau_x}{f} \right), \quad (3)$$

where  $w_E$  is the Ekman vertical velocity,  $\boldsymbol{\tau}$  is computed from ECMWF winds using a bulk formula as reported in Tonani et al. (2008),  $\rho_0$  is the average seawater density taken as 1029 kg m<sup>-3</sup>, and  $f$  is the Coriolis parameter. The  $\beta$  parameter, i.e. the meridional gradient of  $f$ , is computed as  $\beta = 2\Omega \cos \theta / r$ , where  $r$  and  $\Omega$  are the Earth radius and angular velocity, and  $\theta$  is the latitude.

Ekman vertical velocities are defined at the Ekman layer depth  $\delta_E$ , i.e. the e-folding depth of the wind-induced currents. For the Mediterranean Sea  $\delta_E$  is generally between 20 and 40 m. Such an estimate is calculated using the scaling formula  $\delta_E = \pi \sqrt{2A_v / |f|}$  (Pond and Pickard, 1983), where  $A_v$  is the vertical eddy coefficient for momentum, calculated in the model as a function of the Richardson number (Tonani et al., 2008), and normally variable between 10<sup>-2</sup> and 10<sup>-3</sup> m<sup>2</sup> sec<sup>-1</sup>.

We use the daily wind stress fields for the years 1999–2001 to compute the Ekman vertical velocity and the daily current velocity fields of the model output to compute the current vertical velocity. The comparison between the two fields is done on a monthly basis at 30 m depth.

The comparison between the current vertical velocity field and the particle flux data is done by averaging the ocean vertical velocity field on the same time intervals of the particle flux time series, i.e. 10–11 days for Mooring 1 and 14–15 days for Mooring 2.

The obtained fields are then vertically averaged between 100 m and 600 m depth: this interval allows for the surface high frequency signals to be neglected and is deep enough for the nutricline depth to be included at all seasons (Ediger and Yilmaz, 1996; Klein et al., 1999). The vertical velocity fields are then horizontally averaged on rectangular areas of 7381 km<sup>2</sup> and of 7471 km<sup>2</sup> centered on the Urania and Bannock stations respectively. We believe that these areas are large enough for model horizontal phase

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



errors to be taken into account, while conserving the specificity of the circulation features of the two regions.

Correlations between current vertical velocities and particle fluxes are calculated both without a time lag and with a shift between time series of one and two sediment trap rotation intervals, corresponding to a lag of 11 and 22 days for Mooring 1 and to a lag of 15 and 30 days for Mooring 2. This is done in order to assess the delayed effects of ocean vertical velocities on particle fluxes. The statistical significance of the correlation coefficients is tested by applying a correlation t-test (Davis, 2002). The null hypothesis (i.e. that the correlation coefficients are statistically significant) is first of all tested at 90% confidence level: in fact, due to the smallness of the sample, there might be the risk of rejecting the null hypothesis when it should be accepted. In cases where the null hypothesis is accepted at 90% confidence level, we test the null hypothesis also at 95% and 99% confidence levels.

### 3 Results

#### 3.1 Surface circulation in the Ionian Sea

In Fig. 2 a temporal average of the simulated horizontal circulation at 30 m depth for year 1999 is depicted. Modified Atlantic Water entering the Sicily Strait flows mainly eastwards towards the Levantine Sea and separates the Ionian Basin in two regions characterized by an anticyclonic circulation in the south and a weak and prevailing cyclonic circulation in the north. A large meander of the Modified Atlantic Water current, called the Atlantic Ionian Stream (Robinson et al., 1999), is present in the Ionian Sea around 34° N with an anticyclonic curvature and a detached anticyclonic eddy to the north (up to 37° N).

The Ionian Sea was affected in the late 80s by an intensification and northward extension of the anticyclonic circulation (Malanotte-Rizzoli et al., 1999; Demirov and Pinardi, 2002). This circulation change, which is an aspect of the widespread phenomenon

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



known as the Eastern Mediterranean Transient, appears to have recovered since the late 90s (Manca et al., 2003). The weak but cyclonic circulation north of 35° N and the large but limited in extent anticyclonic meander and northward eddy are signatures of such change, which our model succeeds to capture.

### 5 3.2 Vertical velocities in the Ionian Sea

The Ekman Vertical Velocity field (hereafter referred to as EVV) calculated according to Eq. (3), is depicted in Fig. 3 for the months of November 1999 (a), June 2000 (b), November 2000 (c), and February 2001 (d), which are months overlapping with sediment trap data. EVVs have values between  $-0.1$  and  $0.1$   $\text{m day}^{-1}$  on large portions of the Ionian Basin and exhibit a typical large scale structure consistent with the large scale wind field. Positive values, i.e. upward EVVs, are prevalently found in the northern part of the Ionian basin, while the southeastern regions are typically characterized by negative values, i.e. downward EVVs. Seasonal and interannual variability is moreover evident: in February 2001 the Ionian Basin exhibits the largest areas with upward EVV, while June and November 2000 are characterized by prevailing downward EVVs. The upward EVVs are then stronger in the northern part of the basin and in the winter season.

The Current Vertical Velocity field (hereafter referred to as CVV), calculated at 30 m depth according to Eq. (1), is depicted in Fig. 4 for the months of November 1999 (a), June 2000 (b), November 2000 (c), and February 2001 (d). The CVV field is noisier with respect to the EVV field and values generally exceed  $0.1$   $\text{m day}^{-1}$  in absolute value. This is due to the intense mesoscale eddy field that modulates the Ekman vertical velocities. Consistently with the EVV field, a north-south gradient in the CVV sign and a large seasonal variability are evident, with a broadening of the areas of downward CVVs during the summer months and of the areas of upward CVVs during the winter months. The comparison between the CVV field in November 1999 and in November 2000 reveals a significant interannual variability, in agreement with the EVV field.

An overall consistency between the CVV and the EVV fields is therefore observed.

**BGD**

5, 3123–3156, 2008

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



However, the small scale structure of the general circulation leads to a strong variability of current vertical velocities, in both space and time. The Urania and Bannock stations, despite their vicinity, may therefore be characterized by a different circulation regime. This variability does not appear to be correctly captured by the large scale EVV field.

5 We thus believe that the usage of EVVs to compute correlations with sediment trap data is not suitable for our purposes.

### 3.3 Particle fluxes at the Urania and Bannock stations

Total Particle Flux (hereafter referred to as TPF) measured at the Urania site is depicted in Fig. 5 with a light-gray line; current vertical velocity values are superimposed on the same plot with a dark-gray line. Vertical arrows located in the upper part of the panel indicate the timing of the particle flux maxima exceeding  $25 \text{ mg m}^{-2} \text{ day}^{-1}$ .

At 500 m depth (upper panel) TPF is highest at the end of May-beginning of June 2000, when it attains a value of more than  $100 \text{ mg m}^{-2} \text{ day}^{-1}$ . Secondary flux maxima of roughly  $50 \text{ mg m}^{-2} \text{ day}^{-1}$  occur in at the end of September 1999 and in November–December 2000. Fluxes in excess of  $25 \text{ mg m}^{-2} \text{ day}^{-1}$  are observed in November 1999, in March 2000 and in April 2001. TPFs lower than  $2 \text{ mg m}^{-2} \text{ day}^{-1}$  occur throughout the period January–March 2001. At 2800 m depth (lower panel), the highest TPF value is attained at the end of May 2000, with a value of almost  $80 \text{ mg m}^{-2} \text{ day}^{-1}$ , whereas secondary flux maxima in excess of  $25 \text{ mg m}^{-2} \text{ day}^{-1}$  occur in the period of September–October 1999.

TPF at the Urania site is on average  $19 \text{ mg m}^{-2} \text{ day}^{-1}$  at 500 m depth (with a standard deviation of  $28 \text{ mg m}^{-2} \text{ day}^{-1}$ ) and  $15 \text{ mg m}^{-2} \text{ day}^{-1}$  at 2800 m depth (with a standard deviation of  $17 \text{ mg m}^{-2} \text{ day}^{-1}$ ). The large value of the standard deviation indicates that the mean is not a robust estimate of the probability distribution due to the eventful and short time series. However this is the longest available at the moment.

Total Coccolith Flux (hereafter referred to as TCF) measured at the Urania site is depicted in Fig. 6 with a light-gray line; current vertical velocity values are superimposed on the same plot with a dark-gray line. Vertical arrows located in the upper part of the

---

## Particle fluxes and ocean vertical velocities

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



panel indicate the timing of the coccolith flux maxima exceeding  $5 \times 10^8 \text{ nC m}^{-2} \text{ day}^{-1}$ .

TCF at 500 m depth (upper panel) shows a sporadic peak exceeding  $3 \times 10^9 \text{ nC m}^{-2} \text{ day}^{-1}$  at the end of May–beginning of June 2000 and secondary flux maxima in excess of  $5 \times 10^8 \text{ nC m}^{-2} \text{ day}^{-1}$  at the end of September 1999 and in March–April 2000. At 2800 m depth (lower panel) TCF exhibits the largest values (in excess of  $5 \times 10^8 \text{ nC m}^{-2} \text{ day}^{-1}$ ) in the period September–October–November 1999 and at the end of May 2000. TCFs have the same order of magnitude between the upper and deep traps, i.e.,  $10^8 \text{ nC m}^{-2} \text{ day}^{-1}$ : this might be an indication of the good preservation of coccolithophore tests in the descent through the water column. TCF measured at 500 m depth during the second part of the time series is extremely low with respect to the Mooring 1 time series. We suspect that calcite dissolution might have partly occurred in the samples: therefore this part of the time series might not be fully reliable.

We may thus conclude that at the Urania site, fluxes are extremely high at the end of May–beginning of June 2000, i.e., at the beginning of summer, and are generally larger in the autumn and spring seasons.

TPF measured at the Bannock site is depicted in Fig. 7 with a light-gray line; current vertical velocity values are superimposed on the same plot with a dark-gray line. A vertical arrow in the upper part of the panel indicates the particle flux peak exceeding  $25 \text{ mg m}^{-2} \text{ day}^{-1}$ .

TPF is highest at the beginning of June 2000, when it attains a value of almost  $80 \text{ mg m}^{-2} \text{ day}^{-1}$ . There appear to be no other large flux peak: in the remaining part of the time series, TPF remains in fact below  $4 \text{ mg m}^{-2} \text{ day}^{-1}$ , except for the period 7 March–2 June 2000, when it is  $8.84 \text{ mg m}^{-2} \text{ day}^{-1}$ . However, since the particles sinking throughout this period were collected in one single vial, we cannot know whether the higher TPF belongs to one single event or it is due to an overall increase of flux magnitude over the whole period. TPF at the Bannock site at 2800 m depth is on average  $8 \text{ mg m}^{-2} \text{ day}^{-1}$  (with a standard deviation of  $16 \text{ mg m}^{-2} \text{ day}^{-1}$ ).

TPFs measured at the Urania site are totally comparable with those observed in other off-shore sites of the Ionian Sea, while TPFs at the Bannock site are lower. Ziveri

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2000) observe in fact an average TPF of about  $16 \text{ mg m}^{-2} \text{ day}^{-1}$  at 3000 m depth at the Bannock site, while Boldrin et al. (2002) observe an average TPF of about  $17 \text{ mg m}^{-2} \text{ day}^{-1}$  at 2250 m depth in the northern Ionian Sea.

Correlations between TPF and TCF are significant at 99% confidence level at both 500 m and 2800 m depth. This result may suggest an abundance of coccoliths in the total particle flux, as well as a common sinking mechanism between biogenic and lithogenic material. It is moreover evident that some of the flux “events” appear instantaneously in the upper and deep traps. Consistently, correlation coefficients calculated between the 500 m and the 2800 m depth fluxes are significant at 99% confidence level for both TPF and TCF.

Since the sediment traps at different depths are well correlated, we can compute the vertical velocity of the material sinking between 500 m and 2800 m depth. Considering the sampling resolution interval we estimate sinking velocities higher than  $200 \text{ m day}^{-1}$  for both total particles and total coccoliths.

Correlations between the TPFs of the trap at 500 m depth and the trap at 2800 m depth are significant also with 11 days lag (99% confidence level) and with 22 days lag (95% confidence level). We may therefore hypothesize that a range of particle sinking velocities might be occurring in the ocean interior, with values between  $70 \text{ m day}^{-1}$  and more than  $200 \text{ m day}^{-1}$ .

Fluxes of single coccolithophore species at the two investigated depths (Malinverno et al., in preparation) display a pattern that is consistent with this inference. In fact in the 500 m trap there is a rather distinct seasonal signal in assemblage composition, with specific abundance peaks occurring over short time intervals. In contrast in the 2800 m trap the signal is more smoothed and occurs over longer time periods.

Particle sinking speeds of more than  $200 \text{ m day}^{-1}$  are in general agreement with other studies conducted in open areas of the Mediterranean Sea. In the northwestern Mediterranean Sea, Lee et al. (2004) observe, at 200 m depth, a dominant particle sinking regime of  $200 \text{ m day}^{-1}$ . In the northeastern Ionian Sea, Boldrin et al. (2002), estimate particle sinking speeds higher than  $140 \text{ m day}^{-1}$  by computing the time lag

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between sediment trap flux peaks measured at different depths. On the contrary Ziveri et al. (2000) estimate, in the southeastern Ionian Sea, settling rates of 17–25 m day<sup>-1</sup> for coccoliths and of more than 100 m day<sup>-1</sup> for coccospheres by calculating the time lag between the 1979–1985 satellite chlorophyll maximum, and the sediment trap flux peak at 3000 m depth available for the period 1991–1994. However we may argue that at the beginning of the 80s, there could have been different chlorophyll concentrations and phytoplankton community composition, with respect to the beginning of the 90s, because of the Eastern Mediterranean Transient event (Klein et al., 1999).

### 3.4 Correlation between current vertical velocities and particle fluxes

CVVs at the Urania site (Figs. 5 and 6) exhibit positive values exceeding 80 cm day<sup>-1</sup> at the end of September 1999 and in November 2000, whereas negative values lower than -30 cm day<sup>-1</sup> occur in June 2000 and in February 2001. Thus we observe an increase in upward motion in autumn and an increase in downward motion in summer, while winter months exhibit a different behavior between the years 2000 and 2001.

CVVs at the Bannock site (Fig. 7) remain negative during most of the time series. In the period from October 1999 to May 2000, CVVs are on average -59 cm day<sup>-1</sup>. Afterwards they remain negative or quite close to zero except for the period November–December 2001, when an increase of upward vertical velocities up to 50 cm day<sup>-1</sup> occurs.

In both the Urania and Bannock sites, periods of upward vertical velocities are normally associated with higher particle fluxes. This behavior is observed at the Urania site at the end of September 1999, in March 2000 (in the 500 m trap only) and in November 2000. Periods of downward vertical velocities appear instead to coincide with periods of low particle fluxes, as seen in February 2001 at the Urania site and throughout the period October 1999–May 2000 at the Bannock site.

An exception to this pattern occurs at the end of May–beginning of June 2000, when the high particle flux occurring at both the Urania and Bannock sites is not associated with a large increase of upward CVVs. We decided to remove the peak in June 2000

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from the remaining computations: we believe in fact that mechanisms other than ocean vertical velocities are determining the strong particle fluxes observed in this period in both the Urania and Bannock sites. This hypothesis will be discussed in the next section.

5 In Table 2 we illustrate the correlation coefficients – and associated confidence levels – between current vertical velocities and particle fluxes. At the Urania station and during Mooring 1, correlations between coincidental time series of CVV and particle fluxes are significant at 99% confidence level for TCF at 500 m and 2800 m depth and for TPF at 2800 m depth. When the particle flux time series are lagged of one rotation  
10 interval (i.e. 11 days) with respect to the CVV time series, correlations are significant at 99% confidence level for all particle fluxes at 500 m depth, while at 2800 m depth correlations are significant at 95% confidence level. When two temporal shifts are applied (i.e. 22 days), correlations are significant at 99% confidence level for particle fluxes at 2800 m depth, while they are significant at 95% confidence level for TCF at  
15 500 m and not significant for TPF at 500 m depth. During Mooring 2, TPF is correlated with ocean vertical velocities at 95% confidence level for coincidental time series and at 90% confidence level for lagged time series. TCF does not exhibit instead significant correlations (not shown): this behavior might be due to partial calcite dissolution in the samples.

20 At the Bannock station and during Mooring 1, correlations between CVV and TPF at 2800 m depth are significant at 90% confidence level when the time series are shifted of two rotation intervals. During Mooring 2, correlations between CVV and TPF are not significant.

25 We thus conclude that a prevailing positive correlation exists at both the Urania and Bannock stations between particle fluxes and current vertical velocities. Furthermore during Mooring 1 we observe that at 500 m depth correlations are highest with 11 days lag, while at 2800 m depth correlations are highest with 22 days lag. This behavior might suggest a delay process between CVV and particle fluxes, as will be discussed in the next section.

---

**Particle fluxes and  
ocean vertical  
velocities**L. Patara et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## 4 Discussion

### 4.1 Correlation between current vertical velocities and particle fluxes

The positive correlation between upward vertical velocities and particle fluxes might seem at first sight surprising. We might think that downward vertical velocities should enhance particle fluxes through an increase of vertical sinking velocities. To explain this apparent inconsistency, we must first of all consider that the particle sinking rates estimated from the sediment traps are two orders of magnitude higher than the ocean vertical velocities calculated from the model output. We thus have to exclude a direct effect of ocean vertical velocities on deep particle fluxes and we speculate that, in the region analyzed in this study, another sedimentation mechanism is at work.

The high sinking rates estimated in this work might be attained by means of biological aggregation of particles inside fecal pellets produced by macro-zooplankton. Mazzocchi et al. (2003) detect in spring 1999 an increase in abundance of suspension filter feeders (mainly appendicularians, salps, and doliolids) in the eastern Ionian Sea. Pelagic tunicates feed on unicellular phytoplankton by filtering large volumes of water and are adapted to dilute environments where they exhibit a rapid population growth if environmental conditions are favorable (Alldredge and Madin, 1982). Pelagic tunicates produce fecal pellets which sink very rapidly, with a range between 242 and 2700 m day<sup>-1</sup> for salps (Bruland and Silver, 1981; Yoon et al., 1996; Yoon et al., 2001) and between 25 and 504 m day<sup>-1</sup> for appendicularians and doliolids (Turner, 2002 and references therein).

We hypothesize that the following sedimentation mechanism might be occurring in the open areas of the Eastern Mediterranean Sea. An upward vertical velocity can produce a nutricline uplift in the euphotic layer. Coccolithophores would take advantage of the nutrient increase giving rise to enhanced primary production, which would eventually stimulate grazing. The calcitic constituents of coccolithophores would then sink through the water column inside rapidly sinking fecal pellets or other large particles and would be collected by sediment traps (see a schematic representation of this pro-

**BGD**

5, 3123–3156, 2008

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cess in Fig. 8). Considering the sampling resolution of the sediment traps (10–11 days for Mooring 1) we realize that the described mechanism of upwelling-enhanced deep particle fluxes is very fast.

Not only TCFs, but also TPFs are correlated with ocean vertical velocities on coincidental time series. This observation is corroborating the hypothesis of a common sinking mechanism among the various components of the total particle flux, e.g. through biological aggregation inside macro-zooplankton fecal pellets.

Considering the significant correlations between ocean vertical velocities and lagged particle flux time series we envisage a spectrum of particle sinking rates in the ocean interior. This conclusion was also inferred from the correlation values obtained between lagged particle flux time series at 500 m and at 2800 m depth. Associated with the fast sinking regime of more than  $200 \text{ m day}^{-1}$ , there appears to be a slower regime of about  $100 \text{ m day}^{-1}$ , which might be responsible for the significant correlations between ocean vertical velocities and particle fluxes with 11 days lag at 500 m depth and with 22 days lag at 2800 m depth.

The lack of significant correlations at the Bannock site during Mooring 2 suggests that the described model of upwelling-enhanced deep particle fluxes is not always applicable. It is possible that other physical and biological processes are influencing the observed particle fluxes. Nutrient levels in the upper ocean may be also affected by horizontal advection and vertical mixing. In addition seasonal variability of macro-zooplankton abundance might lead to varying sinking regimes throughout the year. Finally we cannot exclude that technical issues related to the trapping efficiency of the sediment traps might have affected the particle flux time series.

#### 4.2 The high sedimentation event of May–June 2000

At the end of May–beginning of June 2000, a strong particle flux peak occurs at both the Urania and Bannock sites (Figs. 5, 6 and 7). This particle flux peak is not associated with an increase of upward CVVs. We suggest that this particle flux peak is due to a strong dust event, which would enhance lithogenic and possibly biogenic flux.

---

## Particle fluxes and ocean vertical velocities

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Particle fluxes and  
ocean vertical  
velocities**

---

L. Patara et al.

Saharan dust transport over the Mediterranean occurs frequently in spring and summer in the form of non-continuous “pulses” (Kubilay et al., 2000). The effect of the Saharan dust particles on biogenic flux may be double: firstly it may represent an important source of nutrients in the euphotic layer (Markaki et al., 2003; Herut et al., 2005). Secondly lithogenic particles may have a ballasting effect which would promote the sinking of the biogenic particles within the ocean interior (Klaas and Archer, 2002).

A time series of SeaWiFS TrueColor satellite images for the period 22 May–13 July 2000 is available (Fig. 9). Images reveal an intense dust event, associated with an atmospheric cyclonic structure, occurring in the period 22–27 May 2000 in the southeastern Ionian Sea. The Urania and Bannock sites are on the direct trajectory of this dust phenomenon. We can therefore reasonably consider this event as responsible for the high flux peak detected in the sediment trap starting from the end of May 2000.

## 5 Conclusions

The mechanisms of sedimentary particle fluxes in the deep Urania and Bannock sites were studied by means of sediment trap records and model simulated vertical velocities.

The current vertical velocities for the area of the Ionian Sea have a large mesoscale signal that is superimposed to a seasonally varying Ekman vertical velocity field at low amplitude. It is found that current vertical velocities at the Urania site are positive (upward motion) in autumn and in spring when pulses of total particle fluxes are found in the water column at 500 m and 2800 m depth.

From sediment trap data the sinking velocities of TPF and TCF are estimated to be 100–200 m day<sup>-1</sup> while current vertical velocities are estimated to be around 1 m day<sup>-1</sup>. This observation brought us to envisage a tight coupling between surface layers and deep ocean through particle aggregation mechanisms. We suggested that fast sinking rates of organic material in the deep ocean are connected to both lithogenic and biological aggregation mechanism that transfer particles rapidly in the deep water column.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In synthesis two emerging fast sinking mechanisms have been captured:

1. Pulses of primary production, triggered by upward current vertical velocities, followed by grazing and macro-zooplankton-related biogenic flux that rapidly conveys the material in the deep ocean.
2. Large Saharan dust events that fertilize the upper ocean and possibly contribute to aggregation of organic material, thus producing large sedimentation fluxes.

If these mechanisms are at work in other open-ocean areas, it is possible to argue that the carbon export to the deep ocean is larger in the cyclonic gyres of the ocean general circulation and during the months of maximum upward vertical velocities. The fast sinking rates estimated in this study are moreover an evidence of the efficiency of the biological pump in sequestering organic carbon from the surface layers of the open areas of the Mediterranean Sea. Further work is however needed to unravel the relevant physical features of the marine environment capable of affecting the deep particle fluxes.

*Acknowledgements.* Sediment trap samples were available for the study of biogenic particles within an Italian-Dutch cooperation agreement. Mooring equipment was made available by NWO/ALW, through the Dutch Marine Research Facilities at NIOZ; ship-time was funded alternatively by NWO and the Italian CoNISMa (National Interuniversity Consortium for Marine Sciences). The study of biogenic particles was performed within the European Project BIODEEP (BIOTEchnologies from the DEEP, EVK3-2000-00042).

## References

- Agostini, V. N. and Bakun, A.: "Ocean triads" in the Mediterranean Sea: physical mechanisms potentially structuring reproductive habitat suitability (with example application to European anchovy, *Engraulis encrasicolus*), *Fish. Oceanogr.*, 11(3), 129–142, 2002.
- Allredge, A. L. and Gotschalk, C.: In situ settling behavior of marine snow, *Limnol. Oceanogr.*, 33(3), 339–351, 1988.

**BGD**

5, 3123–3156, 2008

## Particle fluxes and ocean vertical velocities

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Allredge, A. L. and Madin, L. P.: Pelagic tunicates – unique herbivores in marine plankton, *Bioscience*, 32(8), 655–663, 1982.
- Andersen, V. and Prieur, L.: One-month study in the open NW Mediterranean Sea (DYNAPROC experiment, May 1995): overview of the hydrobiogeochemical structures and effects of wind events, *Deep-Sea Res., Part I, Oceanographic research papers*, 47(3), 397–422, 2000.
- Asper, V. L., Deuser, W. G., Knauer, G. A., and Lohrenz, S. E.: Rapid coupling of sinking particle fluxes between surface and deep ocean waters, *Nature*, 357(6380), 670–672, 1992.
- Boldrin, A., Miserocchi, S., Rabitti, S., Turchetto, M. M. Balboni, V., and Socal, G.: Particulate matter in the southern Adriatic and Ionian Sea: characterization and downward fluxes, *J. Marine Syst.*, 33, 389–410, 2002.
- Bricaud, A., Bosc, E., and Antoine, D.: Algal biomass and sea surface temperature in the Mediterranean Basin – Intercomparison of data from various satellite sensors, and implications for primary production estimates, *Remote Sens. Environ.*, 81(2–3), 163–178, 2002.
- Bruland, K. W. and Silver, M. W.: Sinking rates of fecal pellets from gelatinous zooplankton (salps, pteropods, doliolids), *Mar. Biol.*, 63(3), 295–300, 1981.
- Casotti, R., Landolfi, A., Brunet, C., D’Ortenzio, F., Mangoni, O., Ribera d’Alcalà, M., and Denis, M.: Composition and dynamics of the phytoplankton of the Ionian Sea (eastern Mediterranean), *J. Geophys. Res.-Oceans*, 108(C9), 8116, 2003.
- Davis, J. C.: *Statistics and Data Analysis in Geology*, 3rd edition, Wiley & Sons, New York, 656 pp., 2002.
- Demirov, E. and Pinardi, N.: Simulation of the Mediterranean Sea circulation from 1979 to 1993: Part I. The interannual variability, *J. Marine Syst.*, 33, 23–50, 2002.
- Ediger, D. and Yilmaz, A.: Characteristics of deep chlorophyll maximum in the Northeastern Mediterranean with respect to environmental conditions, *J. Marine Syst.*, 9(3–4), 291–303, 1996.
- Engel, A.: Distribution of transparent exopolymer particles (TEP) in the northeast Atlantic Ocean and their potential significance for aggregation processes, *Deep-Sea Res., Part I, Oceanographic research papers*, 51(1), 83–92, 2004.
- Fowler, S. W., Buat-Menard, P., Yokoyama, Y., Ballestra, S., Holm, E., and Van Nguyen, H.: Rapid removal of Chernobyl fallout from Mediterranean surface waters by biological-activity, *Nature*, 329(6134), 56–58, 1987.
- Herut, B., Zohary, T., Krom, M. D., Mantoura, R. F. C., Pitta, P., Psarra, S., Rassoulzadegan, F., Tanaka, T., and Thingstad, T. F.: Response of East Mediterranean surface water to Saha-

**BGD**

5, 3123–3156, 2008

---

**Particle fluxes and  
ocean vertical  
velocities**L. Patara et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- ran dust: On-board microcosm experiment and field observations, *Deep-Sea Res., Part II, Topical studies in oceanography*, 52(22–23), 3024–3040, 2005.
- Honjo, S.: Coccoliths – Production, Transportation and Sedimentation, *Mar. Micropaleontol.*, 1(1), 65–79, 1976.
- 5 Honjo, S., Doherty, K. W., Agrawal, Y. C., and Asper, V. L.: Direct optical assessment of large amorphous aggregates (marine snow) in the deep ocean, *Deep-Sea Res., Part A, Oceanographic research papers*, 31(1), 67–76, 1984.
- Ignatiades, L.: Scaling the trophic status of the Aegean Sea, eastern Mediterranean, *J. Sea Res.*, 54(1), 51–57, 2005.
- 10 Klaas, C. and Archer, D. E.: Association of sinking organic matter with various types of mineral ballast in the deep sea: Implications for the rain ratio, *Global Biogeochem. Cy.*, 16(4), 1116, 2002
- Klein, B., Roether, W., Manca, B. B., Bregant, D., Beitzel, V., Kovacevic, V., and Luchetta, A.: The large deep water transient in the Eastern Mediterranean, *Deep Sea Res., Part I, Oceanographic Research Papers*, 46(3), 371–414, 1999.
- 15 Knappertsbusch, M. and Brummer, G.-J. A.: A sediment trap investigation of sinking coccolithophorids in the North-Atlantic, *Deep Sea Res., Part I, Oceanographic Research Papers*, 42(7), 1083–1109, 1995.
- Korres, G., Pinardi, N., and Lascaratos, A.: The ocean response to low-frequency interannual atmospheric variability in the Mediterranean Sea, Part I: sensitivity experiments and energy analysis, *J. Climate*, 13(4), 705–731, 2000.
- 20 Kubilay, N., Nickovic, S., Moulin, C., and Dulac, F.: An illustration of the transport and deposition of mineral dust onto the eastern Mediterranean, *Atmos. Environ.*, 34(8), 1293–1303, 2000.
- Lampitt, R. S., Hillier, W. R., and Challenor, P. G.: Seasonal and diel variation in the open ocean concentration of marine snow aggregates, *Nature*, 362(6422), 737–739, 1993.
- 25 Lee, C., Armstrong, R. A., Beck, A., Cochran, J. K., Wakeham, S. G., Peterson, M. L., and Miquel, J.-C.: MEDFLUX: Association of organic matter with ballast minerals in sinking particles, American Geophysical Union Fall Meeting, San Francisco (CA, USA), 13–17 December 2004, OS52A-02, 2004.
- 30 Lévy, M., Mémery, L., and Madec, G.: The onset of a bloom after deep winter convection in the northwestern Mediterranean sea: mesoscale process study with a primitive equation model, *J. Marine Syst.*, 16(1–2), 7–21, 1998.
- Madec, G., Delecluse, P., Imbard, M., and Lévy, C.: OPA version 8.1 Ocean General Circulation

**BGD**

5, 3123–3156, 2008

---

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Model reference manual, Technical Report, Note 11, LODYC/IPSL, Paris, France, 91 pp., 1998.

Malanotte-Rizzoli, P., Manca, B. B., D'Alcalà, M. R., Theocharis, A., Bergamasco, A., Bregant, D., Budillon, G., Civitarese, G., Georgopoulos, D., Michelato, A., Sansone, E., Scarazzato, P., and Souvermezoglou, E.: A synthesis of the Ionian Sea hydrography, circulation and water mass pathways during POEM Phase I, *Prog. Oceanogr.*, 39(3), 153–204, 1997.

Malanotte-Rizzoli, P., Manca, B. B., Ribera d'Alcala, M., Theocharis, A., Brenner, S., Budillon, G., and Oszoy, E.: The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations, *Dynam. Atmos. Oceans*, 29(2–4), 365–395, 1999.

Malinverno, E., Ziveri, P., and Corselli, C.: Coccolithophorid distribution in the Ionian Sea and its relationship to eastern Mediterranean circulation during late fall to early winter 1997, *J. Geophys. Res.-Oceans*, 108(C9), 8115, 2003.

Manca, B. B., Budillon, G., Scarazzato, P., and Ursella, L.: Evolution of dynamics in the eastern Mediterranean affecting water mass structures and properties in the Ionian and Adriatic Seas, *J. Geophys. Res.-Oceans*, 108(C9), 8102, 2003.

Markaki, Z., Oikonomou, K., Kocak, M., Kouvarakis, G., Chaniotaki, A., Kubilay, N., and Mihalopoulos, N.: Atmospheric deposition of inorganic phosphorus in the Levantine Basin, eastern Mediterranean: Spatial and temporal variability and its role in seawater productivity, *Limnol. Oceanogr.*, 48(4), 1557–1568, 2003.

Mazzocchi, M. G., Nervegna, D., D'Elia, G., Di Capua, I., Aguzzi, L., and Boldrin, A.: Spring mesozooplankton communities in the epipelagic Ionian Sea in relation to the Eastern Mediterranean Transient, *J. Geophys. Res.-Oceans*, 108, 8114, 2003.

Molcard, A., Pinardi, N., Iskandarani, M., and Haidvogel, D. B.: Wind driven general circulation of the Mediterranean Sea simulated with a Spectral Element Ocean Model, *Dynam. Atmos. Oceans*, 35(2), 97–130, 2002.

Nittis, K., Pinardi, N., and Lascaratos, A.: Characteristics of the summer 1987 flow-field in the Ionian Sea, *J. Geophys. Res.-Oceans*, 98(C6), 10 171–10 184, 1993.

Pacanowski, R. C. and Gnanadesikan, A.: Transient response in a z-level ocean model that resolves topography with partial cells, *Mon. Weather Rev.*, 126(12), 3248–3270, 1998.

Pinardi, N. and Navarra, A.: Baroclinic wind adjustment processes in the Mediterranean Sea, *Deep Sea Res., Part II, Topical Studies in Oceanography*, 40(6), 1299–1326, 1993.

Pinardi, N. and Masetti, E.: Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: a review, *Palaeogeogr. Palaeoclimatol.*, 158(3–4), 153–174,

**BGD**

5, 3123–3156, 2008

---

## Particle fluxes and ocean vertical velocities

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2000.

Pinardi, N., Arneri, E., Crise, A., Ravaioli, M., and Zavatarelli, M.: The physical, sedimentary and ecological structure and variability of shelf areas in the Mediterranean Sea, in: *The Sea*, Vol. 14, Harvard University Press, 2005.

5 Pond, S. and Pickard, G. L.: *Introductory Dynamical Oceanography*, 2nd edition, Elsevier Butterworth-Heinemann, 349 pp., 1983.

Robinson, A. R., Hecht, A., Pinardi, N., Bishop, J., Leslie, W. G., Rosentroub, Z., Mariano, A. J., and Brenner, S.: Small synoptic mesoscale eddies and energetic variability of the Eastern Levantine Basin, *Nature*, 327(6118), 131–134, 1987.

10 Robinson, A. R., Sellschopp, J., Warn-Varnas, A., Leslie, W. G., Lozano, C. J., Haley, P. J., Anderson, L. A., and Lermusiaux, P. F. J.: The Atlantic Ionian Stream, *J. Marine Syst.*, 20(1–4), 129–156, 1999.

Stemann, L., Gorsky, G., Marty, J. C., Picheral, M., and Miquel, J. C.: Four-year study of large-particle vertical distribution (0–1000 m) in the NW Mediterranean Sea in relation to hydrology, phytoplankton, and vertical flux, *Deep Sea Res., Part II, Topical Studies in Oceanography*, 49(11), 2143–2162, 2002.

Tonani, M., Pinardi, N., Dobricic, S., Pujol, I., and Fratianni, C.: A high-resolution free-surface model of the Mediterranean Sea, *Ocean Sci.*, 4, 1–14, 2008, <http://www.ocean-sci.net/4/1/2008/>.

20 Turner, J. T.: Zooplankton fecal pellets, marine snow and sinking phytoplankton blooms, *Aquat. Microb. Ecol.*, 27(1), 57–102, 2002.

Yoon, W. D., Marty, J. C., Sylvain, D., and Nival, P.: Degradation of faecal pellets in *Pegea confoederata* (Salpidae, Thaliacea) and its implication in the vertical flux of organic matter, *J. Exp. Mar. Biol. Ecol.*, 203(2), 147–177, 1996.

25 Yoon, W. D., Kim, S. K., and Han, K. N.: Morphology and sinking velocities of fecal pellets of copepod, molluscan, euphasiid, and salp taxa in the northeastern tropical Atlantic, *Mar. Biol.*, 139, 923–928, 2001.

Ziveri, P., Rutten, A., de Lange, G. J., Thomson, J., Corselli, C.: Present-day coccolith fluxes recorded in central eastern Mediterranean sediment traps and surface sediments, *Palaeogeogr. Palaeoclimatol.*, 158(3–4), 175–195, 2000.

**BGD**

5, 3123–3156, 2008

---

## Particle fluxes and ocean vertical velocities

L. Patara et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Particle fluxes and ocean vertical velocities**

L. Patara et al.

**Table 1.** Available data from the Urania and Bannock sediment traps, where TPF=Total Particle Flux [ $\text{mg m}^{-2} \text{day}^{-1}$ ] and TCF=Total Coccolith Flux [ $\text{nC m}^{-2} \text{day}^{-1}$ ]. Mooring 1=from 15 September 1999 to 2 June 2000 and Mooring 2=from 30 May 2000 to 9 May 2001.

Sampling period	URANIA station		BANNOCK station
	500 m depth	2800 m depth	2800 m depth
Mooring 1	TPF, TCF	TPF, TCF	TPF
Mooring 2	TPF, TCF		TPF

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Particle fluxes and ocean vertical velocities**

L. Patara et al.

**Table 2.** Correlation coefficients between current vertical velocities and particle fluxes at the Urania and Bannock stations, where TPF=Total Particle Flux [ $\text{mg m}^{-2} \text{day}^{-1}$ ] and TCF=Total Coccolith Flux [ $\text{nC m}^{-2} \text{day}^{-1}$ ]. The degrees of freedom – varying among the different time series – are shown in column 3. Correlation coefficients are computed on coincident time series (no lag) and on time series shifted of one (1 lag) and two (2 lags) sediment trap rotation intervals. The duration of the rotation interval is 10–11 days for Mooring 1 and 14–15 days for Mooring 2. Confidence intervals are depicted below the correlation coefficient values, with \*\*\*=99%, \*\*=95%, \*=90%, and n.s.=not significant.

Correlation coefficients between current vertical velocity and particle fluxes					
Station	Parameter	Degrees of freedom	No lag	1 lag	2 lags
URANIA Station Mooring 1	TPF at 500 m	21	0.4 *	0.63 ***	n.s.
		21	0.75 ***	0.63 ***	0.51 **
	TPF at 2800 m	21	0.55 ***	0.51 **	0.56 ***
		12	0.7 ***	0.61 **	0.67 ***
Mooring 2	TPF at 500 m	19	0.45 **	0.42 *	0.39 *
BANNOCK Station Mooring 1	TPF at 2800 m	7	n.s.	n.s.	0.79 **

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and  
ocean vertical  
velocities

L. Patara et al.



**Fig. 1.** Ionian Sea bathymetry and sediment trap Mooring locations. U=Urania station (latitude 35.2° N, longitude 21.2° E), B=Bannock station (latitude 34.2° N, longitude 20° E).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Year 1999



**Fig. 2.** Annual average for year 1999 of the horizontal currents [ $\text{cm sec}^{-1}$ ] at 30 m depth. The arrow length indicates the  $30 \text{ cm sec}^{-1}$  velocity.

**BGD**

5, 3123–3156, 2008

**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and ocean vertical velocities

L. Patara et al.



**Fig. 3.** Ekman vertical velocities [ $\text{m day}^{-1}$ ]: **(a)** November 1999 **(b)** June 2000 **(c)** November 2000 and **(d)** February 2001. In letters we indicate the position of the Urania (U) and Bannock (B) stations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Particle fluxes and  
ocean vertical  
velocities**

L. Patara et al.



**Fig. 4.** Ocean currents vertical velocities [m day<sup>-1</sup>] at 30 m depth: **(a)** November 1999 **(b)** June 2000 **(c)** November 2000 and **(d)** February 2001. In letters we indicate the position of the Urania (U) and Bannock (B) stations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and ocean vertical velocities

L. Patara et al.



**Fig. 5.** Total Particle Flux (TPF) and Current Vertical Velocity (CVV) at the Urania site for Mooring 1 (15 September 1999 to 2 June 2000) and for Mooring 2 (30 May 2000 to 9 May 2001). Light-gray line: Total Particle Flux [ $\text{mg m}^{-2} \text{day}^{-1}$ ] at 500 m depth (top) and at 2800 m depth (bottom). Dark-gray line: Current Vertical Velocity [ $\text{cm day}^{-1}$ ] integrated on the 100–600 m depth interval. With black arrows we indicate Total Particle Flux peaks exceeding  $25 \text{ mg m}^{-2} \text{day}^{-1}$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Particle fluxes and ocean vertical velocities

L. Patara et al.



**Fig. 6.** Total Coccolith Flux (TCF) and Current Vertical Velocity (CVV) at the Urania site for Mooring 1 (15 September 1999 to 2 June 2000) and Mooring 2 (30 May 2000 to 9 May 2001). Light-gray line: Total Coccolith Flux [ $\text{nC m}^{-2} \text{day}^{-1}$ ]  $\times 10^9$  at 500 m depth (top) and at 2800 m depth (bottom). Dark-gray line: Current Vertical Velocity [ $\text{cm day}^{-1}$ ] integrated on the 100–600 m depth interval. With black arrows we indicate Total Coccolith Flux peaks exceeding  $5 \times 10^8 \text{ nC m}^{-2} \text{day}^{-1}$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and ocean vertical velocities

L. Patara et al.



**Fig. 7.** Total Particle Flux (TPF) and Current Vertical Velocity (CVV) at the Bannock site for Mooring 1 (15 September 1999 to 2 June 2000) and Mooring 2 (30 May 2000 to 9 May 2001). Light-gray line: Total Particle Flux [ $\text{mg m}^{-2} \text{day}^{-1}$ ] at 2800 m depth. Dark-gray line: Current Vertical Velocities [ $\text{cm day}^{-1}$ ] integrated on the 100–600 m depth interval Top: whole time series. Bottom: decreased y-axis scale in the periods of low particle fluxes. The black arrow indicates Total Particle Flux exceeding  $25 \text{ mg m}^{-2} \text{day}^{-1}$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and ocean vertical velocities

L. Patara et al.



**Fig. 8.** Model of upwelling-enhanced deep particle fluxes. A nutricline uplift by means of upward vertical velocities leads to a nutrient increase in the euphotic layer. Primary production and eventually grazing are stimulated. Biological aggregation inside fecal pellets rapidly exports particles in the ocean interior up to the bottom where they are collected by sediment traps.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Particle fluxes and  
ocean vertical  
velocities

L. Patara et al.



**Fig. 9.** TrueColor Satellite images: **(a)** 22 May 2000 at 10:36 a.m., **(b)** 22 May 2000 at 12:14 a.m., **(c)** 24 May 2000 at 12:02 a.m., **(d)** 26 May 2000 at 10:15 a.m., **(e)** 26 May 2000 at 11:51 a.m., **(f)** 27 May 2000 at 10:57 a.m.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

