

**Fatty acid
¹³C-fingerprinting in
Mytilid-bacteria
symbiosis**

V. Riou et al.

**Tracing carbon assimilation in
endosymbiotic deep-sea hydrothermal
vent Mytilid fatty acids by
¹³C-fingerprinting**

V. Riou^{1,2}, S. Bouillon^{1,3}, R. Serrão Santos², F. Dehairs¹, and A. Colaço²

¹Department of Analytical and Environmental Chemistry, Vrije Universiteit Brussel, Brussels, Belgium

²Department of Oceanography and Fisheries, IMAR-University of Azores, Horta, Portugal

³Department of Earth and Environmental Sciences, Katholieke Universiteit Leuven, Leuven, Belgium

Received: 4 May 2010 – Accepted: 5 May 2010 – Published: 10 May 2010

Correspondence to: V. Riou (virginie-riou@hotmail.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Bathymodiolus azoricus mussels thrive at Mid-Atlantic Ridge hydrothermal vents, where part of their energy requirements are met via an endosymbiotic association with chemolithotrophic and methanotrophic bacteria. In an effort to describe phenotypic characteristics of the two bacterial endosymbionts and to assess their ability to assimilate CO₂, CH₄ and multi-carbon compounds, we performed experiments in aquaria using ¹³C-labeled NaHCO₃ (in the presence of H₂S), CH₄ or amino-acids and traced the incorporation of ¹³C into total and phospholipid fatty acids (tFA and PLFA, respectively). 14:0, 15:0, 16:1(n-7)*c+t* and 18:1(n-7)*c+t* PLFA were labeled in the presence of H¹³CO₃⁻ (+H₂S) and ¹³CH₄, while the 12:0 compound became labeled only in the presence of H¹³CO₃⁻ (+H₂S). In contrast, the 16:1(n-9), 16:1(n-8) and (n-6), 18:1(n-8)*c* and (n-7), 20:1(n-7) and 18:2(n-7) PLFA were only labeled in the presence of ¹³CH₄. Some of these symbiont-specific fatty acids also appeared to be labeled in mussel gill tFA when incubated with ¹³C-enriched amino acids, and so were mussel-specific fatty acids such as 22:2(n-7,15). Our results provide experimental evidence for the potential of specific fatty acid markers to distinguish between the two endosymbiotic bacteria, shedding new light on C₁ and multi-carbon compound metabolic pathways in *B. azoricus* and its symbionts.

1 Introduction

Deep-sea hydrothermal vents host peculiar ecosystems fueled by methane, sulfide, iron or even hydrogen (e.g., Perner et al., 2009). The symbiotic association of bacteria with marine invertebrate hosts provides the former with access to the chemical substrates necessary for their metabolism and the latter with a source of organic carbon (and nitrogen). One of the two species of endosymbiotic bathymodioline Mytilids occurring along the Mid-Atlantic Ridge (MAR), *Bathymodiolus azoricus*, is found at

BGD

7, 3453–3475, 2010

¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the northernmost sites Menez Gwen, Lucky Strike and Rainbow (Desbruyères et al., 2001). Transmission electron microscopy has demonstrated the presence of two distinct Gram-negative bacterial endosymbionts inside specialized gill epithelial cells (bacteriocytes). One of these symbionts has centrally stacked intracytoplasmic membranes characteristic for type I methane-oxidising bacteria (MOX; Fiala-Médioni et al., 2002). Analysis of 16S rRNA sequences followed by fluorescence in situ hybridization (FISH) evidenced that *B. azoricus* endosymbiotic 16S rRNA phylotypes cluster with natural symbiotic and cultured sulfide-oxidising (SOX) and type I MOX gamma-proteobacteria (Won et al., 2003; Duperron et al., 2006; Spiridonova et al., 2006). Enzymatic and physiological assays using gills of *B. azoricus* from the Lucky Strike and Menez Gwen vent sites revealed the presence of active enzymes of the metabolic cycles of inorganic C assimilation and sulfide oxidation and of the C₁ carbon assimilation pathway (Fiala-Médioni et al., 2002). Furthermore, Riou et al. (2008) report active ¹³C incorporation from bicarbonate in the presence of sulfide as well as from methane within the gills of *B. azoricus*. This incorporation was followed by carbon transfer to the host's aposymbiotic muscle tissue.

In order to understand the modes of matter and energy transfer to the host, the phenotypes of *B. azoricus* symbionts' need to be characterized. The main difficulty consists in the impossibility to grow the symbionts in pure cultures. However, tracer experiments represent powerful tools enabling the detection of the bacterial signal inside the host. Such experiments can be combined with the analysis of phospholipid ester-linked fatty acids (PLFA) which form about 98% of bacterial membrane lipids as compared to only 50% in eukaryotic cells (Vestal and White, 1989). Classification of PLFA profiles has been proven useful to clarify bacterial genus and species interrelationships established by DNA-based phylogeny (Bodelier et al., 2009). PLFA profiles thus allow to distinguish between different MOX species as evidenced for example by the fact that type I MOX bacteria mainly contain fatty acids with 14 and 16 carbon atoms, while type II MOX bacterial PLFA are mainly composed of 18 carbon atoms (Nichols et al., 1985). In addition, MOX bacteria possess fatty acids that are not found

¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in any other known microorganism – type I: 16:1(n-8) and 16:1(n-5)*t*; type II: 18:1(n-8) – and these compounds therefore represent valuable biomarkers (e.g. Nichols et al., 1985).

In the present study, we investigated the incorporation of ^{13}C -labeled HCO_3^- (in the presence of H_2S) or CH_4 into gill-extracted PLFA from *B. azoricus* in order to assess the metabolic activity of the endosymbionts. This enabled us to establish specific fatty acid (FA) patterns which provide insights into the symbiont phenotypes. The potential for heterotrophic growth by the symbionts or the Mytilid itself was furthermore assessed by tracing the assimilation of amino acids (dissolved into seawater) into gill total fatty acids (tFA). Close examination of the labeled FA patterns helped interpreting physiological processes occurring in *B. azoricus*.

2 Material and methods

2.1 Sampling and aquarium experiments

During the MOMARETO cruise (*R/V Pourquoi Pas?*) in August 2006, a cage was deployed at the Menez Gwen site ($37^\circ 51' \text{N}$ – $32^\circ 31' \text{W}$, 817 m) and loaded with around 400 mussels. The cage was retrieved in May 2007 by the Portuguese vessel *R/V Arquipélago*. *Bathymodiolus azoricus* specimens were transferred to cooled seawater (9°C) for a 14 h transit to Horta, Faial Island (Azores). Their valves were scrubbed clean of visible material, rinsed in chilled seawater and transferred to the Azorean land-based hydrothermal vent laboratory, LabHorta (Colaço and Santos, 2003). Before the start of the experiments, mussels were kept for 38 days in aerated seawater amended with methane and hydrogen sulfide, as described in Riou et al. (2008).

2.2 Mussels incubation with ^{13}C -enriched substrates

The available quantity of labelled methane (25% ^{13}C) allowed running the tracer experiment with $^{13}\text{CH}_4$ for a period of 15 days (Riou et al., 2008). The control experiment

BGD

7, 3453–3475, 2010

^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(with filtered seawater only) and the experiments with $\text{NaH}^{13}\text{CO}_3$ (16% ^{13}C)+ H_2S (Riou et al., 2008) or with ^{13}C -labeled amino acids (98% ^{13}C ; Riou et al., 2010) were continued over a period of 20 days. The amino acid mixture consisted of Gly 20–25%, Ala 15–20%, Tyr 10–15%, Leu 5–10%, Lys 5–10%, Ser 5–10%, Thr 2–5%, Phe 1–5%, Pro 1–5%, Val 1–5%, Met <3%, Trp <1%, Ile <1%, His <1%. At the end of the incubations mussels were transferred into filtered seawater and left for 24 h to allow gut clearance.

Three mussels from each experiment were selected for fatty acid analysis, and dissected into gill, mantle, muscle, and remaining tissues. Mussel tissues were immediately stored at -20°C till they were freeze-dried, a few days after dissection.

2.3 PLFA and tFA preparation

Prior to use, all glassware was precombusted 4 h at 450°C . Lipids were extracted from approximately 100 mg dry tissue (ground to a fine powder right before extraction, using a mortar and pestle) by a modified Bligh and Dyer protocol (Boschker et al., 1998), whereby water in the first extraction step was replaced by Sørensen phosphate buffer pH 7.4 (White et al., 1979). Lipid extracts were fractionated on silicic acid columns into different polarity classes by sequential elution with chloroform, acetone and finally methanol. The phospholipids collected in the methanol fraction were derivatized using mild-alkaline methanolysis (using a methanolic KOH solution) to yield fatty acid methyl esters (hereafter referred to as PLFA since they were derived from the phospholipid fraction).

Total fatty acid methyl esters (tFA) were obtained from 25 mg tissue powder by direct acid methanolysis catalysed for 2 h at 90°C (modified from Lewis et al., 2000) in 3 mL of a fresh solution of methanol/hydrochloric acid 37%/chloroform (10:1:1 volume). After cooling down to room temperature chloroform (1 mL) and water (1 mL) were added to the samples which were then well mixed before the biphasic system was allowed to separate. The chloroform fraction was transferred to a fresh tube and the upper phase was re-extracted with fresh chloroform (1 mL). Chloroform extracts

BGD

7, 3453–3475, 2010

^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



were pooled, cleaned with 2 mL water, dried under a mild nitrogen flow and the tFA were re-suspended in 300 μ L hexane.

2.4 FAME fractionation on Ag^+ -SPE cartridges

PLFA and tFA fractions were resolved further as a function of degree of unsaturation (saturated, *trans*-mono-unsaturated, *cis*-mono-unsaturated, di-unsaturated and poly-unsaturated), using silver ion solid phase extraction columns (Ag^+ -SPE cartridges, Supelco Inc., Bellefonte, PA). The protocol used by Kramer et al. (2008) was modified to recover poly-unsaturated fatty acids (PUFAs) using 5 mL acetone/acetonitrile (60:40 volume), directly after eluting the di-unsaturated fraction.

2.5 FAME ^{13}C content analyses and identification

The individual FAME isotopic compositions were analysed by gas chromatography-combustion-isotope ratio mass spectrometry (GC-c-IRMS) using an HP6890 coupled to a Thermo Finnigan delta+XL via a GC/C III interface. The FAME mix and Ag^+ -SPE fractions were resolved on a fused-silica capillary column (100 m \times 0.25 mm) coated with 0.20 μm CP-Sil 88 (100% cyanopropyl polysiloxane, Varian BV, The Netherlands) after injection in splitless mode at 270 $^{\circ}\text{C}$. Helium was used as a carrier gas at a flow rate of 1 mL min^{-1} and the following temperature program was applied: 4 min at 45 $^{\circ}\text{C}$; an increase at 10 $^{\circ}\text{C min}^{-1}$ to 135 $^{\circ}\text{C}$ and a plateau of 90 min; an increase at 5 $^{\circ}\text{C min}^{-1}$ to 170 $^{\circ}\text{C}$ and a plateau of 25 min; an increase at 10 $^{\circ}\text{C min}^{-1}$ to 195 $^{\circ}\text{C}$ and a plateau of 15 min; an increase at 10 $^{\circ}\text{C min}^{-1}$ to 235 $^{\circ}\text{C}$ and a plateau of 5 min. A careful selection of the type of chromatographic column and the optimization of the temperature program (enabling the detection of FAME 10:0 to 24:0) ensured baseline resolution of the IRMS m/z 44 trace for the majority of the components (including 16:1 and 18:1 positional isomers) and enabled an accurate identification of labeled PLFA. The addition of one methanol carbon per fatty acid molecule during the formation of FAME was corrected as in Abrajano et al. (1994) using the mass balance equation taking a measured

BGD

7, 3453–3475, 2010

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$\delta^{13}\text{C}_{\text{CH}_3\text{OH}}$ value of -40.3‰ :

$$\delta^{13}\text{C}_{\text{FAME}} = [x]\delta^{13}\text{C}_{\text{FA}} + [1 - x]\delta^{13}\text{C}_{\text{CH}_3\text{OH}} \quad (1)$$

Where x is the fractional carbon contribution of the free fatty acid to the methyl ester – e.g.: 18/19 for FA 18:1(n-7).

The weighed $\delta^{13}\text{C}$ signature of fatty acid methyl esters, grouped according to their degree of unsaturation (Fig. 1), was obtained using the following formula (with mono-unsaturated fatty acids – MUFA – taken as an example of fatty acid class):

$$\delta^{13}\text{C}_{\text{MUFA}} = (\% \text{MUFA}_1 * \delta^{13}\text{C}_{\text{MUFA}_1} + \dots + \% \text{MUFA}_z * \delta^{13}\text{C}_{\text{MUFA}_z}) / (\sum \% \text{MUFA}_{1 \text{ to } z}) \quad (2)$$

where $\% \text{MUFA}_x$ is the area obtained from MUFA_x on the chromatogram relative to the sum of the areas obtained from all fatty acids, and $\delta^{13}\text{C}_{\text{MUFA}_x}$ is the carbon isotopic signature of MUFA_x .

FAME identification was achieved by GC-mass spectrometry (GC-MS) using a Thermo Finnigan TRACE GC-MS system, applying the same GC conditions as described for the GC-IRMS analyses. The mass spectrometer was operated under mild conditions of electron impact ionization (EI^+ ; 40 eV) and recorded the mass spectra in the scan mode ($m/z=50-400$). Aliquots of the PLFA or tFA mixture and of the MUFA, DUFA and PUFA fractions were derivatised further by a one step reaction into dimethylloxazolines (DMOX) to locate the unsaturations by GC-MS (Fay and Richly, 1991). We used a temperature gradient similar to the one used for FAME analyses, but since DMOX derivatives are less volatile (see also Precht and Molkenin, 2000) we increased the temperature of the isotherm plateaus at 135°C , 170°C and 195°C for the temperature program described above, by 10°C to 145°C , 180°C and 205°C , respectively. The International Union of Pure and Applied Chemistry PLFA nomenclature used here is described by Guckert et al. (1985): the (n-) notation indicates the position of the carbon-carbon double bond in the FA aliphatic chain starting from the methyl end carbon. The Δ notation, giving the location of FA unsaturation from the carbon at the carboxyl end, is only used in the next paragraphs to discuss enzymatic activities.

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Methane- and sulfide-oxidizing bacteria PLFA fingerprint

Phenotypic characterization is important for accurate taxonomic description of the endosymbionts (Bowman et al., 1993) and interpretation of the compound specific natural stable isotopic signatures. Pond et al. (1998) observed that the 14:0, 18:0, 18:1(n-7), 20:1(n-7), 18:3(n-7), and 20:3(n-7) compounds in the total lipid extract from *Bathymodiolus azoricus* gills were isotopically light and concluded these compounds had been produced by the sulfide-oxidising endosymbiotic bacteria (SOX). Our experimental results contrast with these conclusions and indicate that (i) 14:0, 18:0, 18:1(n-7) and 20:1(n-7) compounds also incorporate C from $^{13}\text{CH}_4$ and (ii) 18:0, 20:1(n-7), 18:3(n-7) and 20:3(n-7) compounds do not incorporate significant amounts of ^{13}C from $\text{H}^{13}\text{CO}_3^-$ in the presence of H_2S .

Our study shows that mussels kept in the presence of $^{13}\text{CH}_4$ or $\text{H}^{13}\text{CO}_3^- + \text{H}_2\text{S}$ incorporate ^{13}C mainly in saturated and mono-unsaturated PLFA (Sat and MUFA, respectively, Fig. 1a). For instance, the weighed average $\delta^{13}\text{C}$ signature of ΣMUFA from the gill PLFA fraction of mussels incubated with $^{13}\text{CH}_4$ varied between +27 and +249‰ (Fig. 1a). From the $^{13}\text{CH}_4$ and $\text{H}^{13}\text{CO}_3^- + \text{H}_2\text{S}$ experiments, three patterns can be distinguished (Fig. 2): (i) PLFA enriched in ^{13}C from both substrates (CH_4 and $\text{HCO}_3^- + \text{H}_2\text{S}$), i.e. 14:0; 15:0; 16:0; 16:1(n-7)*c+t* and 18:1(n-7)*c+t*; (ii) PLFA enriched in ^{13}C only after exposure to $^{13}\text{C}-\text{CH}_4$, i.e., 17:0; 18:0; 16:1(n-9); 16:1(n-8) and (n-6); 18:1(n-13)*c+t* and (n-8); 20:1(n-7) and 18:2(n-7) (note that fatty acid 18:1(n-9) was enriched in only one of the replicate specimens) and (iii) PLFA enriched in ^{13}C only after exposure to $\text{H}^{13}\text{CO}_3^-$ and H_2S : i.e., 12:0. Exposure to $\text{H}^{13}\text{CO}_3^-$ and H_2S also resulted in 17:1(n-8), 20:2(n-9,15) and 20:5(n-3) enrichment in two, and 20:2(n-7,11) in only one out of the three specimens analyzed.

Phylogenetic analyses revealed that the *B. azoricus* methane-oxidising endosymbiont (MOX) is a gamma-proteobacterium related to free living Type I MOX and to MOX

BGD

7, 3453–3475, 2010

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 symbionts from other Bathymodiolids (Duperron et al., 2006; Nakagawa and Takai, 2008; Spiridonova et al., 2006). Although Colaço et al. (2007) did not identify any MOX PLFA biomarkers in wild *B. azoricus* specimens from the Menez Gwen site they did reveal the presence of large amounts of i19:0, 18:1(n-13) and 18:1(n-9), which were attributed to the endosymbionts. Our tracer experiment with $^{13}\text{CH}_4$ revealed that 18:1(n-13) was indeed slightly labeled but this was not the case for 18:1(n-9) (not enriched in the presence of $\text{H}^{13}\text{CO}_3^- + \text{H}_2\text{S}$ either) nor for i19:0 which was not detected in our specimens.

10 As mentioned above, we found that $^{13}\text{CH}_4$ incorporated in *B. azoricus* gill tissue ended up in 14:0, 15:0, 16:0 as well as in *cis* and *trans* 16:1(n-9, 8, 7 and 6) PLFA (Fig. 2). 16:1(n-9) (one of the least abundant 16:1 isomers) also became labeled, although this compound has not been observed previously in PLFA profiles of isolated methanotrophic strains (Bodelier et al., 2009). Overall, labeled PLFA in mussel gill tissue from the $^{13}\text{CH}_4$ experiment represent 95% of the *Methylosphaera hansonii*'s total PLFA content (Table 1; Bodelier et al., 2009). This result is consistent with the study of Pimenov et al. (2002) who could amplify total DNA isolated from *B. azoricus* gill tissue exclusively with 16SrRNA primers targeting the genera *Methylobacter* and *Methylosphaera*.

20 Phylogenetically the *B. azoricus* SOX endosymbiont stands far from any known cultured SOX bacteria. The closest strains such as the obligate chemolithoautotrophic thiodenitrifying gamma-proteobacteria *Thiohalomonas nitratireducens* (Sorokin et al., 2007) only reach around 88% 16S rRNA sequence homology (S. Duperron, personal communication, 2009). Furthermore, the PLFA labeled in our experiments represented only 69% of the fatty acid (FA) extracted from *T. nitratireducens*. It is thus difficult to relate the phenotype of *B. azoricus* SOX symbiont to any free-living bacteria.

25 Conway and MacDowell Capuzzo (1991) suggested that all SOX bacteria desaturate their FA via the anaerobic pathway. The anaerobic desaturation pathway produces long chain MUFA by the elongation of medium chain length *cis*-3-unsaturated intermediates, with 18:1(n-7) as the major end-product and 16:1(n-7) as a secondary product. The

BGD

7, 3453–3475, 2010

^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

O₂-dependent MUFA synthesis pathway is known to produce a large variety of fatty acids (FA) with a double bond insertion occurring mainly in the Δ9 position (Conway and MacDowell Capuzzo, 1991). Note that the delta (Δ) nomenclature which imposes carbon atoms numbering to start from the carboxylic acid end of the acyl chain, is used for describing biochemical reactions. Our results show that 18:1(n-7) and 16:1(n-7), which are the main products of the anaerobic desaturation pathway, were significantly more labeled (−19.8 and +182.2‰, respectively) than the main product of aerobic desaturation, Δ9 MUFA 18:1(n-9) (−29.6‰, Fig. 2). However, the latter fatty acid was less abundant, and therefore incorporation of similar levels of label in the 3 fatty acids could have resulted in a higher δ¹³C signature in PLFA 18:1(n-9). The fact that MUFA 16:1(n-7), which was approximately 3 and 9 fold more abundant than 18:1(n-7) and 18:1(n-9), respectively, was the most labeled compound together with saturated PLFA (12:0, 14:0 and 15:0; see Fig. 2), supports the contention of Conway and MacDowell Capuzzo that *B. azoricus* SOX symbiont desaturates its FA via the anaerobic pathway.

3.2 Amino acid assimilation in tFA

Wright (1982) reports that *Mytilus* and *Modiolus* mussels take up free amino acids dissolved in seawater. Once inside the cell, free amino acids can be incorporated unchanged into proteins (Eccleston and Kelly, 1972) or used for the synthesis of other macromolecules. Excess amino acids (not being incorporated into proteins) cannot be stored in the cells and are generally used as metabolic fuel for the production of FA, ketone bodies or glucose (Berg et al., 2002). The ketogenic amino acids leucine, lysine, isoleucine, phenylalanine, tryptophan and tyrosine (accounting for 41 to 72% of the ¹³C-amino acid mix used in our tracer experiment) can be degraded into acetoacetyl CoenzymeA or acetyl-CoA, which constitutes the basis for de novo FA biosynthesis (e.g., Payne et al., 2001). Alanine, serine and cysteine can be converted to pyruvate which can further turn into acetyl CoA. Tracing the incorporation of amino acid ¹³C into FA biomarkers may thus provide evidence for an osmotrophic activity (feeding on dissolved organics) by *B. azoricus* itself and/or by its endosymbionts. While from the

¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



present study (Riou et al., 2008) it is now clear that *B. azoricus*' symbionts assimilate CH_4 or HCO_3^- in the presence of H_2S , their potential for osmotrophy has never been studied. The fact that gill tissue is the first one showing incorporation of ^{13}C from free amino acids (Riou et al., 2010) could be interpreted as indicating amino acid assimilation by the symbionts hosted in this tissue. However, mussel epithelial cells are the first to access this potential carbon and nitrogen source, since amino acids first enter the gill epithelial cells before they can reach MOX and SOX bacterial cytoplasm.

The experiment with ^{13}C labeled amino acids resulted in significant enrichment of the majority of the total fatty acids (tFA, Fig. 3). Out of the 38 gill tissue tFA only 14 did not show significant incorporation of ^{13}C from the amino acids. The most enriched fatty acid was 12:0 (representing 2% of tFA), followed by 16:1(n-8) (0.3%), 18:1(n-7) (2.7%) and 18:2(n-7) (0.8% of tFA). The unlabelled tFA were the MUFA 16:1(n-9) and 18:1(n-8) (1.6% of the tFA area), 19:1(n-7) (0.5%), and 20:1(n-13) (4.4%); the non-methylene interrupted di-unsaturated (NMID) 18:2(n-7,13) (0.4%), 20:2(n-9,15) (4.7%) and (n-7,15) (6.2%); and the PUFA 18:3(n-7) (11.5%), 18:3(n-5,10,13) (5.3%), 20:3(n-7) (2.3%) and 22:3 isomers (1.5%).

Mollusc lipids contain non-methylene interrupted di-unsaturated (NMID) FA having unusual unsaturation features, since they possess double bonds with more than one methylene group between ethylenic bonds (reviewed in Barnathan, 2009). We observed that the incorporation of ^{13}C from labeled amino acids in gill tissue di-unsaturated tFA ("DUFA" with these being mainly composed of NMI FA; $\delta^{13}\text{C}$ DUFA = -20.9‰) exceeded significantly ^{13}C incorporation from CH_4 or $\text{HCO}_3^- + \text{H}_2\text{S}$ (-32.8 and -35.6‰ , respectively, Fig. 1a and b). In particular, the NMID 22:2(n-7,15) became labeled only when supplying *B. azoricus* with ^{13}C -amino acids (Fig. 3). In bivalves, NMID 22:2(n-9,15) and 22:2(n-7,15) (usually the most abundant in marine invertebrates, see e.g., Dunstan et al., 1993; Kawashima et al., 2003) are synthesized de novo by elongation and $\Delta 5$ desaturation of 18:1n-9 and 16:1n-7 (Zhukova, 1986) and subsequent elongation of 20:2(n-9,15) and 20:2(n-7,15) into the 22:2(n-9,15) and 22:2(n-7,15) acids. The incorporation of ^{13}C label in 22:2(n-7,15) therefore indicates

BGD

7, 3453–3475, 2010

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

that the mollusc host cells have assimilated the amino acids directly in their fatty acids (as opposed to CH_4 and HCO_3^- which are being assimilated by the symbionts). Further evidence for this comes from the observation that MUFA 18:1(n-9) as well as 20:1(n-9) were more enriched in gills of specimens incubated with ^{13}C -amino acids (Fig. 3) than in mussels kept in the presence of $^{13}\text{CH}_4$ or $^{13}\text{CO}_2 + \text{H}_2\text{S}$ (Fig. 2), even though $\sum\text{MUFA}$ in the amino acid experiment was less labeled (+39%, Fig. 1b) than $\sum\text{MUFA}$ for the $^{13}\text{CH}_4$ or $^{13}\text{CO}_2 + \text{H}_2\text{S}$ experiments (+168 and +46, respectively, Fig. 1a). In animal cells, mono-unsaturation of fatty acids is catalyzed by a $\Delta 9$ desaturase which has a maximum affinity for FA 18:0 and yields 18:1 $\Delta 9$ (18:1(n-9)) (Cook, 1996). Since filter-feeding molluscs elongate de novo produced 18:1(n-9) to 20:1(n-9) (Zhukova, 1986), the observed preferential incorporation of ^{13}C from amino acids in 18:1(n-9) as well as 20:1(n-9), indeed points to direct assimilation of amino acids by the Mytilid cell machinery.

As can be noticed from Fig. 3, the FA of the (n-7) series were particularly enriched, with 18:1(n-7) showing the highest labeling followed by 18:2(n-7) and 20:1(n-7). One of the pathways of 18:1(n-7) synthesis is the anaerobic pathway of desaturation, and Tunlid and White (1992) proposed 18:1(n-7) to be a general indicator of the presence of bacteria. However, the highest enrichments from ^{13}C -amino acids observed for 18:1(n-7), 18:2(n-7) and 20:1(n-7) could also indicate preferential use of the labeled acetyl coA resulting from amino acid breakdown to elongate 16:1(n-7), which is one of the most abundant FA (around 10%) in gill tissue. Indeed, had the ^{13}C -amino acids been assimilated by the SOX endosymbionts first, 16:1(n-7) would have ended up being much more labeled than 18:1(n-7), as was observed in the $\text{H}^{13}\text{CO}_3^- + \text{H}_2\text{S}$ experiment. Although it is true that 16:1(n-7) and 18:1(n-7) are the most abundant FA in SOX bacteria (McCaffrey et al., 1989) and that 18:1(n-7) dominates the PLFA profiles of facultative methanotrophic alpha-proteobacteria *Methylocella* species (Dedysh et al., 2005), these particular FA are also biosynthesized by eukaryotic organisms. The elongation of 16:1(n-7) to form 18:1(n-7) by the incorporation of enriched acetyl coA resulting from amino acid degradation might thus be performed by the mussel itself.

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Bacteria possess effective metabolic switch-on/switch-off mechanisms to survive long periods of low food supply and to react rapidly to available suitable nutrients (including low concentrations of dissolved organic matter; Sepers, 1977). The SOX *Beggiatoa* bacteria, for instance, can grow facultatively or mixotrophically on inorganic and soluble organic compounds (Zhang et al., 2005). However, in general, organic compounds do not stimulate the growth of specialist phototrophs, lithotrophs, or methylotrophs. Although some of the isolated bacteria strains using methane and other C₁ compounds also grow on sugars and acids (Patt et al., 1974), many isolates capable of growth on methane have proved incapable of growth on conventional organic media (Whittenbury et al., 1970). Several factors may explain the lack of capacity to assimilate organic compounds such as an inhibition effect by these compounds (although balanced mixtures of amino acids can cancel the inhibition by one or the other amino acid; Smith and Hoare, 1977), the impermeability of the cell (e.g. Eccleston and Kelly, 1972) or the loss of enzymes from the main assimilation pathways (Theisen et al., 2005; Smith and Hoare, 1977). In obligate type I MOX bacteria, the chemical conversion of carbohydrates, lipids and proteins into CO₂ and H₂O for energy production is blocked at the level of the tricarboxylic acid (TCA) cycle, due to the concurrent absence of one of its key enzymes, α -ketoglutarate dehydrogenase, and of isocitrate lyase and malate synthase from the glyoxylate shunt (Trotsenko, 1983). In some cases, succinyl CoA synthetase, another enzyme from the TCA cycle, is also missing, like in *Thiobacillus denitrificans* or *M. capsulatus* (Smith and Hoare, 1977). Our finding that some of the MOX (16:1(n-8) and (n-6)) and SOX (12:0) biomarkers were also labeled after incubation with ¹³C-amino acids might indicate that the symbionts had access to acetyl coA (or acetate) produced during the degradation of the amino acids by the host. Further experiments are needed to ascertain these hypotheses.

4 Conclusions

PLFA and total FA ^{13}C -fingerprinting in *Bathymodiolus azoricus* gill tissues allowed us to trace the assimilation of ^{13}C -enriched HCO_3^- (in the presence of H_2S) and CH_4 into the endosymbionts and host. Based on the selective labeling pattern of PLFA in each of the experimental treatments we could establish qualitative FA profiles of MOX and SOX endosymbiotic bacteria living in association with *B. azoricus*. Additionally, carbon from dissolved free amino acids was found to be incorporated into host specific FA and also into some of the symbiont biomarkers.

The tracer uptake experiments in the present study could only be performed under conditions of atmospheric pressure. It therefore needs to be verified through future experiments whether or not the activated metabolic pathways are the same under high, in situ pressure conditions and atmospheric pressure. While more experimentation is needed to better understand the physiology of the endosymbionts, this experiment has shown its usefulness for positioning the endosymbionts among described strains, as well as for the direct identification of the symbiont biomarkers, for which until now only assumptions had been made.

Acknowledgements. We thank especially C. Leal, M. Laranjo, L. Pires, V. Costa, R. Betten-court, A. J. Cavagna and M. Korntheuer for assistance during experimental and analytical work. Our thanks also go to S. Duperron for his expert advise and to the Victor 6000 team, the crews of “R/V Arquipélago” and “Pourquoi Pas?” and chief scientists of the MoMARETO cruise J. Sarrazin and P. M. Sarradin who were in charge of the cage deployments and recovery. Financial support for this research was provided by the EU project MoMARNET-FP6-RTN/2003/505026 and the Research Council of Vrije Universiteit Brussel. The EU Framework Contract No EVK3-CT1999-00003 (VENTOX) funded the cage system and partially LabHorta with the project DRCT M2.1.1/I/008/2005 – LabHorta – Automation and Up-Grading of LabHorta – Technology for the exploration of marine molecules of deep-sea organisms (PRODESA). IMAR-DOP/UAç research activities are additionally supported through the pluri-annual and programmatic funding schemes of FCT (Portugal) and Azorean Regional Directorate for Science and Technology (DRCT, Azores, Portugal) as Research Unit #531 and Associate Laboratory #9.

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Abrajano, T. A., Murphy, D. E., Fang, J., Comet, P., and Brooks, J. M.: $^{13}\text{C}/^{12}\text{C}$ ratios in individual fatty acids of marine mytilids with and without bacterial symbionts, *Org. Geochem.*, 21, 611–617, 1994.
- 5 Barnathan, G.: Non-methylene-interrupted fatty acids from marine invertebrates: Occurrence, characterization and biological properties, *Biochimie*, 91, 671–678, 2009.
- Berg, J. M., Tymoczko, J. L., and Stryer, L.: Carbon atoms of degraded amino acids emerge as major metabolic intermediates, in: *Biochemistry*, W. H. Freeman and Co., New York, 2002.
- 10 Bodelier, P. L. E., Bär Gillisen, M. J., Hordijk, K., Sinninghe Damste, J. S., Rijpstra, W. I. C., Geenevasen, J. A. J., and Dunfield, P. F.: A reanalysis of phospholipid fatty acids as ecological biomarkers for methanotrophic bacteria, *Int. Soc. Microb. Ecol. J.*, 3, 606–617, 2009.
- Boschker, H. T. S., Nold, S. C., Wellsbury, P., Bos, D., de Graaf, W., Pel, R., Parkes, R. J., and Cappenberg, T. E.: Direct linking of microbial populations to specific biogeochemical processes by ^{13}C -labeling of biomarkers, *Nature*, 392, 801–805, 1998.
- 15 Bowman, J. P., Sly, L. I., Nichols, P. D., and Hayward, A. C.: Revised taxonomy of the methanotrophs: description of *Methylobacter gen. nov.*, emendation of *Methylococcus*, validation of *Methylosinus* and *Methylocystis* species, and a proposal that the family *Methylococcaceae* includes only the group I methanotrophs, *Int. J. Syst. Bacteriol.*, 43, 735–753, 1993.
- 20 Colaço, A. and Santos, R. S.: LABHORTA, A Land-based laboratory for vent studies, in: Towards Planning of Sea-floor Observatory Programs for the MAR region, Proceedings of the II MoMAR Workshop, Arquipélago, Life and Marine Sciences, Supplement 3, XII, edited by: Santos, R. S., Escartin, J., Colaço, A. and Adamczewska, A., 36 p., 2002b.
- Colaço, A., Desbruyères, D., and Guezennec, J.: Polar lipid fatty acids as indicators of trophic associations in a deep-sea vent system community, *Mar. Ecol.*, 28, 15–24, 2007.
- 25 Conway, N., and McDowell, J.: Capuzzo: Incorporation and utilization of bacterial lipids in the *Solemya velum* symbiosis, *Mar. Biol.*, 108, 277–291, 1991.
- Cook, H. W.: Fatty acid desaturation and chain elongation in eukaryotes, in: *Biochemistry of lipids, lipoproteins and membranes*, edited by: Vance, D. E. and Vance, J. E., Elsevier Science, Amsterdam, 129–152, 1996.
- 30 Dedysh, S. N., Knief, C., and Dunfield, P. F.: *Methylocella* species are facultatively methanotrophic, *J. Bacteriol.*, 187(13), 4665–4670, 2005.

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lewis, T., Nichols, P. D., and McMeekin, T. A.: Evaluation of extraction methods for recovery of fatty acids from lipid-producing microheterotrophs, *J. Microbiol. Meth.*, 43, 107–116, 2000.
- McCaffrey, M. A., Farrington, J. W., and Repeta, D. J.: Geochemical implications of the lipid composition of *Thioploca spp.* from the Peru upwelling region –15° S, *Org. Geochem.*, 14, 61–68, 1989.
- 5 Nakagawa, S. and Takai, K.: Deep-sea vent chemoautotrophs: diversity, biochemistry and ecological significance, *FEMS Microbiol. Lett.*, 65, 1–14, 2008.
- Nichols, P. D., Smith, G. A., Antworth, C. P., Hanson, R. S., and White, D. C.: Phospholipid and lipopolysaccharide normal and hydroxy fatty acids as potential signatures for methane-oxidising bacteria, *FEMS Microbiol. Lett.*, 31, 327–335, 1985.
- 10 Patt, T. E., Cole, G. C., Bland, J., and Hanson, R. S.: Isolation and characterization of bacteria that grow on methane and organic compounds as sole sources of carbon and energy, *J. Bacteriol.*, 120, 955–964, 1974.
- Payne, D. J., Warren, P. V., Holmes, D. J., Ji, Y., and Lonsdale, J. T.: Bacterial fatty-acid biosynthesis: a genomics-driven target for antibacterial drug discovery, *Drug Discov. Today*, 6(10), 537–544, 2001.
- 15 Perner, M., Bach, W., Hentscher, M., Koschinsky, A., Garbe-Schönberg, D., Streit, W. R., and Strauss, H.: Short-term microbial and physico-chemical variability in low-temperature hydrothermal fluids near 5° S on the Mid-Atlantic Ridge, *Environ. Microbiol.*, 11(10), 2526–2541, 2009.
- 20 Pimenov, N. V., Kaliuzhnaia, M. G., Khmelenina, V. N., Mitiushina, L. L., and Trotsenko, I. A.: Utilization of methane and carbon dioxide by symbiotrophic bacteria in gills of Mytilidae (*Bathymodiolus*) from the Rainbow and Logachev hydrothermal fields on the Mid-Atlantic Ridge, *Microbiology*, 71(5), 587–594, 2002.
- 25 Pond, D. W., Bell, M. V., Dixon, D. R., Fallick, A. E., and Sargent, J. R.: Stable-Carbon-Isotope composition of fatty acids in hydrothermal vent mussels containing methanotrophic and thiotrophic bacterial endosymbionts, *Appl. Environ. Microbiol.*, 64, 370–375, 1998.
- Precht, D. and Molkenin, J.: Identification and quantitation of *cis/trans* C16:1 and C17:1 fatty acid positional isomers in German human milk lipids by thin-layer chromatography and gas chromatography/mass spectrometry, *Eur. J. Lipid Sci. Tech.*, 2000, 102–113, 2000.
- 30

Won, Y., Hallam, S. J., O'Mullan, G. D., and Vrijenhoek, R. C.: Cytonuclear disequilibrium in a hybrid zone involving deep-sea hydrothermal vent mussels of the genus *Bathymodiolus*, *Mol. Ecol.*, 12, 3185–3190, 2003.

5 Wright, S.: A Nutritional Role for Amino Acid Transport in Filter-Feeding Marine Invertebrates, *Am. Zool.*, 22, 621–634, 1982.

Zhang, C. L., Huang, Z., Cantu, J., Pancost, R. D., Brigmon, R. L., Lyons, T. W., and Sassen, R.: Lipid Biomarkers and Carbon Isotope Signatures of a Microbial (*Beggiatoa*) Mat Associated with Gas Hydrates in the Gulf of Mexico, *Appl. Environ. Microbiol.*, 71(4), 2106–2112, 2005.

10 Zhukova, N. V.: Biosynthesis of Non-methylene-interrupted dienoic Fatty Acids from [¹⁴C] Acetate in Molluscs, *Biochim. Biophys. Acta*, 878, 131–133, 1986.

BGD

7, 3453–3475, 2010

**Fatty acid
¹³C-fingerprinting in
Mytilid-bacteria
symbiosis**

V. Riou et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fatty acid
¹³C-fingerprinting in
 Mytilid-bacteria
 symbiosis**

V. Riou et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Table 1. PLFA displaying significant enrichment after the tracer experiments with H₂S+ ¹³CO₂ (SOX) or with ¹³CH₄ (MOX). X=¹³C labeled. % PLFA = proportion of the PLFA content in cultured strains of *Methylosphaera hansonii* and *Thiohalomonas nitratireducens*.

	MOX	<i>M. hansonii</i> (% PLFA) Bodelier et al., 2009	SOX	<i>T. nitratireducens</i> (% PLFA) Sorokin et al., 2007
12:0			X	3.9
14:0	X	2.5	X	0.5
15:0	X	1.5	X	0.6
16:0	X	14.5	X	25.4
18:0	X	0	X	0.5
16:1(n-6)	X	17.5		
16:1(n-7) <i>c+t</i>	X	20	X	27.3
16:1(n-8)	X	39		
16:1(n-9)	X	0		
17:1(n-8)			X	0.9
18:1(n-7) <i>c+t</i>	X	1.5	X	11.8
18:1(n-8)	X	0		
18:1(n-9)	X	0.5	X	5.1
18:1(n-13) <i>c+t</i>	X	0		
20:1(n-7)	X	0		
18:2(n-7)	X	0		



Fatty acid ¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.




Fig. 1. *Bathymodiolus azoricus* ($n=3$ specimens, mean \pm min-max) gill FAME carbon isotopic signatures. The $\delta^{13}\text{C}$ signatures were corrected for the methyl group added during the transesterification using formula (1), and weighed averages were obtained using formula (2) **(A)**: Gill PLFA after 15 days with $28\ \mu\text{M}$ 25% $^{13}\text{CH}_4$ (black bars) or 20 days with $6\ \mu\text{M}$ $\text{H}_2\text{S} + 16\%$ $\text{H}^{13}\text{CO}_3^-$ (grey bars). **(B)**: Gill total fatty acids (tFA) after 20 days in control conditions (striped bars) or 20 days with $300\ \mu\text{M}$ dissolved 98% ^{13}C -amino acids (white bars). Sat: saturated, MUFA: mono-unsaturated, DUFA: di-unsaturated, NMID: non-methylene interrupted di-unsaturated, PUFA: poly-unsaturated chains.

Fatty acid ^{13}C -fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.




Fig. 2. *Bathymodiolus azoricus* gill tissue PLFA ($n=3$, average \pm min-max) $\delta^{13}\text{C}$ signature after 15 days incubation with $^{13}\text{CH}_4$ (black), or 20 days incubation with $\text{H}^{13}\text{CO}_3^-$ in the presence of H_2S (grey).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Fatty acid ¹³C-fingerprinting in Mytilid-bacteria symbiosis

V. Riou et al.




Fig. 3. *Bathymodiolus azoricus* gill tissue tFA ($n=3$, average \pm min-max) $\delta^{13}\text{C}$ signature after 20 days incubation with ^{13}C -amino acids (white bars), relative to control mussels (striped bars).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion