

This discussion paper is/has been under review for the journal Biogeosciences (BG).  
Please refer to the corresponding final paper in BG if available.

# Tephrostratigraphy and tephrochronology of lakes Ohrid and Prespa, Balkans

R. Sulpizio<sup>1</sup>, G. Zanchetta<sup>2,3</sup>, M. D’Orazio<sup>2</sup>, H. Vogel<sup>4</sup>, and B. Wagner<sup>4</sup>

<sup>1</sup>CIRISIVU, c/o Dipartimento Geomineralogico, Bari, Italy

<sup>2</sup>Dipartimento di Scienze della Terra, Pisa, Italy

<sup>3</sup>INGV sez. Pisa; IGG-CNR, Pisa, Italy

<sup>4</sup>Institute for Geology and Mineralogy, University of Cologne, Köln, Germany

Received: 7 May 2010 – Accepted: 18 May 2010 – Published: 28 May 2010

Correspondence to: R. Sulpizio (r.sulpizio@geomin.uniba.it)

Published by Copernicus Publications on behalf of the European Geosciences Union.

3931

## Abstract

Four cores from Balkans lakes Ohrid and Prespa were studied for recognition of tephra layers and cryptotephra, and the results presented along with the review of data from other two already published cores from Lake Ohrid. The six cores provide a previously unrealised tephrostratigraphic framework of the two lakes, and supply the first detailed tephrochronologic profile (composite) for the Balkans, which spans from the end of the Middle Pleistocene to the end of the Ancient Age (AD 472). A total of 12 tephra layers and cryptotephra were recognised in the cores. One is of Middle Pleistocene age (131 ky) and correlated to the marine tephra layer P-11 from Pantelleria Island. Eight volcanic layers are Upper Pleistocene in age, and encompass the period between ca. 107 ky and ca. 31 ky. This interval contains some of the main regional volcanic markers of the Central Mediterranean area, including X-6, X-5, Y-5 and Y-3 tephra layers. The other layers of this interval have been related to the marine tephra layers C20, Y-6 and C10, while one was for the first time recognised in distal areas and correlated to the Taurano eruption of probable Vesuvian origin. Three cryptotephra were of Holocene age. Two of which have been correlated to Mercato and AD 472 eruptions of Somma-Vesuvius, while the third has been correlated to the FL eruption from Mount Etna. These recognitions provide a link of the Ohrid and Prespa lacustrine successions to other archives of the Central Mediterranean area, like South Adriatic, Ionian, and South Tyrrhenian Seas, lakes of Southern Italy (Lago Grande di Monticchio, Pantano di San Gregorio Magno and Lago di Pergusa) and Balkans (Lake Shkodra).

## 1 Introduction

Owing to the intense explosive volcanic activity that affected the Mediterranean over the last 200 ky (Vezzoli, 1988; Poli et al., 1987; Santacroce, 1987; Rosi and Sbrana, 1987; Keller et al., 1990; Orsi et al., 1996; Pappalardo et al., 1999; Di Vito et al., 2008; Santacroce et al., 2008) application of tephrochronology to volcanology, Quaternary

3932

science, paleoceanography, and archaeology has an exceedingly high potential in this area. In the last 30 y, Quaternary tephra layers have been extensively used to develop a high-resolution event stratigraphy for the late Pleistocene and Holocene across the Central and Eastern Mediterranean (Keller et al., 1978; Paterne et al., 1988, 1990; 5 Narcisi and Vezzoli, 1999; Wulf et al., 2004; Margari et al., 2007; Aksu et al., 2008; Giaccio et al., 2009; Zanchetta et al., 2010). While early work focused mainly on samples from marine cores (Keller et al., 1978; Paterne et al., 1988, 1990; Calanchi et al., 1998), recent work on terrestrial archives (including Italian, Greek, Turkish and Balkan lakes, and Bulgarian, Greeks and Italian cave sites; e.g., St Seymour and Christianis, 10 1995; Narcisi and Vezzoli, 1999; St Seymour et al., 2004; Wulf et al., 2004; Frisia et al., 2008; Giaccio et al., 2008; 2009; Sulpizio et al., 2010; Vogel et al., 2010) have considerably advanced the development of a long, high-resolution tephrostratigraphy, which will link marine and terrestrial records of Pleistocene-Holocene-age across the Mediterranean region and mainland Europe. Nevertheless, there are some areas in 15 the Central Mediterranean, in which volcanologic and Quaternary studies are still at the beginning, and the link to the general tephrostratigraphic Mediterranean network is lacking. The Balkans area across Macedonia, Albania and Montenegro is certainly one, in which tephrostratigraphic and tephrochronologic studies are in their infancy, although some studies on lacustrine settings indicate the area is extremely promising for tephrostratigraphic studies (Wagner et al., 2008; Caron et al., 2010; Lézine 20 et al., 2010; Sulpizio et al., 2010; Vogel et al., 2010). For these areas both tephrostratigraphic and tephrochronologic studies can offer invaluable stratigraphic support for sedimentologic, palaeoclimatic, paleoenvironmental, and volcanological studies. Here we present tephrostratigraphic and tephrochronologic data of lakes Ohrid and Prespa 25 (Fig. 1), which encompass the last 131 ky.

The paper reviews some already published data from the Macedonian side of Lake Ohrid, which are presented along with new data from both Ohrid and Prespa lakes. The recognised tephra layers supply a composite tephrostratigraphy of lakes Ohrid and Prespa, which has been correlated to other archives from the Balkans (Albanian side of

3933

lake Ohrid, Caron et al., 2010; Lézine et al., 2010; Lake Shkodra, Sulpizio et al., 2010), and linked to the regional tephrostratigraphic network of the Central Mediterranean area.

## 2 Study sites and analytical methods

Lake Ohrid (Fig. 1) is a transboundary lake shared by the Republics of Albania and 5 Macedonia. It is located at 693 m a.s.l., and surrounded by high mountain ranges reaching heights up to 2300 m. It is 30 km long, 15 km wide and covers an area of 358 km<sup>2</sup>. The lake basin shows a relatively simple tub-shaped morphology with a maximum water depth of 289 m, an average water depth of 151 m and a total volume of 50.7 km<sup>3</sup> (Popovska and Bonacci, 2007).

10 Lake Ohrid is mainly fed by inflow from karst springs (55%), while the remaining 45% of the hydrological input includes direct precipitation on the lake surface, river and direct surface runoff (Matzinger et al., 2006). The direct watershed of Lake Ohrid covers an area of 1002 km<sup>2</sup> (Popovska and Bonacci, 2007). Surface outflow (60%) through the River Crn Drim to the north and evaporation (40%) are the main hydrological outputs 15 (Matzinger et al., 2006). The average annual precipitation on the Lake Ohrid watershed is 907 mm (Popovska and Bonacci, 2007).

Lake Prespa is located 10 km to the east of Lake Ohrid, at an altitude of 849 m a.s.l. It is a transboundary lake shared between the Republics of Macedonia, Albania, and Greece. The surface area is 254 km<sup>2</sup>, with a catchment area of ca.1300 km<sup>2</sup>, a maximum 20 water depth of 48 m, and a volume of 3.6 km<sup>3</sup>. The total inflow is estimated to 16.9 m<sup>3</sup> s<sup>-1</sup>, with 56% originating from river runoff from numerous small streams, 35% from direct precipitation, and 9% from Lake Mikri Prespa to the south (Matzinger et al., 2006). Water loss derives through evaporation (52%), irrigation (2%) and outflow through karst aquifers (46%).

3934



cryptotephra numbering follows the origin of the core (OT for Ohrid Tephra and PT for Prespa Tephra), the year of core recovery (05-09) and the last two identification numbers of the respective core. Glass shards and micropumice fragments from the different cores were compositionally classified using the silica vs. total alkali diagram (TAS; Le Bas et al., 1986; Fig. 3).

### 3.1 Core Co1200

Core Co1200 contains two discrete tephra layers visible at naked eye. Tephra layer OT0700-1 (40–38 cm) appears light-brown to ochre in colour, and contains highly vesicular, aphyric micro-pumice and glass shards (Fig. 4a). Most of the glass shards show a trachytic composition with only few samples that plot close to the phonolitic field on the TAS diagram (Fig. 3a).

Tephra layer OT0700-2 (120.5–85.5 cm) has a light red colour, and contains aphyric, vesicular micro-pumices and aphyric glass shards with thick septa (Fig. 4b). Glass composition ranges from trachyte to phono-trachyte (Fig. 3a), with three different alkali ratios (Table 2).

### 3.2 Core Co1201

Core Co1201 contains seven discrete tephra layers visible at naked eye inspection and with different thicknesses.

The five tephra layers between 81–78 cm (OT0701-1), 90–89 cm (OT0701-2), at 100 cm (OT0701-3), and between 106–104 cm (OT0701-4) and 126–110 cm (OT0701-5) are light red to rusty-red in colour and comprise aphyric micro-pumice and glass shards. They show identical glass composition (Fig. 3b), and are grouped into three different compositional groups on the basis of the different alkali ratios (Table 2).

Tephra layer OT0701-6 (186–184 cm), appears pink in colour, and contains vesicular, aphyric micro-pumice fragments and glass shards (Fig. 4c). The glass composition is homogeneous phonolitic (Fig. 3b) with alkali ratio greater than 1.5 (Table 2).

3937

Tephra layer OT0701-7 (192–190 cm), is light-brown to red in colour, and contains both aphyric glass shards and poorly vesicular fragments with small pyroxene crystals in the groundmass (Fig. 4d). The glass composition continuously spans between phono-tephryte/shoshonite and phonolite/trachyte (Fig. 3b). Three main compositional groups can be identified on the basis of the alkali ratio and SiO<sub>2</sub> content (Table 2).

### 3.3 Core Co1202

Core Co1202 contains four visible tephra layers and six cryptotephra, which were already described in detail by Vogel et al. (2010a).

Cryptotephra OT0702-1 (77.5–74.5 cm) comprises dark-brown tachylitic fragments with crystalline groundmass (mainly sanidine, clinopyroxene and leucite; Fig. 4e). The composition straddles the foiditic and tephri-phonolitic fields (Fig. 3c).

Cryptotephra OT0702-2 (145.5–144 cm) consists of non-vesicular and blocky fragments with a porphyritic texture. Fragments exhibit mineral inclusions of plagioclase, clinopyroxene and olivine up to some tens of microns in size (Fig. 4f) and frequent occurrences of Fe-Ti oxides. The glass composition is mostly benmoreitic, with few analyses plotting in the trachytic and mugearitic fields on the TAS diagram (Fig. 3c).

Cryptotephra OT0702-3 (277.5–269 cm) comprises mainly aphyric, vesicular micropumice. The bubbles of the micropumices are mainly circular with thick septa (Fig. 4g). The glass composition of the cryptotephra is a fairly homogeneous Na-phonolite (Table 2 and Fig. 3c).

Tephra layer OT0702-4 (620–617 cm) is light brown in colour, and is characterised by normally graded middle to fine sand. Glass shards are mixed with lacustrine sediment up to 2 cm above the tephra layer. The tephra comprises highly vesicular, aphyric micro-pumice and cusped glass shards. Most of the glass shards show a trachytic composition with only few samples that plot close to the phonolitic field on the TAS diagram (Fig. 3c). Trace element distribution (sample 523; Table 3, Fig. 5) shows the less enriched pattern in the four analysed samples, with small negative anomalies in Ba

3938

and Sr and a marked enrichment in Pb (Fig. 5a). The rare earth element (REE) pattern shows a more or less regular decrease passing from light REE (LREE) to heavy REE (HREE; Fig. 5b).

5 Cryptotephra OT0702-5 (696–689 cm) comprises tachilitic particles with a crystal-rich groundmass containing acicular clinopyroxene, plagioclase and sanidine (Fig. 4h). The few glass compositions available range from latite to phonolite when plotted on the TAS diagram (Table 2 and Fig. 3c).

10 Tephra layer OT0702-6 (752–743 cm) is reddish-brown to light-brown in colour, and glass shards are mixed with lacustrine sediments in the overlying 10 cm. Volcanic particles mainly comprise aphyric, vesicular micro-pumices and aphyric glass shards with thick septa. Glass composition ranges from trachyte to phono-trachyte (Fig. 3c), with two different alkali ratio (Table 2). Trace element distribution (sample 565; Table 3, Fig. 5) shows an intermediate enrichment in the analysed samples, with a pronounced negative anomalies in Sr and Eu, moderate anomaly in Ba, and moderate positive anomalies in Th, U and Pb (Fig. 5a). The REE pattern shows a regular decrease in enrichment passing from LREE to HREE, with a pronounced negative Eu anomaly (Fig. 5b).

15 Cryptotephra OT0702-7 (825–822 cm) comprises aphyric, cusped glass shards with thin septa and glassy groundmass. Glass composition ranges from rhyolitic (main) to trachytic (Table 2 and Fig. 3c).

20 Tephra layer OT0702-8 (1146.5–1140 cm) is rusty-red in colour and comprises coarse ash and shows sharp basal and top contacts. Volcanic particles are aphyric micro-pumices and glass shards. Glass composition is mainly phonolitic (Table 2 and Fig. 3c).

25 Tephra layer OT0702-9 (1232.5–1229 cm) is light-brown in colour, and shows sharp basal and top boundaries. Grain size mainly comprises fine to coarse ash. Volcanic particles are highly vesicular, aphyric micro-pumices and glass shards with a large variability in shape and size (Fig. 4i). Glass composition is mainly trachytic, with few analyses that plot into the phonolitic field, and two different alkali ratios (Table 2 and

3939

Fig. 3c). The trace element (sample 1003; Fig. 5, Table 3) distribution shows the most enriched pattern in the four analysed samples, with marked negative anomalies in Ba, Sr and Eu that testify for a feldspar-dominated magma fractionation (Fig. 5a). The REE pattern shows a regular decrease from LREE to HREE, with a marked negative Eu anomaly (Fig. 5b).

5 Cryptotephra OT0702-10 (1447–1440 cm) comprises mainly aphyric cusped glass shards. When plotted on the TAS diagram the glass shards reveal a bimodal chemical composition, which comprises a trachyte and a rhyolite without any compositional trend between (Fig. 3c).

#### 10 3.4 Core Lz1120

Core Lz1120 contains two visible tephra layers and one cryptotephra, which were already described in detail by Wagner et al. (2008).

15 Cryptotephra OT0520-1 (315–310 cm) comprises dark brown, blocky fragments with few spherical or ovoid vesicles. The groundmass comprises small crystals of plagioclase and minor olivine. Single-shard analyses show dispersion between mugearitic and benmoreitic fields (Fig. 3d).

20 Tephra layer OT0520-2 (897–896 cm) comprises light coloured, elongated, highly vesicular fragments. Vesicles are mainly tubular and form channels throughout the entire length of the pyroclastic fragments. Groundmass is glassy, with very few elongated microcrystals of sanidine. Composition of single shards shows a limited variability within the trachytic field (Fig. 3d).

25 Tephra OT0520-3 (1075–1070 cm) is of unknown thickness, as the resistance of this layer prevented penetration of the coring equipment through it. Volcanic fragments comprise mainly light coloured, highly vesicular fragments, with minor dark coloured glass shards and micropumice fragments. In both typologies of volcanic fragments the vesicles are spherical or ovoidal, separated by thin, glassy sets. Groundmass is almost aphyric, even if small sanidine crystals sometimes occur on larger sets among

3940



bubbles. Single shard composition show a trend from the trachytic to the phonolitic fields (Fig. 3d), and can be arranged into three groups on the basis of different alkali ratios (Table 2).

### 3.5 Core Co1205

5 Core Co1204 was recovered from the north-western part of Lake Prespa (Fig. 1). It contains two discrete tephra layers and one cryptotephra.

Tephra layer PT0704-1 (672.5–667.5 cm) has a light grey colour, and contains aphyric, highly vesicular micro-pumice fragments and glass shards with glassy groundmass. The glass composition is homogeneously trachytic (Fig. 3e; Table 2).

10 Cryptotephra PT0704-2 (767.2–764.2 cm) was identified by high Sr count rates through high-resolution XRF-scanning, and contains tachilitic fragments with highly crystalline groundmass. Glass is rare and shows a compositional trend from shoshonites to trachytes (Fig. 3e). Three different compositional groups were identified on the basis of SiO<sub>2</sub>, CaO, and total alkali contents (Table 2).

15 Tephra layer PT0704-3 (879.3–863.3 cm) contains aphyric, highly vesicular micro-pumice and cusped to convolute glass shards with glassy groundmass. Glass composition straddles the phonolitic and trachytic fields (Fig. 3e), it can be split into two or three groups depending on the different alkali ratio (Table 2). The trace element distribution and the REE pattern (sample PR628; Table 3, Fig. 5) are identical to the  
20 sample 565 from core Co1202.

## 4 Discussion

### 4.1 Correlation to proximal deposits and other distal archives

The correlation of a distal tephra layer with proximal counterparts is a critical process, which in many cases implies the contemporaneous use of different data, such as glass

3941

and mineral composition, chronology, lithology, and stratigraphic position. This is because pyroclastic deposits from the same source, with few exceptions, show striking major element composition. Furthermore, pyroclastic deposits from different sources but originating from magmas with similar degree of evolution (e.g. trachytes and rhyolites) are hardly distinguishable on the basis of the sole major element composition. In  
5 this study, cores Lz1120 and Co1202, which contain correlated tephra layers (Wagner et al., 2008; Vogel et al., 2010), can be used as a reference, particularly because core Co1202 contains the largest number of tephra layers in all studied sediment cores from this region (Fig. 2).

10 The youngest volcanic deposit was correlated to the AD 472 (1478 cal. y BP) eruption of Somma-Vesuvius. It occurs as a cryptotephra in core Co1202 (OT0702-1), but has not been recognised in any of the other studied cores (Fig. 2).

The FL (3370±70 cal. y BP; Coltelli et al., 2000) cryptotephra occurs in cores Lz1120 (OT0520-1; Fig. 2; Wagner et al., 2008) and Co1202 (OT0702-2; Fig. 2; Vogel et al.,  
15 2010) from Lake Ohrid, but has not been found in the successions from lake Prespa.

The Mercato (8540±50 cal. y BP; Zanchetta et al., 2010) cryptotephra occurrence is limited to core Co1202 (OT0702-3; Fig. 2; Vogel et al., 2010) from Lake Ohrid.

Due to its thickness and unique, homogeneous trachytic composition (Table 2) the Y-3 tephra layer was recognized in cores Lz1120 (OT0520-2; Wagner et al., 2008),  
20 Co1200 (OT0700-1), and Co1202 (OT0702-4; Vogel et al., 2010) from Lake Ohrid and in core Co1204 (PT0704-1) from Lake Prespa (Fig. 2; Table 2). The ICP-MS analyses of trace elements on sample OT0702-4 (Table 3) reinforces the correlation to the proximal deposits of the SMP1-e eruption (30.67±0.23 cal. ky BP) from the Campi Flegrei caldera, which is indicated as the proximal counterpart of the Y-3 tephra layer  
25 (Di Vito et al., 2008; Zanchetta et al., 2008). The occurrence of the Y-3 tephra layer in four out of six cores (Fig. 2) indicates its usefulness as a stratigraphic marker for lakes Ohrid and Prespa.

The Codola ash (inferred age 33 cal. ky BP; Giaccio et al., 2008 or 34.2 cal. ky BP; Vogel et al., 2010) occurs as a cryptotephra in core Co1202 (OT0702-5; Fig. 2;

3942

Vogel et al., 2010). The Codola fragments have a highly microcrystalline groundmass and a glass composition that straddles the tephri-phonolitic/latitic and the phonolitic/trachytic (Fig. 6a; Di Vito et al., 2008; Giaccio et al., 2008; Santacroce et al., 2008). Based on their stratigraphic position and lithology the tephra layers PT0704-2 in core Co1204 and PT0916-1 in core Co1216 from Lake Prespa are good candidates for correlation to the Codola eruption. The inspection of glass composition illustrates a more complex situation, with some of the analyses from the PT0704-2 and the PT0916-1 samples that plot outside the Codola field (Fig. 6a). In particular, the PT0916-1 analyses define two compositional groups separated by a broad gap in SiO<sub>2</sub> and total alkali content (PT0916-1a and PT0916-1b; Fig. 6a; Table 2). Both groups plot outside the Codola field, either in TAS or SiO<sub>2</sub> vs. CaO diagrams (Fig. 5a, b), being more and less evolved, respectively. The less evolved analyses correlate well to the Taurano composition, which defines an evolutionary trend with the Codola samples in the SiO<sub>2</sub> vs. CaO diagram (Fig. 6b). The Taurano eruption was tentatively assigned to the activity of the Somma-Vesuvius volcano, and approximately dated 36–33 cal. ky BP (Di Vito et al., 2008), however this age range was subsequently changed to 30–33 cal. ky BP on the basis of the correlation to distal tephra layers in the Lago Grande di Monticchio succession (Sulpizio et al., 2008). Proximal deposits comprise porphyritic dark scoriae of shoshonitic/phono-tephritic composition, with a groundmass rich in microlites of clinopyroxene, sanidine and leucite (Di Vito et al., 2008). The evolved group (sample PT0916-1b; Table 2) plots outside the Taurano-Codola trend, and coincides with the Campanian Ignimbrite field in SiO<sub>2</sub> vs. CaO diagram (pinkish area of Fig. 5b). The lithology of the glass shards supports the correlation of this group to the Campanian Ignimbrite deposits, being formed by convolute and cusped shards with glassy groundmass. The occurrence of mixed populations of Taurano and Campanian Ignimbrite ash fragments is not surprising, due to the stratigraphic vicinity of the two eruptions (Di Vito et al., 2008). The analyses of cryptotephra PT0704-2 from core Co1204 can be divided into three groups, on the basis of different contents of SiO<sub>2</sub>, total alkali, and CaO (Fig. 5a, b; Table 2). Among them, only the PT0704-2b samples plot in the Codola

3943

field in both TAS and SiO<sub>2</sub> vs. CaO diagram (Fig. 5a, b). The two analyses of group PT0704-2c are the most evolved, and plot in the Campanian Ignimbrite field in the SiO<sub>2</sub> vs. CaO diagram (Fig. 6b). The PT0704-2a group have shoshonitic/latitic composition (Fig. 6a), and shows a separate compositional trend in the SiO<sub>2</sub> vs. CaO diagram. The correlation of this group with the regional tephrostratigraphy is at present not possible, since tephra layers or cryptotephra with similar lithology, glass composition and age have never been described in the Balkans, in Adriatic/Ionian marine cores, and continental Italy (e.g., Lago Grande di Monticchio succession; Wulf et al., 2004). Based on glass composition, a generic correlation to the activity of Vulcano Island between 53 and 21 ky (De Astis et al., 2000; 2006; Lucchi et al., 2008) is here proposed, although the link to a specific dated eruption is, to date, impossible.

The Campanian Ignimbrite/Y-5 (CI/Y-5; 39.28±0.11 ky; De Vivo et al., 2001) tephra layer has previously been recognized in cores Lz1120 (OT0520-3; Fig. 2; Wagner et al., 2008) and Co1202 (OT0702-6; Fig. 2; Vogel et al., 2010a) from Lake Ohrid. The glass composition of the OT0520-3 and the OT0702-6 tephra layers straddles the trachytic and phonolitic fields (Fig. 3c), and can be organised into two to three different compositional groups on the basis of alkali ratios (Table 2). These compositional groups reflect the involvement in the eruptive processes of three differently evolved trachytic magmas (i.e. from “evolved” to “primitive” trachytes; Fedele et al., 2008), which provides a geochemical fingerprint of the CI/Y-5 deposits also in distal reaches (Civetta et al., 1997; Pappalardo et al., 2002). Based on their lithology, stratigraphic position and major element glass composition the tephra layers found in cores Co1200 (OT0700-2; Fig. 2), Co1201 (OT0701-1, -2, -3, -4, -5; Fig. 2) from Lake Ohrid and in core Co1204 (PT0704-3; Fig. 2) from Lake Prespa can also be correlated to OT0520-3 and OT0702-6 and thus to the CI/Y-5 eruption. ICP-MS analyses of trace elements of samples OT0702-6 and PT0704-3 (Table 3; Fig. 5) reinforce this correlation. If we also consider the glass shards mixed into the PT0916-1 cryptotephra, the CI/Y-5 distal ash occurs in all studied cores from lakes Ohrid and Prespa (Fig. 2). The CI/Y-5 tephra layers is the lowermost volcanic deposit in most of the studied cores, with the only exceptions

3944

of cores Co1201 and Co1202 (Fig. 2).

Cryptotephra OT0702-7 occurs below the CI/Y-5 tephra layer in core Co1202, and was correlated to the Green Tuff eruption from Pantelleria island (Vogel et al., 2010), which corresponds to the Y-6 marine tephra layer (Keller et al., 1978).

5 The tephra OT0701-6 occurs below the CI/Y-5 in core Co 1201, and has a homogeneous trachytic composition (Fig. 3b; Table 2). It has no correspondence in the other cores from lakes Ohrid and Prespa, but can be correlated to the C20 marine tephra layer (dated at 79–80 ky; Paterne et al., 1988), and to the proximal deposits of SA3-b eruption from the Campanian area (Table 3).

10 The tephra layer OT0702-8 occurs in core Co1202 between 825 cm and 822 cm, and was correlated to the X-5 marine tephra layer ( $105 \pm 2$  ky; Keller et al., 1978; Kraml, 1997), which has been tentatively related to the TAU1-b eruption from Campi Flegrei caldera (Di Vito et al., 2008). Due to its chemical composition the OT0701-3 cryptotephra from core Co1201 can tentatively be correlated to the OT0702-8 tephra layer  
15 from core Co1202 and thus to the TAU1-b/X-5 eruption.

Tephra layer OT0702-9 in core Co1202 was correlated to the marine X-6 tephra layer (Keller et al., 1978), of generic Campanian origin. Brauer et al. (2007) quoted an  $^{40}\text{Ar}/^{39}\text{Ar}$  age of  $107 \pm 2$  ky for the X-6 tephra, which is in good agreement with the suggested age of 108.43 ky obtained from the varve-supported chronology of the Lago Grande di Monticchio record. The X-6 tephra layer has not been recognised in the  
20 other studied cores of lakes Ohrid and Prespa (Fig. 2). The evolved trace element composition of this tephra layer (Fig. 5; Table 3) can help in its discrimination from the mess of trachytic tephra layers recognised in the Upper Pleistocene successions.

Cryptotephra OT0702-10 (Fig. 2; Vogel et al., 2010a) in core Co1202 from Lake  
25 Ohrid represents the deepest volcanic deposit recognised in cores from lakes Ohrid and Prespa. It correlates to the marine P11 tephra layer (ca. 131 ka; Paterne et al., 1988, 2008), which sourced from Pantelleria Island.

3945

## 4.2 Composite tephrochronological record and regional correlations

The correlation of the 12 recognised tephra and cryptotephra layers with known proximal deposits allows the reconstruction of a composite stratigraphic and chronological framework for lakes Ohrid and Prespa during the past 131 ky (Fig. 7). Three cryptotephra  
5 occur during Holocene, with the Mercato layer marking the temperature maximum of the Early Holocene. The following seven tephra layers and cryptotephra punctuate the Upper Pleistocene, encompassing about 77 ky (from 30 to 107 ky; Fig. 7). Four markers cluster between 30 and 40 ky (i.e. between the Y-3 and the Y-5 tephra layers), detailing the stratigraphic succession between Heinrich events 3 and 4 (Ton-  
10 That et al., 2001; Zanchetta et al., 2008; Wagner et al., 2010). Particularly significant is the first recognition of the Taurano deposits in a very distal succession, which so far has never been described in nearby Adriatic and Ionian Sea cores. The Y-6 cryptotephra occurs below the CI, at about 49 ky (Fig. 7), linking the Balkans to the Ionian Sea archives. About 30 ky of sedimentation occur below the Y-6 without any tephra  
15 record, being the following tephra layer correlated to the SA3-a eruption and the C-20 marine tephra layer (79–80 ky; Paterne et al., 1988; Fig. 7). The last two tephra layers of the Upper Pleistocene are the X-5 and X-6, dated at 105 and 107 ky, respectively (Fig. 7). The oldest and deepest volcanic deposit recognised is the P-11 tephra layer, which anticipates the inception of the Last Interglacial in the Balkans (Lézine et al.,  
20 2010; Vogel et al., 2010b).

The recognised tephra layers and cryptotephra can be correlated to other published tephrostratigraphic records of the Central Mediterranean area, linking the Balkans with other archives at a regional scale (Fig. 8). Excluding the CI/Y-5 tephra layer, which extensively occurs in the Central Mediterranean area and extends through Aegean Sea and up to Russia (Pyle et al., 2006; Giaccio et al., 2008), the other recognised volcanic  
25 markers have various frequency of occurrence in the published archives (Fig. 8). The most common is the Y-3 tephra layer, which commonly occur in marine cores from South Adriatic (SA), Ionian, and South Tyrrhenian (ST) Seas, and in Lago Grande di

3946





- Italy: implications for volcanic activity in Peninsular Italy and tephrostratigraphy in the Central Mediterranean area, *Quaternary Sci. Rev.*, 28, 2710–2733, 2009.
- Jahns, S. and van den Boogard, C.: New palynological and tephrostratigraphical investigations of two salt lagoons on the island of Mljet, South Dalmatia, Croatia, *Veg. Hist. Archaeobot.*, 7, 219–234, 1998.
- 5 Keller, J., Ryan, W. B. F., Ninkovich, D., and Altherr, R.: Explosive volcanic activity in the Mediterranean over the past 200000 years as recorded in deep-sea sediments, *Geol. Soc. Am. Bull.*, 89, 591–604, 1978.
- Keller, J., Rehren, T., and Stadtbauer, E.: Explosive volcanism in the Hellenic Arc: a summary and review, in: *Thera and the Aegean World III*, edited by: Hardy, D. A., Keller, J., Galanopoulos, V. P., Flemming, N. C., and Druitt, T. H., Thera Foundation, Santorini, Greece, 13–26, 1990.
- 10 Kraml, M.: *Laser-<sup>40</sup>Ar/<sup>39</sup>Ar-Datierungen an distalen marinen tephren des jung-quartären mediterranen Vulkanismus (Ionisches Meer, METEOR-Fahrt 25/4)*, Ph.D. Thesis, Albert-Ludwigs-Universität Freiburg i. Br., 216 pp, 1997.
- Le Bas, M. J., Le Maitre, R. W., Streckeisen, A., and Zanettin, B.: A chemical classification of volcanic rocks based on the Total Alkali–Silica diagram, *J. Petrol.*, 27, 745–750, 1986.
- Lézine, A. M., von Grafenstein, U., Andersen, N., Belmecheri, S., Bordon, A., Caron, B., Cazet, J. P., Erlenkeuser, H., Fouache, E., Grenier, C., Huntsman-Mapila, P., Hureau-Mazaudier, D., Manelli, D., Mazaud, A., Robert, C., Sulpizio, R., Tiercelin, J. J., Zanchetta, G., and Zeqollari, Z.: Lake Ohrid, Albania, provides an exceptional multi-proxy record of environmental changes during the last glacial-interglacial cycle, *Palaeogeogr Palaeoclimatol.*, 287, 116–127, 2010.
- Lindhorst, K., Vogel, H., Krastel, S., Wagner, B., Hilgers, A., Zander, A., Schwenk, T., Wessels, M., and Daut, G.: Stratigraphic analysis of lake level fluctuations in Lake Ohrid: an integration of high resolution hydro-acoustic data and sediment cores, *Biogeosciences Discuss.*, 7, 3651–3689, doi:10.5194/bgd-7-3651-2010, 2010.
- 15 Lucchi, F., Tranne, C. A., De Astis, G., Keller, J., Losito, R., and Morche, W.: Stratigraphy and significance of Brown Tuffs on the Aeolian Islands (Southern Italy), *J. Volcanol. Geoth. Res.*, 177, 49–70, 2008.
- 30 Margari, V., Pyle, D. M., Bryant, C., and Gibbard, P. L.: Mediterranean tephra stratigraphy revisited: results from a long terrestrial sequence on Lesbos Island, Greece, *J. Volcanol. Geoth. Res.*, 163, 34–54, 2007.

3949

- Marianelli, P. and Sbrana, A.: Risultati di misure di standard di minerali e di vetri naturali in microanalisi a dispersione di energia, *Atti Soc. Tosc. Sc. Nat. Mem., Serie A*, 105, 57–63, 1998.
- Matzinger, A., Spirkovski, Z., Patceva, S., and Wüest, A.: Sensitivity of ancient Lake Ohrid to local anthropogenic impacts and global warming, *J. Great Lakes Res.*, 32, 158–179, 2006.
- 5 Narcisi, B. and Vezzoli, L.: Quaternary stratigraphy of distal tephra layers in the Mediterranean – an overview, *Global Planet. Change*, 21, 31–50, 1999.
- Orsi, G., De Vita, S., and Di Vito, M. A.: The restless, resurgent Campi Flegrei nested caldera (Italy): constraints on its evolution and configuration, *J. Volcanol. Geoth. Res.*, 74, 179–214, 1996.
- 10 Pappalardo, L., Civetta, L., D’Antonio, M., Deino, A., Di Vito, M. A., Orsi, G., Carandente, A., De Vita, S., Isaia, R., and Piochi, M.: Chemical and Sr-isotopic evolution of the Phlegraean magmatic system before the Campanian Ignimbrite and the Neapolitan Yellow Tuff eruptions, *J. Volcanol. Geoth. Res.*, 91, 141–166, 1999.
- 15 Pappalardo, L., Civetta, L., De Vita, S., Di Vito, M. A., Orsi, G., Carandente, A., and Fisher, R. V.: Timing of magma extraction during the Campanian Ignimbrite eruption (Campi Flegrei caldera), *J. Volcanol. Geoth. Res.*, 114, 479–497, 2002.
- Paterne, M., Guichard, F., and Labeyrie, J.: Explosive activity of the South Italian volcanoes during the past 80000 years as determined by marine tephrochronology, *J. Volcanol. Geoth. Res.*, 34, 153–172, 1988.
- 20 Paterne, M., Labeyrie, J., Guichard, F., Mazaud, A., and Maitre, F.: Fluctuations of the Campanian explosive volcanic activity (South Italy) during the past 190,000 years as determined by marine tephrochronology, *Earth Planet. Sc. Lett.*, 98, 166–174, 1990.
- Paterne, M., Guichard, F., Duplessy, J. C., Siani, G., Sulpizio, R., and Labeyrie, J.: A 90,000–200,000 yrs marine tephra record of Italian volcanic activity in the Central Mediterranean Sea, *J. Volcanol. Geoth. Res.*, 177, 187–196, 2008.
- 25 Poli, S., Chiesa, S., Gillot, P. Y., and Guichard, F.: Chemistry versus time in the volcanic complex of Ischia (Gulf of Naples, Italy): evidence of successive magmatic cycles, *Contrib. Mineral. Petr.*, 95, 322–335, 1987.
- 30 Popovska, C. and Bonacci, O.: Basic data on the hydrology of lakes Ohrid and Prespa, *Hydrol. Process.*, 21, 658–664, 2007.
- Pyle, D. M., Ricketts, G. D., Margari, V., van Andel, T. H., Sinitsyn, A. A., Praslov, N., and Lisitsyn, S.: Wide dispersal and deposition of distal tephra during the Pleistocene “Campanian

3950

- Ignimbrite/Y5" eruption, Italy, *Quaternary Sci. Rev.*, 25, 2713–2728, 2006.
- Rosi, M. and Sbrana, A. (Eds.): *The Phlaegrean Fields*, Quaderni de "La ricerca Scientifica" 114, Consiglio Nazionale delle Ricerche, Rome, 1987.
- Sadori, L. and Narcisi, B.: The postglacial record of environmental history from Lago di Pergusa, Sicily, *Holocene*, 11, 655–670, 2001.
- 5 Santacroce, R.: *Somma-Vesuvius*, Quaderni de la Ricerca Scientifica, CNR, 114, Progetto Finalizzato Geodinamica, Monografie Finali, 8, CNR, Rome, 1987.
- Santacroce, R., Cioni, R., Marianelli, P., Sbrana, A., Sulpizio, R., Zanchetta, G., Donahue, D. J., and Joron, J. L.: Age and whole rock-glass compositions of proximal pyroclastics from the major explosive eruptions of Somma-Vesuvius: a review as a tool for distal tephrostratigraphy, *J. Volcanol. Geoth. Res.*, 177, 1–18, 2008.
- 10 St Seymour, K. and Christianis, K.: Correlation of a tephra layer in Western Greece with a late Pleistocene eruption in the Campanian province of Italy, *Quaternary Res.*, 43, 46–54, 1995.
- St Seymour, K., Christianis, K., Bouzinos, A., Papazisimou, S., Papatheodorou, G., Moran, E., and Denes, G.: Tephrostratigraphy and tephrochronology in the Philippi Peat Basin, Macedonia, Northern Hellas (Greece), *Quatern. Int.*, 121, 53–65, 2004.
- 15 Sulpizio, R., Caron, B., Wulf, S., Zanchetta, G., Vogel, H., and Wagner, B.: Towards a comprehensive tephrostratigraphic framework for the Upper Pleistocene in the Central Mediterranean region, IAVCEI general assembly, Reykjavik, Iceland, abstract volume, 2008.
- 20 Sulpizio, R., Zanchetta, G., Caron, B., Dellino, P., Mele, D., Giaccio, B., Insinga, D., Paterne, M., Siani, G., Costa, A., Macedonio, G., and Santacroce, R.: The dispersal of ash during Holocene explosive activity of Italian volcanoes insights for volcanic hazard assessment in the Central Mediterranean, *B. Volcanol.*, submitted, 2010.
- Ton-That, T., Singer, B., and Paterne, M.:  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of latest Pleistocene (41 ky) marine tephra in the Mediterranean Sea: implications for global climate records, *Earth Planet. Sc. Lett.*, 184, 645–658, 2001.
- 25 Vezzoli, L.: *Island of Ischia*, CNR, Quaderni della Ricerca Scientifica, vol. 114(10), CNR, Italy, 230 pp, 1988.
- Vogel, H., Zanchetta, G., Sulpizio, R., Wagner, B., and Nowaczyk, N.: A tephrostratigraphic record for the last glacial – interglacial cycle from Lake Ohrid, Albania and Macedonia, *J. Quaternary Sci.*, 25, 320–338, 2010a.
- 30 Vogel, H., Wagner, B., Zanchetta, G., Sulpizio, R., and Rosén, P.: A Paleoclimate record with tephrochronological age control for the last glacial-interglacial cycle from Lake Ohrid, Albania

3951

- and Macedonia, *J. Paleolimnol.*, 44, 295–310, doi:10.1007/s10933-009-9404-x, 2010b.
- Wagner, B., Sulpizio, R., Zanchetta, G., Wulf, S., Wessels, M., and Daut, G.: The last 40 ky tephrostratigraphic record of Lake Ohrid, Albania and Macedonia: a very distal archive for ash dispersal from Italian volcanoes, *J. Volcanol. Geoth. Res.*, 177, 71–80, 2008.
- 5 Wagner, B., Lotter, A. F., Nowaczyk, N., Reed, J. M., Schwab, A., Sulpizio, R., Valsecchi, V., Wessels, M., and Zanchetta, G.: A 40000-year record of environmental change from ancient Lake Ohrid (Albania and Macedonia), *J. Paleolimnol.*, 41, 407–430, 2009.
- Wagner, B., Vogel, H., Zanchetta, G., and Sulpizio, R.: Environmental changes on the Balkans recorded in the sediments from lakes Prespa and Ohrid, *Biogeosciences Discuss.*, 7, 3365–3392, doi:10.5194/bgd-7-3365-2010, 2010.
- 10 Wulf, S., Kraml, M., Brauer, A., Keller, J., Negendank, J. F. W.: Tephrochronology of the 100 ky lacustrine sediment record of Lago Grande di Monticchio (Southern Italy), *Quatern. Int.*, 122, 7–30, 2004.
- Zanchetta, G., Sulpizio, R., Giaccio, B., Siani, G., Paterne, M., Wulf, S., and D'Orazio, M.: The Y-3 Tephra: a Last Glacial stratigraphic marker for the Central Mediterranean Basin, *J. Volcanol. Geoth. Res.*, 177, 145–154, 2008.
- 15 Zanchetta, G., Sulpizio, R., Roberts, N., Cioni, R., Eastwood, W. J., Siani, G., Paterne, M., and Santacroce, R.: Tephrostratigraphy, chronology and climatic events of the Mediterranean Basin during the Holocene: an overview, *Holocene*, in press, 2010.

3952
















3961




3962




**Fig. 4.** SEM photographs of volcanic fragments from selected tephra layers and cryptotephra. (a) Y-3 tephra layer in core Co1200; (b) CI/Y-5 tephra layer from core Co1200; (c) C-20 cryptotephra in core Co1201; (d) X-5 tephra layer in core Co1201; (e) AD 472 cryptotephra in core Co1202; (f) FL cryptotephra in Co 1202 core; (g) Mercato cryptotephra in Co1202 core; (h) Codola cryptotephra in core Co1202; (i) X-6 tephra layer core Co1202.

3963




**Fig. 5.** Primitive-mantle and chondrite-normalised rare earth element (REE) diagrams for the selected samples. Samples 523, 535 and 1033 are from core Co1202, the sample PR628 is from core Co1204.

3964




**Fig. 6.** (a) Total alkali vs. silica diagram for the different groups of OT0702-5 (pale blue dots), PT09-16 (red and green dots), PT0704-2 (purple, orange and black dots); (b) CaO vs. SiO<sub>2</sub> diagram for the same groups of Fig. 5a. 3965



**Fig. 7.** Updated tephrochronology for the last 131 ky of lakes Ohrid and Prespa. Grey boxes indicate the age uncertainties for calibrated <sup>14</sup>C data and for <sup>39</sup>Ar/<sup>40</sup>Ar ages. <sup>1</sup> Coltelli et al. (2000); <sup>2</sup> Zanchetta et al. (2010); <sup>3</sup> Di Vito et al. (2008); <sup>4</sup> Vogel et al. (2010); <sup>5</sup> De Vivo et al. (2001); <sup>6</sup> Paterne et al. (1988); <sup>7</sup> Kraml (1997); <sup>8</sup> Allen et al. (1999); <sup>9</sup> Paterne et al. (2008). The three colours indicate the Middle Pleistocene, Upper Pleistocene and Holocene.





**Fig. 8.** Link of the Ohrid and Prespa tephrostratigraphy to the other archives of the Central Mediterranean area. SA=South Adriatic Sea; ST=South Tyrrhenian Sea.