

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Air-sea exchange of CO₂ at a Northern California coastal site along the California Current upwelling system

H. Ikawa¹, I. Faloona², J. Kochendorfer³, K. T. Paw U², and W. C. Oechel¹

¹Global Change Research Group, San Diego State University, 5500 Campanile Dr., San Diego, CA 92182-4614, USA

²Department of Land, Air, and Water Resources, UC Davis, One Shield Ave, Davis, CA 95616-8627, USA

³Atmospheric Turbulence and Diffusion Division, NOAA, 456 S. Illinois Ave., Oak Ridge, TN 37830, USA

Received: 23 October 2012 – Accepted: 12 November 2012 – Published: 12 December 2012

Correspondence to: H. Ikawa (hikawa@mail.sdsu.edu) and W. C. Oechel (woechel@mail.sdsu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

17707

Abstract

Uncertainty in the air-sea CO₂ exchange (CO₂ flux) in coastal upwelling zones is attributed to high temporal variability, which is caused by changes in ocean currents. Upwelling transports heterotrophic, CO₂ enriched water to the surface and releases CO₂ to the atmosphere, whereas the presence of nutrient-rich water at the surface supports high primary production and atmospheric CO₂ uptake. To quantify the effects of upwelling on CO₂ fluxes, we measured CO₂ flux at a coastal upwelling site off of Bodega Bay, California, during the summer of 2007 and the fall of 2008 using the eddy covariance technique and the bulk method with *p*CO₂ measurements from November 2010 to July 2011. Variations in sea surface temperatures (SST) and alongshore wind speeds suggest that the measurement period in 2007 coincided with a typical early-summer upwelling period and the measurement period in 2008 was during a typical fall relaxation period. A strong source of CO₂ ($\sim 1.5 \pm 7$ SD (standard deviation) g C m⁻² day⁻¹) from the ocean to the atmosphere during the upwelling period was concurrent with high salinity, low SST, and low chlorophyll density. In contrast, a weak source of CO₂ flux ($\sim 0.2 \pm 3$ SD g C m⁻² day⁻¹) was observed with low salinity, high SST and high chlorophyll density during the relaxation period. Similarly, the sink and source balance of CO₂ flux was highly related to salinity and SST during the *p*CO₂ measurement periods; high salinity and low SST corresponded to high *p*CO₂, and vice versa. We estimated that the coastal area off Bodega Bay was likely a source of CO₂ to the atmosphere based on the following conclusions: (1) the overall CO₂ flux estimated from both eddy covariance and *p*CO₂ measurements showed a source of CO₂; (2) although the relaxation period during the 2008 measurements were favorable to CO₂ uptake, CO₂ flux during this period was still a slight source, (3) salinity and SST were found to be good predictors of the CO₂ flux for both eddy covariance and *p*CO₂ measurements, and historical data of daily averaged SST and salinity between 1988 to 2011 show that 99% of the data falls within the range of our observation in May–June 2007, August–September 2008 and November 2010–July 2011 indicating that our data set was representative of the

17708

annual variations in the sea state. Based on the developed relationship between $p\text{CO}_2$ and SST and salinity, the average annual CO_2 flux between 1988 and 2011 was estimated to be $\sim 35 \text{ mol C m}^{-2} \text{ yr}^{-1}$. The peak monthly CO_2 flux of $\sim 7 \text{ mol C m}^{-2} \text{ month}^{-1}$ accounted for about 30 % of the dissolved inorganic carbon in the surface mixed-layer.

5 1 Introduction

Intensive efforts have been underway over the last few decades to better understand and quantify global air-sea CO_2 exchange. These investigations reveal that the oceans absorb approximately one-third of the global anthropogenic CO_2 emissions (Gruber et al., 2009; Takahashi et al., 2009), and have accumulated 118 (± 19) PgC of carbon emitted by fossil fuel combustion, land use change, and cement production between 1800 and 1994 (Sabine et al., 2004). However, current estimates of the global ocean CO_2 uptake still contain considerable uncertainty, and more observations are necessary to better understand this critical carbon reservoir (Doney et al., 2009). One of the largest uncertainties in the global carbon cycle is the role of coastal seas, which potentially account for 30 % of the net global sink of CO_2 from the atmosphere to the ocean (Chen and Borges, 2009). Current global estimates do not account for the variability of the coastal ocean due to the difficulties of quantifying coastal CO_2 fluxes (Takahashi et al., 2009). In order to better understand the dynamics of coastal CO_2 fluxes, we must understand the physical and biological processes unique to coastal waters that control these fluxes.

Coastal upwelling is an important process that conflates the physical and biological influences on the coastal carbon cycle (Dugdale et al., 2006; Gattuso et al., 1998). Upwelling conveys deep nutrient-rich water to the surface and supports diverse coastal ecosystems. Although the surface of coastal upwelling zones covers only 1 % of the global ocean, these areas support approximately 50 % of the fishery industry because of their high primary productivity (Gattuso et al., 1998). Observational studies indicate that upwelling along the California coast has increased over the past 30 yr

17709

(Garcia-Reyes and Largier, 2010). This intensification of upwelling is likely related to increasing greenhouse gases, as the increasing temperature difference between land and sea accelerates coastal upwelling (Bakun, 1990; Diffenbaugh et al., 2004).

The sign of the effects of coastal upwelling on CO_2 fluxes has been debated. Upwelling provides a source of CO_2 from the ocean to the atmosphere because deep water is enriched with respired CO_2 . At the same time, upwelling carries nutrients to the surface that enhance the biological uptake of CO_2 (Borges and Frankignoulle, 2002). Thus, the net CO_2 flux in upwelling zones averaged for inter-seasonal scales can be a sink of CO_2 from the atmosphere to the ocean (Evans et al., 2011; Gago et al., 2003; Hales et al., 2005; Ianson and Allen, 2002). The source and sink pulses of CO_2 flux reported in past studies range from -10 to $30 \text{ mol C m}^{-2} \text{ yr}^{-1}$ (Copin-Montegut and Raimbault, 1994; Friederich et al., 2002; Goyet et al., 1998; Hales et al., 2005; Ianson and Allen, 2002; Ianson et al., 2009; Kelley and Hood, 1971; Lendt et al., 2003; Simpson and Zirino, 1980; Torres et al., 1999), using the sign convention of positive fluxes into the atmosphere from the ocean, and they are frequent two orders magnitude greater than the global average net CO_2 flux of approximately $-0.4 \text{ mol C m}^{-2} \text{ yr}^{-1}$ (Takahashi et al., 2009). This emphasizes the importance of understanding CO_2 fluxes in the complex coastal zones where upwelling occurs. Despite its importance, uncertainty in determining the net CO_2 flux in upwelling zones remains due to the spatial and temporal sparsity of the available measurements.

Another uncertainty in the CO_2 flux of coastal upwelling zones is attributed to the fact that a standard measurement technique for air-sea CO_2 flux is still under development. Quantifications of the spatial and temporal variations of CO_2 flux in upwelling zones were made, for example, by Friederich et al. (2002) for the California coast and Evans et al. (2011) for the Oregon coast. However, these past studies estimated the CO_2 flux using the bulk method based on the partial pressure of CO_2 ($p\text{CO}_2$) measured in surface waters. Currently, the gas transfer coefficient required for the bulk method is determined solely by empirical relationships between gas transfer coefficients and wind speed, which were developed in the open ocean where wind speeds are generally

17710

much higher than coastal areas (Ho et al., 2006; Wanninkhof and McGillis, 1999). Under lower wind speeds, the effects of factors such as surface films and rain may be important as well for determining gas transfer properties (Frew et al., 2004; Jähne et al., 1987), and the effects of breaking waves and bubbles may be especially influential in coastal seas (Keeling, 1993).

The eddy covariance technique utilized in this study is considered to be the most direct CO₂ flux measurement method available. The application of the technique to ocean fluxes is still challenging due to the fact that data processing introduces high uncertainty to the measurement of smaller fluxes (less than 1 gCm⁻² day⁻¹) and the sensors are sensitive to contamination in marine environments (Else et al., 2011; Iwata et al., 2005; Prytherch et al., 2010). However, frequent maintenance and the expected large magnitude of the CO₂ fluxes from coastal upwelling zones enabled us to employ the open path eddy covariance technique.

The objective of this study was to quantify the CO₂ flux from the wind-driven upwelling zone near Bodega Bay, California, and evaluate the physical and biological effects of upwelling on the observed near-shore CO₂ flux. We hypothesized that the sink or source characteristics of the coastal region are determined by a balance between the transports of CO₂ enriched water to the surface and high primary productivity. To test this hypothesis, we measured CO₂ fluxes using the eddy covariance technique and the bulk transfer method with pCO₂ measurements and compared the observed CO₂ fluxes with environmental variables related to upwelling events (e.g. wind speed, salinity, sea temperature) and primary production (e.g. light and nutrient availability and chlorophyll density).

17711

2 Methods

2.1 Site descriptions

The study site was located at the intertidal zone in the vicinity of the Bodega Bay Marine Laboratory (BML) in Northern California (Fig. 1). The coastal sea of Northern California is classically characterized by northerly winds in the summer due to the lateral pressure gradient (Dorman et al., 2000) established by the Pacific High offshore. The predominant alongshore wind causes Ekman transport along with lateral shear vorticity, which generates positive curl in the surface stress, resulting in strong coastal upwelling. Spatial distributions of water temperature indicated that our study area was particularly favorable for wind-driven upwelling events (Day and Faloon, 2009). The region's wind-driven upwelling cycle, nutrient availability and subsequent primary production have been described in CoOP WEST (Wind Events and Shelf Transport) studies (Dugdale et al., 2006; Largier et al., 2006; Wilkerson et al., 2006). The region's typically strong winds are periodically interrupted by a period with weaker winds known as a relaxation period, which lasts for ~ 3–7 days. The repeating cycles of upwelling and relaxation create a unique biological system. Upwelling carries deep water that is relatively cold and nutrient-rich to the surface, but increased turbidity (Day and Faloon, 2009) and lateral exports of nutrients inhibit production, while nutrients brought up to the surface are consumed during the subsequent relaxation period (Largier et al., 2006; Wilkerson et al., 2006).

The year of 2007 and 2008, when the eddy covariance measurements operated, were neutral to El Niño-Southern Oscillation. The winter of 2010, when pCO₂ measurements were operated, was characterized as a moderate El Niño condition.

17712

since 1988, and chlorophyll data was available only from 2005 to 2008. The $p\text{CO}_2$ was measured every hour by a SAMI- $p\text{CO}_2$ (Sun-Burst Sensors, USA) colorimetric sensor mounted on the buoy from November 2010 to July 2011. The measurement was temporarily halted due to a system malfunction in December 2010 through February 2011. The sensor was calibrated by the factory within the range 150–700 μatm . All the data used in this study are summarized in Table 1.

2.4 CO_2 flux estimation with $p\text{CO}_2$ and atmospheric CO_2

CO_2 flux (F_c) was calculated from $p\text{CO}_2$ (C_{pw}), SST, salinity, and atmospheric CO_2 pressure (C_{pa}) with the bulk method for each $p\text{CO}_2$ measurement based on the following equation (e.g. Wanninkhof and McGillis, 1999).

$$F_c = \alpha k [C_{pa} - C_{pw}] \quad (3)$$

The solubility, α , was computed by SST and salinity based on Weiss (1974); the gas transfer coefficient, k , was determined to be the average value calculated following Wanninkhof and McGillis (1999) (hereafter, W&M99), Wanninkhof (1992) (hereafter, W92), Ho et al. (2006) (hereafter, Ho06), Nightingale et al. (2000) (hereafter, N00), and Sweeney et al. (2007) (hereafter, Sw07). The equations for k (md^{-1}) are as follows:

17717

[W&M99]

$$k = 0.0068u^3 \times (Sc/660)^{-0.5}, \quad (4)$$

[W92]

$$k = 0.074u^2 \times (Sc/660)^{-0.5}, \quad (5)$$

[Ho06]

$$k = 0.061u^2 \times (Sc/660)^{-0.5}, \quad (6)$$

[N00]

$$k = [0.053u^2 + 0.024u] \times (Sc/660)^{-0.5}, \quad (7)$$

[Sw07]

$$k = 0.065u^2 \times (Sc/660)^{-0.5}, \quad (8)$$

where, u is wind speed and Sc is Schmidt number (Wanninkhof, 1992).

Data collected monthly by the NOAA/ESRL Carbon Cycle Greenhouse Gases (CCGG) group's aircraft program were used to estimate C_{pa} . The nearest routine sampling location is offshore near Trinidad Head, CA, which is located about 300 km north of Bodega Bay ($\sim 41^\circ 10' \text{N}$, $124^\circ 20' \text{W}$). The lowest altitude data (near 300 m a.s.l.) from each month since September 2003 were used to estimate the average monthly atmospheric CO_2 concentration in the marine boundary layer throughout the region. The CO_2 concentration was converted to the pressure, C_{pa} under the standard atmosphere.

2.5 Effective upwelling and National Data Buoy Center 46013 data

The National Data Buoy Center (NDBC) 46013 buoy is located off of Bodega Bay, California ($38^\circ 14' \text{N}$, $123^\circ 18' \text{W}$). The Effective Upwelling Index (EUI) was introduced by Garcia-Reyes (2011) as a proxy for the cumulative NO_3 . It is estimated using the sea surface temperature measured at the NDBC 46013 buoy (SST_{N13}) from the National

17718

Data Buoy Center (<http://www.ndbc.noaa.gov>). To identify upwelling periods, NO_3 concentrations were estimated based on the NO_3 -SST_{N13} relation (Garcia-Reyes, 2011). EUI was then calculated as a summation of a temporal differential of NO_3 estimated while the daily averaged alongshore wind speed (AWS) was greater than 5 ms^{-1} . EUI was set back to zero when daily AWS was less than 5 ms^{-1} for over 3 days.

2.6 Statistic analysis

2.6.1 For the eddy covariance data set

A t-test was used to test the significance of differences between the 2007 and 2008 CO_2 fluxes and environmental parameters (i.e. SST, salinity, PPF, wind speeds, AWS, friction velocity, chlorophyll density).

2.6.2 For the ρCO_2 data set

Linear correlation analysis was used to investigate relationships between environmental parameters (i.e. SST, salinity, PPF, wind speeds and AWS) and ρCO_2 and CO_2 fluxes calculated from ρCO_2 . Since diurnal variations in ρCO_2 were minimal (less than 10 % of the total variations), these data were averaged for 24-h periods.

2.7 Estimates of dissolved inorganic carbon (DIC) and cumulative monthly CO_2 flux and net primary production (NPP)

To evaluate the monthly net CO_2 exchange between the air-water interface and the monthly net primary production (NPP) and to compare them with the dissolved inorganic carbon (DIC) pool, monthly CO_2 fluxes and NPP were estimated and compared with estimated DIC per unit area within the surface mixed layer.

Since daily averaged ρCO_2 was most correlated with SST and salinity, as will be demonstrated later, daily ρCO_2 from 1988 to 2011 was estimated based on the SST

17719

and salinity data collected from the BML buoy with the following equation:

$$C_{\text{pw}} = -142S_T + 115S_s - 1473 \quad (9)$$

where, S_T is STT and S_s is salinity. CO_2 fluxes were calculated in the same manner as explained above. DIC (C_{dic}) was computed based on the estimated ρCO_2 , SST, salinity and pH (pp 324–325 in Sarmiento and Gruber, 2006):

$$C_{\text{dic}} = C_{\text{pw}}K_0[1 + K_1[\text{H}^+]^{-1} + K_1K_2[\text{H}^+]^{-2}] \quad (10)$$

where, K_0 is the solubility of CO_2 ($\text{mol kg}^{-1} \text{ atm}^{-1}$), K_1 and K_2 are dissociation constants of inorganic carbon species (mol kg^{-1}) determined from temperature and salinity, and $[\text{H}^+]$ is hydrogen ion concentration (mol kg^{-1}) and equal to $10^{-\text{pH}}$. Hauri et al. (2012) reported seasonal cycles of pH off the Central California coast, and based on these results we approximated the pH seasonal cycle with a sine curve. The influence of the seasonal variation of pH on the DIC calculation was minimal (less than 3%). Similarly, the NPP off Central California reported in Kahru and Mitchell (2002) was approximated by a sine curve. The surface mixed-layer depth was approximated by wind stress estimated from wind speeds (Lentz, 1992).

3 Results and discussions

High CO_2 efflux to the atmosphere was observed during the measurement period in 2007. The average CO_2 flux during this period was $1.5 (\pm 7 \text{ SD}) \text{ gC m}^{-2} \text{ day}^{-1}$ with high variability (Fig. 3a, Table 2). Although a relatively strong sink of CO_2 was observed with high wind speeds on 7–12 September 2008, the overall CO_2 flux measured in 2008 was a slight source of CO_2 , with a magnitude of $0.2 (\pm 3 \text{ SD}) \text{ gC m}^{-2} \text{ day}^{-1}$ (Fig. 4a, Table 2), and the average CO_2 flux during the 2008 measurement season was not statistically different from zero (Table 2).

17720

Contrasting differences were found in SST, salinity, and chlorophyll density between the 2007 and 2008 measurement seasons (Table 2). No difference in AWS between the 2007 and 2008 measurement periods was apparent. However, AWS on longer time scales strengthened in late spring to early summer corresponding with a drop in sea surface temperatures (Fig. 5a, b). The AWS then weakened in fall leading to an increase in sea surface temperature (Garcia-Reyes, 2011). A high EUI ($\sim 20 \mu\text{M}$) was still observed during fall 2008, but a negative AWS ($\sim -5 \text{ms}^{-1}$) was frequently observed, creating conditions favorable to relaxation (Fig. 5a, c). The relatively lower SSTs ranging from 10 to 12 °C in May–June and higher SSTs ranging from 15 to 18 °C in August–September shown in Fig. 5b are compatible with the typical seasonal variations in SST, where upwelling intensifies in early summer and relaxes in fall (Garcia-Reyes, 2011). The upwelling season is also characterized by high temporal variations in CO_2 flux, which was also observed in the 2007 measurements, due to the highly dynamic shifts of water chemistry caused by upwelled water with high CO_2 and subsequent CO_2 uptake by primary production (Dugdale et al., 2006; Evans et al., 2011; Gago et al., 2003).

Relaxation periods are favorable to stratification of the upper water column, which was reflected in lower salinity ($\sim 32 \text{psu}$) and higher SST ($\sim 17^\circ\text{C}$) during the 2008 measurement period. Stratification further enhances nutrient depletion on the surface and increases primary production, resulting in higher chlorophyll density. Interestingly, an increasing trend of chlorophyll density was observed during the 2008 measurement period (Fig. 4d). A sink of CO_2 , $\sim 2 \text{gCm}^{-2} \text{day}^{-1}$, was consistently observed on 15 September 2008 when chlorophyll density approached the maximum of 2.3mgm^{-3} . The measurement period in 2008 fell directly between high upwelling events with the peaks of EUI in mid-July and October 2008 (Fig. 5c). Therefore, high nutrient levels were brought to the surface by the preceding upwelling event, increasing the chlorophyll density until the nutrients were depleted. Kahru and Mitchell (2002) reported high NPP of $2.4 \text{gCm}^{-2} \text{day}^{-1}$ (ca. $5 \times 10^{12} \text{gCmonth}^{-1}$ for the area of $6.2 \times 10^4 \text{km}^2$) in summer when nutrient availability was high near our study site. However, the overall CO_2

17721

flux was near zero or a slight source during the 2008 measurement period despite the favorable conditions for CO_2 uptake. This was because the period preceded a high upwelling period that stimulated higher chlorophyll density and probable relaxation conditions, as inferred from the negative AWS.

The overall CO_2 flux estimated from $\Delta p\text{CO}_2$ was also a source of $0.4 (\pm 0.6 \text{SD}) \text{gCm}^{-2} \text{day}^{-1}$ from November 2010 to July 2011 (Fig. 6f). The $p\text{CO}_2$ data were often out of the calibration range (150–700 μatm), and therefore the data may not have been accurate at higher $p\text{CO}_2$ values (Fig. 6e). However, $p\text{CO}_2$ higher than 1000 μatm is likely in our study area (personal communication with John Largier of University of California, Davis, 2007). Correlation analysis showed that $p\text{CO}_2$ was most correlated with SST ($R = -0.54$, $p < 0.001$) followed by salinity ($R = 0.50$, $p < 0.001$), and CO_2 flux was strongly correlated to wind speeds ($R = 0.58$, $p < 0.001$) and AWS ($R = 0.51$, $p < 0.001$) followed by SST ($R = -0.40$, $p < 0.001$) and salinity ($R = 0.34$, $p < 0.001$). Similar to the flux data obtained by the eddy covariance technique, higher salinity and lower SST corresponded with higher $p\text{CO}_2$ and a source of CO_2 (Fig. 7). Salinity and SST buoy data from 1988–2011 showed that salinity reached its maximum level and SST reached its minimum level concurrently, as AWS peaked during the upwelling season in late spring and early summer (Fig. 8). Therefore, although AWS did not reflect the CO_2 flux on short time scales ($\sim 1 \text{h}$), it likely reflected the CO_2 fluxes on a seasonal scale. Interestingly, the seasonal change of the atmospheric CO_2 showed a very similar pattern to AWS, (Fig. 8a) with a higher peak during a high upwelling season in late spring to early summer; however, because the atmospheric changes in CO_2 concentration were so much smaller than those induced in the surface ocean this periodicity did not influence the overall flux significantly.

The CO_2 flux estimated from the eddy covariance technique and from the bulk method both showed that positive CO_2 fluxes were larger than negative fluxes, and that this balance was tightly reflected in salinity and SST. Considering the fact that 99% of the historical data set of salinity and SST fell within the range of our observations, supporting the notion that the data is representative of the region in general,

17722

our data indicates that the overall exchange of CO₂ in this area is likely a net source of CO₂ to the atmosphere. Table 3 lists our CO₂ flux data together with the CO₂ flux in upwelling zones reported in past studies. Although the annual net CO₂ flux in some upwelling zones (e.g. North Pacific and Galician coast) show a sink of CO₂ and high primary production (Evans et al., 2011), particularly in strong downwelling conditions (Ianson et al., 2009), other studies conducted in North-Central California (Feely et al., 2008; Friederich et al., 2002; Wilkerson et al., 2006) further support our conclusion that the upwelling zone off of Bodega Bay is an overall source of CO₂ to the atmosphere. Despite the high chlorophyll density of 11 mg m⁻³ observed in the early summer of 2000 by Wilkerson et al. (2006), their pCO₂ measurements still indicated that the area was a source of CO₂ to the atmosphere (Table 3). Their study also showed that the magnitude of high pCO₂ was similar over the cross-shelf distance of at least 50 km offshore (Wilkerson et al., 2006).

The estimation of CO₂ flux and DIC based on the relationship of pCO₂ with SST and salinity revealed that the monthly CO₂ flux reached its maximum in May (~ 7 mol C m⁻² month⁻¹), which is equal to about 30 % of DIC within the surface mixed-layer (hDIC) (Fig. 9). The annual CO₂ flux was estimated to be about 35 mol C m⁻² yr⁻¹. The fraction of monthly NPP to hDIC became highest (~ 50%) in late summer, and the annual NPP was estimated to be about 40 mol C m⁻² yr⁻¹. This simple estimation suggested that CO₂ flux and NPP were about the same magnitude and together accounted for significant flux compared to the stock of DIC.

4 Summary

The CO₂ flux from the coastal waters of Bodega Bay was measured during a typical upwelling season in 2007 and during a relaxation season in 2008. CO₂ flux was also estimated from pCO₂ measured from November 2010 to July 2011. Contrasting patterns in CO₂ flux and environmental parameters were observed between the two measurement periods in 2007 and 2008. Measurements in 2007 were during a typical

17723

upwelling period, as indicated by the seasonal variations of AWS and EU in early summer, and a source of CO₂ to the atmosphere was observed concurrently with low SST, high salinity, and low chlorophyll density. The measurements in 2008 were, on the contrary, made during a relaxation period with high SST and low salinity that likely led to water stratification and high chlorophyll density. A strong upwelling event preceding the measurements in 2008 likely created favourable conditions for CO₂ uptake. However, the average CO₂ flux over the 2008 measurements was a small source of CO₂ overall. The magnitude of the observed sources CO₂ to the atmosphere estimated by the bulk method were also much higher than that of sinks of CO₂ during the pCO₂ measurements. Similar to the fluxes observed by the eddy covariance technique, those estimated by the pCO₂ measurements were closely related to salinity and SST. The historical data set of salinity and SST (1988–2011) indicated that 99 % of data fell in the range of our observations. Therefore, the area was likely a source of CO₂ to the atmosphere annually. Past studies further support the assertion that the overall conditions are favourable to producing a source of CO₂ in the coastal upwelling area off of Bodega Bay at least within 50 km of the coast. Although AWS did not reflect pCO₂ or CO₂ flux on shorter time scales, high AWS together with low SST and high salinity suggests that AWS likely drives the seasonal cycle of CO₂ flux. Based on the pCO₂ to SST and salinity relationship, the average annual CO₂ flux between 1988 and 2011 was estimated to be 35 mol C m⁻² yr⁻¹, and the amount of CO₂ transferred vertically and consumed by NPP accounted for significant amounts of carbon in comparison to the local dissolved carbon stock.

Acknowledgements. We thank Jackie Sones, Deedee Shideler and logistic supports by Bodega Bay Marine Laboratory, Tessa Hill for pCO₂ data retrieval and Joseph Verfaillie for valuable suggestions on data processing and interpretation. Sea surface temperature, wind speed and wind direction for calculating EU was provided by the National Data Buoy Center. SAMI-pCO₂ data were provided by the Bodega Ocean Acidification Research group (NSF OCE # 0927255 to B. Gaylord), and Bodega Ocean Observing Network (BOON) data were provided by John Largier. Atmospheric CO₂ data from the NOAA/ESRL Carbon Cycle Greenhouse Gases group were provided by Colm Sweeney.

17724

References

- Auble, D. L. and Meyers, T. P.: An open path, fast response infrared absorption gas analyzer for H₂O and CO₂, *Bound.-Lay. Meteorol.*, 59, 243–256, doi:10.1007/BF00119815, 1992.
- Bakun, A.: Global climate change and intensification of coastal ocean upwelling, *Science*, 247, 198–201, doi:10.1126/science.247.4939.198, 1990.
- Baldocchi, D. D., Hicks, B. B., and Meyers, T. P.: Measuring biosphere-atmosphere exchanges of biologically related gases with micrometeorological methods, *Ecology*, 69, 1331–1340, 1988.
- Borges, A. V. and Frankignoulle, M.: Distribution of surface carbon dioxide and air-sea exchange in the upwelling system off the Galician coast, *Global Biogeochem. Cy.*, 16, 1020, doi:10.1029/2000GB001385, 2002.
- Burba, G. G. and Anderson, D. J.: A Brief Practical Guide to Eddy Covariance Flux Measurements: Principles and Workflow Examples for Scientific and Industrial Applications, Li-COR Biosciences, Lincoln, USA, 2010.
- Burba, G. G., McDermitt, D. K., Grelle, A., Anderson, D. J., and Xu, L.: Addressing the influence of instrument surface heat exchange on the measurements of CO₂ flux from open-path gas analyzers, *Glob. Change Biol.*, 14, 1854–1876, doi:10.1111/j.1365-2486.2008.01606.x, 2008.
- Chen, C.-T. A. and Borges, A. V.: Reconciling opposing views on carbon cycling in the coastal ocean: continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO₂, *Deep-Sea Res. II*, 56, 578–590, doi:10.1016/j.dsr2.2009.01.001, 2009.
- Copin-Montegut, C. and Raimbault, P.: The Peruvian upwelling near 15 S in August 1986. Results of continuous measurements of physical and chemical properties between 0 and 200 m depth, *Deep-Sea Res. I*, 41, 439–467, 1994.
- Day, D. A. and Faloon, I.: Carbon monoxide and chromophoric dissolved organic matter cycles in the shelf waters of the Northern California upwelling system, *J. Geophys. Res.*, 114, C01006, doi:10.1029/2007JC004590, 2009.
- Diffenbaugh, N. S., Snyder, M. A., and Sloan, L. C.: Could CO₂-induced land-cover feedbacks alter near-shore upwelling regimes?, *PNAS*, 101, 27–32, doi:10.1073/pnas.0305746101, 2004.

17725

- Doney, S. C., Tilbrook, B., Roy, S., Metzl, N., Le Quééré, C., Hood, M., Feely, R. A., and Bakker, D.: Surface-ocean CO₂ variability and vulnerability, *Deep-Sea Res. II*, 56, 504–511, doi:10.1016/j.dsr2.2008.12.016, 2009.
- Dorman, C. E., Holt, T., Rogers, D. P., and Edwards, K.: Large-scale structure of the June–July 1996 marine boundary layer along California and Oregon, *Mon. Weather Rev.*, 128, 1632–1652, doi:10.1175/1520-0493(2000)128<1632:LSSOTJ>2.0.CO;2, 2000.
- Dugdale, R. C., Wilkerson, F. P., Hogue, V. E., and Marchi, A.: Nutrient controls on new production in the Bodega Bay, California, coastal upwelling plume, *Deep-Sea Res. II*, 53, 3049–3062, doi:10.1016/j.dsr2.2006.07.009, 2006.
- Else, B. G. T., Papakyriakou, T. N., Galley, R. J., Drennan, W. M., Miller, L. A., and Thomas, H.: Wintertime CO₂ fluxes in an Arctic polynya using eddy covariance: evidence for enhanced air-sea gas transfer during ice formation, *J. Geophys. Res.*, 116, C00G03, doi:10.1029/2010JC006760, 2011.
- Evans, W., Hales, B., and Strutton, P. G.: Seasonal cycle of surface ocean pCO₂ on the Oregon shelf, *J. Geophys. Res.*, 116, C05012, doi:10.1029/2010JC006625, 2011.
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for upwelling of corrosive “acidified” water onto the continental shelf, *Science*, 320, 1490–1492, doi:10.1126/science.1155676, 2008.
- Foken, T. and Wichura, B.: Tools for quality assessment of surface-based flux measurements, *Agr. Forest Meteorol.*, 78, 83–105, 1996.
- Frew, N. M., Bock, E. J., Schimpf, U., Hara, T., Haußecker, H., Edson, J. B., McGillis, W. R., Nelson, R. K., McKenna, S. P., Uz, B. M., and Jähne, B.: Air-sea gas transfer: its dependence on wind stress, small-scale roughness, and surface films, *J. Geophys. Res.*, 109, C08S17, doi:10.1029/2003JC002131, 2004.
- Friederich, G. E., Walz, P. M., Burczynski, M. G., and Chavez, F. P.: Inorganic carbon in the Central California upwelling system during the 1997–1999 El Niño-La Niña event, *Prog. Oceanogr.*, 54, 185–203, 2002.
- Gago, J., Gilcoto, M., Pérez, F., and Ríos, A.: Short-term variability of fCO₂ in seawater and air–sea CO₂ fluxes in a coastal upwelling system (Ría de Vigo, NW Spain), *Mar. Chem.*, 80, 247–264, doi:10.1016/S0304-4203(02)00117-2, 2003.
- García-Reyes, M.: Variability in Coastal Upwelling off Central and Northern California, Ph. D. dissertation, University of California, Davis, 2011.

17726

- Garcia-Reyes, M. and Largier, J.: Observations of increased wind-driven coastal upwelling off Central California, *J. Geophys. Res.*, **115**, C04011, doi:10.1029/2009JC005576, 2010.
- Gattuso, J. P., Frankignoulle, M., and Wollast, R.: Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, **29**, 405–434, 1998.
- 5 Goyet, C., Millero, F. J., O'Sullivan, D. W., Eischeid, G., McCue, S. J., and Bellerby, R. G. J.: Temporal variations of $p\text{CO}_2$ in surface seawater of the Arabian Sea in 1995, *Deep-Sea Res. I*, **45**, 609–623, 1998.
- Gruber, N., Gloor, M., Fletcher, S. E. M., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Muller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO_2 , *Global Biogeochem. Cy.*, **23**, GB1005, doi:10.1029/2008GB003349, 2009.
- 10 Hales, B., Takahashi, T., and Bandstra, L.: Atmospheric CO_2 uptake by a coastal upwelling system, *Global Biogeochem. Cy.*, **19**, GB1009, doi:10.1029/2004GB002295, 2005.
- Hauri, C., Gruber, N., Vogt, M., Doney, S. C., Feely, R. A., Lachkar, Z., Leinweber, A., McDonnell, A. M. P., Munnich, M., and Plattner, G.-K.: Spatiotemporal variability and long-term trends of ocean acidification in the California Current System, *Biogeosciences Discuss.*, **9**, 10371–10428, doi:10.5194/bgd-9-10371-2012, 2012.
- 15 Ho, D. T., Law, C. S., Smith, M. J., Schlosser, P., Harvey, M., and Hill, P.: Measurements of air-sea gas exchange at high wind speeds in the Southern Ocean: implications for global parameterizations, *Geophys. Res. Lett.*, **33**, 16611, doi:10.1029/2006GL026817, 2006.
- Ianson, D., Feely, R. A., Sabine, C. L., and Juranek, L. W.: Features of coastal upwelling regions that determine net air-sea CO_2 flux, *J. Oceanogr.*, **65**, 677–687, 2009.
- Iwata, T., Yoshikawa, K., Higuchi, Y., Yamashita, T., Kato, S., and Ohtaki, E.: The spectral density technique for the determination of CO_2 flux over the ocean, *Bound.-Lay. Meteorol.*, **117**, 511–523, doi:10.1007/s10546-005-2773-4, 2005.
- 25 Jähne, B., Münnich, K. O., Dutzi, R. B. A., Huber, W., and Libner, P.: On the parameters influencing air-water gas exchange, *J. Geophys. Res.*, **92**, 1937–1949, doi:10.1029/JC092iC02p01937, 1987.
- Kahru, M. and Mitchell, B. G.: Influence of the El Niño-La Niña cycle on satellite-derived primary production in the California Current, *Geophys. Res. Lett.*, **29**, 1846, doi:10.1029/2002GL014963, 2002.
- 30 Kaimal, J. C., Wyngaard, J. C., Izumi, Y., and Cote, O. R.: Spectral characteristics of surface-layer turbulence, *Quart. J. Roy. Meteorol. Soc.*, **98**, 563–589, 1972.

17727

- Keeling, R. F.: On the role of large bubbles in air-sea gas exchange and supersaturation in the ocean, *J. Mar. Res.*, **51**, 237–271, doi:10.1357/0022240933223800, 1993.
- Kelley, J. J. and Hood, D. W.: Carbon dioxide in the Pacific Ocean and Bering Sea: upwelling and mixing, *J. Geophys. Res.*, **76**, 745–752, doi:10.1029/JC076i003p00745, 1971.
- 5 Largier, J. L., Lawrence, C. A., Roughan, M., Kaplan, D. M., Dever, E. P., Dorman, C. E., Kudela, R. M., Bollens, S. M., Wilkerson, F. P., and Dugdale, R. C.: WEST: a Northern California study of the role of wind-driven transport in the productivity of coastal plankton communities, *Deep-Sea Res. II*, **53**, 2833–2849, 2006.
- Lendt, R., Thomas, H., Hupe, A., and Ittekkot, V.: Response of the near-surface carbonate system of the Northwestern Arabian Sea to the southwest monsoon and related biological forcing, *J. Geophys. Res.*, **108**, 3222, doi:10.1029/2000JC000771, 2003.
- 10 Lentz, S. J.: The surface boundary layer in coastal upwelling regions, *J. Phys. Oceanogr.*, **22**, 1517–1539, 1992.
- Nightingale, P. D., Malin, G., Law, C. S., Watson, A. J., Liss, P. S., Liddicoat, M. I., Boutin, J., and Upstill-Goddard, R. C.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, **14**, 373–387, doi:10.1029/1999GB900091, 2000.
- 15 Papale, D., Reichstein, M., Aubinet, M., Canfora, E., Bernhofer, C., Kutsch, W., Longdoz, B., Rambal, S., Valentini, R., Vesala, T., and Yakir, D.: Towards a standardized processing of Net Ecosystem Exchange measured with eddy covariance technique: algorithms and uncertainty estimation, *Biogeosciences*, **3**, 571–583, doi:10.5194/bg-3-571-2006, 2006.
- Prytherch, J., Yelland, M. J., Pascal, R. W., Moat, B. I., Skjelvan, I., and Neill, C. C.: Direct measurements of the CO_2 flux over the ocean: development of a novel method, *Geophys. Res. Lett.*, **37**, L03607, 2010.
- 25 Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T. H., Kozyr, A., Ono, T., Rios, A. F.: The oceanic sink for anthropogenic CO_2 , *Science*, **305**, 367–371, doi:10.1126/science.1097403, 2004.
- Sachs, L.: *Angewandte Statistik: Anwendung Statistischer Methoden*, Springer, Berlin, 1996.
- 30 Sarmiento, J. L. and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton University Press, 2006.

17728

- Schuepp, P. H., Leclerc, M. Y., MacPherson, J. I., and Desjardins, R. L.: Footprint prediction of scalar fluxes from analytical solutions of the diffusion equation, *Bound.-Lay. Meteorol.*, 50, 355–373, 1990.
- Simpson, J. J. and Zirino, A.: Biological control of pH in the Peruvian coastal upwelling area, *Deep-Sea Res.*, 27, 733–743, 1980.
- Sweeney, C., Gloor, E., Jacobson, A. R., Key, R. M., McKinley, G., Sarmiento, J. L., and Wanninkhof, R.: Constraining global air-sea gas exchange for CO₂ with recent bomb ¹⁴C measurements, *Global Biogeochem. Cy.*, 21, GB2015, doi:10.1029/2006GB002784, 2007.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Kortzinger, A., Steinhoff, T., Hoppema, M., Okafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bate, N. R., and de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO₂, and net sea–air CO₂ flux over the global oceans, *Deep-Sea Res. II*, 56, 554–577, doi:10.1016/j.dsr2.2008.12.009, 2009.
- Torres, R., Turner, D. R., Silva, N., and Rutllant, J.: High short-term variability of CO₂ fluxes during an upwelling event off the Chilean coast at 30° S, *Deep-Sea Res. Pt. I*, 46, 1161–1179, doi:10.1016/S0967-0637(99)00003-5, 1999.
- Wanninkhof, R.: Relationship between wind speed and gas exchange, *J. Geophys. Res.*, 97, 7373–7382, 1992.
- Wanninkhof, R. and McGillis, W. R.: A cubic relationship between air-sea CO₂ exchange and wind speed, *Geophys. Res. Lett.*, 26, 1889–1892, doi:10.1029/1999GL900363, 1999.
- Webb, E. K., Pearman, G. I., and Leuning, R.: Correction of flux measurements for density effects due to heat and water vapour transfer, *Quart. J. Roy. Meteorol. Soc.*, 106, 85–100, doi:10.1002/qj.49710644707, 1980.
- Weiss, R. F.: Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, 2, 203–215, doi:10.1016/0304-4203(74)90015-2, 1974.
- Wilkerson, F. P., Lassiter, A. M., Dugdale, R. C., Marchi, A., and Hogue, V. E.: The phytoplankton bloom response to wind events and upwelled nutrients during the CoOP WEST study, *Deep-Sea Res. II*, 53, 3023–3048, 2006.

17729

Table 1. Data summary.

Parameter	Date range	Location	Position
Eddy covariance tower			
CO ₂ flux			
Friction velocity	28 May–5 Jun 2007,	Bodega Bay,	38°20' N,
Wind speed	31 Aug–25 Sep 2008	CA	123°04' W
Wind direction			
BML meteorological tower			
Air temperature			
Wind speed	1988–2011	Bodega Bay,	38°19' N,
Wind direction		CA	123°04' W
PPFD			
BML buoy			
Sea surface temperature			
Salinity	1988–2011	Near Bodega	38°18' N,
Chlorophyll density	2007–2008	Bay, CA	123°04' W
pCO ₂	Nov 2010–Jul 2011		
NDBC 40613 buoy			
Wind speed			
Wind direction		Near Bodega	38°14' N,
AWS (calculated from WS and WD)	2007–2011	Bay, CA	123°18' W
Sea surface temperature			
Aircraft CO ₂ measurement			
Atmospheric CO ₂	Sep 2003–Jan 2012	Near Trinidad Head, CA	41°05', 124°17' W (~ 4 km in altitude)

17730




Fig. 1. Study site with the approximate 90 % flux footprint, and the wind rose of daily averaged wind directions and speeds (1990–2008) in Bodega Bay, California (38°18'58" N, 123°04'20" W).

17733




Fig. 2. Normalized cospectra of the vertical wind speed and the CO₂ mixing density, $f_{Co} < wC >$ **(a)** and the vertical wind speed and temperature, $f_{Co} < wT >$ **(b)**. The blue line indicates the average cospectra for $z/L > -0.1$ and the red line indicates the average for $z/L < -0.1$. The white line shows $-4/3$ at log-log scale.

17734




Fig. 3. CO₂ flux (a), friction velocity and wind speed (b), SST and PPFD (c), salinity and chlorophyll density (CHL) (d) measured from 28 May to 4 June 2007. All data were plotted every 30 min.

17735




Fig. 4. CO₂ flux (a), friction velocity and wind speed (b) SST and PPFD (c), salinity and chlorophyll density (CHL) (d) measured from 30 August to 24 September 2008. All data were plotted every 30 min.

17736




Fig. 5. Alongshore wind speed calculated from wind speed and direction at the NDBC 46013 buoy, with positive AWS (red dots) and negative AWS (blue dots) differentiated by color **(a)**, sea surface temperature at the NDBC 46013 buoy (SST_{N13}) **(b)**, and effective upwelling index (EUI) **(c)**. All data were averaged daily and plotted. The two boxes indicate the periods when eddy covariance measurements were available.

17737




Fig. 6. Hourly alongshore wind speed (AWS) calculated from wind speed and direction at the NDBC 46013 buoy (black squares), with the 30 point running average (black line) also shown **(a)**, sea surface temperature at the NDBC 46013 buoy (SST_{N13}) **(b)**, and effective upwelling index (EUI) **(c)**, salinity and SST **(d)**, $dpCO_2$ (pCO_2 –atmospheric CO_2) **(e)** and CO_2 flux estimated by the bulk method **(f)** from November 2010 to July 2011. **(a)–(c)** were plotted daily and **(d)–(f)** were plotted hourly.

17738




Fig. 7. Distribution of daily averaged $p\text{CO}_2$ in relation to salinity (psu) and seas surface temperature (SST) ($^{\circ}\text{C}$) measured from November 2010 to July 2011. A multiple linear regression was applied for $p\text{CO}_2$ against both salinity and SST ($R = 0.57$, $p < 0.001$).

17739




Fig. 8. Average seasonal variations of atmospheric CO₂ concentration between September 2003 to August 2012 obtained near Trinidad Head, CA and alongshore wind speed (AWS) between 1988 to 2011 (a) and average seasonal variations of salinity and sea surface temperature (SST) from 1988 to 2011 obtained from the BML buoy (b). Vertical bars show standard errors (SE). The average and SE for each month were calculated from the monthly averages available for all years.

17740




Fig. 9. Monthly averaged seasonal variations of DIC per unit area within the surface mixed layer (hDIC), monthly CO₂ flux ($F_{c,month}$), and monthly NPP (NPP_{month}) **(a)** and the depth of the surface mixed layer depth **(b)** estimated from 1988–2011 data. The vertical lines indicate standard deviations.