

Interactive comment on “Estimates of micro-, nano-, and picoplankton contributions to particle export in the northeast Pacific” by B. L. Mackinson et al.

Anonymous Referee #1

Received and published: 26 September 2014

In this manuscript Mackinson et al. use phytoplankton pigments combined with large-volume pump sampling, ^{234}Th deficiency measurements, and two sediment trap deployments to address the relative contributions of pico-, nano-, and microplankton to passive (sinking) carbon export in the northeast Pacific. This is an important topic, given the hypothesis proposed by Richardson & Jackson (2007) that production stemming from picoplankton may dominate the flux of particulate material in the ocean. To date, there have only been (to my knowledge) three published field studies specifically designed to address this hypothesis, thus this new dataset is quite valuable. There are some methodological issues with the authors' approach (as there will be with any approach to tackling this difficult problem) that I would like to see the authors address

C5484

more directly and succinctly. However, I believe that this is a nice manuscript overall, and is certainly worthy of being published (with moderate revision). Below please find some major and minor issues that I believe should be dealt with:

Major Concerns:

There is no perfect way to address the contribution of picoplankton to particle export, because (1) there is no perfect way to measure export and (2) the source of the exported material is often obscured by grazing, aggregation, physical breakdown, and microbial remineralization processes. The authors have chosen to use a combination of ^{234}Th and pigments as their primary methods for this study. It is very important that they succinctly outline the problems with these methods:

1) ^{234}Th – The two primary methods for measuring vertical carbon fluxes in the field are ^{234}Th and sediment traps. Each has issues (hydrodynamic and degradation for sediment traps; steady-state assumptions and variable C: ^{234}Th ratios for ^{234}Th). For practical reasons, the authors rely very heavily on ^{234}Th measurements for this manuscript (although it is very nice that they have two sediment trap deployments that largely agree with the ^{234}Th -based results). Unfortunately, for their particular question ^{234}Th is inferior to sediment traps. It is incredibly important to note that, when determining the relative contributions of pico-, nano-, and microplankton to export using the authors' approach, the ^{234}Th measurements are COMPLETELY IRRELEVANT. The relative contribution of different size classes to export is completely determined by their pigment ratios in the >50-micron large-volume samples. This needs to be explicitly stated. The authors make the (defensible) assumption that these large particles (likely aggregates) collected by the pump at depth are representative of sinking material. As the authors note at the end of their discussion, however, the pumps do not sample fecal pellets effectively, and fecal pellets may both contribute significantly to export and represent different ratios of micro/nano/picoplankton than the aggregates sampled by the authors.

C5485

2) Pigments – the other half of the authors' primary methodology is pigment analysis to determine the composition of the sinking material. There are a few issues with using pigments for this question (though nucleic acids, the other primary option, may have even greater issues). One issue that the authors have is that indicator pigments do not map perfectly into size classes. It would be nice to see the authors discuss the correlation between different pigments and size-fractionated chlorophyll. Another significant issue is differential pigment degradation. There is no a priori reason to assume that different indicator pigments are degraded at the same rate, especially when considering that picoplankton (primarily grazed by protozoans) and microplankton (largely grazed by mesozooplankton) likely undergo significantly different processes prior to being incorporated in aggregates or fecal pellets. C:pigment ratios may vary significantly with depth and inconsistently between taxa.

Just to reiterate. These are significant issues, but I do not believe that they invalidate the core results of the study. Many of these issues are discussed already at various places in the manuscript. However, given the incredible importance of these issues I believe that they deserve a dedicated section in the discussion. The fact that the proportional role of picoplankton to export (at stations without sediment traps) is determined solely from their ratios in pump samples WITHOUT EVEN NEEDING THE THORIUM MEASUREMENTS should also be explicitly stated in the methods or beginning of the results.

Other Issues:

One of the strengths of this study is that the authors measured export and the contribution of pico-/nano-/microplankton at multiple stations and several different seasons. Given these measurements, it would be nice to see them discuss whether or not there are correlations between export and the contributions of pico/microplankton to surface biomass. Ultimately the Richardson & Jackson hypothesis is important because it pertains to the question of whether or not we would expect export to decrease in a more oligotrophic future ocean. In addition to looking directly at the proportion of picoplank-

C5486

ton in export, the authors can also look at whether or not a picoplankton dominated ocean has less export than a microplankton dominated ocean.

It would be nice to see the authors use pigment:carbon estimates to put together a back-of-the-envelope calculation of the ratio of phytoplankton carbon: total carbon in the deep LV pump samples and sediment trap samples. Do the pigments that the authors measured comprise most of the organic carbon that is being exported or is a significant amount of the sediment trap material unaccounted for?

On a similar note, although the authors do not give any methodological details for their fluorometric chlorophylls (this is an oversight that should be corrected – did they use the acidification method of Strickler & Parsons?) if they used the acidification method, they can get an estimate of phaeopigment concentration in the sediment trap as well. Although chlorophyll is not quantitatively converted to phaeopigments in mesozooplankton guts, phaeopigment concentration can still give an estimate of the proportion of flux that may be due to mesozooplankton fecal pellets and hence likely originating from microplankton but not showing up as microplankton indicator pigments.

Since the authors talk about standing stocks sampled by SV and LV (and these two measurements do not agree) it is very important that they explicitly state when they are using standing stocks derived from SV or LV samples both in the text and in figures.

Figure 8 shows microplankton indicator pigments often dominating even the 1-10 micron size fraction. This should probably be discussed since it clearly illustrates the issues with using phytoplankton pigments as indicators of size.

This paper uses a lot of non-standard abbreviations. I would recommend that the authors add a table at the beginning of the manuscript that lists all their abbreviations so that readers don't have to hunt through the text to find out what mPF or PTh mean. Also, PTh is a strange choice for Th flux, since it could easily be mistaken to mean particulate thorium.

C5487

p. 12633 line 6 – Stukel & Landry 2010, and Lomas & Moran 2011 do not state that picoplankton export is proportional to biomass, but rather that its proportional contribution to export is less than to biomass, but still significant. Amacher et al. (2009, DSR 156(12): 2206-2215) and Stukel et al. (2013, PinO 112-113: 49-59) should probably be cited as other studies that have attempted to directly assess the proportion of picoplankton in export. Amacher used nucleic acids and found a significant role for picos at ESTOC and Stukel used pigments and found a less than proportional role for picos in the Costa Rica Dome.

p. 12636 line 5 – Although there is nothing the authors can do about it at this point (and it probably isn't a huge problem), best practices with thorium involve an acidification step with HNO₃ to bring pH < 2 before spiking with the tracer Th-230 (Pike et al. 2005, Journal of Radioanalytical and Nuclear Chemistry 263(2) 355-360). This brings all the naturally particle-associated Th-234 into the dissolved phase so that it can equilibrate with the added Th-230. Without this step it is possible that the yield of Th-234 (initially bound to naturally occurring particles and colloids) and the yield of Th-230 tracer will be different. Also, no methods for yield analysis are mentioned.

p. 12636 line 10 – “drying over” should be “drying oven”

p. 12636 line 25 - I do not see how mPF, nPF, and pPF are calculated. Is it simply the ratio of the summed indicator pigments that are believed to be responsible for each size class? This seems to be implied by Figure S2 which shows a 1:1 correlation between total indicator pigment and Chl a. This is not, however, the best way to estimate mPF, nPF, and pPF, since different taxa of phytoplankton will have different ratios of indicator pigment : Chl a. A better approach would probably be to multiply each indicator pigment by a pigment:C ratio for the taxa that it represents and then summing these carbon contributions (perhaps using a CHEMTAX approach – e.g. Mackey et al. 1996; MEPS 144: 265-283). Note that this should not change their primary results (the comparison of the proportions of picoplankton to biomass and export), but it would change the total proportion of picoplankton in biomass and export.

C5488

p. 12643 line 23 – The authors have not stated how they determined ²³⁸U concentrations. Did they use the Owens et al. (2011 MarChem 127(1-4):31-39) or Chen et al. (1986, EarthPlanSciLett 80: 241-251) relationships or did they actually measure it directly?

p. 12644 equation 1 – The equation shown neglects the effects of upwelling or downwelling which can (at times) lead to a significant error in simple thorium export models (see Savoye et al. 2006, MarChem 100(3-4): 234-249). Since the authors (like most who study thorium) have no way of estimating upwelling it is acceptable that they have neglected it, however, this term should definitely be included in equation 1 and the rationale behind ignoring it should be given.

p. 12645 line 19 – “decreasee” should be “decrease”

p. 12645 line 25 – the POC/²³⁴Th ratio in traps is substantially higher than the ratio of particles collected by pumps. This is significant since it suggests that there may be a substantial amount of sinking material that is not being collected by the pumps. Such a situation could arise if there is a rapidly sinking particle fraction (perhaps fecal pellets) that has a high C:Th ratio that is similar to the higher bulk C:Th ratio found in surface water as well as a slowly sinking particle fraction that has time to equilibrate with lower bulk C:Th ratios at depth. This should be discussed as it bears on the question of whether or not the pump samples pigment ratios are representative of all sinking material.

p. 12646 line 4 – I cannot find a Fig. 11c

p. 12648 lines 15-20 – the authors point out that much of picoplankton production will be grazed and that the grazing pathway will not show up with their methodology. This is true. However, they then suggest that this may lead to an underestimate of the role of picoplankton by their methodology. This is not true. While picoplankton can certainly be exported by grazing pathways, they are most likely exported after transfer through one (or two) protozoan grazing steps which will degrade a significant fraction

C5489

of their carbon before the protozoans are grazed by large fecal pellet-producing mesozooplankton. Microplankton, by contrast are much more likely to be grazed directly by mesozooplankton, hence contributing a significantly greater fraction of their biomass to export. Thus the grazing pathway (which was not assessed by the methodology used by the authors) is actually much more likely to underestimate the contribution of microplankton than picoplankton. This is particularly important since, as the authors note at the end of the discussion, fecal pellet export is substantially greater than algal aggregate export in their study region.

Table 1 – I find this table to be slightly confusing. It might be easier to read if there were borders around the cells to show which samples go with which cruises.

Table 2 – Please define all abbreviations so readers don't have to hunt through the text.

Figure 2 – I would recommend only showing plots down to 200 m (since I believe that is the deepest depth of the authors' samples) in the interest of making upper water column patterns more visible.

Figure 9 – This is an important figure with a lot of data crammed into it. Is it possible to make it a bit larger so that patterns are more visible?

Figure 10 - legend and figure, there is no symbol for Harrison, 2002

Interactive comment on Biogeosciences Discuss., 11, 12631, 2014.