

Response to Editor:

Main comments

Based on the feedback from two reviewers and my own assessment, I believe the manuscript is worthy of publication. In particular, addressing the concerns of reviewer one about the error in the end member values is important to assessing the uncertainty in the mixing model. However, I have a few additional requests for clarification and editing in the text:

Response: We appreciate the editor's constructive comments and precious time for handling our manuscript. We have thoroughly considered all the comments, and made minor revisions as suggested.

Specific comments:

1. While you note that standard errors have been added, please note what those standard errors are in your response so that I can better evaluate the magnitude of uncertainty (given that it's not included in the current online version of the paper).

Response: We have added the standard errors in our revised version: “Based on our recent studies conducted in the Yanqi Basin (Wang et al., 2015; Zhang, 2013), C:N ratio was 22.1 ± 9.9 and 10 ± 1.8 , and $\delta^{13}\text{C}_{\text{org}}$ value was $-26.4 \pm 1.2\text{\textperthousand}$ and $-23.6 \pm 1.3\text{\textperthousand}$ for the native plants and surface soils around the lake, respectively (page 6, lines 13-16)” and “Lake POC and PON increased from $0.61 \pm 0.04 \text{ mg C L}^{-1}$ and $0.072 \pm 0.005 \text{ mg N L}^{-1}$ in spring to $0.70 \pm 0.16 \text{ mg C L}^{-1}$ and $0.088 \pm 0.02 \text{ mg N L}^{-1}$ in summer, and $\delta^{13}\text{C}_{\text{org}}$ value in POC was $-22.9 \pm 2.56\text{\textperthousand}$ in spring and $-23.5 \pm 0.38\text{\textperthousand}$ in summer (page 6, lines 19-22)”.

2. Introduction, line 21: change "actively" to "active"

Response: We have corrected.

3. Introduction, line 25: remove "the"

Response: Done.

4. Site Description, line 18-19: Please clarify what you mean when you say the max depth was 14 m, but the lake level was 1045 m when sampled (presumably that's relative to sea level?)

Response: This is a good point. We have revised as “It is the largest inland freshwater lake in China, which is about 55 km long from east to west and about 25 km wide from south to north, comprising a total lake surface area of approximately 1005 km^2 , with a maximum water depth of 14 m (Wu et al., 2013). The lake surface was 1045 m above sea level in 2012 when sampling was carried out (page 4, lines 10-14)”.

5. I would like to see a few sentences that state what the potential implications are

for variability in lake surface sediment sources. Does it affect the interpretation of C sources to the lake? Estimates of biological productivity?

Response: Yes, we agree. We have added “Our approach may have uncertainties in determining TOC sources. However, the uncertainties would be small given that the standard errors in $\delta^{13}\text{C}_{\text{org}}$ are small, and C:N ratios diff greatly between sources. Thus, our method will not affect the main conclusion in terms of TOC sources for the lake sediments (page 7, lines 1-4)”.

Reviewer #1:

Main comments

The manuscript by Yu et al. presents a detailed study about spatial distribution of sediment organic matter in Boston Lake, based on which they calculated the contributions of terrestrial plant, soil and lake plankton and evaluate the potential factors responsible for their spatial variability. I think this study address an important issue about widely used geochemical proxies (C/N and $\delta^{13}\text{C}$). Many studies applied C/N and $\delta^{13}\text{C}$ as organic matter source indicators without consideration of other factors such as hydrodynamic and mineral contents. Meanwhile, this manuscript is well written and its topic is suitable for Biogeosciences. I have several concerns, which should be addressed before publish.

Specific comments:

1. Page 7: The authors attributed sediment organic matter to three endmember, high plant, soil and lake plankton. I think it is better to say “terrestrial plants” instead of high plant. High plant (or higher plant) is not an accurate definition because many higher plants such as emerged, floating and submerged plants can be quite abundant in some lakes. In this manuscript, the endmember value for high plant is apparently from land plants.

Response: Yes, we agree. In the revision, we have replaced the “high plant” with “terrestrial plant”.

2. Page 7: for end member values, the authors cited the data from Zhang et al. (2013). I did not check their raw data, but it is kind strange they only provided average values. I believe there are different types of land plants and soils, and therefore, the C/N and $\delta^{13}\text{C}$ should vary with species and sampling sites. In my opinion, those data should be reported with standard errors. Otherwise, the readers can not estimate how much uncertainty of their three end member mixing model. A similar problem exists for the concentrations of POC and PON and $\delta^{13}\text{C}$ values in different seasons. Without SE, we can't judge if those seasonal differences are significant or not.

Response: This is a good point. We agree and have added the standard errors (please see page 6, lines 13-16&19-22).

3. Page 11: delete “as known” since this phrase does not provide any useful information

Response: Done.

4. Figure 2 and other figures: the font size is too small.

Response: Thanks for the comment. We have reproduced all the figures using a larger font size in the revised version.

The list of all relevant changes made in the manuscript:

- 1) We have addressed all the comments made by the editor and reviewers.
- 2) We have updated the authors' information (replaced H. Y. Lan with X.Q. Liu).
- 3) Page 3, line 25: delete “investigation of the dramatic”.
- 4) Page 17, line 2: replace “Doctor of Physical Geography” with “PhD thesis”.
- 5) Page 18, table 1 footnote: delete “and clay, silt and sand fractions (%)".
- 6) Page 25, figure 7 caption: delete “(a)” and “(b)”.
- 7) We have added “the Sino-German project (GZ867) and National pioneer project (XDA05020202)” in the Acknowledgments.
- 8) We also have made some changes the references following the BG requirements for publication.

1 **Spatial distribution and sources of organic carbon in the**
2 **surface sediment of the Boston Lake, China**

4 Z. T. Yu^{1,2,3}, X. J. Wang^{1,4,5}, E. L. Zhang⁵⁶, C. Y. Zhao¹, H. Y. Lan² and X.
5 Q. Liu⁷

6 [1] State¹State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of
7 Ecology and Geography, Chinese Academy of Sciences, 830011 Urumqi, P.R.China+

8 [2]College²College of Resources and Environmental Sciences, Xinjiang University,
9 830046 Urumqi, P.R.China+

10 [3]University³University of Chinese Academy of Sciences, 100049 Beijing, P.R.
11 China+

12 [4]College of Global Change and Earth System Science, Beijing Normal University,
13 100875 Beijing, China

14 [5]Earth System Science Interdisciplinary Center, University of Maryland, College Park,
15 20740 MD, USA+

16 [5]State⁶State Key Laboratory of Lake Science and Environment, Nanjing Institute
17 of Geography and Limnology, Chinese Academy of Sciences, 210008 Nanjing, P.R.
18 China+

19 [7]College of Resource Environment and Tourism, Capital Normal University,
20 100048 Beijing, China

22 Correspondence to: X. J. Wang ([wwwang@essie.umd.edu](mailto:xwang@essie.umd.edu)xwang@bnu.edu.cn)

24 **Abstract**

25 Lake sediment is an important carbon reservoir. However, little is known on the
26 dynamics and sources of sediment organic carbon in the Boston Lake. We collected

1 13 surface (0-2cm) sediment samples in the Boston Lake and analyzed total organic
2 carbon (TOC), total nitrogen (TN), stable carbon isotopic composition in TOC
3 ($\delta^{13}\text{C}_{\text{org}}$) and grain size. We found a large spatial variability in TOC content (1.8-4.4%)
4 and $\delta^{13}\text{C}_{\text{org}}$ value (-26.77‰ to -23.98‰). Using a three end member mixing model
5 with measured TOC:TN ratio and $\delta^{13}\text{C}_{\text{org}}$, we estimated that 54-90% of TOC was
6 from autochthonous sources. Higher TOC content (>3.7%) was found in the east and
7 central-north sections and near the mouth of the Kaidu River, which was attributable
8 to allochthonous, autochthonous plus allochthonous, and autochthonous sources,
9 respectively. The lowest TOC content was found in the mid-west section, which might
10 be a result of high kinetic energy levels. Our study indicated that the spatial
11 distribution of sediment TOC in the Boston Lake was influenced by multiple and
12 complex processes.

13

14 **1 Introduction**

15 Inland water bodies such as rivers and lakes are unique components on the Earth. In
16 spite of their relatively small coverage (Downing et al., 2006), lakes often receive a
17 large amount of terrestrial materials from the watersheds (Battin et al., 2009;Anderson
18 et al., 2013), and store a significant amount of carbon in the sediments (Ferland et al.,
19 2012;Tranvik et al., 2009). Thus, inland lakes may play an important role in the
20 terrestrial carbon cycle. Compared to the oceans, lakes have ~~actively~~active
21 biogeochemical processes with stronger “biological pump”, which often leads to
22 higher sedimentation rates and a large amount of organic carbon (OC) burial at the
23 bottom of lakes (Dean and Gorham, 1998).

24 There have been a number of studies from ~~the~~ North America (Dean and Gorham,
25 1998), West Europe (Bechtel and Schubert, 2009;Woszczyk et al., 2011), East Asia
26 (Khim et al., 2005;Wang et al., 2012) and other regions (Dunn et al., 2008), showing
27 large spatial variability in total organic carbon (TOC) of lake sediment. The
28 magnitude of TOC in surface sediment may depend on many factors, including
29 column water productivity, terrestrial inputs of organic materials, properties of

1 sediment, and rate of microbial activity (Burone et al., 2003;Gireeshkumar et al.,
2 2013). Among them, contributions of autochthonous and allochthonous sources have
3 direct impacts on the spatial distribution, which vary largely across regions (Bechtel
4 and Schubert, 2009;Anderson et al., 2009), partly due to differences in lake
5 productivity and morphology (Barnes and Barnes, 1978). In general, lakes with high
6 productivity have more autochthonous TOC, but lakes with low productivity mainly
7 allochthonous TOC (Dean and Gorham, 1998). There is evidence of littoral sources of
8 TOC in small and shallow lakes, but autochthonous sources, derived from planktonic
9 organisms, in larger and deeper lakes, especially fjord lakes (Shanahan et al.,
10 2013;Sifeddine et al., 2011;Barnes and Barnes, 1978).

11 A number of techniques have been applied to quantify different sources of sediment
12 TOC (Fang et al., 2014;Hanson et al., 2014;Meyers and Ishiwatari, 1993;Bechtel and
13 Schubert, 2009). One of the common approaches is to use two or three end-member
14 mixing models with combined use of TOC to total nitrogen (TN) ratio (C:N) and
15 stable carbon isotope in organic material ($\delta^{13}\text{C}_{\text{org}}$) (Rumolo et al., 2011;Yu et al.,
16 2010;Liu and Kao, 2007). It is well known that there are large differences in C:N ratio
17 and $^{13}\text{C}_{\text{org}}$ value between exogenous and endogenous organic materials (Brodie et al.,
18 2011;Kaushal and Binford, 1999). For example, aqueous organic matters have low
19 C:N ratios (4-10) (Meyers, 2003) whereas vascular land plants have much higher C:N
20 ratios (>20) (Rumolo et al., 2011;Lamb et al., 2004;Sifeddine et al., 2011). On the
21 other hand, due to the difference in isotopic fractionation during photosynthesis,
22 $\delta^{13}\text{C}_{\text{org}}$ value is more negative (ranging from -33‰ to -22‰) in terrestrial C₃ plants
23 (Pancost and Boot, 2004;Wang et al., 2013) and lake plankton (Bertrand et al.,
24 2010;Vuorio et al., 2006) than in C₄ plants (ranging from -16‰ to -9‰) (Pancost and
25 Boot, 2004;Wang et al., 2013).

26 Boston Lake, as the largest lake in Xinjiang of China, is a typical place for studying
27 lake carbon cycle. Previous studies have provided evaluations on water quality (Wu et
28 al., 2013), ~~investigation of the dramatic~~ changes in lake level (Guo et al., 2014), and
29 the controlling factors of carbon and oxygen isotopic composition of surface sediment

1 carbonate (Zhang et al., 2009). A recent study indicated that particulate organic
2 carbon (POC) variability in the water column was affected by allochthonous sources
3 in the Boston Lake (Wang et al., 2014). However, little has been done to assess the
4 dynamics and sources of sediment TOC in the Boston Lake. Therefore, this study was
5 designed to evaluate the spatial distributions of major physical and biogeochemical
6 parameters in the surface sediment, and to quantify the contributions of various
7 sources to the sediment TOC in the Boston Lake.

8

9 **2 Materials and Methods**

10 **2.1 Site description**

11 Boston Lake ($41^{\circ}32' \sim 42^{\circ}14'N$, $86^{\circ}19' \sim 87^{\circ}26'E$) is located in the lowest part of the
12 intermontane Yanqi Basin between the Taklimakan Desert and Tianshan Mountains,
13 Northwest China (Figure 1). It is the largest inland freshwater lake in XinjiangChina,
14 which is about 55 km long from east to west and about 25 km wide from south to
15 north, comprising a total lake surface area of approximately 1005 km^2 , with a
16 maximum water depth of 14 m (Wu et al., 2013). The lake levelsurface was 1045 m
17 above sea level in 2012 when sampling was carried out. The lake lies in the center of
18 the Eurasian Continent and is influenced by a temperate continental climate. The
19 mean annual air temperature is approximately 8.3°C , the mean annual precipitation
20 approximately 65 mm and the mean annual evaporation approximately 1881 mm
21 (Zhang et al., 2009). Winds come mainly from the southwest, indicating dominant
22 influence by the westerly throughout the summer season. Lake water input mainly
23 comes from the Kaidu River that is supplied by melting ice, precipitation and
24 groundwater, whereas water output includes outflow (57%) via the Peacock River and
25 evaporation (43%) (Guo et al., 2014). There are also small seasonal rivers (mainly
26 during flood seasons), e.g., the Huangshui River and Qinhshui River near the
27 northwest of the lake.

1 **2.2 Field sampling and analyses**

2 For the present study, a Kajak gravity corer was used to collect surface sediments
3 from 13 sites in the main section of the Boston Lake in August 2012 (Figure 1). The
4 sampling sites covered most parts of the lake, with water depths ranging from 3 m to
5 14 m. The sediment cores were carefully extruded and the top 2 cm sections were
6 sliced into 1-cm and placed in polyethylene bags that were kept on ice in a cooler
7 during transport and before analyses.

8 Following Liu et al. (2014), each sediment sample (~0.5 g) was pretreated, in a water
9 bath (between 60 and 80 °C), with 10-20 ml of 30% H₂O₂ to remove organic matter,
10 then with 10-15ml of 10% HCl to remove carbonates. The samples were then mixed
11 with 2000 ml of deionized water, and centrifuged after 24 hours of standing. The
12 solids were dispersed with 10 ml of 0.05 M (NaPO₃)₆, then analyzed for grain size,
13 using a Malvern Mastersizer 2000 laser grain size analyzer at the State Key
14 Laboratory of Lake Science and Environment (SKLLSE), Nanjing Institute of
15 Geography and Limnology, Chinese Academy of Sciences (CAS). The Malvern
16 Mastersizer 2000 automatically outputs the median diameter d(0.5) (μm), the diameter
17 at the 50th percentile of the distribution, and the percentages of clay (< 2 μm), silt
18 (2-64 μm) and sand (> 64 μm) fractions.

19 Sediment C and N contents were measured using an Elemental Analyzer 3000 (Euro
20 Vector, Italy) at the SKLLSE, Nanjing Institute of Geography and Limnology, CAS.
21 All samples were freeze-dried and ground into a fine powder, then placed in tin
22 capsules, weighed and packed carefully, according to Eksperiandova et al. (2011). For
23 the analysis of TOC, each sample (~ 0.3 g) was pretreated with 5-10 ml 2M HCl for
24 24h at room temperature to remove carbonate, dried overnight at 40-50 °C, then
25 analyzed for C content using the Elemental Analyzer.

26 For the analyses of δ¹³C_{org}, approximately 0.2 g of the freeze-dried sediment sample
27 was pretreated with 5-10 ml 2M HCl for 24 h at room temperature to remove
28 carbonate, and then rinsed to a pH of approximately 7 with deionized water and dried
29 at 40-50 °C (Liu et al., 2013). The pre-treated samples were combusted in a Thermo

1 elemental analyzer integrated with an isotope ratio mass spectrometer (Delta Plus XP,
2 Thermo Finnigan MAT, Germany). Isotopic data were reported in delta notation
3 relative to the Vienna Pee Dee Belemnite (VPDB).

4 **2.3 Calculations of TOC sources**

5 We applied a three end-member mixing model (Liu and Kao, 2007) to quantify the
6 contributions (f) of three sources (i.e., soil, ~~high terrestrial~~ plant and lake plankton,
7 denoted by 1, 2 and 3, respectively):

$$8 \delta = f_1\delta_1 + f_2\delta_2 + f_3\delta_3 \quad (1)$$

$$9 r = f_1r_1 + f_2r_2 + f_3r_3 \quad (2)$$

$$10 1 = f_1 + f_2 + f_3 \quad (3)$$

11 where δ and r were $\delta^{13}\text{C}_{\text{org}}$ value and C:N ratio, respectively.

12 Given that there were limited crops around the lake and most crops' growing season
13 was less than five months each year, we assumed that native plants, mainly reed
14 (*Phragmites australis* (Cav.) Trin. ex Steud), Manaplant Alhagi (*Alhagi sparsifolia*
15 *Shap*) and Achnatherum splendens (*Achnatherum splendens* (Trin.) Nevski), were
16 responsible for ~~high terrestrial~~ plant's contribution. Based on ~~our~~ recent ~~study~~ studies
17 conducted in the Yanqi Basin (Wang et al., 2015; Zhang, 2013), ~~average~~ C:N ratio was
18 22.1 ± 9.9 and 10 ± 1.8 , and ~~mean~~ $\delta^{13}\text{C}_{\text{org}}$ value was -26.4 $\pm 1.2\%$ and -23.6 $\pm 1.3\%$ for the
19 native plants and surface soils around the lake, respectively. We used ~~these~~ the values
20 as the end-members for the mixing model.

21 We measured POC, particulate organic nitrogen (PON) and $\delta^{13}\text{C}_{\text{org}}$ in POC in the
22 water column of the Boston Lake (Wang et al., 2014). Lake ~~average~~ POC and PON
23 increased from 0.61 ± 0.04 mg C L⁻¹ and 0.072 ± 0.005 mg N L⁻¹ in spring to 0.70 ± 0.16
24 mg C L⁻¹ and 0.088 ± 0.02 mg N L⁻¹ in summer, and $\delta^{13}\text{C}_{\text{org}}$ value in POC was
25 -22.9 $\pm 2.56\%$ in spring and -23.5 $\pm 0.38\%$ in summer. It is reasonable to assume that
26 the seasonal changes were resulted from the production of lake plankton. Accordingly,

1 we estimated that lake plankton (including phytoplankton and zooplankton) would
2 have a C:N ratio of 5.3 and $\delta^{13}\text{C}_{\text{org}}$ value of -27.7‰, and used these values as the
3 end-members for the mixing model.

4 Our approach may have uncertainties in determining TOC sources. However, the
5 uncertainties would be small given that the standard errors in $\delta^{13}\text{C}_{\text{org}}$ are small, and
6 C:N ratios diff greatly between sources. Thus, our method will not affect the main
7 conclusion in terms of TOC sources for the lake sediments.

8 **2.4 Statistical methods and mapping**

9 Correlation analyses were performed using the SPSS Statistics 19 for Windows.
10 Spatial distribution maps were produced using Surfer 9.0 (Golden Software Inc.) and
11 the Kriging method of gridding was used for data interpolation.

12

13 **3 Results**

14 **3.1 Physical characteristics**

15 Figure 2 showed the spatial distributions of the main granulometric variables of the
16 surface sediment. In general, clay content was low (6-17%), showing relatively higher
17 values in the southern part than in the northern part. The highest clay content was
18 found in the southwest, and the lowest in the northwest section. On the other hand, silt
19 content was much high (greater than 80%) with clearly higher values near the mouths
20 of the Kaidu River (southwest) and Huangshui River (northwest). The lowest content
21 of silt was found in the mid-west, between the rivers' mouths, where sand content was
22 highest (Figure 2c). As expected, the spatial distribution of d(0.5) was similar to that
23 of sand, showing the highest values in the mid-west section, indicating strong
24 hydrodynamic effect in this area.

1 **3.2 Spatial distribution of TOC, TN, C:N and $\delta^{13}\text{C}_{\text{org}}$**

2 Concentration of TOC was highly variable, with higher values (4.3-4.4%) found in the
3 northern and eastern sections of the lake (Figure 3a). There was also high
4 concentration of TOC (4.1-4.2%) near the mouth of the Kaidu River (southwest). On
5 the other hand, lower TOC concentration (1.8-2.4%) was observed in the mid-west
6 section. Similarly, TN concentration (ranging from 0.28% to 0.68%) was lowest
7 concentration in the mid-west and highest in the northwest and east sections (Figure
8 3b). Overall, the spatial distribution of TN was similar to that of TOC. The exception
9 was in the northwest area that had high TN value, but low TOC concentration.

10 Figure 4a showed a large spatial variability in the C:N ratio with a range from 4.6 to
11 8.6. In general, C:N ratio was higher in the central part relative to other parts. The
12 highest C:N ratio was found in the mid-west, and the lowest found in the northwest
13 area. The $\delta^{13}\text{C}_{\text{org}}$ values ranged from -26.77‰ to -23.98‰ (Figure 4b). The most
14 negative value was observed in the area of 41.9-42°N and 86.9-87°E, and the least
15 negative value near the mouth of the Huangshui River (northwest). Overall, values of
16 $\delta^{13}\text{C}_{\text{org}}$ were more negative in the eastern and central parts than in the northwestern
17 and southwestern parts.

18 **3.3 Contributions of different sources**

19 Using the three end member mixing model, we calculated the contributions of
20 autochthonous and allochthonous sources to the surface sediment TOC. As shown in
21 Figure 5a, the contribution of lake plankton ranged from 54% to 90%, with the
22 highest in the western shallow lake area, and the lowest in the southern and eastern
23 deep lake area. The contribution of soils varied between 10% and 40%, with the
24 highest in the southeast and central south area (Figure 5b). Apparently, the
25 contribution from native plants was extremely low (< 4%), with only a few sites
26 showing values of 10-12% (Figure 5c). On average, the contributions from lake
27 plankton, soils and native plants were 66%, 30% and 4%, respectively.

1 There were large differences in the spatial distributions of TOC between the
2 autochthonous and allochthonous sources. Autochthonous TOC revealed highest value
3 (~3.5%) near the mouth of the Kaidu River and lowest (~1.5%) in the mid-west of the
4 lake (Figure 6a). For the area east of 87°E, autochthonous TOC showed a clear
5 increase from south to north. On the other hand, there was an apparent elevation in the
6 allochthonous TOC, from 0.5% in the west to 1.9% in the east (Figure 6b).

7

8 **4 Discussion**

9 The concentration of TOC in the surface sediment of the Boston Lake ranged from
10 1.8-4.4%, which was relatively higher than those (0.2-2%) in the Tibetan Plateau
11 (Lami et al., 2010; Wang et al., 2012) and Yangtze floodplain (Wu et al., 2007; Dong et
12 al., 2012), but much lower than those (5-13%) in the lakes of the Yunnan-Guizhou
13 Plateau (Zhu et al., 2013; Wu et al., 2012). Low TOC contents in the Tibetan Plateau
14 lakes were a consequence of low biological productivity owing to the high altitude
15 and low temperature (Lami et al., 2010). Although lakes in the Yangtze floodplain had
16 higher productivity in the water column due to eutrophication (Qin and Zhu, 2006),
17 most of them were shallow lakes that were subject to frequent turbulence and
18 resuspension of sediments (Qin et al., 2006). In addition, warm-humid climate in the
19 Yangtze floodplain could promote decomposition of POC in the water column and
20 TOC in the sediments (Gudasz et al., 2010), which led to less TOC storage in the
21 surface sediments. On the other hand, lakes in the Yunnan-Guizhou Plateau were deep
22 with higher lake productivity, which had favorable TOC burial conditions (Jiang and
23 Huang, 2004).

24 Sediment organic compounds are either of terrestrial origins or derived from
25 phytoplankton and zooplankton remains and feces (Meyers, 2003; Meyers and
26 Ishiwatari, 1993; Barnes and Barnes, 1978). A number of studies have demonstrated
27 that TOC in small and shallow lakes are attributable to allochthonous sources, but
28 TOC in larger and deeper lakes to autochthonous sources that are derived from
29 planktonic organisms (Shanahan et al., 2013; Sifeddine et al., 2011; Barnes and Barnes,

1 1978). Our analyses showed that the majority of TOC was autochthonous in the
2 surface sediment of the Boston Lake. We also found a significant negative relationship
3 between TOC and dry bulk density (Table 1), confirming that higher TOC (with
4 lighter weight) would be a result of sedimentation of non-terrestrial organic materials.
5 Our study demonstrated large spatial variability in the TOC of the surface sediment in
6 the Boston Lake, with higher values in the central north and east sections and near the
7 mouth of the Kaidu River, but lower values in the west section and mid-south section
8 (Figure 3a). Further analyses showed that the highest autochthonous TOC was found
9 near the mouth of the Kaidu River and the highest allochthonous TOC in the east
10 section (Figure 6). There is evidence of high productivity near the sources of nutrients,
11 such as estuaries owing to extra nutrient input from riverine (Deng et al., 2006; Lin et
12 al., 2002). Nutrient conditions in the Boston Lake may be largely affected by the
13 transportation of the Kaidu River, which has a significant decline from the mouth to
14 the east section. Similar finding was also observed in the Nam Co Lake (Wang et al.,
15 2012).

16 | ~~As known,~~ TOC burial in sediments is a result of sedimentation of POC. Here, we
17 compared the spatial pattern of autochthonous TOC in the 0-1 cm sediment with the
18 summer POC reported by Wang et al. (Wang et al., 2014), which showed the highest
19 values of both variables near the mouth of the Kaidu River (Figure 7). Statistical
20 analysis indicated that the correlation was not significant ($r = 0.14$, $P > 0.1$, Table 1)
21 between these two variables, which might be due to the mismatch in the locations of
22 the lowest values. As shown in Figure 2&3, coarse particle components were
23 dominant in the mid-west section where TOC was the lowest. Table 1 also illustrated
24 that TOC had a negative relationship with sand content and $d(0.5)$. Usually, in a
25 relatively close hydraulic equivalence, coarser sediment particles indicated a stronger
26 water energy environment (Jin et al., 2006; Molinaroli et al., 2009). These analyses
27 indicated that the relative lower TOC values in the mid-west section of the Boston
28 Lake were attributable to both the lower POC in the water column and higher kinetic
29 energy level.

1 The magnitudes and spatial distribution of TOC in lake sediment may reflect multiple,
2 complex processes (Sifeddine et al., 2011;Woszczyk et al., 2011;Dunn et al.,
3 2008;Wang et al., 2012). Our analyses showed a significant negative relationship
4 between the $\delta^{13}\text{C}_{\text{org}}$ value and water depth (Table 1), implying that the shallow
5 sections in the Boston Lake accumulated more allochthonous TOC (with less negative
6 $\delta^{13}\text{C}$). Apart from the lake own characteristics (such as lake current and depth), other
7 factors may have influences on the dynamics of TOC. For example, land use changes
8 such as agricultural development and fertilization would enhance the riverine input of
9 nutrients, leading to changes in lake productivity and subsequently altering TOC
10 burial in the sediment (Rumolo et al., 2011;Lami et al., 2010;Lamb et al., 2006).
11 There has been evidence of climate change and human activities over the past decades
12 in the surrounding region, which has caused remarkable lake level changes in the
13 Boston Lake (Guo et al., 2014). All these changes would have impacts on the
14 production of POC and TOC burial. Further studies are needed to assess the spatial
15 and temporal variations in the water column biological production to better
16 understand the dynamics of OC in the Boston Lake and the impacts of human activity
17 and climate change.

18

19 **Acknowledgments**

20 This study is financially supported by the National Key Basic Research Program
21 (2013CB956602), ~~and~~the Special Environmental Research Funds for Public Welfare
22 of the State Environmental Protection Administration (~~grant 201309041~~201309041),
23 the Sino-German project (GZ867) and National pioneer project (XDA05020202).

24

25 **References**

- 26 Anderson, N. J., D'Andrea, W., and Fritz, S. C.: Holocene carbon burial by lakes in
27 SW Greenland, Global Glob. Change Biology, 15, 2590-2598, 2009.
28 Anderson, N. J., Dietz, R. D., and Engstrom, D. R.: Land-use change, not climate,
29 controls organic carbon burial in lakes, Proceedings of the Royal Society B

- 1 [Biological Sciences](#), [280P. Roy. Soc. B-Biol. Sci.](#), [280: 20131278](#),
2 [doi:10.1098/rspb.2013.1278](#), 2013.
- 3 Barnes, M. A., and Barnes, W. C.: Organic Compounds in Lake Sediments, in: Lakes,
4 edited by: Lerman, A., Springer New York, 127-152, 1978.
- 5 Battin, T. J., Luyssaert, S., Kaplan, L. A., Aufdenkampe, A. K., Richter, A., and
6 Tranvik, L. J.: The boundless carbon cycle, [Nature Geoscience](#), [Nat. Geosci.](#), [2](#),
7 [598-600](#), 2009.
- 8 Bechtel, A., and Schubert, C. J.: A biogeochemical study of sediments from the
9 eutrophic Lake Lugano and the oligotrophic Lake Brienz, Switzerland, [Organic](#)
10 [Geochemistry](#), [Org. Geochem.](#), [40](#), [1100-1114](#), 2009.
- 11 Bertrand, S., Sterken, M., Vargas-Ramirez, L., De Batist, M., Vyverman, W., Lepoint,
12 G., and Fagel, N.: Bulk organic geochemistry of sediments from Puyehue Lake and
13 its watershed (Chile, 40 °S): Implications for paleoenvironmental reconstructions,
14 [Palaeogeography](#), [Palaeoclimatology](#), [Palaeoecology](#), [Palaeogeogr. Palaeocl.](#), [294](#),
15 [56-71](#), 2010.
- 16 Brodie, C. R., Leng, M. J., Casford, J. S. L., Kendrick, C. P., Lloyd, J. M., Yongqiang,
17 Z., and Bird, M. I.: Evidence for bias in C and N concentrations and $\delta^{13}\text{C}$
18 composition of terrestrial and aquatic organic materials due to pre-analysis acid
19 preparation methods, [Chemical Geology](#), [Chem. Geol.](#), [282](#), [67-83](#), 2011.
- 20 Burone, L., Muniz, P., PIRES-VANIN, A., MARIA, S., and RODRIGUES, M.:
21 Spatial distribution of organic matter in the surface sediments of Ubatuba Bay
22 (Southeastern-Brazil), [Anais da Academia Brasileira de Ciências](#), [An. Acad. Bras.](#)
23 [Cien.](#), [75](#), [77-80](#), 2003.
- 24 Dean, W. E., and Gorham, E.: Magnitude and significance of carbon burial in lakes,
25 reservoirs, and peatlands, [Geology](#), [26](#), [535-538](#), 1998.
- 26 Deng, B., Zhang, J., and Wu, Y.: Recent sediment accumulation and carbon burial in
27 the East China Sea, [Global Biogeochemical Cycles](#), [20](#), [GB3014](#), [Biogeochem. Cy.](#)
28 [20](#), [GB3014](#), [doi:10.1029/2005 GB 002559](#), 2006.
- 29 Dong, X., Anderson, N. J., Yang, X., and Shen, J.: Carbon burial by shallow lakes on
30 the Yangtze floodplain and its relevance to regional carbon sequestration,

- 1 ~~Global~~Glob. Change ~~Biology~~Biol., 18, 2205-2217, 2012.
- 2 Downing, J. A., Prairie, Y., Cole, J., Duarte, C., Tranvik, L., Striegl, R., McDowell, W.,
3 Kortelainen, P., Caraco, N., and Melack, J.: The global abundance and size
4 distribution of lakes, ponds, and impoundments, ~~Limnology and~~
5 ~~Oceanography~~Limnol.Oceanogr., 51, 2388-2397, 2006.
- 6 Dunn, R. J. K., Welsh, D. T., Teasdale, P. R., Lee, S. Y., Lemckert, C. J., and Meziane,
7 T.: Investigating the distribution and sources of organic matter in surface sediment
8 of Coombabah Lake (Australia) using elemental, isotopic and fatty acid biomarkers,
9 ~~Continental~~Cont. Shelf ~~Research~~Res., 28, 2535-2549, 2008.
- 10 Eksperiandova, L. P., Fedorov, O. I., and Stepanenko, N. A.: Estimation of
11 metrological characteristics of the element analyzer EuroVector EA-3000 and its
12 potential in the single-reactor CHNS mode, ~~Microchemical Journal~~Microchem. J.,
13 99, 235-238, 2011.
- 14 Fang, J., Wu, F., Xiong, Y., Li, F., Du, X., An, D., and Wang, L.: Source
15 characterization of sedimentary organic matter using molecular and stable carbon
16 isotopic composition of n-alkanes and fatty acids in sediment core from Lake
17 Dianchi, China, ~~Science of The Sci.~~ Total ~~Environment~~Enviro., 473, 410-421,
18 2014.
- 19 Ferland, M.-E., del Giorgio, P. A., Teodoro, C. R., and Prairie, Y. T.: Long-term C
20 accumulation and total C stocks in boreal lakes in northern Québec, Global
21 ~~Biogeochemical Cycles~~Biogeochem. Cy., 26, GB0E04,
22 doi:10.1029/2011GB004241, 2012.
- 23 Gireeshkumar, T. R., Deepulal, P. M., and Chandramohanakumar, N.: Distribution and
24 sources of sedimentary organic matter in a tropical estuary, south west coast of
25 India (Cochin estuary): A baseline study, ~~Marine Pollution Bulletin~~Mar. Pollut.
26 Bull., 66, 239-245, 2013.
- 27 Gudasz, C., Bastviken, D., Steger, K., Premke, K., Sobek, S., and Tranvik, L. J.:
28 Temperature-controlled organic carbon mineralization in lake sediments, Nature,
29 466, 478-481, 2010.
- 30 Guo, M., Wu, W., Zhou, X., Chen, Y., and Li, J.: Investigation of the dramatic changes

- 1 in lake level of the Boston Lake in northwestern China, *Theor. Appl. Climatol.*, 111,
2 2014., 119, 341-351, 2015.
- 3 Hanson, P. C., Buffam, I., Rusak, J. A., Stanley, E. H., and Watras, C.: Quantifying
4 lake allochthonous organic carbon budgets using a simple equilibrium model,
5 *Limnol. Oceanogr.*, 59, 167-181, 2014.
- 6 Jiang, J., and Huang, Q.: Distribution and variation of lakes in Tibetan Plateau and
7 their comparison with lakes in other part of China, *Water Resources Protection*, 4,
8 24-27, 2004.
- 9 Jin, Z., Li, F., Cao, J., Wang, S., and Yu, J.: Geochemistry of Daihai Lake sediments,
10 Inner Mongolia, north China: Implications for provenance, sedimentary sorting,
11 and catchment weathering, *Geomorphology*, 80, 147-163, 2006.
- 12 Kaushal, S., and Binford, M.: Relationship between C:N ratios of lake sediments,
13 organic matter sources, and historical deforestation in Lake Pleasant, Massachusetts,
14 USA, *J. Paleolimnol.*, 22, 439-442, 1999.
- 15 Khim, B.-K., Jung, H. M., and Cheong, D.: Recent variations in sediment organic
16 carbon content in Lake Soyang (Korea), *Limnology*, 6, 139-139, 2005.
- 17 Lamb, A. L., Leng, M. J., Umer Mohammed, M., and Lamb, H. F.: Holocene climate
18 and vegetation change in the Main Ethiopian Rift Valley, inferred from the
19 composition (C/N and $\delta^{13}\text{C}$) of lacustrine organic matter, *Quaternary Science*
20 *Reviews, Sci. Rev.*, 23, 881-891, 2004.
- 21 Lamb, A. L., Wilson, G. P., and Leng, M. J.: A review of coastal palaeoclimate and
22 relative sea-level reconstructions using $\delta^{13}\text{C}$ and C/N ratios in organic material,
23 *Earth-Science Reviews, Sci. Rev.*, 75, 29-57, 2006.
- 24 Lami, A., Turner, S., Musazzi, S., Gerli, S., Guilizzoni, P., Rose, N., Yang, H., Wu, G.,
25 and Yang, R.: Sedimentary evidence for recent increases in production in Tibetan
26 plateau lakes, *Hydrobiologia*, 648, 175-187, 2010.
- 27 Lin, S., Hsieh, I. J., Huang, K.-M., and Wang, C.-H.: Influence of the Yangtze River
28 and grain size on the spatial variations of heavy metals and organic carbon in the
29 East China Sea continental shelf sediments, *Chemical Geology, Chem. Geol.*, 182,
30 377-394, 2002.

- 1 Liu, K.-K., and Kao, S.-J.: A Three End-Member Mixing Model Based on Isotopic
2 Composition and Elemental Ratio, *Terrestrial, Atmospheric & Oceanic Sciences*,
3 18, [1067-1075](#), 2007.
- 4 Liu, W., Li, X., An, Z., Xu, L., and Zhang, Q.: Total organic carbon isotopes: A novel
5 proxy of lake level from Lake Qinghai in the Qinghai–Tibet Plateau, China,
6 [Chemical Geology, Chem. Geol.](#), 347, 153-160, 2013.
- 7 Liu, X., Herzschuh, U., Wang, Y., Kuhn, G., and Yu, Z.: Glacier fluctuations of
8 Muztagh Ata and temperature changes during the late Holocene in westernmost
9 Tibetan Plateau, based on glaciolacustrine sediment records, [Geophysical Research Letters, Geophys. Res. Lett.](#), 41, 2014GL060444, [doi:10.1002/2014GL060444](#),
10 2014.
- 12 Meyers, P. A., and Ishiwatari, R.: Lacustrine organic geochemistry—an overview of
13 indicators of organic matter sources and diagenesis in lake sediments, [Organic](#)
14 [geochemistry, Org. Geochem.](#), 20, 867-900, 1993.
- 15 Meyers, P. A.: Applications of organic geochemistry to paleolimnological
16 reconstructions: a summary of examples from the Laurentian Great Lakes, [Organic](#)
17 [Geochemistry, Org. Geochem.](#), 34, 261-289, 2003.
- 18 Molinaroli, E., Guerzoni, S., De Falco, G., Sarretta, A., Cucco, A., Como, S., Simeone,
19 S., Perilli, A., and Magni, P.: Relationships between hydrodynamic parameters and
20 grain size in two contrasting transitional environments: The Lagoons of Venice and
21 Cabras, Italy, [Sedimentary Geology, Sediment. Geol.](#), 219, 196-207, 2009.
- 22 Pancost, R. D., and Boot, C. S.: The palaeoclimatic utility of terrestrial biomarkers in
23 marine sediments, [Marine Chemistry, Mar. Chem.](#), 92, 239-261, 2004.
- 24 Qin, B., Yang, L., Chen, F., Zhu, G., Zhang, L., and Chen, Y.: Mechanism and control
25 of lake eutrophication, [Chin. Sci. Bull.](#), 51, 2401-2412, 2006.
- 26 Qin, B., and Zhu, G.: The nutrient forms, cycling and exchange flux in the sediment
27 and overlying water system in lakes from the middle and lower reaches of Yangtze
28 River, *Science in China Series D*, 49, 1-13, 2006.
- 29 Qin, B., Yang, L., Chen, F., Zhu, G., Zhang, L., and Chen, Y.: Mechanism and control
30 of lake eutrophication, [Chin. Sci. Bull.](#), 51, 2401-2412, 2006.


- 1 Rumolo, P., Barra, M., Gherardi, S., Marsella, E., and Sprovieri, M.: Stable isotopes
2 and C/N ratios in marine sediments as a tool for discriminating anthropogenic
3 impact, *Journal of Environmental Monitoring, J. Environ. Monitor.*, 13, 3399-3408,
4 2011.
- 5 Shanahan, T. M., McKay, N., Overpeck, J. T., Peck, J. A., Scholz, C., Heil Jr, C. W.,
6 and King, J.: Spatial and temporal variability in sedimentological and geochemical
7 properties of sediments from an anoxic crater lake in West Africa: Implications for
8 paleoenvironmental reconstructions, *Palaeogeography, Palaeoclimatology,*
9 *Palaeoecology, Palaeogeogr., Palaeocl.*, 374, 96-109, 2013.
- 10 Sifeddine, A., Meyers, P., Cordeiro, R., Albuquerque, A., Bernardes, M., Turcq, B.,
11 and Abrão, J.: Delivery and deposition of organic matter in surface sediments of
12 Lagoa do Caçó (Brazil), *J. Paleolimnol.*, 45, 385-396, 2011.
- 13 Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T.
14 J., Dillon, P., Finlay, K., Fortino, K., and Knoll, L. B.: Lakes and reservoirs as
15 regulators of carbon cycling and climate, *Limnology and Oceanography, Limnol.*
16 *Oceanogr.*, 54, 2298-2314, 2009.
- 17 Vuorio, K., Meili, M., and Sarvala, J.: Taxon-specific variation in the stable isotopic
18 signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of lake phytoplankton, Freshwater *Biology, Biol.*, 51,
19 807-822, 2006.
- 20 Wang, X., Cui, L., Xiao, J., and Ding, Z.: Stable carbon isotope of black carbon in
21 lake sediments as an indicator of terrestrial environmental changes: An evaluation
22 on paleorecord from Daihai Lake, Inner Mongolia, China, *Chemical Geology, Chem.*
23 *Geol.*, 347, 123-134, 2013.
- 24 Wang, X., Fang, C., Yu, Z., Wang, J., Peng, D., and Jingjing, K.: Seasonal variations
25 of particulate and dissolved organic carbon in Boston Lake, Xinjiang, Journal of
26 Lake Sciences, 26, 552-558, 2014.
- 27 Wang, X., Wang, J., Xu, M., Zhang, W., Fan, T. and Zhang, J.: Carbon accumulation
28 in arid croplands of northwest China: pedogenic carbonate exceeding organic
29 carbon, *Sci. Rep-UK*, 5:11439, 10.1038/srep11439, 2015.
- 30 Wang, Y., Zhu, L., Wang, J., Ju, J., and Lin, X.: The spatial distribution and

- 1 sedimentary processes of organic matter in surface sediments of Nam Co, Central
2 Tibetan Plateau, Chin. Sci. Bull., 57, 4753-4764, 2012.
- 3 Woszczyk, M., Bechtel, A., Gratzer, R., Kotarba, M. J., Kokociński, M., Fiebig, J.,
4 and Cieśliński, R.: Composition and origin of organic matter in surface sediments
5 of Lake Sarbsko: A highly eutrophic and shallow coastal lake (northern Poland),
6 Organic Geochemistry, Org. Geochem., 42, 1025-1038, 2011.
- 7 Wu, F., Xu, L., Sun, Y., Liao, H., Zhao, X., and Guo, J.: Exploring the relationship
8 between polycyclic aromatic hydrocarbons and sedimentary organic carbon in three
9 Chinese lakes, J Soils Sediments, Soil. Sediment., 12, 774-783, 2012.
- 10 Wu, J., Huang, C., Zeng, H., GerhardH, S., and Rick, B.: Sedimentary evidence for
11 recent eutrophication in the northern basin of Lake Taihu, China: human impacts on
12 a large shallow lake, J. Paleolimnol., 38, 13-23, 2007.
- 13 Wu, J., Ma, L., and Zeng, H.: Water Quality and Quantity Characteristics and Its
14 Evolution in Lake Boston, Xinjiang over the Past 50 Years, Scientia Geographica
15 Sinica, 33, 231-237, 2013.
- 16 Yu, F., Zong, Y., Lloyd, J. M., Huang, G., Leng, M. J., Kendrick, C., Lamb, A. L., and
17 Yim, W. W. S.: Bulk organic $\delta^{13}\text{C}$ and C/N as indicators for sediment sources in the
18 Pearl River delta and estuary, southern China, Estuarine, Coastal and Estuar. Coast.
19 Shelf Science, 87, 618-630, 2010.
- 20 Zhang, C., Mischke, S., Zheng, M., Prokopenk, A., Guo, F., and Feng, Z.: Carbon and
21 Oxygen Isotopic Composition of Surface-Sediment Carbonate in Boston Lake
22 (Xinjiang, China) and its Controlling Factors, Acta Geologica Sinica English
23 Edition, Geol. Sin.-Engl., 83, 386-395, 2009.
- 24 Zhang, J.: Impacts of land use on soil organic matter in Yanqi Basin, Doctor of
25 Physical Geography Ph.D. thesis, Xinjiang Institute of Ecology and Geography,
26 Chinese Academy of Sciences, Urumqi, Xinjiang, P.R. China, 2013.
- 27 Zhu, Z., Chen, J. a., and Zeng, Y.: Abnormal positive delta C-13 values of carbonate
28 in Lake Caohai, southwest China, and their possible relation to lower temperature,
29 Quaternary International, Quatern. Int., 286, 85-93, 2013.


1 Table 1. Correlation coefficient (r) between various variables for the sediments.

Variables	WD	DBD	d(0.5)	Cla y	Silt	Sand	TOC	$\delta^{13}\text{C}_{\text{org}}$
TOC	0.50	-0.58 ^{*5} <u>8^a</u>	-0.74 ^{**7} <u>1^b</u>	0.1 8	0.77 ^{**7} <u>7^b</u>	-0.76 ^{**7} <u>6^b</u>		-0.15
TN	0.07	-0.82 ^{**8} <u>3^b</u>	-0.60 ^{*6} <u>0^a</u>	-0.0 5	0.79 ^{**7} <u>9^b</u>	-0.72 ^{**7} <u>2^b</u>	0.74 ^{**7} <u>1^b</u>	0.45
C:N	0.50	0.50	0.01	0.2 5	-0.19	0.11	0.14	-0.82 ^{**8} <u>2^b</u>
$\delta^{13}\text{C}_{\text{org}}$	-0.66 ^{*6} <u>6^a</u>	-0.46	-0.13	0.0 3	0.21	-0.20	-0.15	
POC				-0.2				
	-0.42	-0.41	0.11	9	0.11	-0.02	0.14	0.22


2 WD = water depth (m), DBD = dry bulk density (g cm⁻³), d(0.5) = median diameter
3 (µm) and clay, silt and sand fractions (%) from the 0-2 cm sediments. Significance of
4 Pearson correlation is marked with ^{*a} (p<0.05) and ^{**b} (p<0.01) asterisks.



4 Figure 1. Map of the Boston Lake with the water depth and the 13 sampling stations
 5 (red dots). Bathymetric was measured in 2008 by Wu et al. (2013) and bathymetric
 6 contours were plotted by using software ArcGIS 9.3 and Corel DRAW X3.





1




2


Figure 2. Distributions of (a) clay, (b) silt, (c) sand and (d) the median diameter ($d(0.5)$, μm) in the 0-1 cm (color map) and 1-2 cm (dashed lines). The spatial distribution maps (Figure 2-7) were produced using Surfer 9.0 (Golden Software Inc.) and the interpolated data in the maps was made using the Kriging method of gridding.






2 Figure 3. Spatial distributions of (a) total organic carbon (TOC)
 3 and (b) total nitrogen (TN) in the 0-1 cm (color map) and 1-2 cm (dashed lines).





2 Figure 4. Spatial distribution of (a) C:N ratio and (b) carbon stable isotope ($\delta^{13}\text{C}_{\text{org}}$) of
3 TOC in the 0-1 cm (color map) and 1-2 cm (dashed lines).








Figure 5. Spatial patterns of the relative contributions for TOC in the 0-1cm (color map) and 1-2 cm (dashed lines) sediments. (a) TOC from lake plankton (TOC_{lp}), (b) TOC from surface soils (TOC_{ss}), and (c) TOC from native plants (TOC_{np}).





2 Figure 6. Spatial distributions of (a) autochthonous TOC (TOC_{auto}) and (b)
3 allochthonous sources TOC (TOC_{allo}) in the 0-1 cm (color map) and 1-2cm (dashed
4 lines) sediments.




Figure 7. Spatial distributions of POC concentrations in summer (color map) and autochthonous TOC in the 0-1 cm sediment ($\text{TOC}_{\text{auto},0-1\text{cm}}$, dashed lines). POC data were from Wang et al. (2014).