

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Age structure, carbonate production and shell loss rate in an Early Miocene reef of the giant oyster *Crassostrea gryphoides*

M. Harzhauser¹, A. Djuricic², O. Mandic¹, T. A. Neubauer¹, M. Zuschin³, and N. Pfeifer²

¹Natural History Museum Vienna, Geological-Paleontological Department, Vienna, Austria

²TU Wien, Department of Geodesy and Geoinformation, Vienna, Austria

³University of Vienna, Department of Paleontology, Vienna, Austria

Received: 22 July 2015 – Accepted: 9 September 2015 – Published: 28 September 2015

Correspondence to: M. Harzhauser (mathias.harzhauser@nhm-wien.ac.at)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

12, 15867–15900, 2015

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We present the first analysis of population structure and cohort distribution in a fossil oyster reef based on more than 1121 shells of the giant oyster *Crassostrea gryphoides* (Schlotheim, 1813). Data derive from Terrestrial Laser Scanning of a Lower Miocene shell bed covering 459 m². Within two transects, individual shells were manually outlined on a digital surface model and cross-checked based on high-resolution orthophotos, resulting in accurate information on center line length and area of exposed shell surface. A growth model was calculated, revealing this species as the fastest growing and largest *Crassostrea* known so far. Non-normal distribution of size, area and age data hints at the presence of at least four distinct recruitment cohorts. The rapid decline of frequency amplitudes with age is interpreted to be a function of mortality and shell loss. The calculated shell half-lives range around few years, indicating that oyster reefs were geologically short-lived structures, which could have been fully degraded on a decadal scale.

Crassostrea gryphoides reefs were widespread and common along the Miocene circum-Tethyan coasts. Given its enormous growth performance of ~ 150 g carbonate per year this species has been an important carbonate producer in estuarine settings. Yet, the rapid shell loss impeded the formation of stable structures comparable to coral reefs.

1 Introduction

The genus *Crassostrea* Sacco, 1897 comprises numerous commercially exploited species. The modes of growth and population structures of extant *Crassostrea* species are of paramount importance for oyster fishery (FAO, 2015). Consequently, a wealth of data exists on the ontogeny, biology and ecological requirements. These data, in turn are a valuable base for the interpretation of the autecology of fossil congeners. Extant *Crassostrea* are sessile bivalves adapted to estuarine and intertidal environments

BGD

12, 15867–15900, 2015

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where they have to cope with high environmental stress. Whilst physicochemical stress is managed by genetic response (Zhang et al., 2012), the formation of thick shells is a strategy against predation (Lombardi et al., 2013; Robinson et al., 2014). The largest and fastest growing *Crassostrea* species flourished during the Miocene and Early Pliocene and became replaced by comparatively smaller and thinner species thereafter (Kirby and Jackson, 2004). Among these, *Crassostrea gryphoides* (Schlotheim, 1813) is the largest, attaining shell lengths of up to 80 cm and individual ages of more than 40 years (Harzhauser et al., 2010). The Pleistocene–Holocene extirpation of large and thick-shelled *Crassostrea* species was explained by a shift from shallow-marine towards estuarine-intertidal habitats to escape from predation (Kirby, 2000, 2001). At least for *C. gryphoides* this model does not fit because the oyster lived as secondary soft-bottom dweller in the intertidal zone of estuaries along the circum-Tethyan coasts (Laurain, 1980; Schultz, 2001; Mandic et al., 2004). This species evolved during the Late Oligocene in the European Paratethys Sea and became ubiquitous in the Paratethys Sea, the proto-Mediterranean Sea and the eastern Atlantic throughout the Early to Late Miocene. It might even have entered the Indian Ocean during the Miocene (Newton and Smith, 1912) and reached the North Sea during the Middle Miocene (Schultz, 2001). Pliocene records from the eastern Atlantic and North Africa, however, may need verification (see Schultz, 2001 for a detailed list of occurrences). This species became extinct around the Miocene/Pliocene boundary or with the onset of the Pliocene cooling at the latest.

Studies on growth in fossil *Crassostrea* species and the resulting carbonate production were published by Chinzei (1995), Kirby (2001) and Kirby and Jackson (2004). These studies are based on collection material comparing species and specimens from different stratigraphic horizons. No study, however, tried to capture the size and age structures of a fossil *Crassostrea* reef, presumably representing a real population of coeval specimens. The lack of such studies is clearly linked to the fact that fossil shell beds usually represent time-averaged assemblages (Kidwell, 1986, 1991), which only vaguely reflect original community structures. Although some in-situ preserved

(Chatterji et al., 1985; Nagi et al., 2011). Nevertheless, the Miocene species is closely related to the extant Asian–Pacific *Crassostreinae* species, which show a large genetic difference from Atlantic species (Littlewood, 1994; Ó Foighil et al., 1995; Wang et al., 2004; Ren et al., 2010; Salvi et al., 2014). Both groups differ considerably in mitochondrial genes, nuclear genome and chromosome structures (Wang et al., 2004; Ren et al., 2010; Salvi et al., 2014). The divergence between both groups seems to have happened already in Cretaceous times and the diversification of the Asian Pacific group started during the Eocene (Ren et al., 2014). Based on this genetic evidence and biogeographic separation, Salvi et al. (2014) introduced the genus *Magallana* Salvi, Macali and Mariottini, 2014 for the Asian Pacific species group with *Ostrea gigas* Thunberg, 1793 as type species. Salvi et al. (2014), however, did not provide a formal “description or definition that states in words characters that are purported to differentiate the taxon” (ICNZ Art. 13.1.1). The authors only refer to a description of the type species. Therefore, *Magallana* is formally unavailable (see also Marshall, 2015) and we use *Crassostrea* in its traditional sense.

2 Geological setting and paleoenvironment

The investigated oyster reef is excavated at Stetten in Lower Austria (48°22′03.33″ N, 16°21′33.22″ E). It is part of the about 600-m-thick lower Miocene (upper Burdigalian, Karpatian) siliciclastic succession of the Korneuburg Basin, which is a 20 km long and 7 km wide halfgraben within the Alpine–Carpathian thrust-belt (Fig. 2) (Wessely, 1998). The basin fill represents a quick alternation of shoreface and tidal flat deposits, which formed in an embayment of the Paratethys Sea (Zuschin et al., 2014). More than 650 species-level taxa have been described from the area (Sovis and Schmid, 1998, 2002) and the paleoenvironments are fairly well understood (Harzhauser et al., 2002; Zuschin et al., 2014). The oyster reef flourished in an estuary fringed by salt marshes, Taxodiaceae swamps and scattered *Avicennia* mangroves (Harzhauser et al., 2002; Kern et al., 2010). The pollen record of the section documents a warm subtropical climate

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with marked seasonality (Kern et al., 2010). A warm and wet summer season with c. 204–236 mm precipitation during the wettest month was opposed by a rather dry winter season with precipitation of c. 9–24 mm during the driest month. The mean annual temperature ranged between 15.7–20.8 °C, with about 9.6–13.3 °C during the cold season and 24.7–27.9 °C during the warmest month. These data suggest similarities with the modern “Cwa” climate of Koeppen (1936). Today this climate covers parts of northern India extending into south-eastern Asia (south Nepal, Myanmar, northern Thailand) to East China and in central south Africa (east Angola, Zambia, north Zimbabwe, north Mozambique) (Peel et al., 2007; Kern et al., 2010). The distinct seasonality was also revealed in sclerochronologic analysis of one of the *Crassostrea* shells collected from the reef (Harzhauser et al., 2010). This shell exhibits a regular annual rhythm of at least 11 seasons with a temperature range of 9.8 °C. Thus, the paleoclimatic and paleoenvironmental frame of the *C. gryphoides* reef is comparable to the settings of modern *Crassostrea* reefs in the subtropical parts of the Asian Pacific.

Harzhauser et al. (2015) describe the complex taphonomy of the shell bed, which was formed by a tsunami or an exceptional storm and represents an event deposit *sensu* Einsele et al. (1991) and Kidwell (1991). Despite the bias of post-event processes, when the exhumed shells were exposed for few years on the seafloor below fair weather base, the rapid subsequent burial preserved most of the original distribution patterns.

3 Materials and methods

3.1 Data acquisition by Terrestrial Laser Scanning and orthophotos

The shell bed was excavated manually with steel graters and brushes; no water was used for cleaning. The surface of the shell bed was flat during deposition but has now an undulate relief of few decimeters due to post sedimentary tectonics. Terrestrial Laser Scanning was applied to document the site as georeferenced 3-D point cloud (Otepka

et al., 2013). A Faro Focus Laser scanner with a nominal point measurement accuracy of 1 mm (SD) in each coordinate and a sampling distance of approximately 1 mm was used. The individual point clouds of each scan were transformed first into one common coordinate system and then georeferenced by control points to Universal Transverse Mercator (UTM) coordinates (resolution below 2 mm). A robust filter (pre-processing) was applied to reduce measurement noise while preserving surface structures like sharp edges (Nothegger and Dorninger, 2009). The surface triangulation is based on the Poisson surface reconstruction method (Kazhdan et al., 2006). The points of this triangulation are used for interpolating a regular grid of heights above the plane of the shell bed using the scientific software OPALS (Pfeifer et al., 2014). In addition, a Canon 60D with a Canon EF 20 mm f2.8 was used to capture more than 300 photos from a moving platform. The camera was placed approximately orthogonal to the fossil bed. From the photos with a nominal ground resolution of approximately $0.6 \text{ mm pixel}^{-1}$ an orthophoto mosaic was generated with a resolution of $0.5 \text{ mm pixel}^{-1}$. To detect patterns in the distribution and composition of shells two transects (N–S, W–E) were defined, each represented by 42 m^2 with a central overlap (Fig. 1). All objects within this area were manually outlined on the digital surface model and cross checked based on the high-resolution orthophotos.

As discussed by Harzhauser et al. (2015) the assemblage is not monospecific but contains about 46 molluscan species of which *Crassostrea* predominates (79.4 %). A detailed description of the composition and distribution of the other taxa is given in Harzhauser et al. (2015). Herein, we only focus on *Crassostrea gryphoides*. In total, 10284 objects were defined. Of these, 8169 objects were identified as *Crassostrea gryphoides* of which 1121 are complete shells (see Supplement Table); 86 % of the specimens display various degrees of fragmentation and are excluded from the size-frequency analysis. The fragments usually show sharp fractures and therefore, the main cause for fragmentation seems to be predatory and hydrodynamic breakage. The ratio between left and right valves is balanced (0.98).

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Shell length and area

Crassostrea gryphoides shows a very broad range of morphologies, ranging from elongate shells to strongly curved and sigmoidal shapes (Fig. 3). Therefore, measuring shell length as straight line, as done in other extant and fossil *Crassostrea* species, is inadequate. To overcome this problem, we evaluated shell length based on the 2-D center line, which is an imaginary curved line spanning the maximum growth-length of the shell. For the automatic determination of the center line we used the shell margins, which comprise about 1000 points on average. For easier calculation the outline point number was reduced to 100 and then filtered to points with close to even spacing. In the next step, a Delaunay triangulation was calculated between the filtered outline points (Delaunay, 1934), constrained by the edges between the outline points. To find the center line for each oyster outline, the Voronoi diagram was formed (Voronoi, 1908) from the triangulation. The edges between neighboring Voronoi vertices within the boundary are the medial axis transform (MAT) for the oyster outline (Aichholzer et al., 1996). The longest 2-D path in this tree was found using Dijkstra's algorithm between MAT end points (Kirk, 2015).

The center line lengths of 1121 complete *Crassostrea* shells, rounded to the nearest mm, range from 48 to 602 mm with a mean of 237 mm ($\sigma = 89$ mm) (Fig. 4a). The data distribution displays a positive skewness of 0.52 and the Shapiro–Wilk test excludes normal distribution for raw data and log₁₀-transformed measurements. Area data range from 1708 to 56 755 mm² with a mean of 16 983 mm² ($\sigma = 8414$ mm²) (Fig. 4b). These data show also a positive skewness (0.83) and normal distribution is rejected by the Shapiro–Wilk test.

Based on the manual outlines, the exposed shell area can be deduced directly. Area data are slightly underestimated because shells are not always exposed parallel to the bedding plane but may be somewhat oblique. Despite the fact, that area data are somewhat biased by oblique shells, the correlation between center line lengths and

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



areas is highly significant (raw data: $r = 0.92$, $p < 0.001$; log 10-transformed: $r = 0.93$, $p < 0.001$) (Fig. 5).

3.3 Length frequency data

Non-normal distribution of length–frequencies is a common pattern in extant *Crassostrea* reefs (e.g. Coakley, 2004; Baqueiro Cárdenas and Aldana Aranda, 2007; Harding et al., 2008; Nurul Amin et al., 2008; Ross and Luckenbach, 2009; Goslier et al., 2014). It results from seasonal and/or annual recruitment with distinct cohorts. For instance, the (sub)tropical *C. madrasensis* and *C. rhizophorae* display a distinct annual recruitment peak (Nurul Amin et al., 2008; Mancera and Mendo, 1996). In multiannual communities this results in a right-skewed distribution due to the loss of old specimens by natural mortality and shell loss. For extant *Crassostrea* reefs, the analysis of the cohorts is routinely performed using Bhattacharya’s model or the EM-Algorithm of Dempster et al. (1977), which tries to detect normal distributions within the length–frequency data. Consequently, in order to test for cohort mixing, lengths of *C. gryphoides* were subjected to mixture analysis, a maximum-likelihood method for estimating the parameters of two or more univariate normal distributions, based on a pooled univariate sample (Hammer, 2015). Statistical analyses were performed in PAST versions 2.17c and 3.06 (Hammer et al., 2001). Akaike’s Information Criterion (AIC) was used to test the goodness of fit of the maximum likelihood estimates to the length–frequency data.

In log10-transformed length frequency diagrams, the maximum likelihood based analysis reveals lowest AIC values for four or five cohorts. Similarly, log-transformed area data have lowest AIC values if four or five cohorts are detected. Assuming more groups does not lower the AIC or the computed cohorts comprise unrealistic narrow cohort ranges, which are nested within larger ones.

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.4 Growth model

Kirby et al. (1998), Kirby (2000, 2001) and Kirby and Jackson (2004) used ligamental increments of fossil and recent *Crassostrea* species to estimate individual life spans, assuming that increments are formed annually. The ligamental area of these *Crassostrea* species is typically structured by perpendicular convex ridges and concave furrows corresponding to phases of rapid and low calcification. The specimens from the Stetten site lack such well-defined convex tops. In all specimens of *C. gryphoides* analyzed by Harzhauser et al. (2010) for stable isotopes, the counting of increments would have resulted in a large overestimation of the life spans. Similarly, Alam and Das (1999) documented a clear misfit between growth increments and age for the extant *C. madrasensis*. Therefore, we restrict our age estimates solely on growth rates of *C. gryphoides* deduced from stable isotope profiles published by Harzhauser et al. (2010). According to these authors, a shell from Stetten attained 43 cm in length at an age of 11 years and a second one from a slightly younger site was 63 cm long at an age of 16 years. These values document an average growth rate of $\sim 3.9 \text{ cm year}^{-1}$ and might serve as base for rough age estimates for the complete adult shells from the Stetten biostrome. For juvenile shells, this estimate would be wrong due to the non-linear mode of growth known from extant *Crassostrea* species, which show very high initial growth rates (Kennedy et al., 1996). Similarly, growth rates of *Crassostrea gryphoides* seem to decline in very old and large specimens as shown for a 78 cm long and 41 year old shell from the late Langhian of Austria (Harzhauser et al., 2010). To cope with the non-linear growth, growth curves of extant *Crassostrea* species are routinely calculated with the von Bertalanffy equation (von Bertalanffy, 1934). This equation is $SL_t = SL_{\max} (1 - e^{-k(t-t_0)})$, where SL_t is shell length at time t , SL_{\max} is the asymptotic shell length, t_0 is the size at time 0, and k is a rate constant. Herein, we used the length of the center line of each shell as SL_t as this measure captures the real growth length of the partly strongly curved or sigmoid specimens. For SL_{\max} we used the size-to-age data of the 78 cm long shell, which is the largest individual known so far.

3.5 Ratio between chalky and foliate layers

The calcitic *Crassostrea* shells consist of two structures: thin but densely spaced foliate layers separated by thick layers of light-weight chalky material (Stenzel, 1971; Higuera-Ruiz and Elorza, 2009). This fast growing structure is interpreted to be a major adaptive advantage of *Crassostrea* to impede drilling predation and to prevent from sinking in the soft bottom (Seilacher, 1984; Chinzei, 1995; Kirby, 2001; Vermeij, 2014). In fossil shells the chalky layer is completely recrystallized and has the same density as the foliate layer. Nevertheless, it is optically easily recognized by its lighter color and the nearly opaque appearance. A polished longitudinal section of an articulated *C. gryphoides* shell (providing data for left and right shells) was scanned and the ratio between both shell structures was quantified by image analysis (Fig. 6). This method is only an approximation to the true value, as the ratio may vary locally (Durve and Bal, 1960; Chinzei, 1995), but it is clearly an improvement compared to former studies that used only linear transects or sectors within the shell (Durve and Bal, 1960; Chinzei, 1995; Kirby, 2001).

Image analysis of the cross-section documents proportions of 64 % chalky to 36 % foliate layers for the right shell and of 61 to 39 % for the left shell. The density of the chalky layer when wet ranges around $1.15\text{--}1.32\text{ g cm}^{-3}$ (Chinzei, 1995) and the density of the foliate layer ranges around $2.2\text{--}2.5\text{ g cm}^{-3}$ (Chinzei, 1995; Yoon et al., 2003) and has a clear upper limit by the density of calcite (2.7 g cm^{-3}). Using 1.2 g cm^{-3} for the chalky layer and 2.2 g cm^{-3} for the foliate layer as rough estimates result in a mean density of 1.84 and 1.81 g cm^{-3} for the right and left valves, respectively. Shell density in *Crassostrea* species is independent from age and size (Lombardi et al., 2013) and therefore the density estimates can be applied to the entire data set.

3.6 Shell volume

The volume of nine individual shells was determined using close range laser scanning technology, which provides high resolution models with sub-mm resolution. The spec-

BGD

12, 15867–15900, 2015

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



imens were selected from the collections of the Natural History Museum and vary in center length from 141 to 406 mm; four of these shells represent left-right shell pairs of an individual (Fig. 3). The data were captured with a measuring arm (METRIS MCA, 3600 M7). A hand-help triangulation laser scanner (a laser plane and a camera) was mounted at the end of two arms of fixed length with flexible joints. The laser scanner takes measurements with a maximum scan rate of 80 stripes per second with a strip-width of about 200 mm; the camera has a resolution of 1000 dots strip⁻¹.

In a first step, more than half of each shell was scanned and in a second step the other half. The overlap between both parts was more than 70 %, which was sufficient for successful registration of the scanned parts. During this registration process, the geometric transformation is determined, which puts the two 3-D laser point clouds together based on the points in the overlapping part. This procedure is done using the iterative closest point (ICP) algorithm (Glira et al., 2015). The resulting point cloud is analyzed in order to reduce noise and thus improve the surface description. Outliers (wrongly determined points not on the surface) were manually eliminated. Additionally, the raw point cloud (over 1.5 million points shell⁻¹) was uniformly sub-sampled to allow interactive handling. The final resolution is better than 0.18 mm (i.e. around 25 points mm⁻²). For volume calculation the point cloud has to be transformed into a closed mesh. Remaining holes, non-manifold surfaces and additional not connected components were identified and removed. Finally, the surface area of the mesh and its volume were computed using the algorithm of Mirtich (1996).

Based on the 9 measured shells a relation between center line length and volume can be deduced. The largest shell measured is 406 mm long but no empirical volume data are available for larger shells, because these cannot be removed from the protected site. Therefore, the von Bertalanffy equation would not be applicable for shells larger than ~ 40 cm. Consequently, we chose a logistic function to approximate the inverse von Bertalanffy equation:

$$V = \frac{3.2439 \times 10^6}{1 + 118.86 e^{-0.0099889 SL}},$$

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

where v is the volume in mm^3 and SL the center line length in mm (Fig. 7). Applying this equation to all shells results in a total volume of $393\,273\text{cm}^3$ with a mean shell volume of 350.8cm^3 ($\sigma = 313.7$). These values do not change significantly if a non-linear Gompertz growth model is assumed as frequently done for *Crassostrea* (Lopez et al., 2013; Ginger et al., 2013). The respective equation

$$v = 1.978 \times 10^8 e^{-9.1401} e^{-0.0013762x}$$

results in a total volume of $398\,474\text{cm}^3$. Applying the above discussed average shell gravity of 1.82g cm^{-3} results in a total carbonate mass of $\sim 715\text{kg}$ (logistic) to $\sim 725\text{kg}$ (Gompertz) for all 1121 shells. Thus, based on the age models of the shells, the annual carbonate production per shell can be calculated, which ranges from 74gyr^{-1} ($\sigma = 2.9$) (Gompertz) to 83gyr^{-1} ($\sigma = 2.8$) (logistic), accounting for $\sim 150\text{gyr}^{-1}$ living $^{-1}$ oyster individual (= two valves).

4 Discussion

4.1 From lengths to cohorts

Based on the assumption that the size-frequency cohorts represent age classes it is apparent to establish length-at-age relationships. Oysters, especially subtropical to tropical species, can have extended spawning seasons resulting in more than one recruitment event per year. In protracted spawning situations the cohort signature in multiple year class situations is that of broad rather than narrow peaks in the demographic data, which fits to our data. Therefore, the assignment to certain years is only tentative. Applying the von Bertalanffy equation to the length data reveals a strongly right-skewed distribution with 50% of the shells ranging between 3 and 6 years (Fig. 8a). The frequency of specimens between 6 and 9 years decreases rapidly and the contribution by shells older than 9 years is subordinate although outliers with up to 16 years occur.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Again the non-normal distribution of the von Bertalanffy growth model data suggests cohort mixing and the mixture analysis assumes at least four significant cohorts with low AIC value. Due to the rareness of large and aged shells, the fourth cohort displays a rather low amplitude and we assume that at least two natural cohorts may be amalgamated in this group. The pattern suggests more or less continuous recruitment accentuated by very successful settlement peaks every 2 or 3 years. Similarly, comparable patterns in extant *Crassostrea* reefs are linked to fluctuating mortality rates and changing recruitment success (Southworth et al., 2010). The data show that old and large shells are rare. The reason for this may be a generally high mortality during the early years of growth, resulting in low survival rates and few old specimens. The high amount of fragments of large shells, however, suggests that a distinctly higher proportion of large shells existed but became successively destroyed.

4.2 Shell loss and mortality

The rapid decline of old shells is a combination of two processes. First, natural mortality will result in a tail of large shells. Second, shells of died-off *Crassostrea* are known to have surprisingly short half-lives ranging from few years to few decades (Powell et al., 2006, 2015; Waldbusser et al., 2011). Natural degradation processes, such as fragmentation, dissolution and hydrodynamic export may account for 30 % loss of shells per year (Southworth et al., 2010). The high amount of fragments with sharp edges suggests that fragmentation is a major factor in our case. The importance of hydrodynamic export cannot be evaluated due to the limited outcrop area. Dissolution, however, is a minor factor as the well preserved shell-surfaces lack any signs of chemical degradation. Based on the declining amplitudes of the cohorts the total shell loss can be computed as exponential decay function (Fig. 8b) revealing initial half-lives of less than 4 years. The high proportion of fragmented, abraded and/or bioeroded oyster shells (fragments: $n = 7047$) would balance the “missing frequencies” easily and points to fragmentation as key factor. We have no evidence for age-specific shell destruction and assume that all age classes were equally affected by this process.

Overall, under the environmental conditions as present in the Early Miocene estuary, a died-off *Crassostrea gryphoides* reef would have been fully degraded within one or two decades if not buried below the taphonomically active zone. In respect to the high sedimentation rates in the rapidly subsiding basin with rates around 0.6 m kyr^{-1} (Zuschin et al., 2014), a rapid burial of the reef is not unlikely, thus capturing the population structure.

4.3 *Crassostrea* as carbonate factory?

The image analysis based estimation of chalky and foliate layers shows that the values for chalky layers are slightly lower than reported for other fast-growing *Crassostrea* species. The amount of chalky layers in left shells of the extant *Crassostrea* “*gryphoides*” (sensu Newton and Smith, 1912) is $\sim 70\%$ (Durve and Bal, 1960). The shells of the Miocene *Crassostrea titan* (Conrad, 1853) and *C. gravitesta* (Yokoyama, 1926) were reported to comprise even up to $\sim 90\%$ of chalky deposits (Chinzei, 1995; Kirby, 2001) but this may be a slight overestimation, neglecting the high amount of dense calcite in the hinge areas. A re-evaluation of the illustrated specimen of *C. gravitesta* in Chinzei (1995) using our method reveals a somewhat lower but still high value of 77% chalky layer. The gravity of the fast growing *C. titan*, *C. gravitesta* and *C. cahobasensis* (Pilsbry and Brown, 1917) was estimated to range around $1.35\text{--}1.40 \text{ g cm}^{-3}$ (Chinzei, 1995; Kirby and Jackson, 2004). These values are very close to the empirical data on the fast growing *C. ariakensis* (Fujita, 1913) ($1.44 \pm 0.12 \text{ g cm}^{-3}$) measured by Lombardi et al. (2013). Slightly higher values are given for *C. gigas* ($1.63 \pm 0.35 \text{ g cm}^{-3}$, mean = 1.58) by Chinzei (1995). Generally, species with low amounts of chalky layer have much higher densities; e.g., *C. virginica* shells range around $2.18\text{--}2.35 \text{ g cm}^{-3}$ (Kirby and Jackson, 2004; Lombardi et al., 2013). Our gravity estimate of $\sim 1.82 \text{ g cm}^{-3}$ for *C. gryphoides* is thus somewhat higher than expected for such a growth type. In fact, large parts of the shells are very light-weighted and fit well to the patterns discussed by Chinzei (1995) and Kirby (2001). The main difference is the large proportion of heavy shell material in the huge umbos and hinges.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A comparison of the growth curve of *C. gryphoides* with the von Bertalanffy growth models of fossil and extant *Crassostrea* species (Fig. 9) reveals this oyster as outstandingly fast growing species. Thus, *C. gryphoides* was an important carbonate producer in Neogene estuaries and lagoons where it lived as secondary soft-bottom dweller in dense colonies in a mixed mode of shell-supported reclining and mud sticking (sensu Seilacher et al., 1985; Seilacher and Gishlick, 2014). Therefore, dense populations with more than 100 specimens per m^2 can be expected. Even within the shell bed, which is clearly not in-situ, the average density is 129 shells (~ 64 individuals) per m^2 (including also moderately fragmented shells). This would point to a hypothetic annual carbonate production of up to 15 kg m^{-2} with the oyster reef. Although this calculation is just a very rough estimate, it indicates that the carbonate production is in the range of fast growing coral reefs with productions of $6\text{--}10 \text{ kg m}^{-2} \text{ yr}^{-1}$ (Montaggioni, 2005; Jones et al., 2015). A major difference, however, is the rapid shell loss in *Crassostrea* reefs, which prevents the formation of rigid and stable structures comparable to coral reefs.

5 Conclusions

Crassostrea gryphoides was the fastest growing and largest Crassostreinae species known so far. Despite the fact, that this species could attain outstanding individual ages of four decades (Harzhauser et al., 2010), the bulk of specimens analyzed herein lived less than 10 years, typically growing up to about 300 mm in length.

The non-normal distribution in the size, area and age frequency data are best explained by the presence of distinct recruitment cohorts, comparable to modern oyster reefs. About four cohorts are detected by mixture analysis and the rapidly decreasing amplitudes of frequency of these cohorts is interpreted to reflect the combined effect of mortality, respectively the declining life expectancy with age, and the shell loss by biotic and physical factors. As no accumulation of large and aged shells occurred, whilst the amount of fragments is high, we assume that shell loss is an important factor to explain the strongly right-skewed distribution. Shell half-lives ranged around 2–4 years

and within less than two decades the seemingly rigid and persisting structure of a *Crasostrea gryphoides* reef could have been completely degraded. This may explain the rareness of in-situ *C. gryphoides* reefs in the fossil record although the shells are frequent and ubiquitous.

The significant growth performance is clearly boosted by the formation of up to 64 % percentage of fast-growing and lightweight chalky material. The subtropical climate with warm winter temperatures above c. 10 °C and the nutrient-rich setting in an estuary will additionally have supported the excessive growth.

Due to its fast growth and large shells, the carbonate production of *C. gryphoides* is outstanding. Dense colonies might have produced around 15 kg m⁻² yr⁻¹ of carbonate, which is within the range of fast growing coral reefs. Therefore, this oyster may have been a major carbonate producer in the circum-Tethyan area throughout the Miocene. In contrast to coral reefs, however, the high shell loss rates did not allow to form stable persistent structures.

The Supplement related to this article is available online at doi:10.5194/bgd-12-15867-2015-supplement.

Acknowledgements. The study was financed by the Austrian Science Fund (FWF project no. P 25883-N29 “Smart-Geology für das größte fossile Austernriff der Welt”). We would like to thank Florian Rist from the Institute of Art and Design, Vienna University of Technology for providing access to the photo studio and for his support during data acquisition.

References

- Afsar, N., Siddiqui, G., and Roberts, D.: Parasite inspection in five commercially important oyster species (Mollusca: Bivalvia) of Pakistan, *J. Basic Appl. Sci.*, 10, 220–225, 2014.
- Aichholzer, O., Aurenhammer, F., Alberts, D., and Gärtner, B.: A novel type of skeleton for polygons, *J. Univ. Comp. Sci.*, 1, Springer, Berlin, Heidelberg, 752–761, 1996.

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Alam, M. D. and Das, N. G.: Growth and age determination of an intertidal cupped oyster *Crassostrea madrasensis* (Preston) (Bivalvia: Ostreidae) around Moheshkhali Channel, Bay of Bengal, Indian J. Mar. Sci., 28, 329–331, 1999.

Baqueiro Cárdenas, E. R. and Aldana Aranda, D.: Differences in the exploited oyster (*Crassostrea virginica* (Gmelin, 1791)) populations from different coastal lagoons of the Gulf of Mexico, Transitional Waters Bull., 2, 21–35, 2007.

Berthome, J. P., Prou, J., and Bodoy, A.: Performances de croissance de l'huître creuse, *Crassostrea gigas* (Thunberg) dans le bassin d'élevage de Marennes-Oléron entre 1979 and 1982, Haliotis, 15, 183–192, 1986.

Chatterji, A., Ansari, Z. A., Ingole, B. S., and Parulekar, A. H.: Length–weight relationship of giant oyster *Crassostrea gryphoides* (Schlotheim), Mahasagar – Bull. Nat. Inst. Oceanogr., 18, 521–524, 1985.

Chávez-Villalba, J., López-Tapia, M., Mazón-Suástegui, J., and Robles-Mungaray, M.: Growth of the oyster *Crassostrea corteziensis* (Hertlein, 1951) in Sonora, Mexico, Aquac. Res., 36, 1337–1344, 2005.

Chinzei, K.: Adaptive significance of the lightweight shell structure in soft bottom oysters, Neues Jahrb. Geol. P.-A., 195, 217–227, 1995.

Chinzei, K.: Adaptation of oysters to life on soft substrates, Hist. Biol., 25, 223–231, 2013.

Coakley, J. M.: Growth of the eastern oyster, *Crassostrea virginica*, in Chesapeake Bay, Thesis, Faculty of the Graduate School of the University of Maryland, College Park, available at: <http://drum.lib.umd.edu/bitstream/1903/1471/1/umi-umd-1590.pdf> (last access: 24 September 2015), 1–263, 2004.

Conrad, T. A.: Notes on shells with descriptions of new species, P. Acad. Nat. Sci. Phila., 6, 199–200, 1853.

Delaunay, B.: Sur la sphere vide, Otdelenie Matematicheskikh i Estestvennykh Nauk, 7, 793–800, 1934.

Dellmour, R. and Harzhauser, M.: The Iván Canyon, a large Miocene canyon in the Alpine–Carpathian Foredeep, Mar. Petrol. Geol., 38, 83–94, 2012.

Dempster, A. P., Laird, N. M., and Rubin, D. B.: Maximum likelihood from incomplete data via the EM algorithm, J. Roy. Stat. Soc. B, 39, 1–38, 1977.

Durve, V. S.: Malacological differences between the oysters *Crassostrea gryphoides* (Schlotheim) and *Crassostrea madrasensis* Preston, Indian J. Fish., 202, 624–625, 1974.

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Durve, V. S. and Bal, D. V.: Some observations on shell-deposits of the oyster *Crassostrea gryphoides* (Schlotheim), P. Indian Acad. Sci. B, 54, 45–55, 1960.
- Einsele, G., Ricken, W., and Seilacher, A.: Cycles and Events in Stratigraphy, Springer-Verlag, Berlin, 1991.
- 5 FAO: Aquaculture Fact Sheets, Food and Agriculture Organization of the United Nations, Fisheries and Aquaculture Department, available at: <http://www.fao.org/fishery/culturedspecies/search/en> (last access: 14 May 2015), 2015.
- Fujita, T.: Nihon Suisan Dobutsugaku (Aquatic Zoology in Japan), Shokabo, Tokyo, Japanese aquatic fisheries animals, vol 2., Shokabu, Tokyo, 1913.
- 10 Ginger, K. W. K., Vera, C. B. S., Dineshram, R., Dennis, C. K. S., Adela, L. J., Yu, Z., and Thiagarajan, V.: Larval and post-larval stages of Pacific oyster (*Crassostrea gigas*) are resistant to elevated CO₂, PLoS ONE, 8, e64147, doi:10.1371/journal.pone.0064147, 2013.
- Glira, P., Pfeifer, N., Briese, C., and Ressler, C.: A correspondence framework for ALS strip adjustments based on variants of the ICP Algorithm, J. Photogram., Remote Sensing, Geoinf. Process., accepted for publication, 2015.
- 15 Gmelin, J. F.: Caroli a Linnei systema natura per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, disserentis, synonymis, locis etc. Editio decima tertia, aucta, reformata, cura J. F. Gmelin, 1. Vermes testacea, G. E. Beer, Lipsiae, 3021–4120, 1791.
- 20 Groslier, T., Toft Christensen, H., Davids, J., Dolmer, P., Elmedal, I., Holm, M. W., and Hansen, B. W.: Status of the pacific oyster *Crassostrea gigas* (Thunberg, 1793) in the western Limfjord, Denmark – five years of population development, Aquatic Invasions, 9, 175–182, 2014.
- Hammer, Ø.: PAST, Paleontological Statistics Version 3.06, Reference manual, Natural History Museum, University of Oslo, available at: <http://folk.uio.no/ohammer/past/past3manual.pdf> (last access: 24 September 2015), 225 p., 2015.
- 25 Hammer, Ø., Harper, D. A. T., and Ryan, P. D.: PAST: Paleontological Statistics Software Package for education and data analysis, Palaeontol. Electron., 4, 1–9, 2001.
- Harding, J. M., Mann, R., and Southworth, M. J.: Shell length–at-age relationships in James River, Virginia oysters (*Crassostrea virginica*) collected four centuries apart, J. Shellfish Res., 27, 1109–1115, 2008.
- 30

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Harzhauser, M., Böhme, M., Mandic, O., and Hofmann, C.-C.: The Karpatian (Late Burdigalian) of the Korneuburg Basin – a palaeoecological and biostratigraphical synthesis, *Beitr. Paläont.*, 27, 441–456, 2002.

Harzhauser, M., Piller, W. E., Müllegger, S., Grunert, P., and Micheels, A.: Changing seasonality patterns in Central Europe from Miocene Climate Optimum to Miocene Climate Transition deduced from the *Crassostrea* isotope archive, *Global Planet. Change*, 76, 77–84, 2010.

Harzhauser, M., Djuricic, A., Mandic, O., Zuschin, M., Dorninger, P., Nothegger, C., Székely, B., Puttonen, E., Molnár, G., and Pfeifer, N.: Disentangling the history of complex multi-phased shell beds based on the analysis of 3-D point cloud data, *Palaeogeogr. Palaeoclimatol.*, 437, 165–180, doi:10.1016/j.palaeo.2015.07.038, 2015.

Hertlein, L. G.: Descriptions of two new species of marine pelecypods from west Mexico, Southern Calif. Acad. Sci. Bull., 50, 68–75, 1951.

Higuera-Ruiz, R. and Elorza, J.: Biometric, microstructural, and high-resolution trace element studies in *Crassostrea gigas* of Cantabria (Bay of Biscay, Spain): anthropogenic and seasonal influences, *Estuar. Coast. Shelf S.*, 82, 201–213, 2009.

Hoşgör, I.: Presence of *Crassostrea gryphoides* (Schlotheim) from the lower-middle Miocene sequence of Kahramanmaraş Basin (SE Turkey); its taxonomy, paleoecology and paleogeography, *Min. Res. Explor. Bull.*, 136, 17–28, 2008.

Jones, N. S., Ridgwell, A., and Hendy, E. J.: Evaluation of coral reef carbonate production models at a global scale, *Biogeosciences*, 12, 1339–1356, doi:10.5194/bg-12-1339-2015, 2015.

Kazhdan, M., Bolitho, M., and Hoppe, H.: Poisson surface reconstruction, in: P. fourth Eurographics Sym. Geom. Proc., 26–28 June 2006, Cagliari, Sardinia, 7, 61–70, 2006.

Kennedy, V. S., Newell, R. I. E., and Eble, A. F.: The Eastern Oyster *Crassostrea virginica*, Maryland Sea Grant College Publication UM-SG-TS-96-01, College Park, Maryland, USA, 1–750, 1996.

Kern, A., Harzhauser, M., Mandic, O., Roetzel, R., Ćorić, S., Bruch, A. A., and Zuschin, M.: Millennial-scale vegetation dynamics in an estuary at the onset of the Miocene Climate Optimum, *Palaeogeogr. Palaeoclimatol.*, 304, 247–261, 2010.

Kidwell, S. M.: Models for fossil concentrations: paleobiologic implications, *Paleobiology*, 12, 6–24, 1986.

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Kidwell, S. M.: The stratigraphy of shell concentrations, in: Taphonomy, Releasing the Data Locked in the Fossil Record, edited by: Allison, P. A. and Briggs, D. E. G., Plenum Press, New York, 211–290, 1991.

Kirby, M. X.: Paleoecological differences between Tertiary and Quaternary *Crassostrea* oysters, as revealed by stable isotope sclerochronology, *Palaios*, 15, 132–141, 2000.

Kirby, M. X.: Differences in growth rate and environment between Tertiary and Quaternary *Crassostrea* oysters, *Paleobiology*, 27, 84–103, 2001.

Kirby, M. X. and Jackson, J. B. C.: Extinction of a fastgrowing oyster and changing ocean circulation in Pliocene tropical America, *Geology*, 32, 1025–1028, 2004.

Kirby, M. X., Soniat, T. M., and Spero, H. J.: Stable isotope sclerochronology of Pleistocene and recent oyster shells (*Crassostrea virginica*), *Palaios*, 13, 560–569, 1998.

Kirk, J.: Advanced Dijkstra's minimum path algorithm, available at: <http://www.mathworks.com/matlabcentral/fileexchange/20025-advanced-dijkstras-minimum-path-algorithm> (last access: 2 April 2015), 2015.

Koepfen, W.: Das geographische System der Klimate, in: Handbuch der Klimatologie, edited by: Koepfen, W. and Geiger, R., Gebrüder Bornträger, Berlin, 1–44, 1936.

Laurain, M.: *Crassostrea gryphoides* et *C. gingensis* (Schlotheim, 1813) deux expressions morphologiques d'une même espèce (Miocène, Bivalvia), *Geobios*, 13, 21–43, 1980.

Littlewood, D. T. J.: Molecular phylogenetics of cupped oysters based on partial 28S rDNA gene sequences, *Mol. Phylogenet. Evol.*, 3, 221–229, 1994.

Lombardi, S. A., Chon, G. D., Jin-Wu Lee, J., Lane, H. A., and Paynter, K. T.: Shell hardness and compressive strength of the eastern oyster, *Crassostrea virginica*, and the Asian oyster, *Crassostrea ariakensis*, *Biol. Bull.*, 225, 175–183, 2013.

Lopes, G. R., Araujo de Miranda Gomes, C. H., Tureck, C. R., and de Melo, C. M. R.: Growth of *Crassostrea gasar* cultured in marine and estuary environments in Brazilian waters, *Pesqui. Agropecu. Bras.*, 48, 975–982, 2013.

Mahar, M. A. and Awan, K. P.: Cultivation of oyster *Crassostrea gryphoides* (Schlotheim) through rafts at Ambra creek coastal belt of Arabian sea, Sindh Pakistan, *Sindh Univ. Res. J. (Sci. Ser.)*, 44, 119–124, 2012.

Mancera, E. and Mendo, J.: Population dynamics of the oyster *Crassostrea rhizophorae* from the Ciénaga Grande de Santa Marta, Colombia, *Fish. Res.*, 26, 139–148, 1996.

Mandic, O., Harzhauser, M., Schlaf, J., Piller, W. E., Schuster, F., Wielandt-Schuster, U., Nebelsick, J. H., Kroh, A., Rögl, F., and Bassant, P.: Palaeoenvironmental Reconstruction of an

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Powell, E. N., Kraeuter, J. N., and Ashton-Alcox, K. A.: How long does oyster shell last on an oyster reef?, *Estuar. Coast. Shelf S.*, 69, 531–542, 2006.
- Powell, E. N., Klinck, J. M., Guo, X., Ford, S. E., and Bushek, D.: The potential for oysters, *Crassostrea virginica*, to develop resistance to Dermo disease in the field: evaluation using a gene-based population dynamics model, *J. Shellfish Res.*, 30, 685–712, 2011.
- Powell, E. N., Mann, R., Ashton-Alcox, K. A., Kim, Y., and Bushek, D.: The allometry of oysters: spatial and temporal variation in the length–biomass relationships for *Crassostrea virginica*, *J. Mar. Biol. Assoc. UK*, doi:10.1017/S0025315415000703, 2015.
- Preston, H. B.: Report on a collection of Mollusca from the Cochin and Ennur Backwaters, *Rec. Indian Mus.*, 12, 27–39, 1916.
- Ragaini, L. and Di Celma, C.: Shell structure, taphonomy and mode of life of a Pleistocene ostreid from Ecuador, *Boll. Soc. Paleontol. I.*, 48, 79–87, 2009.
- Reece, K. S., Cordes, J. F., Stubbs, J. B., Hudson, K. L., and Francis, E. A.: Molecular phylogenies help resolve taxonomic confusion with Asian *Crassostrea* oyster species, *Mar. Biol.*, 153, 709–721, 2008.
- Ren, J., Liu, X., Jiang, F., Guo, X., and Liu, B.: Unusual conservation of mitochondrial gene order in *Crassostrea* oysters: evidence for recent speciation in Asia, *BMC Evol. Biol.*, 10, 394, doi:10.1186/1471-2148-10-394, 2010.
- Robinson, E. M., Lunt, J., Marshall, C. D., and Smee, D. J.: Eastern oysters *Crassostrea virginica* deter crab predators by altering their morphology in response to crab cues, *Aquat. Biol.*, 20, 11–118, 2014.
- Ross, P. G. and Luckenbach, M. W.: Population Assessment of Eastern Oysters (*Crassostrea virginica*) in the Seaside Coastal Bays, Final Report, Virginia Coastal Zone Management Program, College of William and Mary, Wachapreague, available at: <http://www.deq.virginia.gov/Portals/0/DEQ/CoastalZoneManagement/task10-02-07.pdf>, 101 pp., 2009.
- Sacco, F.: I molluschi dei terreni terziarii del Piemonte e della Liguria. Parte XXIII. Pelecypoda (Ostreidae, Anomiidae e Dimyidae), Carlo Clausen, Torino, 1–46, 1897.
- Salvi, D., Macali, A., and Mariottini, P.: Molecular phylogenetics and systematics of the bivalve family Ostreidae based on rRNA sequence-structure models and multilocus species tree, *PLoS ONE*, 9, e108696, doi:10.1371/journal.pone.0108696, 2014.
- Schultz, O.: *Bivalvia neogenica* (Nuculacea – Unionacea), in: *Catalogus Fossilium Austriae*, edited by: Piller, W. E., Akademie der Wissenschaften, Wien, 1/1, 1–379, 2001.

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Seilacher, A.: Constructional morphology of bivalves: evolutionary pathways in primary versus secondary soft-bottom dwellers, *Palaeontology*, 27, 207–237, 1984.
- Seilacher, A. and Gishlick, A. D.: *Morphodynamics*, CRS Press, London, 2014.
- Seilacher, A., Matyja, B. A., and Wierzbowski, A.: Oyster beds: morphologic response to changing substrate conditions, *Lect. Notes Earth Sci.*, 1, 421–435, 1985.
- Siddiqui, G. and Ahmed, M.: Oyster species of the sub tropical coast of Pakistan (northern Arabian Sea), *Indian J. Mar. Sci.*, 31, 108–118, 2002.
- Southworth, M., Harding, J. M., Wesson, J. A., and Mann, R.: Oyster (*Crassostrea virginica*, Gmelin 1791) population dynamics on public reefs in the Great Wicomico River, Virginia, USA, *J. Shellfish Res.*, 29, 271–290, 2010.
- Sovis, W. and Schmid, B.: Das Karpat des Korneuburger Beckens, Teil 1, *Beitr. Paläont.*, 23, 1–413, 1998.
- Sovis, W. and Schmid, B.: Das Karpat des Korneuburger Beckens, Teil 2, *Beitr. Paläont.*, 27, 1–467, 2002.
- Sowerby, G. B.: Monograph of the genus *Ostraea*, *Conchologia Iconica*, 18, 6–33, 1871.
- Stenzel, H. B.: Oysters, in: *Treatise on Invertebrate Paleontology*, edited by: Moore, R. C., Geological Society of America and the University of Kansas Press, Lawrence, N/3, 953–1224, 1971.
- Thunberg, C. P.: Tekning och Beskrifning på en stor Ostronsort ifrån Japan, *Kongliga Vetenskaps Akademiens Nya Handlingar*, Stockholm, 14, 140–142, 1793.
- Trivedi, S., Aloufi, A. A., Ansari, A. A., and Ghosh, S. K.: Molecular phylogeny of oysters belonging to the genus *Crassostrea* through DNA barcoding, *J. Entomol. Zool. Stud.*, 3, 21–26, 2015.
- Vermeij, G.: The oyster enigma variations: a hypothesis of microbial calcification, *Paleobiology*, 40, 1–13, 2014.
- von Bertalanffy, L.: Untersuchungen über die Gesetzmäßigkeiten des Wachstums. I. Allgemeine Grundlagen der Theorie; mathematische und physiologische Gesetzmäßigkeiten des Wachstums bei Wassertieren, *Arch. Entwicklungsmech. Org.*, 131, 613–652, 1934.
- von Schlotheim, E. F.: Beiträge zur Naturgeschichte der Versteinerungen in geognostischer Hinsicht, in: *Leonhard's Taschenbuch für die gesammte Mineralogie mit Hinsicht auf die neuesten Entdeckungen*, edited by: Leonhard, C. C., Series 1, 7, Frankfurt a. M., Johann Christian Hermann, 1–134, 1813.

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Voronoi, G.: Nouvelles applications des paramètres continus à la théorie des formes quadratiques. Deuxième mémoire. Recherches sur les paralléloèdres primitifs, *J. Reine Angew. Math.*, 133, 97–178, 1908.

Waldbusser, G. G., Steenson, R. A., and Gren, M. A.: Oyster shell dissolution rates in estuarine waters: effects of pH and shell legacy, *J. Shellfish Res.*, 30, 659–669, 2011.

Wang, Y., Xus, Z., and Guo, X.: Differences in the rDNA-bearing chromosome divide the Asian–Pacific and Atlantic species of *Crassostrea* (Bivalvia, Mollusca), *Biol. Bull.*, 206, 46–54, 2004.

Wessely, G.: Geologie des Korneuburger Beckens, *Beitr. Paläont.*, 23, 9–23, 1998.

Yokoyama, M.: Fossil Mollusca from the oil-fields of Akita, *J. Fac. Sci., Imperial Univ. Tokyo, Sect. 2*, 1/9, 377–389, 1926.

Yoon, G.-L., Kim, B.-T., Kim, B.-O., and Han, S.-H.: Chemical–mechanical characteristics of crushed oyster-shell, *Waste Manage.*, 23, 825–834, 2003.

Zhang, G., Fang, X., Guo, X., Li, L., Luo, R., Xu, F., Yang, P., Zhang, L., Wang, X., Qi, H., Xiong, Z., Que, H., Xie, Y., Holland, P. W., Paps, J., Zhu, Y., Wu, F., Chen, Y., Wang, J., Peng, C., Meng, J., Yang, L., Liu, J., Wen, B., Zhang, N., Huang, Z., Zhu, Q., Feng, Y., Mount, A., Hedgecock, D., Xu, Z., Liu, Y., Domazet-Lošo, T., Du, Y., Sun, X., Zhang, S., Liu, B., Cheng, P., Jiang, X., Li, J., Fan, D., Wang, W., Fu, W., Wang, T., Wang, B., Zhang, J., Peng, Z., Li, Y., Li, N., Wang, J., Chen, M., He, Y., Tan, F., Song, X., Zheng, Q., Huang, R., Yang, H., Du, X., Chen, L., Yang, M., Gaffney, P. M., Wang, S., Luo, L., She, Z., Ming, Y., Huang, W., Zhang, S., Huang, B., Zhang, Y., Qu, T., Ni, P., Miao, G., Wang, J., Wang, Q., Steinberg, C. E., Wang, H., Li, N., Qian, L., Zhang, G., Li, Y., Yang, H., Liu, X., Wang, J., Yin, Y., and Wang, J.: The oyster genome reveals stress adaptation and complexity of shell formation, *Nature*, 490, 49–54, 2012.

Zuschin, M., Harzhauser, M., Hengst, B., Mandic, O., and Roetzel, R.: Long-term ecosystem stability in an Early Miocene estuary, *Geology*, 42, 1–4, 2014.




Figure 1. Part of the digital surface model of the shell bed; the white cross indicates the area, within which all objects were digitally outlined and evaluated. It contains 1121 complete shells of *C. gryphoides* and 7047 fragments of that species (a). Outcrop pictures of the shell bed illustrating the density and extent of the shell bed (b–c), (b) width c. 2 m; (c) see workers as scale.

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 2. Geographic position of the Stetten site within the Korneuburg Basin north of Vienna in Austria; shaded areas represent pre-Miocene basement; dashed lines are major faults (modified from Dellmour and Harzhauser, 2012).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Figure 3. Models of 6 shells based on high resolution laser scanning data of shells from the collections of the Natural History Museum. These specimens document the broad range of morphologies and were used for volume calculations; L = left shell, R = right shell.

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Size frequency diagrams for center line length and area data (log transformed) with cohorts (dashed lines) as detected by mixture analysis (a–b). Box-plot illustrating the strongly right-skewed distribution for fragments ($n = 7047$) and a clear separation from the size distribution pattern of complete shells ($n = 1121$) (c).




Figure 5. Regression analysis revealing a significant correlation between length and area of complete shells.

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



BGD

12, 15867–15900, 2015

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 6. Longitudinal section through *C. gryphoides* from the oyster reef site showing the high amount of chalky layers (grey) and the low amount of foliate layers (black).

BGD

12, 15867–15900, 2015

Age structure,
carbonate production
and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 7. Logistic function showing the relation between center line length and shell volume based on empirical measurements of 9 shells (dots); L = left shell, R = right shell; shell pairs are linked; numbers correspond to specimens illustrated in Fig. 3.




Figure 9. Comparison of von Bertalanffy growth model of *Crassostrea gryphoides* with selected fossil and extant *Crassostrea* species. 1: *C. titan* (Conrad, 1853), Miocene, California, USA (Kirby, 2001), 2: *C. madrasensis* (Preston, 1916), recent, Bangladesh (Nurul Amin et al., 2008); 3: *C. virginica* (Gmelin, 1791), recent and Pleistocene Virginia, USA (Kirby, 2001; Powell et al., 2011); 4: *C. corteziensis* (Hertlein 1951), recent, Mexico (Chávez-Villalba et al., 2005); 5: *C. gigas* (Thunberg, 1793), recent, Marennes-Oleron, France (Berthome et al., 1986).

BGD

12, 15867–15900, 2015

Age structure, carbonate production and shell loss rate

M. Harzhauser et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

