

Dear Dr. Jack Middelburg,

We submit the revised version of the manuscript:

Deep ocean mass fluxes in the coastal upwelling off Mauritania from 1988 to 2012: variability on seasonal to decadal timescales

by Fischer and co-authors for publication in BG.

Our response includes replies and comments to the reviewers' criticisms and remarks, and the explanations for the changes we made in the revised version. In addition, several smaller changes were made to the text, tables and figures. The major revised parts in the text are marked in red and comments are given (see marked-up version).

We thank the Editorial Team of BG for considering our paper for publication. We appreciate the reviewers' comments and suggestions and acknowledge their help to improve the ms. We hope that this revised version will meet your expectations and the journals' requirements and that it will find your positive consideration,

Yours sincerely,

Gerhard Fischer

1 **Changes according to Anonymous Referee #1 (RC)**

2
3 RC

4 Fischer et al. present 25 years of sediment trap fluxes collected off Cape Blanc in the area of
5 permanent upwelling. This is a very impressive data set and it is quite obvious that it is dominated by
6 interannual variations and not by a clear seasonality. It is therefore consistent to investigate the
7 relationship between fluxes and indices of climate oscillations of multiannual or decadal scale (AMO,
8 NAO, ENSO). The authors manage to present this complex matter in a comprehensive and concise
9 way. The paper is well written and well organized and the evaluation of the results is critical and
10 refrains from over-interpretation. Tables and Figures are of good quality and present the results
11 appropriately.

12
13 I have some questions regarding the data used in the paper which need to be clarified. The authors
14 state that fluxes in the upper and lower traps are very similar and therefore use the upper fluxes to fill
15 in gaps in the lower record. The paper cited to show this (Fischer et al., 2009b) presents only one
16 record of upper and lower traps. The authors should show that this can be done for the whole data set
17 despite the vertical distance of about 2000m between the upper and lower traps.

18
19 AC

20 The reviewer is right in mentioning this problem and stating that the paper by Fischer et al. (2009b)
21 presented only one annual time series (deployment CB-13, 2002-2003, see also Table 1 of the
22 submitted ms). Fischer et al. (2009b) showed a good match between upper and lower trap fluxes
23 regarding absolute values and timing of peaks, but also an increase in flux with depth during winter-
24 spring due to an additional lateral source, which is located closer to coastal upwelling east of the CB
25 trap location. Due to this process, the lower CB traps are better suited to record any long-term
26 changes in coastal upwelling intensity.

27
28 We admit, however, that we cannot be sure that the upper traps mimic the temporal and absolute flux
29 pattern of the lower traps in all seasons in each year in all details. For our study, the winter fluxes may
30 be critical due to their potential relationship to the winter NAO (Figs. 6, 7, 9a). When plotting all
31 available lower and upper trap total mass fluxes for winter, a close correspondence is observed
32 ($R^2=0.84$ for $N=10$), with slightly higher fluxes in the deeper trap due to lateral particle advection
33 processes (described in detail by Fischer et al., 2009b, and modelled by Karakas et al., 2006, 2009).
34 However, considering the entire record presented here, it seems that the upper trap fluxes of the
35 winter seasons 1998 and 2004 may be critical due to a smaller filament area (Fig. 7b). The area with
36 elevated chlorophyll and high particle concentrations may not have reached the upper offshore trap.
37 Because of lateral advection from the east and the larger catchment area of the deeper traps (Siegel
38 and Deuser, 1997), particle fluxes might have been higher in the deeper water column in winter 1998
39 and 2004. A possible alternative is eliminating the two data points (winter 1998 and 2004) in Figure 7.
40 This slightly improves the correlation between winter fluxes and the NAO, without significantly
41 influencing our findings and conclusions. However, we keep the two data points in Figure 7 also in the
42 revised version, mentioning in the caption and in the corresponding text passages that the upper trap
43 data from winter 1998 and 2004 might have been different from the deeper traps. Additionally, we
44 added the argumentation presented here to chapter 3.2 (mass fluxes) in the revised version.

45
46 RC

47 Moreover, the authors integrate their data for spring, summer, fall and winter and thus lose some of
48 the original high resolution sampling. The authors should explain their reasons.

49
50 AC

51 The reason for this is that the focus of our MS is the longer-term variability on timescales of seasons
52 and beyond (as stated by reviewer #1, page C7709, second sentence). Due to logistical reasons, we
53 have very different time resolutions in our trap samplings (from a few days to several weeks), which
54 may sometimes limit comparisons between specific intervals. Furthermore, represented data points
55 from intervals with a high sampling resolution (e.g. a few days) can hardly be seen in the figures. By
56 doing so, this would partly lead to overloaded figures and may be confusing for the reader. This is now
57 explained in the revised version of chapter 3.2. (mass fluxes).

58
59 RC

60 Page 17653, line 14: incomplete sentence

61
62 AC

63 Sentence deleted

64

65 RC

66 Page 17657, lines 17-20: This part on the time delay is too short and difficult to understand. The
67 motivation/results should be explained in more detail.

68

69 AC

70 This section/sentence indeed needs more explanation. We rephrased this accordingly and explained
71 our arguments on/ideas concerning a possible time delay between NAO forcing in winter (DJFM) and
72 flux response in the deep ocean in spring in the revised version of the discussion section 5.2 in more
73 detail.

74

Comments to Anonymous Referee # 2 (RC)

RC

General Comments

The authors present a >20 yr record of mass fluxes in the North Atlantic – this is a rare and valuable data set. The authors are trying to write a synthesis of a huge data set, and this is a worthwhile, if very challenging task. Understandably, the authors look for correlations between flux data and climate metrics. However, the manuscript suffers from an apparent listing of potential hypotheses that are at best modestly supported by the data in sections where there is not space to fully develop these ideas, especially the Introduction. Given the restrictions on space, the manuscript would be better served by articulating the working hypothesis that is best supported by the data in the beginning, i.e., that the organic carbon and BSi fluxes are mostly highly correlated with dust deposition, and discuss the alternative hypotheses in the discussion section. A table of flux (including total, lithogenic, Corg, and BSi) correlations with different climate statistics, i.e., NAO, ENSO Index, sea surface temperature, sea surface pressure, might be a more coherent and easily digestible way to present this data.

AC

We understand the argumentation of reviewer #2 to make the manuscript easier to digest. Our major motivation was to investigate the variability of deep ocean mass fluxes in the NE Atlantic and link them to major climate phenomena, such as the NAO. From a statistical point of view, we see no impact of the NAO on the deep ocean mass fluxes (BSi and Corg). This is due to local episodic perturbations, e.g. by dust deposition, which is assumed to weaken the relationship between fluxes and the NAO (as pointed out in the abstract). We rephrased the introduction and focus now more on the flux correlations and the NAO index (reviewer #2: 'the working hypotheses that is best supported by the data'). We tried to further clarify and focus the Introduction by slight modifications such that it now follows a very clear structure on the major factors influencing our study area as discussed in earlier publications and motivating us to analyze our records in the light of these earlier findings (1: upwelling trends ('Bakun hypothesis'), 2: ENSO, 3: NAO). In the discussion section, we discussed other possible drivers (e.g. ENSO, AMO) as suggested by reviewer #2 (see also below). The AMO is now mainly addressed in chapter 5.5. We did not want to over-interpret the results and the number of realizations of AMO cycles in our dataset is too low for statistically robust statements.

Compiling a large matrix of correlation coefficients between all kinds of climate statistics and fluxes of all parameters will mostly show statistically insignificant values ($r^2 < 0.1$), except for the NAO and BSi/Corg. Additionally, we believe that it makes little sense to compare i.e. lithogenic fluxes (=dust: dependent on aridity, winds on land, etc.) with i.e. SST in the ocean. Instead, it makes more sense to try to relate the NAO (influence on coastal upwelling and biomass) to BSi (=diatom, primary producers) fluxes or organic carbon fluxes, which is presented in the respective figures. Another problem is the potential effect of time lags (e.g. from ENSO, say we use a 3.4. index) from the Pacific Ocean to the NE Atlantic coastal upwelling. Potential time lags of zero months up to 1-2 years are discussed in the literature. It would be difficult to capture all these timescales without enlarging the discussion and the paper significantly.

Nevertheless, we focus on the flux relationships with NAO in the revised abstract and introduction section and, as suggested by reviewer #2, move other influences more to the discussion section. We added a few sentences in the revised introduction section, which will make it more clear why we focused on specific parameters. Discussing all possible relationships is well beyond the scope of this paper. We therefore selected those parameters, which may be interpreted in the light of earlier findings in the literature and that are related, e.g. to upwelling of nutrient rich deep water and the production of biogenic silica.

RC

I also expect that the study would benefit by comparisons of their data with data collected in similar locations, i.e., the long-term study sites off the California coast, e.g., CalCOFI data sets and data sets from the San Pedro and Santa Monica Basins. There have been a number of studies of those California sites looking at relationships between productivity, water column oxygenation, winds, and upwelling. It would be interesting to see if there are similar trends observed in the Eastern Pacific and Eastern Atlantic basins.

AC

We briefly mentioned the other EBUEs in the revised manuscript and added a few papers from the California Current System (see chapters 2.3., 3.2., 5.3., and new reference list). We did not intend to

make an inter-comparison study of fluxes of the Canary Current with other Eastern Boundary Upwelling Systems such as the California Current System. An inter-comparison between both systems would have further enlarged the manuscript. Both systems are dominated by different climate forcings on the longer-term (different relative importance of NAO, ENSO etc.), making a comparison rather complicated and clear trends less obvious. Furthermore, a high and continuous dust supply is less pronounced in the California Current System compared to the CC system. An inter-comparison needs to include the Benguela and the Humboldt Current Systems as well and could be the topic of a separate study.

RC

p. 17660: How does the fact that particles analyzed in this study were < 1 mm (because of sample filtration) relate to the predominant grain size of dust particles being between 10 and 20 nm? Are those dust grains broken down in size between the surface ocean and deep traps? If the authors are invoking mineral dust as the primary driver of the sinking flux in this manuscript, but the samples exclude particles > 1 mm, is that consistent with the dominant grain size of dust being much larger than the filter size?

AC

There seems to be some misunderstanding. Dust particles of 10-20 μm may easily pass through a filter of 1 mm. The point is that these fine-grained dust particles can only sink at higher rates (10-100 meters per day) when being incorporated into larger organic-rich particles such as marine snow aggregates (in our case off Cape Blanc around 1mm or less in size, Nowald et al., 2015) or fecal pellets (several 100 μm large). Organic carbon and dust together with other particles sink to depth and form the deep ocean flux. However, within the sampling cups, those particles mostly disintegrate due to their fragile nature. Consequently, we do not exclude any particles except some large carcasses from crustaceans, larger pteropods and rare 'swimmers' which are in the size fraction > 1mm. We slightly modified the method section accordingly.

RC

Specific Comments

In abstract, the discussion of AMO is confusing – since there is no or only a weak correlation, I recommend dropping the discussion of AMO and focus on the positive relationship with dust deposition. It may worth mentioning that you looked for a relationship between BSi and AMO in the discussion, but it is distracting in the abstract. Abstract would benefit from not discussing the hypotheses that were not supported by the data – only focus on the hypothesis supported by the data, and the data that supported the hypothesis.

AC

See AC comment above. It has been deleted in the abstract and is referred to only in the revised Discussion section. In the revised abstract, we will mainly refer to the NAO, the flux relationships and the 'Bakun hypothesis'.

RC

P. 3 is confusing – a list of hypotheses that are sometimes supported by the data and sometimes not – unsure what message we are to take away from this other than that there is no statistically significant correlation w/ anything? If this is not the message, the Intro needs to be restructured around a single, coherent message. I appreciate that the authors are trying to look for correlations between their data and climate indices, but this information is better suited to the discussion.

AC

Indeed, there is no a single coherent message in the literature concerning the influence of larger scale climatic oscillations and reactions within the NW African coastal upwelling system (see ACs above). As mentioned above, we have changed the focus in the revised Introduction on the potentially primary driver, i.e. the NAO. We rephrased and slightly restructured the introduction in order to make the different motivations and hypotheses available in the literature and the main points addressed in our study more clear.

RC

Section 2.3, line 20-25: How do the authors evaluate and quantify the relative strength/magnitude of correlation between climate variables and flux metrics? Which statistics are used? Section 2.3 did not convince me that there were meaningful correlations between climate indices and upwelling and/or flux metrics at the trap location.

AC

The temporal correlations shown in Fig. 3 indicate a linear relationship of sea level pressure variations at each gridpoint with the respective climate index (NAO, ENSO, AMO). Sea level pressure variations are a direct indicator for wind variations and therefore also for wind-driven (coastal) upwelling. The patterns in Fig. 3 clearly show that wind changes in our study area are part of large-scale teleconnection patterns (also supported by the literature given in the Introduction) and that consequently, wind-driven upwelling changes may be linked to the respective phase of the climate oscillations.

However, the sediment trap fluxes will always record the superimposed influence of the teleconnection patterns. At this stage, it is difficult to disentangle the relative importance of NAO, ENSO and decadal fluctuations for the fluxes based on the very limited number of realizations of the respective combinations (e.g. El Niño coinciding with a NAO-phase etc.). Longer sediment trap records will potentially help in clarifying this issue. In this sense, our sentence on p.17650 l.25 should be taken as a cautionary note which gives a perspective for future work in this respect.

RC

p. 17652 line 1: The text says total nitrogen was measured, but it is not reported in the tables or figures. This would be a very valuable set of data to include. If the authors chose not to include the total nitrogen flux data they should not report that it was measured.

AC

For further information, the nitrogen data are now included in Tables 2 and 3. However, we did not discuss them in detail.

RC

p. 17652, line 13: Please describe the factor of 2 that the Corg is multiplied by p. 17654, Results: Please specify whether differences in the bulk fluxes are statistically significant higher in winter and summer than fall and spring.

AC

To estimate organic matter composed mainly of C, N, H and O, we used a conversion factor of two as about 50-60% of marine organic matter is constituted by organic carbon (e.g. Hedges et al., 2002). Fig. 4 does not show significant differences between seasons, therefore our focus is on the longer-term and interannual changes of fluxes.

RC

The Results section would benefit from stating the ranges of the total, Corg, BSi, CaCO₃ fluxes.

AC

We presented the values and some ranges in the revised Results section.

RC

p. 17655, Results, lines 12-19: reporting the slopes together with the correlations would be valuable.

AC

We now present the slopes and show them in Table 3, together with the number of data points and the statistical significance

RC

p. 17658: Please discuss how the analysis of Corg fluxes in trap samples collected >1000 m affects interpretations relative to fluxes of BSi, CaCO₃, and lithogenic fluxes, that do not experience flux attenuation with depth the way Corg fluxes do, and whether this is expected to affect a correlation with remote sensing data of sea surface chl

AC

Biological/bacterial degradation of Corg is much faster than dissolution of CaCO₃, BSi and mineral/lithogenic components. Surface chlorophyll should be better related to Corg than to lithogenic and biogenic minerals – the phytoplankton pigments will also be degraded as the total Corg. Therefore, we did not compare CaCO₃ and lithogenic fluxes to surface chlorophyll from satellite

imagery. If the Corg decay over time is constant (which is not precisely known), we may relate it to surface chlorophyll produced by phytoplankton.

RC

p. 17661: Doesn't an increased mass flux with La Nina conditions contradict other text where the authors state that fluxes are not correlated with the strength of upwelling?

AC

No, there is no contradiction. We provided a description of a single ENSO event 1997-1999 and the resulting consequences for the NW upwelling as known from literature. We did not argue that fluxes are not correlated with the strength of upwelling. We speculate that the reviewer possibly had in mind part of the introduction (p. 17647 ll 14-16), where we cautiously note that there does not necessarily have to be a simple link between changes in upwelling and deep ocean fluxes.

RC

Table 3: Does important mean statistically significant? If so, how significant?

AC

No, it does not mean "statistically significant". We deleted this sentence. Instead, we added the number of data points, the slopes and indicate the statistical significance (r^2 , at 99.9% confidence level).

RC

Table 4: Similarly, what is meant by "important"?

AC

We meant 'important' for our study. We deleted 'important'.

RC

Figure 2: What is implied by "strong changes"? It is not clear what the reader should note happening over the past four decades. Is there something unusual? If so, unusual relative to what?

AC

We removed this sentence in the caption.

RC

Figure 5: Not clear to mean if the gray shaded area is the data from the shallower trap? If so, it appears that there is more data from the upper than lower trap, and so the figure should be about data from the upper and not lower trap.

AC

More data from the lower trap than from the upper one are available and the former is better suited to record coastal upwelling. We changed the respective figures and the captions. Light/white bars denote sampling gaps where upper trap data have been used to fill the gaps of the lower trap record.

RC

Figures 5 and 6: Not clear what the shading of El Nino/La Nina represents, since those colors don't appear elsewhere in the figures nor are they described in the figure captions

AC

This is now better described in the revised figure captions and the shading is shown in the other figures as well. Blue means the cold and red color means the warm phase of the 1997-1999 ENSO cycle.

RC

Technical Comments Comma usage and grammar are problematic throughout the manuscript

AC

A native speaker checked grammar in the revised version.

RC

Page 17653, line 14: incomplete sentence

AC

Deleted

RC

Figure 4: The font size of the y-axis labels is too small to read – you could replace with “Mass flux”, “Corg flux”, “BSi flux”, “CaCO₃ flux”, and “lithogenic flux” and note in the figure caption that all are mean seasonal values.

AC

done

RC

Figure 5 caption: polynome should be polynomial?

AC

done

2 **Deep ocean mass fluxes in the coastal upwelling off Mauritania from 1988 to 2012:**
3 **variability on seasonal to decadal timescales**

4
5
6 *Fischer, G.,^{1,2*} Romero, O.², Merkel, U.^{1,2}, Donner, B.², Iversen, M.^{2,3}, Nowald, N.², Ratmeyer, V.²,*
7 *Ruhland, G.², Klann, M.² and G. Wefer²*

8
9
10 ¹ Geosciences Department, University of Bremen, Klagenfurter Str,
11 28359 Bremen, Germany

12
13 ² Marum Center for Marine and Environmental Sciences, Leobener Str,
14 28359 Bremen, Germany

15
16 ³ Alfred Wegener Institute for Polar and Marine Research, Am Handelshafen,
17 27570 Bremerhaven, Germany

18
19 * *corresponding author*
20
21

22 **Abstract**

23 A more than two-decadal sediment trap record from the Eastern Boundary Upwelling Ecosystem (EBUE) off
24 Cape Blanc, Mauritania, is analysed with respect to deep ocean mass fluxes, flux components and their
25 variability on seasonal to decadal timescales. The total mass flux revealed interannual fluctuations which
26 were superimposed by fluctuations on decadal timescales ~~possibly linked to the Atlantic Multidecadal~~
27 ~~Oscillation (AMO)~~. High winter fluxes of biogenic silica (BSi), used as a measure of marine production
28 (mostly by diatoms) largely correspond to a positive North Atlantic Oscillation (NAO) index ~~during boreal~~
29 ~~winter~~ (Dec.-March). However, this relationship is weak. The highest positive BSi anomaly was in winter
30 2004-2005 when the NAO was in a neutral state. More episodic BSi sedimentation events occurred in several
31 summer seasons between 2001 and 2005, when the previous winter NAO was neutral or even negative. We
32 suggest that distinct dust outbreaks and deposition in the surface ocean in winter ~~and occasionally but also~~ in
33 summer/fall enhanced particle sedimentation and carbon export on ~~rather~~ short timescales via the ballasting
34 effect ~~_ thus leading to these episodic sedimentation events~~. Episodic perturbations of the marine carbon
35 cycle by dust outbreaks (e.g. in 2005) ~~might have~~ weakened the relationships between fluxes and large scale
36 climatic oscillations. As phytoplankton biomass is high throughout the year ~~in our study area~~, any dry (in
37 winter) or wet (in summer) deposition of fine-grained dust particles is assumed to enhance the efficiency of
38 the biological pump by ~~being~~ ~~incorporating~~ ~~ed~~ ~~dust~~ into dense and fast settling organic-rich aggregates. A
39 good correspondence between BSi and dust fluxes was observed for the dusty year 2005, following a period
40 of rather dry conditions in the Sahara/Sahel region. Large changes of all ~~bulk~~ fluxes occurred during the
41 strongest El Niño-Southern Oscillation (ENSO) in 1997-1999 where low fluxes were obtained for almost one
42 year during the warm El Niño and high fluxes in the following cold La Niña phase. ~~For decadal timescales,~~
43 Bakun (1990) suggested an intensification of coastal upwelling due to increased winds ('Bakun upwelling
44 intensification hypothesis'; Cropper et al., 2014) and global ~~climate~~ change. We did not observe an increase
45 of any flux component off Cape Blanc during the past two and a half decades which might support this
46 ~~hypothesis~~. Furthermore, fluxes of mineral dust did not show any positive or negative trends over time which
47 ~~would have~~ ~~might~~ ~~suggested~~ enhanced desertification or 'Saharan greening' during the last few decades.

48

49

50 1. Introduction

51 Eastern Boundary Upwelling Ecosystems (EBUEs; Freon et al., 2009) cover only about 1% of the total
 52 ocean area but contribute with about 15% to total marine primary production (Carr, 2002; Behrenfeld and
 53 Falkowski, 1997). Roughly, 20% of the marine global fish catch is provided by the four major EBUEs (Pauly
 54 and Christensen, 1995), the Benguela, the Canary, the Californian and the Humboldt Current Systems.
 55 Continental margins may be responsible for >more than 40% of the carbon sequestration in the ocean
 56 (Muller-Karger et al., 2005) and are thus, highly relevant for the global carbon cycle. In the literature,
 57 multiple factors with potential influence on upwelling systems have been mentioned. To discuss all of them,
 58 would be beyond the scope of this paper and we therefore focus on three major factors.

59 In the 1990s, a discussion began whether global warming may lead to intensified coastal upwelling in the
 60 EBUEs (e.g. Bakun, 1990: ‘Bakun upwelling intensification hypothesis’; Cropper et al., 2014). Since then,
 61 various studies showed contradicting results, depending on the timescales regarded, the area studied and the
 62 methods applied. The longer-term time series analysis of wind stress and sea surface temperature (SST) by
 63 Narayan et al. (2014) from coastal upwelling areas seems to support the ‘Bakun upwelling intensification
 64 hypothesis’, but correlation analysis showed ambiguous results concerning the relationships of upwelling to
 65 the North Atlantic Oscillation (NAO). With some modification, the ‘Bakun hypothesis’ is supported for the
 66 ~~NW-African Canary Current (CC)~~ coastal upwelling system by Cropper et al. (2014). ~~These authors found~~
 67 ~~indications of a relationship between upwelling and NAO (mainly in winter), but no signs of teleconnections~~
 68 ~~between upwelling and the El Niño Southern Oscillation (ENSO) or the Atlantic Multidecadal Oscillation~~
 69 ~~(AMO) were observed.~~ Using an upwelling index derived from SSTs and remote sensing wind stress,
 70 Marcello et al. (2011) obtained increased offshore spreading of upwelled waters off Cape Blanc from 1987 to
 71 2006. Other authors, however, found a warming trend of the Canary Current (CC)-System (e.g. Aristegui et
 72 al., 2009). Bode et al. (2009) observed a continuous decrease in upwelling intensity in the northern CC
 73 around the Canary Islands during the past 40 years, associated with ~~a-the~~ warming of ~~the~~ surface waters, a
 74 decrease in zooplankton abundance, and, locally, in phytoplankton abundance. Studying a sediment core off
 75 Cape Ghir, Morocco, a cooling of the northern Canary Current in the 20th century was inferred (McGregor et
 76 al. 2007).

77 ~~The An~~ influence of tropical Pacific interannual variability on EBUEs has also been ~~already~~ proposed earlier.
 78 A link between the cold La Niña period (1997-1999 ENSO cycle) and the Mauritanian upwelling via a
 79 strengthening of the north-easterly (NE) trade winds in fall and winter was described by Pradhan et al.
 80 (2006). Helmke et al. (2005) correlated these anomalous events in-with deep-ocean carbon fluxes at the
 81 mesotrophic Cape Blanc study site. Using ocean colour data, Fischer et al. (2009) showed a large extension
 82 of the Cape Blanc filament from fall 1998 to spring 1999 when comparing it to the rest of the record
 83 ~~(spanning from 1997_ to 2008). Persistent La Niña conditions between summer 1998 and summer of 1999~~
 84 ~~caused nutrient driven increases of net marine primary production (NPP) and in the Mauritanian upwelling~~
 85 ~~plume as well (Behrenfeld et al., 2001).~~ Using remote sensing data, Nykjaer and Van Camp (1994) found a

86 weak northwest ~~NW African~~ upwelling south of 20°N during and after the strong 1982-1983 El Niño event,
87 ~~(end of 1982 to early 1984).~~

88 The NW African margin and the low-latitude North Atlantic are heavily influenced by Saharan dust
89 transport, deposition (e.g. Kaufman et al., 2005) and sedimentation (Brust et al., 2001). Dust particles
90 influence the earth's radiation balance, ~~and~~ supply micro-nutrients ~~(e.g. such as iron)~~ and macro-nutrients to
91 the ocean surface waters (e.g. Jickells et al., 2005; Neuer et al., 2004). Additionally, dust acts as ballast
92 mineral ~~both for total flux~~ (Armstrong et al., 2002; Klaas and Archer, 2002) ~~and~~ for organic carbon-rich
93 particles (e.g. Fischer et al., 2009, a, b; Bory and Newton, 2000; Iversen and Ploug, 2010; Iversen et al.,
94 2010; Bressac et al., 2014). Dunne et al. (2007) suggested that dust may be the major carrier for organic
95 carbon to the seafloor. A clear coupling between atmospheric dust occurrence and deep-sea lithogenic
96 particle fluxes at 2000 m water depths was observed in the subtropical north Atlantic (33°N, 22°W; Brust et
97 al., 2011). Fischer and Karakas (2009) proposed that high dust supply may increase particle settling rates by
98 ballasting and result in relatively high organic carbon fluxes in the Canary Current NW African upwelling
99 system compared to other EBUEs. Wintertime African dust transport is suggested to be affected by the NAO
100 (Chiapello et al. 2005; Hsu et al., 2012). As dust plays a major role in the Cape Blanc area with respect to
101 deep ocean fluxes and the intensity of coastal upwelling is affected by the NAO as well, the major focus of
102 this long-term study will be on the relationship between deep ocean mass fluxes and NAO forcing.

103 From the mesotrophic Cape Blanc study site CB_{meso} located about 200 nm off the coast (Fig. 1a), we
104 obtained an almost continuous sediment trap record of ~~export~~ fluxes (mostly from about 3500 m water depth)
105 for the past 25 years (1988-2012, ~~only~~ interrupted between 1992 and 1993). Long time series of particle
106 fluxes are rare, in particular from coastal upwelling sites with high productivity. Although SSTs and wind
107 data analysis over longer time scales (e.g. decades) for the NW African upwelling system and other
108 EBUEs are very important to test the 'Bakun upwelling intensification hypothesis' (Bakun, 1990; Cropper et
109 al., 2014), any potential increase of upwelling intensity does not necessarily result in an increase of
110 phytoplankton standing stock and/or productivity and/or deep ocean mass fluxes (e.g. Ducklow et al., 2009).
111 Hence, for studying the potential changes of the biological pump and carbon sequestration in the deep ocean
112 over decades and over a larger area, sediment traps are a primary and probably the best choice. As deep
113 ocean sediment traps have a rather large catchment area for particles formed in the surface and subsurface
114 waters (e.g. Siegel and Deuser, 1997), they integrate rather local and small-scale effects, events and
115 processes in the highly dynamic EBUE off Mauritania.

116

117 **2. Study area**

118 *2.1 Oceanographic and biological setting*

119 The sediment trap mooring array CB_{meso} is deployed in the Canary Current ~~(CC)~~ System within one of the
120 four major EBUEs (Freon et al., 2009) (Fig. 1a). Coastal upwelling is driven there by alongshore trade
121 winds, leading to offshore advection of surface waters, which are replaced by colder and nutrient-rich

122 | subsurface waters. Around 21°N off Cape Blanc (~~Fig. 1a~~), a prominent cold filament leads to offshore
 123 | streaming of cold and nutrient-rich waters from the coast to the open ocean up to about 450 km offshore
 124 | (~~Fig. 1a~~). This cold tongue is named the ‘giant Cape Blanc filament’ (Van Camp et al., 1991), being one of
 125 | the largest filaments within all EBUEs.

126 | The relationship between the coastal winds, SST and the biological response (~~e.g.~~ changes in chlorophyll) off
 127 | Mauritania seems to be strong and almost immediate (Mittelstaedt, 1991; Pradhan et al., 2006). Trade winds
 128 | persist throughout the year and intensify in late winter to reach their highest intensity in spring (Barton et al.,
 129 | 1998; Nykjaer and Van Camp, 1994; Meunier et al., 2012). According to Lathuilière et al. (2008), our study
 130 | area is located within the Cape Blanc inter-gyre region (19-24°N) which is characterized by a weaker
 131 | seasonality (peaks in winter-spring and fall). Following the definition by Cropper et al. (2014), our study
 132 | area is situated on the southern rim of the strong and permanent coastal upwelling zone (21°-26°N) (Fig. 1a).

133 | The cold and nutrient-rich southward flowing CC departs from the coastline south of Cape Blanc, later
 134 | forming the North Equatorial Current (NEC) (Fig. 1a). South of about 20°N, a recirculation gyre drives a
 135 | poleward coastal current fed by the North Equatorial Counter Current (NECC) during summer. The
 136 | Mauritanian Current (MC) flows northward along the coast to about 20°N (Fig. 1a; Mittelstaedt, 1991),
 137 | bringing warmer surface water masses from the equatorial realm into the study area. Where the CC departs
 138 | from the coast, a NE-SW orientated salinity front in the subsurface waters is observed, the Cape Verde
 139 | Frontal Zone (CVFZ, Zenk et al., 1991) (Fig. 1a), which separates the salty and nutrient-poor North Atlantic
 140 | Central Water (NACW) from the nutrient-richer and cooler South Atlantic Central Water (SACW). Both
 141 | water masses may be upwelled and mixed laterally and frontal eddies develop off Cape Blanc (Meunier et
 142 | al., 2012) (Fig. 1a). Lathuilière et al. (2008) offered a comprehensive overview of the physical background,
 143 | i.e. the ocean circulation off NW Africa.

144 | Fig. 1.

145 |

146 | ***2.2 Importance of dust supply and Sahel rain fall for the study area***

147 | Dust supply from land to the low-latitude North Atlantic Ocean is not only dependent on the strength of the
 148 | transporting wind systems (NE trade winds at lower levels and Saharan Air Layer above) but also on the
 149 | rainfall and dryness in the multiple source regions in West Africa (~~see Nicholson, 2013;~~ Goudie and
 150 | Middleton, 2001; Nicholson, 2013). During long periods of droughts (e.g. in the 1980s), dust loadings over
 151 | the Sahel experienced extraordinary increases (N’Tchayi Mbourou et al., 1997). As mass fluxes and settling
 152 | rates of larger marine particles (i.e. marine snow) are assumed to be influenced by mineral dust particles via
 153 | the ballasting effect (Armstrong et al., 2002; Fischer et al., 2007a, 2010; Iversen and Ploug, 2010; Bressac
 154 | et al., 2014; Dunne et al., 2007; Thunell et al., 2007), climatic conditions on land need to be considered. The
 155 | contribution of dust to the settling particles in the deep ocean off Cape Blanc amounts to one-third on
 156 | average of the total mass flux (Fischer et al., 2010), but it may be as high as 50% during particular flux
 157 | events (Nowald et al., 2015). As shown by Jickells et al. (2005), modelled dust fluxes from the Saharan

158 region and their variability may be influenced by ENSO and NAO cycles (see also Goudie and Middleton,
 159 2001; Chiapello et al., 2005; Hsu et al., 2012; Diatta and Fink, 2014). During the time period of this study
 160 (1988-2012, Fig. 2), the wintertime (Dec-Jan-Feb-Mar = DJFM) NAO index after Hurrell (Hurrell, 1995) is
 161 characterized by switches from extremely positive (e.g. 1989, 1990) to extremely negative values (e.g. in
 162 1996, 2010) (Fig. 2).

163 Climate over West Africa is also influenced by the continental Inter-Tropical Convergence Zone (ITCZ; also
 164 named Intertropical Front, Nicholson, 2013). This low-pressure zone separates the warm and moist SW
 165 monsoon flow from the dry NE trade winds coming from the Sahara. The tropical rainbelt in the Atlantic
 166 realm originates from the convergence of the NE and SE trade wind systems and ~~latitudinally~~ migrates
 167 roughly between $\sim 3^{\circ}\text{S}$ (boreal winter) and $\sim 15^{\circ}\text{N}$ (boreal summer) in the course of the year (Lucio et al.,
 168 2012). On longer timescales, severe Sahel drought intervals occurred in the 1980s (Chiapello et al., 2005;
 169 Nicholson, 2013). ~~According to Shanahan et al. (2009), changing Atlantic SSTs (AMO) exert a control on~~
 170 ~~the persistent droughts in West Africa in the 1980s and 1990s.~~ Recent evidence shows that Sahel rainfall
 171 may have recovered during the last two decades and that the region is now ‘greening’ (Fontaine et al., 2011;
 172 Lucio et al., 2012).

173 Fig. 2.

174

175 *2.3 Large-scale teleconnections affecting the study area*

176 Ocean-atmosphere dynamics at our study site is influenced by large-scale atmospheric teleconnections and
 177 climate modes. Here, such teleconnections are illustrated based on results from a long-term present-day
 178 climate control run which was performed using the Comprehensive Climate System Model version 3
 179 (CCSM3; Collins et al., 2006; Yeager et al., 2006). Atmospheric sea-level pressure (SLP) patterns describe
 180 the near-surface air flow which affects ocean upwelling and currents as well. We therefore correlated
 181 simulated SLP with prominent teleconnection indices such as the NAO SLP index (Hurrell, 1995) and the
 182 Niño3 area-averaged (150°W - 90°W , 5°S - 5°N) SST index, both calculated from the model results (Fig. 3).
 183 Boreal winter is the season where the NAO is strongest and where tropical Pacific SST anomalies associated
 184 with ENSO events tend to peak.

185 Correlations during winter show that NAO and ENSO may have opposite effects on the NW African/eastern
 186 Atlantic realm (Fig. 3 a,b), for instance on wind fields, and consequently on upwelling with potential
 187 implications for deep ocean mass fluxes. A positive phase of the NAO is associated with anomalous high
 188 pressure in the Azores high region (Fig. 3a) and stronger northeasterly winds along the NW African coast. In
 189 contrast, a positive phase of ENSO (El Niño event) goes along with a weakening of the northeasterlies in the
 190 study area (Fig. 3 b). It should be noted, however, that the magnitude of correlation in our study area is larger
 191 for the NAO than for ENSO. This should be taken into account when disentangling the relative importance
 192 of these climate modes. Apart from seasonal-to-interannual timescales, low-frequent climate variability may
 193 impact on our study area as well and is probably linked to Atlantic ~~SST~~ sea surface temperature variations on

194 | decadal-to-interdecadal timescales, e.g. the [Atlantic Multidecadal Oscillation \(AMO\)](#). The correlation of SLP
 195 | with area-averaged (0°-70°N, 60°-10°W) SST fluctuations over periods above 10 years highlights a centre of
 196 | action in the tropical Atlantic with SLP reductions (weaker northeasterly winds) along with higher Atlantic
 197 | basin-wide SST during a positive AMO phase (Fig. 3c). This shows the potential importance of longer-term
 198 | Atlantic basin-scale SST variations for alongshore winds and upwelling (trends) at our trap location

199 | ENSO related teleconnections in the NW African upwelling system have been described by several authors
 200 | (Behrenfeld et al., 2001; Pradhan et al, 2006; Zeeberg et al., 2008) and can be illustrated by the negative
 201 | correlation of SLP with eastern tropical Pacific SST (Fig. 3b, chapter 3.3.). Fischer et al. (2009b) showed
 202 | that the size of the Cape Blanc filament was small in winter-spring 1997-1998 and unusually high from fall
 203 | 1998 to spring 1999 (Fig. 7b; Fig. 1e). This is documented by reduced (warm El Niño) and elevated (cold La
 204 | Niña) deep ocean mass fluxes of all components. In certain years, the filament [area](#) was more than twice [as](#)
 205 | [large](#) in spring [as than](#) in fall (e.g. 1999 La Niña Event). [Tropical Pacific variability on interannual ENSO](#)
 206 | [timescales is also an important factor in driving ecosystem variability in the California Current System \(for a](#)
 207 | [summary see Checkley and Barth, 2009\).](#)

208 | Fig. 3.

209

210 | **3. Material and Methods**

211 | **3.1 Sediment traps and moorings**

212 | We used deep-moored (>1000 m), large-aperture time-series sediment traps of the Kiel and Honjo type with
 213 | 20 cups and 0.5 m² openings, equipped with a honeycomb baffle (Kremling et al., 1996). Mooring and
 214 | sampling dates are given in Table 1. As the traps were moored in deep waters (mostly [→ below](#) 1000m),
 215 | uncertainties with the trapping efficiency due to strong currents (e.g. undersampling, Yu et al., 2001;
 216 | Buesseler et al., 2007) and/or due to the migration and activity of zooplankton migrators ('swimmer
 217 | problem') are assumed to be minimal. Prior to the deployments, the sampling cups were poisoned with
 218 | HgCl₂ (1 ml of conc. HgCl₂ per 100ml of filtered seawater) and pure NaCl was used to increase the density
 219 | in the sampling cups to 40‰. Upon recovery, samples were stored at 4°C and wet-split in the home
 220 | laboratory using a rotating McLane wet splitter system. [Larger](#) swimmers [such as crustaceans](#) were picked
 221 | by hand with forceps and were removed by filtering carefully through a 1 mm sieve and all flux data here
 222 | refer to the size fraction of <1 mm. In almost all samples, the fraction of particles >1 mm was negligible,
 223 | [only in a few samples larger pteropods were found.](#)

224 | **3.2 Mass fluxes**

225 | Analysis of the fraction < 1 mm, using ¼ or 1/5 wet splits, was performed according to Fischer and Wefer
 226 | (1991). Samples were freeze-dried and the homogenized samples were analyzed for bulk (total mass),
 227 | organic carbon, total nitrogen, carbonate and biogenic opal (BSi = [biogenic silica](#)). Organic carbon, [nitrogen](#)
 228 | and calcium carbonate were measured by combustion with a CHN-Analyser (HERAEUS). Organic carbon
 229 | was measured after removal of carbonate with 2 N HCl. Overall analytical precision based on internal lab

standards was better than 0.1% ($\pm 1\sigma$). Carbonate was determined by subtracting organic carbon from total carbon, the latter being measured by combustion without pre-treatment with 2N HCl. BSi was determined with a sequential leaching technique with 1M NaOH at 85°C (Müller and Schneider, 1993). The precision of the overall method based on replicate analyses is mostly between ± 0.2 and $\pm 0.4\%$, depending on the material analyzed. For a detailed table of standard deviations for various samples we refer to Müller and Schneider (1993). Lithogenic fluxes or the non-biogenic material was estimated according to:

$lithogenic\ material = dust = total\ mass - carbonate - opal - 2xC_{org}$ (~~= organic matter~~).

We estimated organic matter by multiplying organic carbon by a factor of two as about 50-60% of marine organic matter is constituted by organic carbon (Hedges et al., 1992). Some studies have shown a clear linear relationship between lithogenic fluxes and particulate aluminum (e.g. Ratmeyer et al., 1999a), the latter being derived from clay minerals as part of the lithogenic (non-biogenic) component. Grains size studies from Ratmeyer et al. (1999a, b) and further microscopic analysis provide evidence that most of the lithogenic material in the study area was derived from quartz grains in the fine silt fraction (10-30 μm ; see also Friese et al., 2016). ~~We h~~Here we attribute ~~all~~ the lithogenic flux to dust-derived material (=mineral dust flux) as no large rivers ~~suppl~~ies suspended material to the study area off Cape Blanc.

Due to logistical reasons, we had very different time resolutions of the sediment trap collections (a few days to several weeks) which limits comparisons between specific intervals and years. Seasonal fluxes were calculated and shown to allow comparison between the seasons mainly with respect to interannual variability. Seasons were defined ~~using~~ the dates of opening and closure of the sampling cups closest to the start of the astronomical seasons (March 21, June 21, September 23, December 21) (Table 2). Where lower trap data (around 3500 m) were not available, the upper trap data (around 1000 m) were used, which mostly match the lower trap fluxes with respect to seasonality (Fischer et al., 2009b). ~~As the deeper traps have a higher collection area due to the 'statistical funnel' (Siegel and Deuser, 1997), they might have collected slightly more material, in particular in the winter season. When plotting all available lower and upper trap total mass fluxes for winter, a close correspondence is observed ($r^2=0.84$, $N=10$), with slightly higher fluxes in the deeper trap due to lateral particle advection processes (Fischer et al., 2009b, Karakas et al., 2006, 2009). However, considering the entire record presented here, it seems that the upper trap fluxes of the winter seasons 1998 and 2004 may be critical due to smaller filament areas. As a consequence, the area/filament with elevated chlorophyll and high particle concentrations may not have reached the upper offshore trap. Because of lateral particle advection from the east (Karakas et al., 2006) and the larger catchment area of the deeper traps (Siegel and Deuser, 1997), particle fluxes might have been higher in the deeper water column in winter 1998 and 2004. However,~~ ~~in general,~~ the seasonal patterns and the composition of the particle fluxes were rather similar between the upper and lower traps (Fischer et al., 2009b). The long-term means and standard deviations were calculated using only the available deeper trap flux values. The seasonal anomalies of the bulk fluxes were calculated using the deviations from the mean values of the respective seasons.

Kommentar [g1]: Chapter on major comment of reviewer #1

268 **3.3 Carbonate producers**

269 To determine the major carbonate producers, the trap material was carefully wet-sieved with a 1 mm-screen
 270 and split into aliquots by a rotary liquid splitter. Generally a 1/5 split of the < 1 mm-fraction was used to pick
 271 planktonic foraminifers and pteropods from the wet solution. Foraminifers and pteropods were picked by
 272 hand with a pipette under a ZEISS Stemi 2000 microscope and rinsed with fresh water for three times and
 273 dried at 50°C overnight and counted. The mass fluxes of total carbonate producers expressed as mg m⁻² day⁻¹
 274 are mainly constituted of planktonic foraminifera, pteropods and nannofossils/coccolithophorids. Masses of
 275 foraminifera and pteropods were determined with a Sartorius BP 211D analytical balance.

276 **3.4 Additional web-based data:**

277 To put our flux results from the deep ocean into a broader context ~~to the surface water properties~~, we used
 278 several observational datasets available from several [the](#) websites given below. For ocean colour, time series
 279 from the MODIS or SeaWiFS sensors based on a 1°x1° box from 20°N-21°N and 201°-240°W (9 km
 280 resolution) slightly to the east of the study site CB have been chosen due to the generally prevailing E-W
 281 directed current system, transporting particles to the west (Helmke et al., 2005). Larger boxes, e.g. 2°x2° or
 282 4°x4°, revealed similar results. For the aerosol optical thickness (AOT, 869 nm, 9 km resolution), a 1°x1°
 283 box was chosen from the SeaWiFS and MODIS data.

284 *Ocean colour from MODIS (9 km resolution):*

285 http://oceancolor.gsfc.nasa.gov/cgi/l3?ctg=Standard&sen=A&prd=CHL_chlor_a&per=SN&date=21Jun2002&res=9km&num=24

287 *Ocean colour from SeaWiFS (9 km resolution):*

288 http://oceancolor.gsfc.nasa.gov/cgi/l3/S19972641997354.L3m_SNAU_CHL_chlor_a_9km.png?sub=img

289 *GIOVANNI-derived time series AOT (Aerosol Optical Thickness) and chlorophyll from SeaWiFS and MODIS:*

291 http://gdata1.sci.gsfc.nasa.gov/daac-bin/G3/gui.cgi?instance_id=ocean_month

292 *AOD (Aerosol Optical Depths) and dust and rainfall pattern (animation):*

293 http://earthobservatory.nasa.gov/GlobalMaps/view.php?d1=MODAL2_M_AER_OD&d2=TRMM_3B43M

294 *NAO (North Atlantic Oscillation) index based on station data of sea level pressure:*

295 <http://climatedataguide.ucar.edu/guidance/hurrell-north-atlantic-oscillation-nao-index-station-based>

296 *ENSO (El Niño-Southern Oscillation) Niño3.4 SST index:*

297 http://iridl.ldeo.columbia.edu/filters/.NINO/SOURCES/.NOAA/.NCEP/.EMC/.CMB/.GLOBAL/.Reyn_SmithOlv2/.monthly/.ssta/NINO34/T

299 *AMO (Atlantic Multidecadal Oscillation) SST index:*

300 <http://www.esrl.noaa.gov/psd/data/correlation/amon.us.data>

301 Table 1.

302

303

304 4. Results

305 On the long-term, seasonal bulk fluxes were highest in boreal winter and summer and slightly lower in
 306 spring and fall (Figs. 4, 5, 6a; Table 2). Total bulk fluxes reached 23.6 and 23.1 g m⁻² in winter and summer,
 307 respectively (Table 2). For spring and fall, total mass fluxes were as high as 19.6 and 21.1 g m⁻², respectively
 308 (Table 2). However, the seasonal differences in the bulk fluxes are not statistically significant. Along with
 309 the highest mass fluxes, winter and summer seasons also exhibit the highest standard deviations (Fig. 4),
 310 pointing to a high interannual variability ~~during these seasons.~~ In general, this interannual variability is
 311 clearly higher than the seasonal differences in bulk fluxes. Only the lithogenic components, i.e. the mineral
 312 dust particles, did not show an increase during summer and only peaked in winter (up to 7.4 g m⁻²) when dust
 313 plumes were most frequent (Goudie and Middleton, 2001). High summer fluxes of up to 16.9 g m⁻² were
 314 mostly due to high carbonate sedimentation (Fig. 4), both of primary (coccolithophoresids) and secondary
 315 producers (foraminifera and pteropods). Organic carbon and BSi showed a rather similar pattern (Fig. 4, 6a)
 316 with a maximum in winter (up to 1.1 and 1.7 g m⁻², respectively) and a secondary maximum in summer/fall.
 317 This is reflected in the close correspondence between both flux components for these seasons (Table 3).
 318 Highest mass fluxes coinciding with highest positive flux anomalies lasting for several seasons occurred in
 319 1988-89, 1998-99, and 2005-2006 (Fig. 5).

320 Following the strong ENSO cycle 1997-1999, total flux anomalies were low or negative over a longer period
 321 (fall 1999 to fall 2004), only interrupted by an episodic peak in summer 2002 (Fig. 5a, b). ~~More~~ Other
 322 episodic peaks in sedimentation were found in winter/spring 1996-1997 and in the winter seasons 2004-
 323 2005, 2006-2007 and 2009-2010 (Fig. 5a, b). Longer intervals (several seasons) of negative flux anomalies
 324 were obtained in 1997-98 and 2009-11 (Fig. 5b). Total fluxes decreased from 1988 to 1991, from spring
 325 2007 to 2010, later increasing from 2010 to 2012 (Fig. 5).

326 In general, the major bulk flux components followed the total flux and were well inter-correlated, except that
 327 the relationship between organic carbon and carbonate was weak in summer (Table 3). However, the
 328 regression-based relationships (i.e. the slope) varied interannually (e.g. Fischer et al., 20079a). The matrix in
 329 Table 3 shows the correlation coefficients between organic carbon and nitrogen, BSi, carbonate, and
 330 lithogenic fluxes for the four seasons (lower traps only). Organic carbon and BSi (mainly diatoms) were
 331 highly correlated during the major upwelling events in winter ($R^2=0.70$, $N=16$) and summer/fall, whereas the
 332 relationship between organic carbon and total carbonate in summer was weak ($R^2=0.16$, $N=19$; Table 3).
 333 Dust fluxes peaked together with organic carbon, preferentially in winter and fall ($R^2=0.63$ ($N=16$) and 0.67
 334 ($N=18$), respectively Table 3). The tight coupling between organic carbon and nitrogen is not surprising as
 335 both elements constitute organic matter formed during photosynthesis, which is later degraded in the upper
 336 water column forming sinking phytodetritus. The slope is almost constant (0.13-0.11) and the reciprocal
 337 value reflects the Redfield Ratio (Redfield et al., 1963) of the sinking organic-rich particles (Table 3). The
 338 molar C:N varied seasonally between 8.9 and 10.6, typical for sinking detritus collected in deep sediment
 339 traps.

340 On the long-term, the composition of settling particles in the deeper traps off Cape Blanc consisted of
 341 roughly 56% carbonate, ca. 30% lithogenic particles, ~~and about~~ 4% ~~of~~ organic carbon, 0.5% nitrogen and
 342 5% BSi (Fig. 4). BSi contained mostly a mixture of coastal and open-ocean diatoms (Romero et al. 1999,
 343 2002, and unpubl. data). The BSi flux pattern (Fig. 6) was influenced by switches from a positive to a
 344 negative NAO index which were reflected in decreasing winter opal fluxes, e.g. from 1989 to 1991, 1995-
 345 1996 and 2007 to 2010. From 2001 through 2006, NAO variability was rather low and the index was around
 346 zero or slightly negative (Fig. 6c; Table 4). Nevertheless, BSi fluxes varied considerably and showed
 347 episodic peaks in the summer seasons 2001, 2002 and 2003 (Fig. 6a, b). ~~Except in spring 2005~~, BSi flux was
 348 high and showed (positive anomalies) ~~in the entire year 2005~~, except for spring 2005 (Fig. 6a, b; Table 4).

349 The general flux pattern of BSi (Fig. 6a, b) with values from almost zero to 1.91 g m⁻² did not match the
 350 SeaWiFS ocean colour time series trend which showed an overall decrease in chlorophyll from 1997 to 2010
 351 (Fig. 6d). The organic carbon flux pattern (not shown, values from almost zero to 1.1 g m⁻²) did not follow
 352 the ocean colour data from MODIS/SeaWiFS either as well. Peak chlorophyll values were observed mostly
 353 during spring, except in 1998 (fall maximum) and 2007 (summer maximum). The MODIS ocean colour
 354 values generally mimicked the SeaWiFS pattern, except for the discrepancy in summer 2010 (Fig. 6d).

355 Tables 2-3; Figs. 4-6.

356

357 5. Discussion

358 5.1 Particle transport processes in the water column

359 Mass fluxes and particle transport processes off Cape Blanc (Mauritania) have been described by
 360 summarizing articles of Fischer et al. (2009b) and Karakas et al. (2006). Common flux patterns were the
 361 increase of fluxes in late winter-spring and late summer of all components at both trap levels. This matched
 362 the seasonal intensification of coastal upwelling (e.g. Meunier et al., 2012) due to wind forcing and a
 363 stronger offshore streaming of the Cape Blanc filament (e.g. Fischer et al., 2009). The increase of fluxes in
 364 late summer to fall was mostly due to enhanced biogenic carbonate sedimentation (Fig. 4d), associated with
 365 the northward flowing warm MC coming from tropical regions (Mittelstaedt, 1991). In the Canary
 366 Current/NW African upwelling system, which is dominated by carbonate producers ~~(Fischer et al., 2007)~~,
 367 particle settling rates are rather high (around 300 m d⁻¹), compared to EBUEs dominated by BSi
 368 sedimentation (Fischer and Karakas, 2009; Fischer et al., 2009a). As suggested by Fischer et al. (2009a), the
 369 relatively high organic carbon flux in the deep ocean off NW Africa may be due to the high availability of
 370 mineral ballast, i.e. from coccolithophorids and fine-grained mineral dust (Iversen et al., 2010; Iversen and
 371 Ploug, 2010; Ploug et al., 2008). Direct evidence for the influence of the deposition of dust particles on the
 372 settling rates of larger particles and the flux attenuation in the epi- and mesopelagic has been found on short
 373 time-scales, i.e. days. This was observed during a severe dust outbreak in January 2012 (Iversen, unpubl.
 374 observations) by deploying drifting traps before and after the dust outbreak (Fig. 8a, insert image).

375

Kommentar [g2]: Flux ranges and values are now given (reviewer #2)

376 5.2 Influence of the NAO on biogenic silica sedimentation

377 The NAO both affects coastal upwelling and productivity off Mauritania through wind forcing (upwelling)
 378 and dust/nutrient supply (Chiapello et al., 2005), mainly during winter (DJFM) (Goudie and Middleton,
 379 2001; Cropper et al., 2014). Indeed, we observed an increase of both the winter NAO index and associated
 380 winter BSi fluxes (Fig. 6, 7a, ~~$R^2=0.18$, $N=20$~~), the latter known to be indicative of coastal upwelling
 381 strengths and productivity. When plotting winter BSi fluxes versus the Azores pressure alone (DJFM Ponta
 382 Delgada SLP, 1989-2002), the relationship improves slightly ($R^2=0.19$ $N=11$, not shown) but remains
 383 statistically insignificant. Since upwelling is wind-driven and large-scale wind patterns in the study area are
 384 positively correlated to NAO variability (Fig. 3a), a close linkage between a positive (negative) NAO and
 385 higher (lower) BSi fluxes can be expected. Organic carbon flux showed less correspondence to the winter
 386 NAO index (not shown). No clear relationship can be seen between the winter (DJFM) NAO index and
 387 ~~spring~~ BSi and organic carbon fluxes later in spring, if we consider a time delay of a few weeks between
 388 wind forcing of coastal upwelling, high chlorophyll standing stock, particle formation and in-spring and
 389 sedimentation and, finally, the documentation of increasing fluxes in the deep traps in spring.

Kommentar [g3]: Reviewer #1

390 From 2001 to 2006 when the winter NAO index became close to zero (Fig. 6c), the BSi flux showed rather
 391 unusual (episodic) peaks either in summer, fall ~~and or~~ in winter 2004-2005 (Fig. 6a, b). This suggests
 392 increasing coastal upwelling in summer and fall (e.g. Cropper et al., 2014) and/or a strengthening of the
 393 northward flowing and warmer MC, combined with an enhanced supply of a nutrient- and Si-rich source
 394 water (SACW instead of NACW). We favour the latter scenario as there is evidence of unusual warm surface
 395 water conditions (SST anomalies of +3°C) related to weak trade wind intensity between 2002 and 2004
 396 (Zeeberg et al., 2008, Alheit et al., 2014). These conditions might have led to a stronger influence of the
 397 northward flowing MC and the silicate-rich SACW which mixes into the Cape Blanc upwelling filament
 398 and, thus, contributed to higher BSi productivity and sedimentation. Such a scenario was proposed by
 399 Romero et al. (2008) to explain the extraordinary high content of BSi in Late Quaternary sediments
 400 deposited off Cape Blanc during Heinrich Event 1 and Younger Dryas following the Last Glacial Maximum.

401 The 2004-2005 winter BSi flux clearly falls off the regression line of winter BSi flux versus the winter NAO
 402 index (Fig. 7a). Exceptional conditions in 2005 are also indicated when plotting the area with high
 403 chlorophyll (> 1 mg Chl_a m⁻³) covered by the Cape Blanc filament (Fischer et al., 2009a) versus the BSi
 404 fluxes (Fig. 7b). In general, A a larger (smaller) Cape Blanc filament area has been associated with higher
 405 (lower) BSi fluxes (Fig. 7b) and also with higher total mass fluxes (not shown) and BSi fluxes. However, in
 406 winter of 2004-2005 (a relatively cold season with negative SST anomalies), the filament area was smaller
 407 and chlorophyll standing stock was lower (Fig. 6d, 7b). Nevertheless, BSi fluxes were the highest of the
 408 entire record. ~~It appears that the area of the filament with high chlorophyll biomass is not related linearly to~~
 409 ~~deep ocean mass fluxes~~. The seasonal variability of chlorophyll from the entire SeaWiFS record (1997-2010,
 410 Fig. 6d) ~~also~~ indicates no relationship between the chlorophyll standing stock and deep ocean BSi flux (or
 411 organic carbon flux, not shown). These observations point to additional regulators for organic carbon and
 412 BSi export to the deep sea. Ocean colour imagery even revealed a decreasing trend from 1997 to 2010 (Fig.

413 | 6d), which suggests a decrease in upwelling. This is ~~not in concert~~ consistent with the ‘Bakun upwelling
 414 | intensification hypothesis’ (Bakun, 1990) nor with studies from Kahru and Mitchell (2008). Throughout
 415 | 2005, however, the positive BSi flux anomalies corresponded well with positive dust flux anomalies (Fig. 6b,
 416 | 8b). As seen from Aerosol Optical Thickness (AOT, Fig. 8c), dust availability was rather high in 2005 and
 417 | corresponded to high dust sedimentation in summer and fall 2005 (Fig. 8; see chapter 6.5.3). We suggest,
 418 | therefore, that the linear relationship between the NAO index and BSi fluxes may be biased in years of
 419 | anomalous dust input into the surface ocean.

420 | Fig. 7.

421 |

422 | **5.3 Interaction between mineral dust and the biological pump**

423 | Fischer and Karakas (2009) stated that particle settling rates ~~off NW Africa~~ and organic carbon fluxes in the
 424 | Canary Current system were unusually high compared to other EBUEs. This was mainly attributed to
 425 | particle loading by dust particles ~~off NW Africa~~ (see also Fischer et al., 20079a, b, Iversen et al., 2010). ~~In~~
 426 | ~~January 2012 (RV Poseidon cruise 425) when chlorophyll increased in the study area, a two-day Saharan~~
 427 | ~~dust storm (Fig. 8a, insert image) led to increased carbon fluxes at 100 and 400 m water depth as measured~~
 428 | ~~by free drifting traps prior and after the storm event (Iversen, unpubl.).~~ BSi and lithogenic (mineral dust)
 429 | fluxes point to a close linear relationship ($R^2=0.78$ ~~(linear)~~ ~~or 0.9 (power law)~~; $N=1921$, Fig. 9a) mainly in
 430 | winter where dust availability and deposition is high (Goudie and Middleton, 2001), but not in summer (~~Fig.~~
 431 | ~~9b~~; $R^2=0.3156$, $N=23$, Fig. 9b). High supply of dust into the surface ocean is often associated with dry
 432 | conditions in the Sahel/Sahara in the previous year (Engelstaedter et al., 2006; Prospero and Lamb, 2003;
 433 | Moulin and Chiapello, 2004). Indeed, the interval 2002-2004 in particular is known to have been much
 434 | warmer and drier during summer/autumn on land and in the ocean (Zeeberg et al., 2008; Alheit et al., 2014).
 435 | These conditions might have allowed the later wind-induced mobilization of larger amounts of dust particles
 436 | by upward moving winds into the atmosphere and lead to a dust-enriched atmosphere during the entire year
 437 | 2005, combined with elevated deep ocean mass fluxes.

438 | Typically, highest dust flux off Cape Blanc occurs in winter, whereas part of the summer dust load (Fig. 8) is
 439 | transported further westward and deposited in the Caribbean Sea (Goudie and Middleton, 2001; Prospero and
 440 | Lamb, 2003). However, the rainfall pattern exhibits elevated precipitation in summer and fall 2005 when the
 441 | tropical rainbelt was far north; this might have led to unusual wet deposition of dust in summer over our
 442 | study site (Friese et al., ~~ms~~ 2016). As shown earlier, BSi fluxes show positive anomalies in summer and fall
 443 | 2005 (Fig. 6b), pointing to a stronger dust-influenced biological pump.

444 | In contrast to BSi, winter sedimentation of mineral dust did not show any common trend with the winter
 445 | NAO index (not shown). Using satellite-derived AOT, Chiapello et al. (2005) suggested a close relationship
 446 | of atmospheric dust content and the NAO index. High AOT, however, does not necessarily correspond with
 447 | high dust deposition into the ocean. Moreover, dust deposition into the ocean surface does not unavoidably
 448 | and directly result in particle export and transfer to the deep ocean. Dust deposition is not only controlled by

449 wind strength and direction in the trap area but also by source region conditions and precipitation over the
 450 trap site. Consequently, considering the NAO as the only controlling factor for dust deposition and
 451 sedimentation even if the correlation between SLP (and thus winds) and NAO is strong in the study area
 452 (Fig. 3a), would be an oversimplification.

453 Another explanation for the missing relationship could be that fine-grained dust accumulates in surface
 454 waters until the biological pump produces sufficient organic particles to allow the formation of larger
 455 particles which then settle into the deep ocean (Bory et al., 2002; Ternon et al., 2010, Nowald et al., 2015).
 456 Cape Blanc dust particles have predominant grain sizes between 10 and 20 μm (Ratmeyer et al., 1999a, b;
 457 [Friese et al., 2016](#)) and, thus, would sink too slowly to build a deep ocean flux signal. We propose that only
 458 the close coupling between the organic carbon pump, dust particles and the formation of dense and larger
 459 particles led to elevated export and sedimentation (Bory et al., 2002; Fischer et al., 2009a, Fischer and
 460 Karakas, 2009); ~~Iversen, unpubl.~~ [Thunell et al. \(2007\)](#) found that organic carbon fluxes strongly correlated
 461 [with mineral fluxes in other upwelling-dominated continental margin time series such as the Santa Barbara](#)
 462 [Basin located within the California Current System](#). However, the detailed processes [and interaction](#) between
 463 different [groups types](#) of phytoplankton and types of ballast minerals (e.g. quartz versus clay minerals etc.)
 464 are largely unknown and need clarification. Laboratory experiments with different ballast minerals (e.g.
 465 Iversen and Roberts, 2015) and measurements of organic carbon respiration and particle settling rates
 466 suggest a significant influence of ballast minerals on particle settling rates, carbon respiration and flux
 467 (Ploug et al., 2008; Iversen and Ploug, 2010). In a time series study with optical measurements, addressing
 468 particle characteristics (e.g. sizes) and using fluxes at the nearby eutrophic sediment trap off Cape Blanc
 469 (CB_{eu}), Nowald et al. (2015) suggested an influence of dust outbreaks on particle sedimentation down to
 470 1200 m. Interestingly, settling organic-rich particles off Cape Blanc were only around 1 mm in size (~~marine~~
 471 ~~snow is >0.5 mm by definition~~) during the two-year deployment from 2008 to 2010 (Nowald et al., 2015).
 472 Higher fluxes were mostly attributed to higher numbers of small particles rather than to larger particle sizes
 473 during blooms in the Cape Blanc area (Nowald et al., 2015). ~~This demonstrates the need for seasonal studies~~
 474 ~~on particle characteristics and particle settling rates, together with process studies to get further insights into~~
 475 ~~the links between mineral dust input into the ocean and the biological pump.~~

476 Figs. 8, 9.

477

478 **5.4 Carbonate fluxes and potential ENSO teleconnections**

479 Deep ocean total mass and carbonate fluxes (Figs. 5 and 10) showed elevated values over more than a year
 480 from summer 1998 to fall 1999 during a La Niña event, whereas BSi and dust fluxes showed positive
 481 anomalies of shorter duration (fall 1998 to spring 1999) (Fig. 6b). Investigating SeaWiFS-derived ocean
 482 colour in the Mauritanian upwelling region, Pradhan et al. (2006) obtained a link between the multivariate
 483 ENSO index, the strength of upwelling and the chlorophyll standing stock (250% increase) during the 1998-
 484 1999 La Niña. They also observed that during the mature La Niña phase in the Pacific Ocean, NW African

485 trade winds increased in winter-spring. Coincidentally, Helmke et al. (2005) obtained a more than doubling
486 of the deep ocean organic carbon fluxes in fall 1998 to summer 1999 during the major La Niña phase.

487 We obtained positive carbonate flux anomalies with a longer duration in summer 1998 to fall 1999 and
488 summer 2005 to spring 2006 (Fig. 10b). During fall 1998 (La Niña phase), the area of the Cape Blanc
489 filament was unusually large compared with fall 1997 (El Niño phase) (Fig. 1 d, e). The contribution of
490 major carbonate producers to total carbonate flux varied both on seasonal and interannual timescales (Fischer
491 et al., 2007a). These authors observed that nannofossils contributed almost 95% to carbonate sedimentation
492 in 1991 (a relatively cold year) but only 64% in 1989 (a relatively warm year). On the long-term,
493 nannofossils showed a rather low seasonality. Among the calcareous microorganisms, pteropods had the
494 strongest seasonal signal which did not quite match the pattern of carbonate flux (Fig. 10a, c). As previously
495 observed by Kalberer et al. (1993), a possible explanation is the high pteropod flux (mostly *Limacina inflata*)
496 in summer 1989 due to unusual high SSTs. In our record, we found distinct pulses of pteropods in the
497 summer seasons of 1998, 2002 and 2004 (Fig. 10c). In particular, the peaks in 2002 and 2004 can be
498 attributed to anomalously warm conditions in the study area (Zeeberg et al., 2008; Alheit et al., 2014). Here,
499 a period of near-neutral NAO together with an almost permanent El Niño phase during 2002-2004
500 might have acted in concert towards weakening trade winds which allows a stronger influence of
501 the warm and northward flowing MC, supplying high amounts of pteropods from tropical waters. In
502 summary, ENSO may impact differently on different flux components. Whereas an increase in
503 pteropod fluxes is found during the El Niño phase, La Niña induces an increase in total carbonate
504 flux.

505 Fig. 10.

506

507 ***5.5 Decadal variability and potential trends in mass fluxes***

508 Our records allow a first estimate of deep-ocean mass flux variations beyond seasonal-to-interannual
509 timescales. The 'Bakun upwelling intensification hypothesis' (Bakun, 1990) has been supported by other
510 studies using long-term SSTs, wind stress records or upwelling indices (e.g. Cropper et al., 2014, Narayan
511 et al., 2010). Kahru and Mitchell (2008) applied satellite derived chlorophyll time series from SeaWiFS to
512 conclude that chlorophyll standing stock in major upwelling regions of the world oceans had increased since
513 September 1997. However, these records are rather short (1997-2006) and started in an unusual period with
514 the strongest ENSO ever reported (1997-1998). In our record, no long-term trend in any mass flux
515 component from 1988 through 2012 is seen, which indicates a long term increase or decrease in the strength
516 of coastal upwelling off Cape Blanc. The 1997-2010 chlorophyll time series from SeaWiFS (Fig. 6d) shows
517 a decreasing standing stock, which might indicate a decrease in the strength of coastal upwelling in the Cape
518 Blanc area. The upwelling indices used by Cropper et al. (2014) showed a ~~downward-decreasing~~ trend from
519 1980 to 2013 for the Mauritanian-Senegalese upwelling zone (12-19°N), while observing some interdecadal
520 variability. All these observations together point to regional differences within the upwelling system along

521 the NW African coast (Cropper et al., 2014) with respect to long-term trends in upwelling and chlorophyll
 522 standing stock. According to these findings, only the southernmost weak permanent upwelling zone (21-
 523 26°N) would be in concert with the ‘Bakun upwelling intensification hypothesis’. Another implication is that
 524 trends detected from near-surface data/indices are not necessarily reflected in changes of deep-ocean mass
 525 fluxes and organic carbon sequestration. No evidence of decreasing dust fluxes from the Sahara/Sahel is seen
 526 in our lithogenic (dust) flux -record (Fig. 8a), which might indicate ‘Saharan greening’ and reduced dust
 527 plumes during the past two decades (Zhao et al., 2010; Fontaine et al. 2011). Thus, mass flux patterns might
 528 be partly independent from chlorophyll standing stock or the size of the Cape Blanc filament.

529 Long-term model simulations under present-climate boundary conditions allow to study the linkages within
 530 the climate system on decadal timescales and beyond. Climate modes such as the AMO are operating in this
 531 frequency band, and a correlation between large-scale patterns of SLP and North Atlantic SST (AMO) index
 532 (both lowpass-filtered for periods above 10 years, Fig. 3c) suggests that even on these long timescales,
 533 climate modes such as the AMO might impact on climate variables such as SLP, SST and wind patterns,
 534 specifically through a weakening of the trade winds over the eastern Atlantic during the AMO
 535 warm phase (Fig. 3c). This response of the winds to low-frequent SST variations is consistent with
 536 earlier findings on interdecadal Atlantic SST variability (Kushnir, 1994; Alexander et al., 2014),
 537 and could influence the main characteristics of particle fluxes at our study site (Fig. 5). However, as
 538 current particle flux records from sediment traps only cover a few decades and cannot resolve AMO cycles
 539 with statistical robustness, continuation of trap experiments are essential to capture all relevant timescale
 540 variations. They will help to understand modern particle settling rates and the interpretation of marine
 541 sediment records used in paleoclimate reconstructions.

542

543 6. Summary and Conclusions

544 In our study, we presented a sediment trap record from the Eastern Boundary Upwelling Ecosystem area off
 545 Cape Blanc (Mauritania) for the period 1988-2012. Our major findings can be summarized as follows (also
 546 see Table 4):

547 ~~We made the following major findings which are in part summarized in Table 4:~~

548 ~~1. Fluxes from 1988 to 2012 point to a long term decadal variability, probably related to the Atlantic~~
 549 ~~Multidecadal Oscillation (AMO)-record is~~

550 ~~2.1. Winter BSi fluxes showed a trend of increasing values with an increasing NAO Hurrell Index (Table 4)~~
 551 ~~and the increasing Azores SLP as well. However, both relationships ~~is~~are statistically insignificant.~~

552 ~~3.2. Episodic BSi flux peaks occurred between 2000 and 2005 when the NAO was neutral or negative (Table~~
 553 ~~4. Dust outbreaks, followed by dry (winter) and wet (summer) deposition (e.g. in 2005) into the ocean, might~~
 554 ~~have modified the efficiency of the biological pump and resulted in increased downward fluxes (e.g. of BSi~~
 555 ~~or organic carbon) which were not related to any large scale forcings ~~such as the NAO.~~~~

556 | ~~43.~~ Only the extreme 1997-2000 ENSO was documented clearly in the record, with low fluxes for almost a
 557 | year during the warm El Niño phase, followed by high fluxes of almost a year during the following cold La
 558 | Niña phase (~~Table 4~~).

559 | ~~54.~~ In addition to episodic BSi fluxes, episodic peaks of pteropods occurred in the summers 2002 and 2004
 560 | (Fig. 10c, Table 4). This ~~occurred during was due to~~ a neutral NAO ~~phase~~ and weakening trade winds,
 561 | allowing a stronger influence from tropical surface waters from the south via the Mauritanian Current (MC)
 562 | and an entrainment of Si-rich subsurface waters (~~SACW~~).

563 | ~~65.~~ Teleconnections from ENSO and the NAO may have opposite effects on the NW African
 564 | upwelling (~~Fig. 3~~) with potential implications for deep ocean mass fluxes. In particular, ENSO
 565 | might confound the relationship between the NAO and BSi fluxes.

566 | ~~6. Fluxes from 1988 to 2012 point to a long-term decadal variability, probably related to the Atlantic~~
 567 | ~~Multidecadal Oscillation. However, the time series record is too short to reproduce AMO cycles with~~
 568 | ~~statistical robustness.~~

569 | ~~7. No long-term trend of any flux component was observed in the Mauritanian upwelling off Cape Blanc and~~
 570 | ~~therefore does not which might~~ support the ‘Bakun upwelling intensification hypothesis’ (Bakun, 1990;
 571 | Cropper et al., 2014).

572 | ~~8. We found no evidence of an increasing/decreasing supply of dust and its deposition off Mauritania~~
 573 | ~~between 1988 and 2012.~~

574 | Table 4

575 |

576 | The long-term flux record allows insights into the influences of major climatic oscillations such as the NAO
 577 | and ~~and AMO~~ on ~~the~~ particle export and transfer of particles to the deep ocean and might help to evaluate
 578 | how ~~the~~ ~~is~~ ecosystem ~~off Mauritania~~ could develop in the future. We have some indications that the
 579 | relationships between major Northern Hemisphere climate oscillations (e.g. the NAO) and deep ocean mass
 580 | fluxes are weakened by short-term ecosystem perturbations, e.g. due to dust outbreaks, the latter probably
 581 | leading to episodic sedimentation pulses into the deep ocean. The complex processes of the interaction of
 582 | non-biogenic particles (e.g. different minerals within dust, e.g. Iversen and Roberts, 2015) with organic
 583 | materials produced by photosynthesis, aggregate formation and disintegration in the epi- and mesopelagic,
 584 | particle characteristics (e.g. Nowald et al., 2015), settling rates and remineralization require further process
 585 | studies, combined with laboratory experiments and different modelling approaches (e.g. ~~particle (dis-)~~
 586 | ~~aggregation~~, Karakas et al., 2009).

587 | Additionally, our record provides information on potential long-term changes or trends of mass fluxes which
 588 | point to ecosystem changes or an intensification/weakening of the NW African upwelling system in the study
 589 | area. Considering the present ~~and limited~~ record of bulk fluxes of more than two decades, we have no

590 | indication of any long-term trend which might suggest a fundamental ecosystem change or a regime shift
591 | (step-wise change) in this important coastal upwelling ecosystem ~~off NW Africa~~.

592

593 | *Author contribution*

594 | G. Fischer prepared the ms with contributions from the co-authors. O. Romero investigated the diatom
595 | producers and contributed to the discussion, U. Merkel ~~performed~~ the model simulations and the analysis, B.
596 | Donner studied the carbonate producers, M. Iversen and his group did the dust experiments and provided
597 | unpublished results/observations, N. Nowald and V. Ratmeyer performed the optical observations and
598 | analysis of particles, G. Ruhland and M. Klann designed the sediment trap experiments and analysed the
599 | sediment trap samples, G. Wefer planned the entire program and contributed to the discussion.

600

601 | *Acknowledgements*

602 | We are greatly indebted to the masters and crews of many expeditions ~~which were listed in~~ (Table 2). Many
603 | thanks also go to the chief scientists of the expeditions for their support during the cruises and for the
604 | planning activities and cooperations. We also would like to thank the Mauritanian, Moroccan and German
605 | authorities for their help during the planning phases of the expeditions. This work was only possible because
606 | of the long-term funding by the Deutsche Forschungsgemeinschaft through the SFB 261 (The South Atlantic
607 | in the Late Quaternary: Reconstruction of Mass Budget and Current Systems, 1989-2001) and the Research
608 | Center Ocean Margins. During about the last decade, the study is supported by the Marum Excellence
609 | Cluster, ‘The Ocean in the Earth System’. The model simulation done by U. Merkel has been performed at
610 | the supercomputer of the Norddeutscher Verbund für Hoch- und Höchstleistungsrechnen (HLRN),
611 | Hannover, Germany. We thank the two anonymous reviewers for helpful, fair and constructive comments
612 | and the associate editor for handling the manuscript.

613

614

615 **References**

- 616 Alexander, M.A., Kilbourne, K.H., and Nye, J.A.: Climate variability during warm and cold phases of the
617 Atlantic Multidecadal Oscillation (AMO) 1871–2008. *J. Marine Syst.*, Vol. 133, 14–26, 2014.
- 618 Alheit, J., Licandro, P., Coombs, S., Garcia, A., Giráldez, A., Santamaría, M.T.G., Slotte, A. and Tsikliras,
619 A.C.: Atlantic Multidecadal Oscillation (AMO) modulates dynamics of small pelagic fishes and
620 ecosystem regime shifts in the eastern North and Central Atlantic. *J. Marine Syst.*, 133, 88–102, 2014.
- 621 Arístegui, J., Barton, E. C., Álvarez-Salgado, X. A., Santos, A. M. P., Figueiras, F. G., Kifani, S.,
622 Hernández-León, S., Mason, E., Machú, E., and Demarcq, H.: Sub-regional ecosystem variability in the
623 Canary Current upwelling, *Prog. Oceanogr.*, 83, 33–48., 2009.
- 624 | Armstrong, R.A., Lee, C., Hedges, J.I., Honjo, S., and Wakeham, S.G.: A new mechanistic model of organic
625 carbon fluxes in the ocean based on the quantitative association of POC with ballast minerals, *Deep-Sea*
626 *Res. II*, 49, 219–236, 2002.
- 627 Bakun, A.: Global climate change and intensification of coastal ocean upwelling. *Science* 247, 198–201,
628 1990.
- 629 Barton, E. D., Arístegui, J., Tett, P., Cantón, M., García-Braun, J., Hernández-León, S., Nykjaer, L.,
630 Almeida, C., Almunia, J., Ballesteros, S., Basterretxea, G., Escáñez, J., García-Weill, L., Hernández-
631 Guerra, A., López-Laatzén, F., Molina, P., Montero, M.F., Navarro-Pérez, E., Rodríguez, J. M., van
632 Lenning, K., Vélez, H., and Wild, K.: Eastern Boundary of the North Atlantic: northwest Africa and
633 Iberia, in: *The Global Coastal Ocean*, Vol. 11, edited by: Robinson, A. R. and Brink, K., John Wiley and
634 Sons, New York, Chichester, Weinheim, Brisbane, Singapore, Toronto, 29–67, 1998.
- 635 Behrenfeld, M. J., Randerson, J. T., McClain, C. R., Feldman, G. C., Los, S. O., Tucker, C. J., Falkowski, P.,
636 Field, C. B., Frouin, R., Esaias, W. E., Kolber, D. D., and Pollack, N. H.: Biospheric primary production
637 during an ENSO transition, *Science*, 291, 2594–2597, 2001.
- 638 | Behrenfeld, M.J., [and](#) Falkowski, P.G.: Photosynthetic rates derived from satellite based chlorophyll
639 concentration. *Limnol. Oceanogr.*, 42 (1), 1–20, 1997.
- 640 | Bode, A., Alvarez-Ossorio, M.T., Cabanas, J.M., Miranda, A., [and](#) Varela, M.: Recent trends in plankton and
641 upwelling intensity off Galicia (NW Spain). *Prog. Oceanogr.*, 83, 1–4, 342–350, 2009.
- 642 | Bory, A.J.M., [and](#) Newton, P.P.: Transport of airborne lithogenic material down through the water column in
643 two contrasting regions of the eastern subtropical North Atlantic Ocean. *Global Biogeochem. Cycles* 14
644 (1), 297–315, 2000.
- 645 | Bory, A., Dulac, F., Moulin, C., Chiapello, I., Newton, P.P., Guelle, W., Lambert, C.E., [and](#) Bergametti, G.:
646 Atmospheric and oceanic dust fluxes in the northeastern tropical Atlantic ocean: how close a coupling?
647 *Ann. Geophys.*, 20, 2067–2076, 2002.
- 648 | [Bressac, M., Gieu, C., Doraxan, D., Bourrin, F., Desboeufs, K., Leblond, N., and C. Ridame: Quantification](#)
649 [of the lithogenic pump following a dust deposition event, *Biogeosciences*, 11, 1007-1020, 2014.](#)
- 650 | Brust, J., Schulz-Bull, D.E., Leipe, T., Chavagnac, V., [and](#) Wanick, J.J.: Descending particles: from the
651 atmosphere to the deep ocean: A time series study in the subtropical NE Atlantic. *Geophys. Res. Lett.*, 38,
652 L06603. <http://dx.doi.org/10.1029/2010GL045399>, 2011.
- 653 Buesseler, K. O., Antia, A. A., Chen, M., Fowler, S. W., Gardner, W. D., Gustafsson, O., Harada, K.,
654 Michaels, A. F., Rutgers van der Loeff, M., Sarin, M., Steinberg, D. K., and Trull, T.: An assessment of
655 the use of sediment traps for estimating upper ocean particle fluxes, *J. Mar. Res.*, 65, 345–416, 2007.
- 656 Carr, M.-E.: Estimation of potential productivity in Eastern Boundary Currents using remote sensing. *Deep-*
657 *Sea Res. I*, 49, 59–80, 2002.
- 658 | [Checkley, D.M.Jr., and Barth, J.A.: Patterns and processes in the California Current Systems. *Prog.*](#)
659 [Oceanogr.](#), 83, 49-64, 2009.
- 660 | Chiapello, I., Moulin, C., [and](#) Prospero, J.M.: Understanding the long-term variability of African dust
661 transport across the Atlantic as recorded in both Barbados surface concentrations and large-scale Total
662 Ozone Mapping Spectrometer (TOMS) optical thickness. *J. Geophys. Res.*, 110, D18S10,
663 doi:10.1029/2004JD005132, 2005.

- 664 Collins, W. D., Bitz, C.M., Blackmon, M.L. Bonan, G.B., Bretherton, C.S., Carton, J.A., Chang, P., Doney,
665 S.C., Hack, J.J., Henderson, T.B., Kiehl, J.T., Large, W.G., McKenna, D.S., Santer, B.D., and Smith,
666 R.D.: The Community Climate System Model Version (CCSM3), *J. Climate*, 19, 2122-2143. 2006.
- 667 Cropper, T.E., Hanna, [and](#) E., Bigg, G.R.: Spatial and temporal seasonal trends in coastal upwelling off
668 Northwest Africa, 1981–2012. *Deep-Sea Res. II*, 86, 94–111, 2014.
- 669 Diatta, S., and Fink, A. H.: Statistical relationship between remote climate indices and West African
670 monsoon variability, *Int. J. Climatol.*, 34, 3348–3367, doi:10.1002/joc.3912, 2014.
- 671 [Ducklow, H.W., Doney, S.C., and Steinberg, D.K.: Contributions of long-term research and time-series
672 observations to marine ecology and biogeochemistry. *Annu. Rev. Mar. Sci.*, 1, 279-302, 2009.](#)
- 673 Dunne, J. P., Sarmiento, J.L., and Gnanadesikan, A.: A synthesis of global particle export from the surface
674 ocean and cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cycles*, 21,
675 GB4006, doi:10.1029/2006GB002907, 2007.
- 676 Engelstaedter, S., Tegen I., and Washington, R.: North African dust emissions and transport. *Earth Sci. Rev.*,
677 79, 73-100, 2006.
- 678 Fischer, G., and Wefer, G.: Sampling, preparation and analysis of marine particulate matter. In: *The Analysis
679 and Characterization of Marine Particles*, edited by Hurd, D.C., and Spencer, D.W., *Geophysical
680 Monograph Series*, No. 63, 391-397, 1991.
- 681 Fischer, G., Donner, B., Ratmeyer, V., Davenport, R., [and](#) Wefer, G.: Distinct year-to-year particle flux
682 variations off Cape Blanc during 1988–1991: Relation to delta $\delta^{18}\text{O}$ -deduced sea-surface temperatures
683 and trade winds. *J. Marine Res.*, 54 (1), 73–98, 1996.
- 684 [Fischer, G., Wefer, G., Romero, O., Dittert, N., Ratmeyer, V. and Donner, B.: Transfer of particles into the
685 deep Atlantic and the global Ocean: control of nutrient supply and ballast production. In: *The South
686 Atlantic in the Late Quaternary: Reconstruction of material budget and current systems*, edited by Wefer,
687 G., Mulitza, S. and Ratmeyer, V., Springer, Berlin, Heidelberg, New York, 21-46, 2003.](#)
- 688 Fischer, G., Karakas, G., Blaas, M., Ratmeyer, V., Nowald, N., Schlitzer, R., Helmke, P., Davenport, R.,
689 Donner, B., Neuer, S., and Wefer, G.: Mineral ballast and particle settling rates in the coastal upwelling
690 system off NW Africa and the South Atlantic. 98, 281–298, doi:10.1007/s00531-007-0234-7, 2009a.
- 691 [Fischer, G., Reuter, C., Karakas, G., Nowald, N., and Wefer, G.: Offshore advection of particles within the
692 Cape Blanc filament, Mauritania: Results from observational and modelling studies. *Prog. Oceanogr.*, 83,
693 322–330, 2009b.](#)
- 694 Fischer, G., [and](#) Karakas, G.: Sinking rates and ballast composition of particles in the Atlantic Ocean:
695 implications for the organic carbon fluxes to the deep ocean. *Biogeosciences* 6, 85–102, 2009b.
- 696 [Fischer, G., Reuter, C., Karakas, G., Nowald, N., and Wefer, G.: Offshore advection of particles within the
697 Cape Blanc filament, Mauritania: Results from observational and modelling studies. *Prog. Oceanogr.*, 83,
698 322–330, 2009.](#)
- 699 Fischer, G., S. Neuer, R. Davenport, O. Romero, V. Ratmeyer, B. Donner, T. Freudenthal, H. Meggers and
700 G. Wefer. The Northwest African Margin. In: *Carbon and Nutrient Fluxes in Continental Margins: A
701 Global Synthesis*, edited by Liu, K. K., Atkinson, L., Quinones, R., Talaue-McManaus, L., IGBP Book
702 Series, Springer, Berlin, 77-103. 2010.
- 703 Fontaine, B., Roucou, P., Gaetani, M., and Marteau, R.: Recent changes in precipitation, ITCZ convection
704 and northern tropical circulation over North Africa (1979-2007). *Int. J. Climatol.*, 31, 633-648, 2011.
- 705 Fréon, P., Barange, M., and Aristegui, J.: Eastern Boundary Upwelling Ecosystems: integrative and
706 comparative approaches. *Prog. Oceanogr.*, 83, 1–14, 2009.
- 707 Friese, C., van der Does, M., Merkel, U., Iversen, M., Fischer, G., and Stuu, J.-B.: Environmental factors
708 controlling the seasonal variation in particle size of modern Saharan dust deposited offshore Cape, Blanc.
709 *Aeolian Res.*, [in press submitted](#), 2016.
- 710 Goudie, A.S., and Middleton, N.J.: Saharan dust storms: nature and consequences. *Earth Sci. Rev.*, 56, 179-
711 204, 2001.

- 712 | [Hedges, J.I., Baldock, J.A., Gelin, Y., Lee, C., Peterson, M.L., and Wakeham, S.G.: The biochemical and](#)
713 | [elemental compositions of marine plankton: a NMR perspective. *Marine Chemistry* 78 \(1\), 47–63, 2002.](#)
- 714 | Helmke, P., Romero, O., and Fischer, G.: Northwest African upwelling and its effect on off-shore organic
715 | carbon export to the deep sea, *Global Biogeochem. Cy.*, 19, GB4015, doi:10.1029/2004GB002265, 2005.
- 716 | Hsu, N.C., Gautam, R., Sayer, A.M., Bettenhausen, C., Li, C., Jeong, M.J., Tsay, S.-C., and Holben, B.N.:
717 | Global and regional trends of aerosol optical depth over land and ocean using SeaWiFS measurements
718 | from 1997-2010. *Atmos. Chem. Phys.*, 12, 8037-8053, 2012.
- 719 | Hurrell, J.W.: NAO Index Data provided by the Climate Analysis Section, NCAR, Boulder, USA, Updated
720 | regularly. Accessed 01 January 2012, 1995.
- 721 | Iversen, M.H., and Ploug, H.: Ballast minerals and the sinking carbon flux in the ocean: carbon-specific
722 | respiration rates and sinking velocities of marine snow aggregates. *Biogeosciences* 7, 2613–2624.
723 | <http://dx.doi.org/10.5194/bg-7-2613-2010>, 2010.
- 724 | Iversen, M.H., Nowald, N., Ploug, H., Jackson, G.A., and Fischer, G.: High resolution profiles of vertical
725 | particulate organic matter export off Cape Blanc, Mauritania: degradation processes and ballasting
726 | effects. *Deep Sea Res. I*, 57, 771–784, 2010.
- 727 | Iversen, M.H., and Robert, M.L.: Ballasting effects of smectite on aggregate formation and export from a
728 | natural plankton community. *Marine Chem.*, 175, 18-27, 2015.
- 729 | Jickells, T.D., An, Z.S., Andersen, K.K., Baker, A.R., Bergametti, G., Brooks, N., Cao, J.J., Boyd, P.W.,
730 | Duce, R.A., Hunter, K.A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P.S., Mahowald, N., Prospero,
731 | J.M., Ridgwell, A.J., Tegen, I., and Torres, R.: Global iron connections between desert dust, ocean
732 | biogeochemistry, and climate. *Science*, 308 (5718), 67–71, 2005.
- 733 | Kahru, M., and Mitchell, B. G.: Ocean colour reveals increased blooms in various parts of the world ocean.
734 | *EOS* 89, 170, 2008.
- 735 | Kalberer, M., G. Fischer, J. Pätzold, B. Donner, M. Segl and Wefer, G.: Seasonal sedimentation and stable
736 | isotope records of pteropods off Cape Blanc. *Marine Geol.*, 113, 305-320, 1993.
- 737 | Karakas, G., Nowald, N., Blaas, M., Marchesiello, P., Frickenhaus, S., and Schlitzer, R.: High-resolution
738 | modeling of sediment erosion and particle transport across the northwest African shelf. *J. Geophys. Res.*,
739 | 111. C06025, <http://dx.doi.org/10.1029/2005JC003296>, 2006.
- 740 | Karakas, G., Nowald, N., Schäfer-Neth, C., Iversen, M.H., Barkmann, W., Fischer, G., Marchesiello, P., and
741 | Schlitzer, R.: Impact of particle aggregation on vertical fluxes of organic matter. *Prog. Oceanogr.* 83,
742 | 331–341, 2009.
- 743 | Kaufman, Y. J., Koren, I., Remer, L. A., Tanré, D., Ginoux, P., and Fan, S.: Dust transport and deposition
744 | from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) spacecraft over the Atlantic
745 | Ocean, *J. Geophys. Res.*, 110, D10S12, doi:10.1029/2003JD004436, 2005.
- 746 | Klaas, C., and Archer, D.E.: Association of sinking organic matter with various types of ballast in the deep
747 | sea: Implications for the rain ratio, *Global Biogeochem. Cycles*, 16(4), 1116,
748 | doi:10.1029/2001GB001765, 2002.
- 749 | Kremling, K., Lentz, U., Zeitzschell, B., Schulz-Bull, D.E., and Duinker, J.C.: New type of time-series
750 | sediment trap for the reliable collection of inorganic and organic trace chemical substances. *Rev. Scient.*
751 | *Instr.*, 67, 4360–4363, 1996.
- 752 | Kushnir, Y.: Interdecadal variations in North Atlantic sea surface temperature and associated
753 | atmospheric conditions. *J. of Climate*, 7(1), 141-157, 1994.
- 754 | Lathuilière, C., Echevin, V., and Levy, M.: Seasonal and intraseasonal surface chlorophyll-a variability along
755 | the northwest African coast. *J. Geophys. Res., Oceans*, 13 (C5), C05007, doi:10.1029/2007/JC004433,
756 | 2008.
- 757 | Lucio, P.S., Baldicero Molion, L.C., de Avial-Valadão, C.E., Conde, F.C., Malheiro Ramos, A., and Dias de
758 | Melo, M.L.: Dynamical outlines of the rainfall variability and the ITCZ role over the West Sahel.,
759 | *Atmospheric Climate Sci.*, 2, 337-350, 2012.

- 760 Mahowald, N., Kohfeld, K., Hansson, M., Balkanski, Y., Harrison, S.P., Prentice, I.C., Schulz, M., and
 761 Rodhe, H.: Dust sources and deposition during the last glacial maximum and current climate: a
 762 comparison of model results with paleodata from ice cores and marine sediments. *J. Geophys. Res.,*
 763 *Atmospheres*, 104 (D13), 15895–15916, 1999.
- 764 Marcello, J., Hernandez-Guerra, A., Eugenio, F., and Fonte, A.: Seasonal and temporal study of the
 765 northwest African upwelling system. *Internat. J. Remote Sens.*, 32, 7, 1843-1859, 2011.
- 766 McGregor, H.V., Dima, M., Fischer, H.W., and Mulitza, S.: Rapid 20th century increase in coastal upwelling
 767 off Northwest Africa. *Science*, 315, 637-639, 2007.
- 768 Meunier, T., Barton, E.D., Barreiro, B., and Torres, R.: Upwelling filaments off Cape Blanc: interaction of
 769 the NW African upwelling current and the Cape Verde frontal zone eddy field? *J. Geophys. Res., Oceans*,
 770 117 (C8), C08031, doi:10.1029/2012JC007905, 2012.
- 771 Mittelstaedt, E.: The ocean boundary along the northwest African coast: Circulation and oceanographic
 772 properties at the sea surface: *Prog. Oceanogr.*, 26, 307–355, 1991.
- 773 Moulin, C. and Chiapello, I.: Evidence of the control of summer atmospheric transport of African dust over
 774 the Atlantic by Sahel sources from TOMS satellites (1979-2000). *Geophys. Res. Lett.*, 31, L02107,
 775 doi:10.1029/2003GL019031, 2004.
- 776 Müller P.J. and Schneider, R.: An automated leaching method for the determination of opal in sediments and
 777 particulate matter. *Deep Sea Res. I*, 40, 425-444, 1993.
- 778 Muller-Karger, F. E., Varela, R., Thunell, R., Luerssen, R., Hu, C., and Walsh, J. J.: The importance of
 779 continental margins in the global carbon cycle, *Geophys. Res. Lett.*, 32, L01602,
 780 doi:10.1029/2004GL021346, 2005.
- 781 Narajan, N., Paul, A., and Schulz, M.: Trends in coastal upwelling intensity during the late 20th century.
 782 *Ocean Sci.*, 6, 815-823, 2014.
- 783 Neuer, S., Torres-Padron, M.E., Gelado-Caballeo, M.D., Rueda, M.J., Hernandez-Brito, J., Davenport, R.,
 784 and Wefer, G.: Dust deposition to the eastern subtropical North Atlantic gyre: Does ocean's
 785 biogeochemistry respond? *Global Biogeochem. Cy.*, 18, GB4020,
 786 <http://dx.doi.org/10.1029/2004GB002228>, 2004.
- 787 Nicholson, S.E.: The West African Sahel: A review of recent studies on the rainfall regime and its
 788 interannual variability. *ISRN Meteorology*, 2013, 453521, doi:10.1155/2013/453521, 2013.
- 789 Nowald, N., Iversen, M.H., Fischer, G., Ratmeyer, V., and Wefer, G.: Time series of in-situ particle
 790 properties and sediment trap fluxes in the coastal upwelling filament off Cape Blanc, Mauritania. *Prog.*
 791 *Oceanogr.*, 137, Part A, 1–11, 2015, doi:10.1016/j.pcean.2014.12.015, 2015.
- 792 N'Tchayi Mbourou, G., Berrand, J.J., and Nicholson, S.E.: The diurnal and seasonal cycles of wind-borne
 793 dust over Africa north of the equator. *J. Appl. Meteorology*, 36, 868-882, 1997.
- 794 Nykjaer, L., and Van Camp, L.: Seasonal and interannual variability of coastal upwelling along northwest
 795 Africa and Portugal from, 1981 to 1991. *J. Geophys. Res.*, 99, 197–207, 1994.
- 796 Pauly, D., and Christensen, V.: Primary production required to sustain global fisheries. *Nature*, 374, 255-257,
 797 1995.
- 798 Ploug, H., Iversen, M.H., and Fischer, G.: Ballast, sinking velocity, and apparent diffusivity within marine
 799 snow and zooplankton fecal pellets: implications for substrate turnover by attached bacteria. *Limnol.*
 800 *Oceanogr.*, 53 (5), 1878–1886, 2008.
- 801 Pradhan, Y., Lavender, S.J., Hardman-Mountford, N.J., and Aiken, J.: Seasonal and inter-annual variability
 802 of chlorophyll-a concentration in the Mauritanian upwelling: observation of an anomalous event during
 803 1998–1999. *Deep Sea Res. II*, 53, 1548–1559, 2006.
- 804 Prospero, J.M.: Mineral-aerosol transport to the North Atlantic and North Pacific: The impact of African and
 805 Asian sources. In: *The long range atmospheric transport of natural and contaminant substances*, edited by
 806 Knap, A.H., Dordrecht, Mathematical and Physical Sciences. Kluwer Academic Publishers, 19–52, 1990.
- 807 Prospero, J.M., and Lamb, P.J.: African droughts and dust transport to the Caribbean: climate change
 808 implications. *Science*, 302, 1024-1027, 2003.

- 809 Ratmeyer, V., Fischer, G. and Wefer, G.: Lithogenic particle fluxes and grain size distributions in the deep
810 ocean off northwest Africa: Implications for seasonal changes of aeolian dust input and downward
811 transport, *Deep-Sea Res. II*, 46, 1289-1337, 1999a.
- 812 Ratmeyer, V., Balzer, W., Bergametti, G., Chiapello, I., Fischer, G., and Wyputta, U.: Seasonal impact of
813 mineral dust on deep-ocean particle flux in the eastern subtropical Atlantic Ocean, *Mar. Geol.*, 159, 241–
814 252, 1999b.
- 815 [Redfield, A. C., Ketchum, B.H., and Richards, F.A.: The influence of organisms on the composition of
816 seawater. In: *The Sea*, edited by Hill, M.N., Vol. 2, Wiley and Sons, Chichester, 26-77, 1963.](#)
- 817 [Romero, O.E., Lange, C.B., Swap, R.J., and Wefer, G.: Eolian-transported freshwater diatoms and phytoliths
818 across the equatorial Atlantic record temporal changes in Saharan dust transport patterns. *J. Geophys.
819 Res.*, 104, 3211-3222, 1999.](#)
- 820 [Romero, O.E., Lange, C.B., and Wefer, G.: Interannual variability \(1988-1991\) of siliceous phytoplankton
821 fluxes off northwest Africa. *J. Plank. Res.*, 24 \(10\), 1035-1046, doi:10.1093/plankt/1024.1010.1035.
822 2002.](#)
- 823 Romero, O. E., Kim, J.-H. , and Donner, B.: Submillennial-to-millennial variability of diatom production off
824 Mauritania, NW Africa, during the last glacial cycle, *Paleoceanography*, 23, PA3218,
825 doi:10.1029/2008PA001601, 2008.
- 826 Shanahan, T. M., Overpeck, J. T., and Anchukaitis, K. J., Beck, J. W., Cole, J. E., Dettman, D. L., Peck, J.
827 A., Scholz, C. A., and King, J. W.: Atlantic Forcing of persistent drought in West Africa, *Science*, 324,
828 377–380, 2009.
- 829 Siegel, D.A., [and](#) Deuser, W.G.: Trajectories of sinking particles in the Sargasso Sea: modeling of statistical
830 funnels above deep-ocean sediment traps. *Deep-Sea Res. I*, 44 (9–10), 1519–1541, 1997.
- 831 Ternon, E., Guieu, C., Loye-Pilot, M.-D., Leblond, N., Bosc, E., Gasser, B., Miquel, J.C., [and](#) Martin, J.: The
832 impact of Saharan dust on the particulate export in the water column of the North Western Mediterranean
833 Sea. *Biogeosciences* 7, 809–826, 2010.
- 834 [Thunell, R., Benitez-Nelson, C., Varela, R., Astor, Y., and Muller-Karger, F.: Particulate organic carbon
835 fluxes along upwelling-dominated continental margins: rates and mechanisms. *Glob. Biogeochem. Cy.*,
836 21, GB1022, doi:10.1029/2006GB002793, 2007.](#)
- 837 Van Camp, L., Nykjær, L., Mittelstaedt, E., and Schlittenhardt, P.: Upwelling and boundary circulation off
838 Northwest Africa as depicted by infrared and visible satellite observations. *Prog. Oceanogr.*, 26, 357–402,
839 1991.
- 840 Yeager, S. G., Shields, C.A., Large, W.G., and Hack, J.J. : The Low-Resolution CCSM3, *J. Climate*, 19,
841 2545-2566, 2006.
- 842 Yu E. F., Francois R., Honjo S., Fleer A.P., Manganini S.J., Rutgers van der Loeff, M.M., and Ittekkot V.:
843 Trapping efficiency of bottom-tethered sediment traps estimated from the intercepted fluxes of ²³⁰Th and
844 ²³¹Pa. *Deep-Sea Res. I*, 48, 865-889, 2001.
- 845 Zeeberg, J., Corten, A., Tjoe-Awie, P., Coca, J., and Hamady, B.: Climate modulates the effects of
846 *Sardinella aurita* fisheries o_ Northwest Africa, *Fish. Res.*, 89, 65–75, 2008.
- 847 Zenk, W., Klein, B., and Schroder, M.: Cape Verde Frontal Zone. *Deep Sea Res. I*, 38, 505–530, 1991.
- 848 Zhao, T. X.-P., Laszlo, I., Guo, W., Heindinger, A., Cao, C., Jelenak, A., Tarpley, D., and Sullivan, J.: Study
849 of long-term trend in aerosol optical thickness observed from operational AVHRR satellite instrument, *J.*
850 *Geophys. Res.*, 113, D07201, doi:10.1029/2007JD009061, 2008.
- 851

852 **Figure Captions**

853 Fig. 1. General setting of the study area: a: Oceanographic setting in the area of the long-term mooring site
 854 Cape Blanc (CB_{meso}) within the Cape Blanc filament (green arrow), dissolving into eddies (indicated as
 855 circles with arrows) further offshore. The Cape Verde Frontal Zone (CVFZ) separating the subsurface water
 856 masses of the NACW and the SACW (Zenk et al., 1991) is shown. Upwelling zones are marked according to
 857 Cropper et al. (2014). Ocean colour map (chlorophyll, 9 km resolution) from MODIS is shown for two
 858 extreme years, winter 2006-2007 (b: NAO+) and winter 2009-2010 (c: NAO-). SeaWiFS ocean colour
 859 during two contrasting situations for the strongest ENSO cycle 1997-1999: fall 1997 during the warm El
 860 Niño phase (d), and fall 1998 during the cold La Niña event (e). The study site CB_{meso} is indicated by a
 861 square box in the ocean colour pictures. green arrow indicates the Cape Blanc filament, yellow arrows the
 862 major dust transport. *MC=Mauritanian Current, CC=Canary Current, NEC=North Equatorial Current*.

863
 864 Fig. 2. The NAO Hurrell index (DJFM, station-based, Lisbon-Reykjavik; Hurrell, 1995) plotted from 1864 to
 865 2012. Note the strong changes during the last four decades. Grey shading indicates the time period covered
 866 by the long-term flux record off Cape Blanc, Mauritania. A 5-point running mean is shown by the thick line.

867
 868 Fig. 3: Teleconnections affecting the study site off Cape Blanc. Correlation of simulated sea-level pressure
 869 (SLP) with a: the NAO SLP index after Hurrell (Hurrell, 1995; boreal winter season), b: the Nino3 SST
 870 index (boreal winter season), and c: North Atlantic SST (low pass-filter applied considering periods above
 871 10 years). Analysis based on the last 100 model years of a present-day control simulation using the CCSM3
 872 model.

873
 874 Fig. 4. Long-term sSeasonal means of major bulk fluxes of the lower traps only (a: total, b: organic carbon,
 875 c: nitrogen, d: biogenic silica (=BSi), e: carbonate, and f: lithogenic=mineral dust) and the respective
 876 standard deviations (1 s.d.), which reflect interannual variability. Relative contributions (%) of BSi, organic
 877 carbon, nitrogen, carbonate and lithogenic materials to total mass in the respective seasons are indicated by
 878 numbers below the bars.

879
 880 Fig. 5. a: Total mass fluxes of the lower traps (grey-shaded). Gaps were filled with upper trap data (light grey
 881 barslight-stippled). Deviations of the seasonal total mass fluxes from the long-term seasonal means
 882 (anomalies), fitted with a 9-order polynomial (eial (b). c: Atlantic Multidecadal Oscillation (AMO) Index based
 883 on monthly SST fitted with a 9th-order polynomial fit (dashed blue line). The strong ENSO cycle 1997-1999
 884 with a warm El Niño and a cold La Niña phase is indicated.

885
 886 Fig. 6. a: Seasonal flux of biogenic silica (BSi, green) with gaps filled from the upper trap data (light green
 887 barslight-green). Deviations of the long-term seasonal means (anomalies, b). c: The NAO Hurrell index
 888 (DJFM). d: Seasonal chlorophyll concentration both from the MODIS (light green) and the SeaWiFS (dark
 889 green) sensors at 9 km resolution. Note that high chlorophyll biomass is generally occurring in spring but

890 sometimes in summer/fall as well (e.g. in 1998, 2007). SeaWiFS chlorophyll reveals a downward trend from
 891 1997 to 2010, not mimicked in any flux data. The strong ENSO cycle 1997-1999 with a warm El Niño and a
 892 cold La Niña phase is indicated.

893
 894 Fig. 7. a: The NAO Hurrell index (DJFM, Hurrell, 1995) plotted against winter BSi fluxes from Fig. 6. Note
 895 the increase of BSi with increasing NAO index. However, the ~~relationship is weak~~~~correlation coefficient is~~
 896 ~~low~~ due to unusual sedimentation events in the years 1998-99, 2002, 2004, and, in particular in 2005. When
 897 omitting the data point from 2005, the correlation coefficient increases, but remains low ($R^2=0.14$, $N=20$).
 898 Upper trap flux data from winter 1998 and 2004 may be too low as the filament with elevated chlorophyll
 899 was small and the particles did not reach the upper trap (see text). Omitting these two data point would
 900 slightly improve the relationship.

Kommentar [g4]: See comment reviewer #1

901 b: The size of the Cape Blanc filament (Fischer et al., 2009) during winter months (DJFM) versus winter
 902 BSi fluxes shows higher fluxes with larger filament size. When omitting the ~~BSi flux from~~ winter ~~2005~~~~BSi~~
 903 ~~flux from 2005~~, a statistically significant relationship between filament size and fluxes is obtained ($R^2=0.63$,
 904 $N=10$). Years given in the figure denote the respective winter seasons (e.g. 1999 = Dec 1998 – Mar 1999).

905
 906 Fig. 8. a: Seasonal flux of lithogenic ~~partieles~~ (=mineral dust) particles (orange) with gaps filled from the
 907 upper trap data (light orange bars~~light orange~~). Deviations from the long-term seasonal means (anomalies, b).
 908 Note the large positive anomalies with longer duration in 1988-89, 1997-2000 and 2005-2006. From about
 909 2000 to 2004-2005, lithogenic fluxes remain rather low. In 2005, dust sedimentation and BSi flux (Fig. 6b)
 910 were high throughout the year. c: The AOT from the SeaWiFS (brown) and MODIS (light brown) sensors
 911 shows repeatedly high values in summer, but not in winter when dust sedimentation is highest in the study
 912 area. A typical short-term (2-day) dust storm in January 2012 ~~with a duration of about 2 days~~ is shown as
 913 insert in the upper right. The strong ENSO cycle 1997-1999 with a warm El Niño and a cold La Niña phase
 914 is indicated.

915
 916 Fig. 9. Relationships between BSi and lithogenic (= mineral dust) fluxes for the winter (a) and summer (b)
 917 seasons. Note the high correspondence in winter (~~linear~~- $R^2=0.78$, $N=21$); a ~~much~~ lower coefficient is found
 918 for the summer season. During the outstanding year of 2005 (see Fig. 7), both points for winter and summer
 919 are close to the linear regression line.

920
 921 Fig. 10. a: Seasonal flux of total carbonate (blue) with gaps filled from the upper trap data (light blue bars).
 922 Deviations from the long-term seasonal means (anomalies, b). c: Seasonal flux of pteropods. During the
 923 strongest ENSO cycle 1997-2000, longer periods of low and high carbonate fluxes occurred. Note the
 924 episodic sedimentation pattern of pteropods with maxima e.g. in summer 1998, 2002 and 2004. The strong
 925 ENSO cycle 1997-1999 with a warm El Niño and a cold La Niña phase is indicated.

926
 927

928 Table 1. Deployment data of the moorings and traps at the mesotrophic sediment trap site CB, Cape Blanc,
 929 Mauritania. Associated ships' cruises and references to earlier publications on fluxes are indicated.

Trap	LAT	LONG	water depth	trap depth	sampling		no	remark,	relevant cruise recovery/
name	N	W	m	m	start	end	of samples	reference to fluxes	GeoB no.
CB-1 lower	20°45.3'	19°44.5'	3646	2195	22.03.88	08.03.89	13	<i>Fischer et al. 1996, 2003</i>	Meteor 9/4/ GeoB 1121-4
CB-2 lower	21°08.7'	20°41.2'	4092	3502	15.03.89	24.03.90	22	<i>Fischer et al. 1996, 2003</i>	Meteor 12/1/ GeoB 1230-1
CB-3 lower	21°08.3'	20°40.3'	4094	3557	29.04.90	08.04.91	17	<i>Fischer et al. 1996, 2003, 2010</i>	Polarstern ANT IX/4
CB-4 lower	21°08.7'	20°41.2'	4108	3562	03.03.91	19.11.91	13	<i>Fischer et al. 1996, 2003, 2010</i>	Meteor 20/1/ GeoB 1602-1
CB-5 lower	21°08.6'	20°40.9'	4119	3587	06.06.94	27.08.94	19		Meteor 29/3/ GeoB 2912-1
CB-6 upper	21°15.0'	20°41.8'	4137	771	02.09.94	25.10.95	20	<i>Fischer et al. 2010</i>	Polarstern ANT XIII/1
CB-7 lower	21°15.4'	20°41.8'	4152	3586	20.11.95	29.01.97	20		Meteor 38/1/ GeoB 4302-7
CB-8 upper	21°16.3'	20°41.5'	4120	745	30.01.97	04.06.98	20	<i>Fischer et al. 2010</i>	Meteor 41/4/ GeoB 5210-2
CB-9 lower	21°15.2'	20°42.4'	4121	3580	11.06.98	07.11.99	20	<i>Helmke et al. 2005</i>	Metero 46/1/ GeoB 6103-3
CB-10 lower	21°17.2'	20°44.1'	4125	3586	10.11.99	10-10-00	3	mostly no seasonal sampling	Polarstern ANT XVIII/1
CB-11 upper	21°16.8'	20°43.0'	4113	1003	11.10.00	30.03.01	20		Poseidon 272/ GeoB 7401-1
CB-12 lower	21°16.0'	20°46.5'	4145	3610	05.04.01	22.04.02	14		Meteor 53/1c/ GeoB 7917-1
CB-13 lower	21°16.8'	20°46.7'	4131	3606	23.04.02	08.05.03	20	<i>Fischer et al., 2009</i> <i>Fischer and Karkas, 2009</i>	Meteor 58/2b/ GeoB 8628-1
CB-14 upper	21°17.2'	20°47.6'	4162	1246	31.05.03	05.04.04	20		Poseidon 310/ no number
CB-15 lower	21°17.9'	20°47.8'	4162	3624	17.04.04	21.07.05	20		Meteor 65/2/ no number
CB-16 lower	21°16.8'	20°47.8'	4160	3633	25.07.05	28.09.06	20		Poseidon 344/1/ GeoB 11401-1
CB-17 lower	21°16.4'	20°48.2'	4152	3614	24.10.06	25.03.07	20		Merian 04/b/ GeoB 11833-1
CB-18 lower	21°16.9'	20°48.1'	4168	3629	25.03.07	05.04.08	20		Poseidon 365/2/ GeoB 12907-1
CB-19 lower	21°16.2'	20°48.7'	4155	3617	22.04.08	22.03.09	20		Merian 11/2/ GeoB 13616-4
CB-20 upper	21°15.6'	20°50.7'	4170	1224	03.04.09	26.02.10	19		Poseidon 396/ GeoB 14201-3
CB-21 lower	21°15.6'	20°50.9'	4155	3617	28.02.10	04.04.11	20		Merian 18/1/ GeoB 15709-1
CB-22 lower	21°16.1'	20°50.9'	4160	3622	05.05.11	11.01.12	15		Poseidon 425/ GeoB 16101-1
CB-23 lower	21°15.8'	20°52.4'	4160	3622	20.01.12	22.01.13	18		Poseidon 445/ GeoB 17102-5

931 | Table 2. Seasonal flux data [and percentages of major bulk components of total flux](#) at the mesotrophic sediment trap site CB from 1988 to 2012.

CB meso	interval		sample no. of trap	season	year	duration days	remark	TTL mass	BSibieg-opal	org. carbon	nitrogen	carbonate	lithogenic	BSi	org. carbon	nitrogen	carb.	lith.
	start	end						g m ⁻²	%	%	%	%	%					
CB-1 lower	22.03.88	11.06.88	#1-3	spring	1988	81		15,64	1,91	0,59	0,069	4,89	7,66	12,23	3,77	0,44	31,25	48,96
	11.06.88	27.09.88	#4-7	summer		108		23,01	1,57	1,07	0,135	10,83	8,47	6,81	4,66	0,59	47,07	36,81
	27.09.88	17.12.88	#8-10	fall		81		14,12	0,86	0,51	0,056	6,78	5,46	6,11	3,60	0,40	48,00	38,70
	17.12.88	08.03.89	#11-13	winter	1989	81		11,50	0,89	0,45	0,055	5,09	4,63	7,70	3,92	0,48	44,21	40,23
CB-2 lower	15.03.89	25.06.89	#1-6	spring		102		12,91	0,52	0,40	0,051	6,92	4,68	4,03	3,10	0,40	53,60	36,25
	25.06.89	18.09.89	#7-11	summer		85		13,62	0,49	0,39	0,046	7,48	4,87	3,60	2,86	0,34	54,92	35,76
	18.09.89	29.12.89	#12-17	fall		102		16,29	0,67	0,48	0,056	8,21	6,45	4,11	2,95	0,34	50,40	39,59
	29.12.89	24.03.90	#18-22	winter	1990	85		14,06	0,75	0,46	0,055	6,81	5,58	5,33	3,27	0,39	48,44	39,69
CB-3 lower	29.04.90	03.07.90	#2-4	spring		64,5		12,68	0,49	0,39	0,047	6,78	4,64	3,87	3,04	0,37	53,49	36,55
	03.07.90	27.09.90	#5-8	summer		86		13,20	0,40	0,39	0,044	7,35	4,67	3,05	2,98	0,33	55,64	35,35
	27.09.90	22.12.90	#9-12	fall		86		9,76	0,40	0,44	0,052	4,02	4,46	4,08	4,52	0,53	41,21	45,66
	22.12.90	18.03.91	#13-16	winter	1991	86		10,89	0,58	0,73	0,091	4,50	4,34	5,33	6,73	0,84	41,35	39,87
CB-4 lower	18.03.91	22.06.91	#17 + #1-5	spring		71,5	gap	4,87	0,24	0,38	0,053	2,09	1,77	4,89	7,83	1,09	43,02	36,36
	22.06.91	20.09.91	# 6-14	summer		90		9,06	0,48	0,83	0,110	3,66	3,25	5,30	9,16	1,21	40,40	35,87
	20.09.91	19.11.91	# 15-20	fall		60		2,67	0,13	0,17	0,023	1,31	0,89	4,87	6,37	0,86	49,06	33,33
							no sampling											
CB-5 lower	06.06.94	23.06.94	#1-4	spring	1994	17		1,76	0,05	0,05	0,007	1,27	0,35	2,84	2,84	0,40	72,16	19,89
	23.06.94	27.08.94	#5-19	summer		65		7,30	0,16	0,14	0,023	5,97	0,89	2,19	1,92	0,32	81,78	12,19
CB-6 upper	24.09.94	21.12.94	# 2-5	fall		88		11,58	0,26	0,70	0,104	6,82	3,10	2,27	6,02	0,90	58,92	26,74
	21.12.94	19.03.95	# 6-9	winter	1995	88		12,44	0,96	0,74	0,113	5,89	4,12	7,69	5,91	0,91	47,33	33,14
	19.03.95	15.06.95	# 10-13	spring		88		3,50	0,22	0,24	0,042	1,59	1,20	6,29	6,86	1,20	45,43	34,29
	15.06.95	11.09.95	# 14-17	summer		88		0,24	0,00	0,00	0,001	0,00	0,00	0,00	0,00	0,42	0,00	0,00
CB-7 lower	20.11.95	19.12.95	#1	fall		29		2,91	0,18	0,12	0,015	1,22	1,26	6,26	4,26	0,52	42,06	43,33
	19.12.95	16.03.96	#2-5	winter	1996	88		8,02	0,37	0,34	0,044	3,80	3,16	4,55	4,28	0,55	47,40	39,48
	16.03.96	12.06.96	#6-9	spring		88		9,55	0,63	0,61	0,080	4,76	2,94	6,58	6,38	0,84	49,83	30,82
	12.06.96	30.09.96	#10-14	summer		110		7,44	0,20	0,29	0,036	4,72	1,95	2,66	3,90	0,48	63,39	26,15
	30.09.96	27.12.96	#15-18	fall		88		8,59	0,38	0,40	0,049	3,73	3,69	4,40	4,64	0,57	43,38	42,91

Table 2. continued

CB meso	interval		sample no. of trap	season	year	duration days	remark	TTL mass	BSi ^{bio-epal}	org. carbon	nitrogen	carbonate	lithogenic	BSi	org. carbon	nitrogen	carb.	lith.
	start	end						g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	%	%	%	%	%
CB-7/8 lower	27.12.96	20.03.97	#19-20 + #1-2	winter	1997	82		14,24	0,78	0,77	0,097	5,37	6,55	5,46	5,41	0,68	37,70	46,00
CB-8 upper	20.03.97	20.06.97	# 3-6	spring		98		17,72	0,62	1,05	0,131	9,69	5,30	3,50	5,94	0,74	54,68	29,92
	20.06.97	02.10.97	# 7-10	summer		98		4,25	0,04	0,20	0,026	2,91	0,90	0,92	4,66	0,61	68,50	21,20
	02.10.97	14.12.97	# 11-13	fall		73,5		0,49	0,01	0,04	0,006	0,25	0,12	2,86	7,14	1,22	51,84	24,08
	14.12.97	22.03.98	# 14-17	winter	1998	98		1,68	0,05	0,15	0,024	0,84	0,45	3,21	8,87	1,43	50,12	26,49
	22.03.98	18.06.98	#18-20 + #1	spring		81	gap	1,57	0,01	0,06	0,008	1,21	0,20	0,45	3,70	0,51	76,80	12,62
CB-9 lower	18.06.98	09.09.98	# 2-4	summer		82,5		17,67	0,61	0,58	0,074	12,57	3,34	3,45	3,29	0,42	71,11	18,87
	09.09.98	28.12.98	# 5-8	fall		110		17,06	1,07	0,77	0,086	9,31	5,15	6,25	4,48	0,50	54,59	30,19
	28.12.98	20.03.99	# 9-11	winter	1999	82,5		16,33	1,19	0,62	0,073	7,18	6,71	7,27	3,81	0,45	43,99	41,11
	20.03.99	11.06.99	# 12-14	spring		82,5		19,55	1,08	0,65	0,083	11,77	5,40	5,53	3,33	0,42	60,17	27,61
	11.06.99	29.09.99	# 15-18	summer		110		16,88	0,51	0,57	0,068	11,80	3,43	3,02	3,35	0,40	69,92	20,35
CB-9/10 lower	29.09.99	16.12.99	#19-20+ #1-2	fall		75	gap	2,20	0,09	0,09	0,010	1,11	0,75	4,26	3,90	0,45	50,36	34,07
	16.12.99	21.03.00	#3	winter	2000	94		8,92	0,14	0,45	0,076	7,26	0,93	1,59	5,04	0,85	81,39	10,46
	21.03.00	21.06.00	#3	spring		92		8,74	0,14	0,44	0,075	7,11	0,91	1,59	5,05	0,86	81,33	10,45
	21.06.00	21.09.00	#3	summer		92		8,74	0,14	0,44	0,075	7,11	0,91	1,59	5,05	0,86	81,33	10,45
CB-11 upper	11.10.00	18.12.00	#3 + # 1-8	fall		87		8,32	0,41	0,56	0,087	5,13	1,68	4,93	6,73	1,05	61,66	20,19
	18.12.00	22.03.01	#9-19	winter	2001	93,5		6,51	0,39	0,60	0,093	3,57	1,34	6,01	9,25	1,43	54,84	20,58
CB-12 lower	05.04.01	27.06.01	# 1-4	spring		83		6,50	0,32	0,25	0,034	2,91	2,76	4,92	3,85	0,52	44,77	42,46
	27.06.01	01.10.01	# 5-9	summer		96,25		12,49	1,03	0,63	0,091	6,47	3,75	8,25	5,04	0,73	51,80	30,02
	01.10.01	17.12.01	# 10-13	fall		77		7,90	0,53	0,43	0,050	3,72	2,79	6,71	5,44	0,63	47,09	35,32
	17.12.01	21.03.02	# 14 sammel	winter	2002	94,25		0,88	0,05	0,04	0,009	0,75	0,02	5,68	4,20	1,02	85,23	2,27
CB-13 lower	23.04.02	19.06.02	#1-3	spring		57		6,03	0,27	0,23	0,029	3,53	1,78	4,46	3,78	0,48	58,42	29,55
	19.06.02	22.09.02	#4-8	summer		95		23,10	1,03	0,62	0,085	16,85	3,98	4,44	2,69	0,37	72,94	17,23
	22.09.02	26.12.02	#9-13	fall		95		9,51	0,42	0,32	0,042	5,53	2,92	4,42	3,36	0,44	58,16	30,66
	26.12.02	31.03.03	#14-18	winter	2003	95		11,41	0,55	0,35	0,050	6,78	3,39	4,78	3,07	0,44	59,37	29,72
	31.03.03	08.05.03	#19-20	spring		38		7,71	0,69	0,27	0,036	3,68	2,79	8,92	3,54	0,47	47,73	36,23

Table 2. continued

CB meso	interval		sample no. of trap	season	year	duration days	remark	TTL mass	BSi	org. carbon	nitrogen	carbonate	lithogenic	BSi	org. carbon	nitrogen	carb.	lith.
	start	end						g m ⁻²	%	%	%	%	%					
CB-14 upper	15.06.03	16.09.03	#2-7	summer		93		11,35	1,26	0,83	0,104	5,77	2,67	11,06	7,32	0,92	50,80	23,52
	16.09.03	18.12.03	#8-13	fall		93		8,28	0,84	0,45	0,061	3,99	2,56	10,16	5,48	0,74	48,14	30,87
	18.12.03	20.03.04	#14-19	winter	2004	93		0,58	0,03	0,03	0,005	0,29	0,16	5,39	5,22	0,87	49,91	27,48
CB-15 lower	17.04.04	25.06.04	#1-3	spring		69		12,49	0,66	0,45	0,059	7,58	3,36	5,30	3,58	0,47	60,64	26,90
	25.06.04	25.09.04	#4-7	summer		92		15,21	0,43	0,39	0,053	10,75	3,25	2,80	2,54	0,35	70,64	21,34
	25.09.04	26.12.04	#8-11	fall		92		8,34	0,48	0,36	0,043	4,22	2,93	5,72	4,25	0,52	50,60	35,14
	26.12.04	28.03.05	#12-15	winter	2005	92		23,56	1,69	1,12	0,152	12,18	7,44	7,18	4,76	0,65	51,69	31,59
CB-16 lower	28.03.05	28.06.05	#16-19	spring		92		7,72	0,24	0,28	0,041	5,00	1,93	3,04	3,65	0,53	64,72	24,94
	28.06.05	27.09.05	#20 + 1-3	summer		87,5	gap	18,23	1,12	0,63	0,078	10,46	5,40	6,13	3,43	0,43	57,38	29,62
	27.09.05	22.12.05	#4-7	fall		86		15,87	1,19	0,63	0,074	6,98	6,45	7,51	3,94	0,47	43,97	40,63
	22.12.05	18.03.06	#8-11	winter	2006	86		14,90	0,72	0,46	0,056	8,71	4,54	4,82	3,11	0,38	58,46	30,48
	18.03.06	12.06.06	#12-15	spring		86		15,16	0,92	0,66	0,085	9,04	3,87	6,09	4,37	0,56	59,64	25,51
CB-17 lower	12.06.06	28.09.06	#16-20	summer		107,5		6,07	0,45	0,24	0,031	3,55	1,58	7,38	3,97	0,51	58,51	26,11
	24.10.06	23.12.06	#1-8	fall		60		4,34	0,14	0,14	0,021	2,81	1,09	3,28	3,30	0,48	64,87	25,24
	23.12.06	23.03.07	#9-20	winter	2007	90		19,89	1,00	0,84	0,112	12,42	4,78	5,03	4,23	0,56	62,47	24,03
CB-18 lower	25.03.07	25.06.07	#1-5	spring		92		11,22	0,38	0,48	0,061	7,05	2,83	3,40	4,24	0,54	62,87	25,24
	25.06.07	28.09.07	#6-10	summer		95		8,57	0,29	0,33	0,040	4,79	2,83	3,43	3,83	0,47	55,94	32,96
	28.09.07	13.12.07	#11-14	fall		76		7,19	0,39	0,28	0,033	4,05	2,20	5,38	3,87	0,46	56,31	30,56
	13.12.07	17.03.08	#15-19	winter	2008	95		10,58	0,64	0,50	0,061	5,43	3,51	6,03	4,69	0,58	51,37	33,22
CB-19 lower	17.03.08	23.06.08	#20 + 1-4	spring		81	gap	5,49	0,24	0,22	0,029	4,04	0,76	4,43	4,03	0,53	73,67	13,92
	23.06.08	16.09.08	#5-9	summer		85		12,59	0,82	0,63	0,072	8,95	1,58	6,51	4,99	0,57	71,12	12,58
	16.09.08	27.12.08	#10-15	fall		102		9,01	0,47	0,44	0,045	4,64	3,03	5,17	4,87	0,50	51,45	33,60
	27.12.08	22.03.09	#16-20	winter	2009	85		9,51	0,63	0,42	0,050	6,56	1,47	6,60	4,44	0,53	69,04	15,47

938

Table 2.continued

CB meso	interval		sample no. of trap	season	year	duration days	remark	TTL mass	BSi biog. opal	org. carbon	nitrogen	carbonate	lithogenic	BSi	org. carbon	nitrogen	carb.	lith.
	start	end						g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	g m ⁻²	%	%	%	%	%
CB-20 upper	03.04.09	30.06.09	#1-5	spring		88		9,74	0,23	0,44	0,060	8,63	0,07	2,36	4,56	0,62	88,59	0,67
	30.06.09	28.09.09	#6-10	summer		90		3,25	0,09	0,16	0,021	2,63	0,21	2,74	4,95	0,65	80,90	6,43
	28.09.09	21.12.11	#11-(15)	fall		84		0,26	0,01	0,02	0,002	0,16	0,06	2,31	6,92	0,77	61,54	22,31
	21.12.11	26.02.10	#(11)- 19	winter	2010	67,5		18,77	0,66	0,95	0,086	10,78	5,42	3,54	5,07	0,46	57,45	28,85
CB-21 lower	20.03.10	28.06.10	#2-6	spring		100		7,34	0,24	0,31	0,047	4,78	1,33	3,27	4,26	0,64	65,12	18,12
	28.06.10	16.09.10	#7-10	summer		80		7,72	0,27	0,27	0,040	6,01	0,69	3,50	3,48	0,52	77,85	8,94
	16.09.10	25.12.10	#11-15	fall		100		9,81	0,26	0,50	0,041	6,00	2,55	2,65	5,13	0,42	61,16	25,99
	25.12.10	15.03.11	#16-19	winter	2011	80		4,94	0,20	0,20	0,029	3,44	0,89	4,05	4,13	0,59	69,64	18,02
CB-21/22 lower	15.03.11	21.06.11	#20 + 1-3	spring		67	gap	4,90	0,18	0,21	0,028	3,93	0,46	3,61	4,23	0,57	80,22	9,39
	21.06.11	14.09.11	#4-8	summer		85		10,45	0,28	0,54	0,048	7,86	1,23	2,63	5,16	0,46	75,22	11,77
	14.09.11	25.12.11	#9-14	fall		102		12,52	0,46	0,56	0,057	8,92	2,02	3,65	4,43	0,46	71,25	16,13
CB-22/23 lower	25.12.11	24.03.12	#15 +1+3	winter	2012	81,5 (90,5)	gap	17,91	0,87	0,60	0,086	10,08	5,74	4,86	3,35	0,48	56,28	32,05
	24.03.12	18.06.12	#4-7	spring		86		13,54	0,51	0,56	0,064	5,93	5,97	3,77	4,17	0,47	43,80	44,09
	18.06.12	12.09.12	#8-11	summer		86		12,90	0,27	0,31	0,046	8,67	3,35	2,09	2,37	0,36	67,21	25,97
	12.09.12	29.12.12	#12-16	fall		107,5		21,10	0,98	0,73	0,097	10,96	7,71	4,62	3,45	0,46	51,94	36,54

939

940

941


942 Table 3. Correlation coefficients between organic carbon flux and major bulk flux components for the four
 943 different seasons (lower trap data only). Number of data points (N) and the slopes (s) for the regression lines
 944 are given as well. Statistically significant values for R^2 at a 99.9% confidence level are indicated in bold.

organic carbon	<i>winter</i>	<i>spring</i>	<i>summer</i>	<i>fall</i>
nitrogen	0.96 N=16 s=0.13	0.93 N=19 s=0.12	0.92 N=19 0.12	0.93 N=18 0.11
BSi	0.70 N=16 s=1.3	0.46 N=19 s=1.7	0.63 N=19 s=1.4	0.75 N=18 s=1.4
carbonate	0.56 N=15 s=8.9	0.56 N=19 s=10.9	0.16 N=19 s=6.0	0.82 N=18 s=13.2
lithogenic (=mineral dust)	0.63 N=16 s=6.9	0.53 N=19 s=8.5	0.43 N=19 s=5.6	0.67 N=18 s=8.6


945
 946 Table 4. Summary of important flux changes between 1988 and 2012 ~~at site CB_{meso}~~ which are related to large
 947 scale climate modes such as NAO and ENSO. The record is divided into six major periods, including the
 948 outstanding year 2005 ([see text](#)).

Period/years	1988-1991	1997-1999 El Niño - La Niña	2001-2005/6	2005	2007-2010	2010-2012
FORCING:						
NAO	decreasing	increasing	negative or neutral	neutral	decreasing	increasing
ENSO		strongest ENSO	weak ENSOs	neutral		
FLUX RESPONSE:						
BSi	decreasing	first decreasing, then increasing	episodic peaks	High throughout , except spring	decreasing	increasing
Carbonate	decreasing	generally high, pteropod peaks	major episodic pteropod peaks			
Lithogenic (dust)	decreasing	first decreasing, then increasing		high throughout , except spring	decreasing	increasing

949




950
951
952 Fig. 1
953



954
955
956 Fig. 2.
957
958

959




960

961 Fig. 3.

962

963




964

965


966 Fig. 4.

967

CB-1 to CB-23
upper and lower traps



968

969 Fig. 5.



970
971
972
973


Fig. 6.



974
975
976
977

Fig. 7.

978




979

980

981 Fig. 8.


982



983
984
985
986

Fig. 9.

987



988

989

990 Fig. 10.