

*Biogeosciences Discussions* is the access reviewed discussion forum of *Biogeosciences*

# Imminent ocean acidification projected with the NCAR global coupled carbon cycle-climate model

M. Steinacher<sup>1</sup>, F. Joos<sup>1,2</sup>, T. L. Frölicher<sup>1</sup>, G.-K. Plattner<sup>1,\*</sup>, and S. C. Doney<sup>3</sup>

<sup>1</sup>Climate and Environmental Physics, Physics Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

<sup>2</sup>Oeschger Centre for Climate Change Research, University of Bern, Erlachstrasse 9a, 3012 Bern, Switzerland

<sup>3</sup>Dept. of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA 02543-1543, USA

\*now at: Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, Universitätstrasse 16, 8092 Zürich, Switzerland

Received: 22 September 2008 – Accepted: 23 September 2008

– Published: 14 November 2008

Correspondence to: M. Steinacher (steinacher@climate.unibe.ch)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Ocean acidification from the uptake of anthropogenic carbon is simulated for the industrial period and IPCC SRES emission scenarios A2 and B1 with a global coupled carbon cycle-climate model. Earlier studies identified seawater saturation state with respect to aragonite, a mineral phase of calcium carbonate, as a key variable governing impacts on corals and other shell-forming organisms. Globally in the A2 scenario, water saturated by more than 300%, considered suitable for coral growth, vanishes by 2070 AD ( $\text{CO}_2 \approx 630$  ppm), and the ocean volume fraction occupied by saturated water decreases from 42% to 25% over this century. The largest simulated pH changes worldwide occur in Arctic surface waters, where hydrogen ion concentration increases by up to 185%. Projected climate change amplifies the decrease in Arctic surface mean saturation and pH by more than 20%, mainly due to freshening and increased carbon uptake in response to sea ice retreat. Modeled saturation compares well with observation-based estimates along an Arctic transect and simulated changes have been corrected for remaining model-data differences in this region. Aragonite undersaturation in Arctic surface waters is projected to occur locally soon and to become more widespread as atmospheric  $\text{CO}_2$  continues to grow. The results imply that surface waters in the Arctic Ocean will become corrosive to aragonite, with potentially large implications for the marine ecosystem, if anthropogenic carbon emissions are not reduced and atmospheric  $\text{CO}_2$  not kept below 450 ppm.

## 1 Introduction

The continued emissions of  $\text{CO}_2$  by human activities causes atmospheric  $\text{CO}_2$  to rise, climate to warm, and the ocean to become less alkaline (e.g. Feely et al., 2008; German Advisory Council on Global Change, 2006; Royal Society, 2005; Orr et al., 2005; Caldeira and Wickett, 2003). Today's atmospheric  $\text{CO}_2$  concentration of 384 ppm is far above the natural range reconstructed for the last 800 000 years (172 to 300 ppm; Lüthi

**BGD**

5, 4353–4393, 2008

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2008; Siegenthaler et al., 2005) and probably the last 20 million years (Prentice et al., 2001). An important aspect of the human perturbation is its rate of change, as it codetermines the impacts of global warming and elevated CO<sub>2</sub> on natural and socioeconomic systems and their capabilities to adapt. The 20th century increase in CO<sub>2</sub> occurred more than an order of magnitude faster than any sustained change during at least the past 22 000 years (Joos and Spahni, 2008). Carbon cycle and climate change projections based on the latest set of mitigation and reference scenarios from the integrated assessment modelling community indicate that CO<sub>2</sub> and global mean surface temperature will continue to rise over the 21st century and decadal-scale rates of change in temperature and CO<sub>2</sub> will remain high for several decades even for stringent mitigation scenarios (Van Vuuren et al., 2008; Strassmann et al., 2008).

The purpose of this paper is to investigate the consequences of rising atmospheric CO<sub>2</sub> and climate change on ocean chemistry and the ocean's acid-base state with the Climate System Model CSM1.4-carbon of the U.S. National Center for Atmospheric Research (NCAR). We investigate how the saturation state of seawater with respect to aragonite, a mineral phase of calcium carbonate (CaCO<sub>3</sub>), will evolve over this century under two IPCC non-intervention greenhouse gas emission scenarios. A particular emphasis is on the Arctic Ocean, a region that was not analyzed for saturation state changes in a recent assessment (Orr et al., 2005), but where large changes in CaCO<sub>3</sub> saturation (Gangstø et al., 2008; Steinacher, 2007; McNeil and Matear, 2007), fresh-water balance, and sea ice (Stroeve et al., 2007; Holland et al., 2006) are expected for the coming decades. We also analyze how the ocean volume fraction of different saturation regimes is changing with increasing CO<sub>2</sub> and quantify seasonal and interannual variability worldwide.

A particular concern of rising atmospheric CO<sub>2</sub> is the impact on marine organisms and marine ecosystems. Recent work indicates that ocean acidification from the uptake of anthropogenic CO<sub>2</sub> has adverse consequences for many marine organisms via decreased CaCO<sub>3</sub> saturation, affecting calcification rates, and via disturbance to acid-base physiology (Fabry et al., 2008; Hoegh-Guldberg, 2005; Langdon, 2002). Particu-

**Imminent ocean acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



larly vulnerable appear organisms that build shells or other structures of calcium carbonate ( $\text{CaCO}_3$ ) in the relatively soluble mineral forms of aragonite or high-magnesium  $\text{CaCO}_3$ . Examples are pteropods (Orr et al., 2005), a major zooplankton group in high latitudes, corals (Kleypas et al., 2006; Langdon and Atkinson, 2005), and sea urchins (Fabry et al., 2008). Experimental evidence suggests a range of responses to ocean acidification (Fabry et al., 2008, Table 1) such as a reduction in foraminifera shell mass, shell dissolution for pteropods in water undersaturated with respect to aragonite, reduced calcification rates, reduced growth rates, reduced metabolisms, and increased mortality in molluscs or reduced fertilization for a number of groups. Responses to acidification are likely to vary across species and life stage. Scleractinian corals were found to survive and recover after experiencing decalcification in acidic water (Fine and Tchernov, 2007). Currently experimental evidence is however limited with regard to the number of species tested at relevant  $\text{CO}_2$  concentrations.

Uptake of anthropogenic carbon by the ocean changes the acid-base state of seawater. Currently, about a quarter of the anthropogenic  $\text{CO}_2$  emitted into the atmosphere is taken up by the ocean. The hydrolysis of  $\text{CO}_2$  in seawater lowers ocean pH, making the oceans less alkaline (Caldeira and Wickett, 2003). Carbonate ion concentration (Feely et al., 2004) and the saturation state of water with respect to calcium carbonate ( $\text{CaCO}_3$ ) is decreasing through the reaction:



The  $\text{CaCO}_3$  saturation state,  $\Omega$ , is the product of the concentrations of calcium ions ( $\text{Ca}^{2+}$ ) and carbonate ions ( $\text{CO}_3^{2-}$ ) divided by the apparent stoichiometric solubility product  $K'_{\text{sp}}$ :

$$\Omega = [\text{Ca}^{2+}][\text{CO}_3^{2-}]/K'_{\text{sp}} \quad (2)$$

The solubility product ( $K'_{\text{sp}} = [\text{Ca}^{2+}]_{\text{sat}}[\text{CO}_3^{2-}]_{\text{sat}}$ ) is either defined for aragonite or calcite, two mineral phases of  $\text{CaCO}_3$  secreted by marine organisms. The  $\text{CO}_3^{2-}$  saturation concentration is about 50% higher for the mineral phase calcite ( $\Omega_{\text{calc}}$ ) than for the

**Imminent ocean acidification**

M. Steinacher et al.

less stable aragonite ( $\Omega_{\text{arag}}$ ). The concentration of  $\text{Ca}^{2+}$  is linearly related to salinity and nearly constant. It is thus convenient to express the saturation state as deviation of the  $\text{CO}_3^{2-}$  concentration from its saturation concentration:  $\Delta[\text{CO}_3^{2-}] = [\text{CO}_3^{2-}] - [\text{CO}_3^{2-}]_{\text{sat}}$ . Undersaturated seawater ( $\Omega < 1$  or  $\Delta[\text{CO}_3^{2-}] < 0$ ) is corrosive to  $\text{CaCO}_3$  in the absence of protective mechanisms.

The  $\text{CaCO}_3$  saturation state  $\Omega_{\text{arag}}$  is a key variable for assessing biological impacts of ocean acidification. Available experimental and observational evidence suggests that the biogenic production and dissolution of  $\text{CaCO}_3$  is mainly controlled by the ambient saturation state, rather than by pH or other factors. This has been shown for corals, coralline algae, coccolithophorids, foraminifera, echinoderms, mesocosm coral reef communities and natural coral reef ecosystems (Kleypas et al., 2006; Schneider and Erez, 2006; Hoegh-Guldberg, 2005; Orr et al., 2005; Gattuso et al., 1998). Tropical coral reef development was associated with  $\Omega_{\text{arag}} \geq 4.0$  by Kleypas et al. (1999), and Langdon and Atkinson (2005) project a 40–50% decrease of coral calcification rates by 2065 ( $\Omega_{\text{arag}} = 3.1$ ) relative to  $\Omega_{\text{arag}} = 4.6$ , which corresponds to preindustrial conditions (year 1880) in the tropical ocean in their study.

A range of responses is found for pelagic calcifiers under changing  $\Omega_{\text{arag}}$  (Fabry, 2008). Recently Iglesias-Rodriguez et al. (2008) reported laboratory evidence that calcification and net primary production per cell in the coccolithophore species *Emiliani huxleyi* are significantly increased by high  $\text{CO}_2$  partial pressures. Yet the growth rates were found to be significantly lower at higher  $\text{CO}_2$  levels. Given that coccolithophores are a major contributor to the open-ocean carbonate pump, this could have important implications for the overall community production of  $\text{CaCO}_3$  relative to organic carbon, under increasing  $\text{pCO}_2$ . However, the implied feedback on atmospheric  $\text{CO}_2$  has been estimated to be small in global modelling studies (Heinze, 2004).

The main controls on 21st century  $\Omega_{\text{arag}}$  are the uptake of anthropogenic carbon and the subsequent increase in dissolved inorganic carbon (DIC) (Orr et al., 2005). Second order changes in saturation state arises from alterations in the distribution of DIC and alkalinity (Alk) due to a reorganization of the marine biological cycle and

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ocean circulation as well as ocean warming (decreasing CO<sub>2</sub> solubility) and changes in the freshwater fluxes. Changes in sea ice affect  $\Omega_{\text{arag}}$  by changing the air-sea gas exchange as well as the freshwater balance. The direct influence of temperature and salinity changes on the CaCO<sub>3</sub> solubility product ( $K'_{\text{sp}}$ ) is negligible.

5 Large changes in  $\Omega_{\text{arag}}$  and pH have been projected using ocean-only models (Kley-  
pas et al., 1999; Caldeira and Wickett, 2003; Gangstø et al., 2008). Recently, the  
impact of climate change on ocean acidification also has been considered (Cao et al.,  
2007; McNeil and Matear, 2007; Orr et al., 2005). In most regions, a relatively small  
negative climate change feedback is found on changes in  $\Omega_{\text{arag}}$  ( $\approx 10\%$ ). An exception  
10 is the Arctic, where larger changes in  $\Omega_{\text{arag}}$  are found for a simulation including climate  
change compared to one without climate change (McNeil and Matear, 2007).

Here, we complement earlier studies by using the fully coupled NCAR CSM1.4-  
carbon model to investigate the evolution of  $\Omega_{\text{arag}}$  over the 21st century for the SRES  
A2 and B1 scenarios. A particular emphasis is on the Arctic Ocean, a region for which  
15 changes in saturation state have not been discussed in detail. Recently, oceanographic  
data have become available to quantify the current saturation state in the Arctic (Jut-  
terström and Anderson, 2005) and that provide now a basis for our modeling efforts.  
In the next section, the model and experimental details are described. Then, model  
results are compared with observations, providing also statistical measures of data-  
20 model agreement. In the results section, projected global and regional changes in sat-  
uration state are described and climate feedbacks quantified. The change in Arctic sat-  
uration state is quantified both from direct model results and by combining observation-  
based estimates of the current saturation state (Jutterström and Anderson, 2005) with  
modeled changes following a similar approach as Orr et al. (2005). Before concluding,  
25 the projected anthropogenic changes are compared to modeled natural seasonal and  
interannual variations.

**Imminent ocean  
acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Methods

### 2.1 The NCAR CSM1.4-carbon model

Simulations were performed with the coupled climate-carbon cycle model NCAR CSM1.4-carbon (Doney et al., 2006; Fung et al., 2005). The core of the model used in this study is a modified version of the NCAR CSM1.4 coupled physical model, consisting of ocean, atmosphere, land and sea ice physical components integrated via a flux coupler without flux adjustments (Boville and Gent, 1998). The atmospheric model CCM3 is run with a spectral truncation resolution of  $\approx 3.75^\circ$  (T31 grid) and 18 levels in the vertical (Kiehl et al., 1998). The ocean model is the NCAR CSM Ocean Model (NCOM) with 25 levels in the vertical and a resolution of  $3.6^\circ$  in longitude and  $0.8^\circ$  to  $1.8^\circ$  in latitude (T31 $\times$ 3 grid) (Gent et al., 1998). The sea ice component (CSIM) simulates the dynamic and thermodynamic evolution of ice concentration, thickness, temperature, velocity, and snow cover in response to forcing by the atmosphere and ocean (Bettge et al., 1996). The model is known to simulate too much ice in the Northern Hemisphere mainly due to too extensive ice cover in the North Pacific and North Atlantic. There is some contribution to the excess ice volume in the Arctic Ocean through larger-than-observed summer sea ice concentrations. Further, maximum thickness in the Arctic occurs against the Bering Strait, not against the Canadian Archipelago as observed (Weatherly et al., 1998).

The CSM1.4-carbon model combines the NCAR Land Surface Model (LSM) (Bonan, 1996) with the Carnegie-Ames-Stanford Approach (CASA) biogeochemical model (Randerson et al., 1997) and a derivative of the OCMIP-2 biotic carbon model (Najjar et al., 2007), where the nutrient uptake has been changed from a restoring to a prognostic formulation and an iron cycle has been added (Doney et al., 2006). A constant export ratio of  $\text{CaCO}_3$  to organic carbon of 0.07 and constant Redfield ratios were used (Anderson and Sarmiento, 1994). There is no feedback between changes in saturation state, pH or other carbonate chemistry variables and the biological carbon cycle implemented.

**BGD**

5, 4353–4393, 2008

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean  
acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A sequential spin-up procedure including 1000 years of fully coupled simulation under preindustrial conditions was employed by Doney et al. (2006). Starting from this nearly steady state, four simulations over the period 1820 AD to 2100 AD were performed: (1) a 280 year long control simulation, two simulations with historical emissions over the industrial period followed by the (2) A2 IPCC emission scenario and (3) B1 IPCC emission scenario, and (4) a SRES A2 “no-warming case” simulation, where the concentrations of greenhouse gases and other radiative agents were kept at their preindustrial values in the radiation module of the model. Differences in results between this no-warming simulation and the standard SRES A2 simulation are taken to reflect the impact of climate change on carbonate chemistry and other parameters.

The model was driven by emissions of CO<sub>2</sub> from fossil fuel and land use change (Fig. 1) as well as by anthropogenic sulfur emissions. Concentrations of the non-CO<sub>2</sub> greenhouse gases CH<sub>4</sub>, N<sub>2</sub>O, CFC-11, CFC-12 are prescribed according to reconstructions for 1820 to 2000 AD and as projected from emissions for 2000 to 2100 AD. The CFC-11 concentration is scaled to include the forcing of other halogenated species and SF<sub>6</sub>. Emissions of non-CO<sub>2</sub> greenhouse gases were translated to concentration using formulations by Forster et al. (2007). A recurring annual cycle of ozone and natural sulfate aerosol were also included. Over the historical period, solar irradiance changes were prescribed following Wang et al. (2005) and spatially-explicit sulfate aerosols from explosive volcanic eruptions after Ammann et al. (2007). After year 2000, the solar irradiance was kept constant and no volcanic eruptions were assumed. Further details are given by Frölicher et al. (submitted).

## 2.2 Carbonate chemistry and tracer data

For model evaluation, the gridded data from the Global Data Analysis Project (GLODAP) (Key et al., 2004) for DIC and Alk, and from the World Ocean Atlas 2001 (WOA01) (Conkright et al., 2002) for temperature (*T*), salinity (*S*), phosphate (PO<sub>4</sub>) and silicate (SiO<sub>4</sub>) were used. The two data sets were regridded by multi-axis linear interpolation onto the model grid. The gridded GLODAP data have annual resolution and do not

include the Arctic.

Carbonate chemistry, pH, carbonate ion concentration, and the saturation state have been calculated offline from modeled or observation-based quantities using the standard OCMIP carbonate chemistry routines (Orr et al., 2008). In these routines, based on work by Dickson (2002) and Millero (1995), the total pH scale (Lueker et al., 2000) is utilized. The observation-based carbonate variables were computed for each month, which corresponds to the time resolution of the model output. Due to lack of monthly data, annual means of Alk and DIC from GLODAP have been used. Also, annual means of silicate and phosphate from WOA01 at depths below 500 m, as well as seasonal means of temperature and salinity from WOA01 at depths below 1500 m have been used. From the monthly model output, the carbonate variables were computed for each month from 1990 to 1999. Annual averages from the monthly data are then used for data-model comparison. Because the CSM model does not include  $\text{SiO}_4$ , a seasonal cycle of observation-based  $[\text{SiO}_4]$  from WOA01 has been used in all calculations of simulated pH and calcium carbonate saturation state.

For the Arctic, observation-based estimates for  $\Omega_{\text{arag}}$  and related variables were calculated from  $T$ ,  $S$ , Alk, DIC,  $\text{PO}_4$ , and  $\text{SiO}_4$  measurements using the OCMIP chemistry routines and data from the Arctic Ocean Expedition 1991 (ODEN-91, cruise 77DN1991072), Arctic Ocean Section 1994 (AOS-94, cruise 18SN19940726), and the Arctic Climate System Study 1996 (ARCSYS-96, cruise 06AQ19960712) (Jutterström and Anderson, 2005). These data are available from the preliminary CARINA database ([http://cdiac.ornl.gov/oceans/CARINA/Carina\\_table.html](http://cdiac.ornl.gov/oceans/CARINA/Carina_table.html)). Where not referring to this specific section, the Arctic Ocean is defined by the basin index of the CSM1.4 model ( $\text{rmask}=10$ ), i.e. all waters north of  $65^\circ \text{N}$ , except the Labrador Sea and the Greenland, Iceland, and Norwegian (GIN) Seas ( $<80^\circ \text{N}$  and  $35^\circ \text{W}-18^\circ \text{E}$ ).

**BGD**

5, 4353–4393, 2008

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3 Results

#### 3.1 Comparison of modeled aragonite saturation and $\text{CO}_3^{2-}$ concentration with observation-based estimates

The modeled values and spatial pattern for  $\Omega_{\text{arag}}$  and  $[\text{CO}_3^{2-}]$  (averaged over the years 1990–1999 in the transient simulation) correspond reasonably well with the data-based estimates in all major ocean basins (Fig. 2). The correlation coefficient  $r$  between modeled and data-based  $\Omega_{\text{arag}}$  is between 0.90 and 0.95 for the Atlantic, Pacific, and Global Ocean. The model represents the amplitude of variations in  $\Omega_{\text{arag}}$  well in all major basins, as indicated by a relative standard deviation close to unity in the Taylor diagram (Fig. 2). Correlation coefficients are somewhat smaller (0.86 to 0.91) for  $[\text{CO}_3^{2-}]$  than for  $\Omega_{\text{arag}}$ . Model biases in  $[\text{CO}_3^{2-}]$  are particularly large in the Pacific where the relative standard deviation is smaller than indicated by observations.

The present distribution of  $\Delta[\text{CO}_3^{2-}]$  (Fig. 3) and  $\Omega_{\text{arag}}$  is governed by the distribution of DIC and Alk. Surface  $[\text{CO}_3^{2-}]$  is approximately proportional to the difference Alk–DIC. Consequently, the nutrient and carbon rich water of the North Pacific thermocline is undersaturated with respect to aragonite, while the relatively nutrient and carbon poor water of the North Atlantic thermocline is oversaturated. The increase of the solubility product  $K'_{\text{sp}}$  with pressure (depth) leads to a decrease of the saturation state in the deep ocean, although the carbonate ion concentration is relatively homogeneous below 1000 m.

Deviations in modeled and observation-based zonal mean  $[\text{CO}_3^{2-}]$  and  $\Delta[\text{CO}_3^{2-}]$  are less than  $20 \mu\text{mol/l}$  in the deep Atlantic, the Southern Ocean, and the southern Pacific (Fig. 3). Larger deviations are found in the North Pacific and in the thermocline of the tropical Atlantic.  $[\text{CO}_3^{2-}]$  is underestimated in the tropical Atlantic related to too high DIC concentration in the model caused by nutrient trapping. On the other hand, the model underestimates export and remineralisation of organic matter in the North Pacific, causing too low DIC concentration and too high  $[\text{CO}_3^{2-}]$  in the thermocline and at

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



depth (Frölicher et al., submitted). Correspondingly, the modeled aragonite saturation horizon in the North Pacific is too deep compared with observation-based estimates.

In the Arctic Ocean, the modeled distribution of  $\Omega_{\text{arag}}$  corresponds reasonably well to the observation-based estimates (years 1991–1996; Fig. 4). The observations suggest an oversaturation of around 50% in the top five hundred meter, around 25% at 1000 m depth and undersaturation below 2000 m. Horizontal gradients are relatively small along the transect. The model tends to underestimate  $\Omega_{\text{arag}}$  at the surface (up to 0.3), while below 500 m  $\Omega_{\text{arag}}$  is overestimated by 0.2 to 0.4. Correspondingly, the simulated saturation horizon is deeper than the observation-based horizon.

### 3.2 Projected global mean changes

Figure 1 provides projected global mean changes for a few key variables that broadly characterize the investigated scenarios. Anthropogenic carbon emissions increase to almost 30 GtC yr<sup>-1</sup> by the end of the century for the high-emission SRES A2, whereas emissions peak at around 12 GtC yr<sup>-1</sup> and decline thereafter in the more optimistic B1 scenario. Atmospheric CO<sub>2</sub> reaches 840 ppm and 538 ppm by the end of the century for the A2 and B1 scenario, respectively. Ocean surface DIC content increases on average by 8% (0.17 mmol kg<sup>-1</sup>) and by 5% (0.10 mmol kg<sup>-1</sup>). Mean aragonite saturation of surface water decreases from a preindustrial  $\Omega_{\text{arag}}$  value of 3.4 to 1.8 (A2) and to 2.3 (B1) by the end of the century. Concomitant reductions in pH are 0.40 and 0.23 units. Global mean surface temperature increase is 1.97 and 0.94°C, reflecting the low climate sensitivity of the CSM1.4 (Friedlingstein et al., 2006).

Simulated climate change feedbacks on  $\Omega_{\text{arag}}$  and pH are relatively small on the global scale. The CO<sub>2</sub> rise is slightly smaller in the no-warming case than in the standard SRES A2 simulation (Fig. 1), consistent with the low climate-carbon cycle feedback of the NCAR CSM1.4 (Fung et al., 2005). The increase in ocean surface DIC by the end of the century is about 8% larger without climate-feedback and consequently the decrease in surface  $\Omega_{\text{arag}}$  is also slightly enhanced (by about 4%). In contrast, the decrease in pH is about 4% smaller in the no-warming simulation as found earlier (Cao

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2007).

Changes in DIC,  $\Omega_{\text{arag}}$ , and pH are not linearly related to the atmospheric CO<sub>2</sub> increase (Thomas et al., 2007). The Revelle factor defined as the percentage change in CO<sub>2</sub> partial pressure divided by the percentage change in DIC increases with increasing CO<sub>2</sub>. Doubling the atmospheric CO<sub>2</sub> from 280 to 560 ppm causes global mean surface DIC to increase by 6% (11 mmol kg<sup>-1</sup>), whereas increasing CO<sub>2</sub> by a further 280 ppm to 840 ppm causes an additional DIC increase of only 3%. Therefore, a higher fraction of the emissions stays in the atmosphere for the high emission A2 scenario than for the low emission B1 scenario. On the other hand, surface mean aragonite saturation decreases by 1.1 unit for a doubling of CO<sub>2</sub>, but only by 0.5 units for the additional increase from 560 to 840 ppm. Similarly, pH decreases by 0.24 unit for the first 280 ppm increase and by 0.15 unit for the next 280 ppm.

### 3.3 Regional changes in saturation at the surface

It is projected that  $\Omega_{\text{arag}}$  of surface waters will continue to decrease rapidly in all regions and that high-latitude waters will become undersaturated (Figs. 5 and 6). Arctic surface waters are projected to become undersaturated with respect to aragonite within a few decades only, when following business as usual scenarios. The averaged surface annual mean  $\Omega_{\text{arag}}$  becomes less than unity in the model when atmospheric CO<sub>2</sub> exceeds 490 ppm (2040 AD in A2 and 2050 AD in B1). The Southern Ocean (65 to 80° S) becomes undersaturated on average about 20 years later in the A2 scenario, when atmospheric CO<sub>2</sub> concentration exceeds 580 ppm, consistent with Orr et al. (2005). By the end of the century, at 840 ppm, high-latitude surface waters are projected to be undersaturated poleward of about 50° for the A2 scenario.

The biggest absolute changes in  $\Omega_{\text{arag}}$  are found in tropical and subtropical surface waters. In the Tropics (30° N–30° S)  $\Omega_{\text{arag}}$  is projected to drop from 4.2 (1820) to 2.3 in A2. For atmospheric CO<sub>2</sub> higher than 585 ppm, surface waters will have a zonal mean  $\Omega_{\text{arag}}$  of less than three everywhere. These conditions are expected to have significant impacts on the ability of coral reef ecosystems to maintain their structures against the

---

**Imminent ocean acidification**

M. Steinacher et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



forces of erosion and dissolution (Kleypas et al., 2006).

Turning to the B1 scenario surface changes in  $\Omega_{\text{arag}}$  are similar as for the A2 scenario until 2050, as both scenarios follow very similar pathways of atmospheric  $\text{CO}_2$ . After 2050, the more gradual increase of atmospheric  $\text{CO}_2$  in B1 slows down the decrease in  $\Omega_{\text{arag}}$ . By 2100, undersaturation at the surface is limited to the Arctic Ocean and surface  $\Omega_{\text{arag}}$  will stabilize at 2.5 to 3.0 in tropical and subtropical regions. The Southern Ocean surface remains slightly supersaturated throughout the B1 simulation.

### 3.3.1 Changes at depth and in the volume of supersaturated waters

Saturation is decreasing not only at the surface but also in the thermocline and the deep ocean as anthropogenic carbon continues to invade the ocean. This is illustrated by Hovmöller diagrams of  $\Delta[\text{CO}_3^{2-}]$  for selected regions (Fig. 7). The thermocline changes in  $[\text{CO}_3^{2-}]$  are relatively modest over the past 200 years. However, the modeled  $\text{CO}_3^{2-}$  concentrations decrease rapidly over this century and the saturation horizon, separating over- and undersaturated waters, shoals in most regions.

For the A2 scenario and over this century, the anthropogenic perturbation is projected to penetrate the top 3500 m in the North Atlantic, leading to a shoaling of the saturation horizon from 3250 m to 1300 m in mid and high latitudes and an extension of the undersaturated watermasses in the low latitude Atlantic thermocline. In the Arctic and in the Southern Ocean supersaturation is already small today and the top 300 and 400 m will become undersaturated over the coming decades in the A2 scenario (Fig. 7). In the tropical Pacific the anthropogenic perturbation extends down to the depth of about 1000 m and the saturation horizon remains at 1500 m.

The anthropogenic perturbation causes a substantial loss of habitat for calcifying organisms. We introduce five classes of saturation levels: (i) more than 400% saturated ( $\Omega_{\text{arag}} > 4$ ) as considered optimal for coral growth, (ii) 300 to 400% saturated, considered as adequate for coral growth, (iii) 200 to 300% saturated and (iv) 100 to 200% saturated, both considered as marginal to inadequate for coral growth (Langdon and

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean  
acidification**

M. Steinacher et al.

Atkinson, 2005; Kleypas et al., 2006), and finally (v) undersaturated water ( $\Omega_{\text{arag}} < 1$ ) considered to be unsuitable for pteropods (Feely et al., 2004; Orr et al., 2005). The volume of water that is more than 400% saturated is projected to vanish in the next three decades for the A2 and B1 scenario (Fig. 8). Water with  $\Omega_{\text{arag}}$  larger than three is projected to vanish by 2070 ( $\text{CO}_2 \approx 630$  ppm) in the A2 scenario and to be reduced to 11% of the respective preindustrial volume in the B1 scenario. Further, the volumes of waters with  $2 < \Omega_{\text{arag}} < 3$  and  $1 < \Omega_{\text{arag}} < 2$  are projected to decrease by about 75% and 30% over the industrial period and this century for the A2 scenario; corresponding changes are about half as large for B1 than for A2.

Overall the volume occupied by water oversaturated with respect to aragonite decreases from about 42% to 25% of the total ocean volume. We note that the model correctly simulates the total volume occupied by the first three classes, whereas the volume with an aragonite saturation between 100 and 200% is overestimated (crosses in Fig. 8). Therefore, these projections might be somewhat too optimistic.

### 3.3.2 Changes in the Arctic Ocean and climate feedbacks

Our projection that the Arctic ocean will become undersaturated with respect to aragonite in a few decades calls for a more detailed analysis. We combine observation-based estimates of  $\Omega_{\text{arag}}$  with the modeled changes to obtain an alternative estimate of the saturation state in the Arctic in 2100:  $\Omega(t) = \Omega_{\text{obs.}}(t_0) + \Omega_{\text{model}}(t) - \Omega_{\text{model}}(t_0)$ . In this way,  $\Omega_{\text{arag}}$  is adjusted for known model biases similar to the approach of Orr et al. (2005), who applied simulated changes in DIC to observation-based estimates. The difference between adjusting  $\Omega$  directly and calculating  $\Omega$  from adjusted DIC and Alk concentrations is small ( $< 1\%$ ). In 2100 and for A2, the entire water column is projected to be undersaturated except some small regions at depths between 1000 m and 2000 m (Fig. 4). For comparison, the model-only projection also show a distinct undersaturation in the top few hundred meters ( $\Omega_{\text{arag}}$  between 0.25 and 0.75), whereas water remains oversaturated between 1000 and 3500 m in the model only projection as the modeled  $\text{CO}_3^{2-}$  concentration is biased high. In any case, the emerging undersatura-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tion of the surface Arctic ocean is a robust feature and independent of these model biases.

Undersaturation occurs first in small regions and over short periods of the year. Its spatial and seasonal occurrence increases with rising atmospheric CO<sub>2</sub> as summarized in Table 1 for the investigated Arctic transect and for the model-data approach. 10% of the surface area is projected to become undersaturated with respect to aragonite in at least one month of the year when atmospheric CO<sub>2</sub> has reached 409 ppm at year 2016 in the A2 and at year 2018 in the B1 scenario. At 428 ppm (2023 AD in A2, 2025 AD in B1) 10% of the surface is projected to become undersaturated in the annual mean, and at 450 ppm (2029 AD in A2, 2034 AD in B1) throughout the year. More than 50% of the surface waters of that section will become undersaturated at CO<sub>2</sub> levels of 534 ppm (2050 AD in A2, 2085 AD in B1), and at 765 ppm (2090 AD in A2) the entire water column is projected to become undersaturated.

The largest pH changes over the 21st century (−0.45 in A2, −0.23 in B1) are simulated in the Arctic Ocean. This turns the Arctic Ocean from a region with relatively high pH (0.06 units above the global average) to a region with very low pH (0.09 units below the global average by 2100 in A2). [H<sup>+</sup>] increases from 7.3 nmol/l to 20.8 nmol/l in A2 and to 12.35 nmol/l in B1, respectively.

Next the role of climate change and of different processes for the projected Arctic changes is addressed. Warming in the Arctic region is much higher than on global average (polar amplification) and summer sea ice cover is projected to be significantly reduced. The Arctic Ocean is projected to become fresher in response to sea ice melt, altered evaporation and precipitation, and river input. Small climate feedbacks on  $\Omega_{\text{arag}}$  and pH have been reported for the global ocean. However, climate change amplifies the decrease in average annual mean Arctic surface  $\Omega_{\text{arag}}$  and pH by 22% and 27%, respectively (Fig. 9). In the simulation without global warming, alkalinity remains roughly constant in the Arctic, while the dissolved inorganic carbon concentration increases as a result of anthropogenic carbon uptake. Both surface Alk and DIC decrease in the simulation with global warming (Fig. 10). Additional freshwater input into the Arctic Ocean

**Imminent ocean acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



dilutes the tracer concentrations. The simulated decrease in DIC is smaller than that for Alk, mainly because the uptake of anthropogenic carbon tends to increase DIC. Overall, the difference between Alk and DIC, governing the  $\text{CO}_3^{2-}$  concentration, decreases more in the standard global warming simulation than in the simulation without climate change, thereby amplifying the changes in saturation state.

We can quantitatively attribute the changes of Alk and DIC, and thus  $[\text{CO}_3^{2-}]$ , to different mechanisms. A budget of the Alk and DIC fluxes entering and leaving the Arctic surface ocean (top 200 m) is established (Fig. 11) for the A2 scenario. The changes in the difference between Alk and DIC is taken as a proxy for changes in the  $\text{CO}_3^{2-}$  concentration. This approximation works well in the Arctic surface ocean (inset of Fig. 11). We note that the regression slope is slightly different in the no-warming simulation ( $0.43$ ,  $r^2=0.94$ ) than in the standard global warming simulation ( $0.38$ ,  $r^2=0.90$ ). In the no-warming simulation, the freshwater balance, ocean circulation, the marine biological cycle and Alk remain mostly unperturbed. We note a small perturbation of the global hydrologic cycle, which arises most likely due to the interaction of increasing atmospheric  $\text{CO}_2$  with the terrestrial biosphere. However, the most relevant changes are the uptake of anthropogenic (excess) carbon by gas exchange through the air-sea interface and the export of excess carbon out of the top 200 m by surface-to-deep and lateral (to the North Atlantic and through the Bering Strait) exchange. The resulting storage of anthropogenic carbon in the top 200 m increases DIC and decreases (Alk-DIC) by  $0.10$  mmol/l by 2100. In the standard global warming simulation, additional freshwater input dilutes both Alk and DIC. The decrease in Alk and DIC by freshwater input is proportional to their concentration and thus larger for Alk than for DIC. As a result, (Alk-DIC) is reduced by  $0.035$  mmol/l on average. The dilution of DIC by freshwater lowers the  $\text{CO}_2$  partial pressure which in combination with an increase in ice-free area enhances uptake of anthropogenic carbon through gas exchange. Increased uptake is further promoted by the somewhat higher atmospheric  $\text{CO}_2$  concentration in the warming than in the no-warming simulation. On the other hand the dilution of Alk increases the  $\text{CO}_2$  partial pressure, which tends to decrease the uptake of anthropogenic

**Imminent ocean acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean  
acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



carbon. The net effect is an increase in  $\text{CO}_2$  uptake by about 40% in the warming simulation relative to the no-warming simulation. The export of excess carbon out of the top 200 m by physical transport is also enhanced in the warming simulation. Further, the projected changes in environmental conditions enhance marine biological productivity and the export of organic material and  $\text{CaCO}_3$  out of the Arctic surface. As a result, surface (Alk-DIC) is increased by 0.013 mmol/l. Thus, the biological feedbacks partly counter the physically-driven climate feedback. Taken all fluxes together, (Alk-DIC) decreases by 0.036 mmol/l (38%) more in the warming than in the no-warming simulation. This may be compared to a 0.013 mmol/l (34%) higher decrease in  $[\text{CO}_3^{2-}]$ .

Freshwater input at the ocean surface in the Arctic is increased due to melting of sea ice, as well as increased precipitation and reduced evaporation. In the CSM 1.4, rivers deliver pure freshwater to the ocean, with DIC and alkalinity set to zero. Land-ice is absent from the simulations. The annual mean sea ice volume in the Arctic Ocean is reduced by 69% in the A2 simulation from  $3.48 \times 10^4 \text{ km}^3$  (preindustrial) to  $1.07 \times 10^4 \text{ km}^3$  (2100). The summer sea ice cover is reduced by 74% from  $8.7 \times 10^6 \text{ km}^2$  (preindustrial) to  $2.3 \times 10^6 \text{ km}^2$  (2100), which correspond to a cover of 88% and 23% of the total Arctic Ocean surface ( $9.9 \times 10^6 \text{ km}^2$ ), respectively. The reduced sea ice cover allows gas exchange to occur in a larger area and the availability of light is increased in the ocean surface layer. The summer sea ice extent in the CSM is 10–30% higher than observed during the last 25 years, but the reduction in sea ice cover is at the lower end of the range of projections for the 21st century. Holland et al. (2006) project abrupt reduction in summer Arctic sea ice and near ice-free conditions by 2040 for the SRES A1B scenario, whereas in our SRES A2 simulation summer sea ice cover remains above 70% of the total Arctic Ocean surface until about 2050. Freshwater input and warming also cause an increase in stratification, here defined by the density difference between the surface and 200 m depth.

Modelled marine biological production in the Arctic increases due to a number of factors (Steinacher, 2007). Most importantly, enhanced availability of photosynthetic active radiation in response to reduced sea ice cover (and changes in cloud cover)

**Imminent ocean  
acidification**

M. Steinacher et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and ocean surface warming stimulate productivity. Simulated shortwave irradiance increases by 200% at the air-water interface and sea surface temperature increases by 0.31°C in scenario A2 by 2100. On the other hand, iron and phosphate concentration decrease only by a few percent. We note that production is limited by temperature and light, but not by phosphate and iron. The averaged mixed layer depth remains roughly unchanged.

Outside the Arctic, the impact of climate change on ocean acidification is relatively small. It globally affects  $\Omega_{\text{arag}}$  and pH equally (about 4%), but the decrease in  $\Omega_{\text{arag}}$  is reduced, whereas the decrease in pH is enhanced by climate change. This is consistent with the analysis of Cao et al. (2007) for carbon emission scenarios. McNeil and Matear (2007) found a climate-change feedback reducing the change in  $\Omega_{\text{arag}}$  by about 15% but almost no feedback on pH; they applied a scenario with prescribed CO<sub>2</sub> concentration and not a scenario with prescribed carbon emission as done here.

The climate feedback on  $\Omega_{\text{arag}}$  is negative in most regions of the oceans. Exceptions are the Arctic, the Labrador Sea and the high-latitude North Atlantic, the Antarctic polar zone and a few coastal regions (Fig. 9). The modeled high latitude changes are linked to sea ice as discussed for the Arctic; we note that the model simulates too much sea ice in the North Atlantic and North Pacific (Weatherly et al., 1998) and the magnitude of the positive feedback in the high latitude North Atlantic may not be realistic. The climate feedback on pH is positive (or only slightly negative) in all regions.

### 3.4 Seasonal and interannual variability of $\Omega_{\text{arag}}$

The simulated seasonal and interannual variability of  $\Omega_{\text{arag}}$  is small compared to the projected anthropogenic changes discussed above. The zonal mean seasonal cycle of  $\Omega_{\text{arag}}$  shows the typical quadrupole space-time pattern (Fig. 12a and b) with largest positive seasonal deviations in summer and largest negative deviations in winter in both hemispheres and at mid-latitudes. This seasonal cycle is driven by low summer DIC and high winter DIC values linked to thermally driven air-sea exchange and nutrient drawdown during the spring and summer growing season, as well as winter convection

**Imminent ocean acidification**

M. Steinacher et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and mixing. The seasonal variability of simulated  $\Omega_{\text{arag}}$  is small in the high latitude and in the tropical surface ocean ( $\pm 0.1$  in zonal mean  $\Omega_{\text{arag}}$ ; less than 5%). In Northern Hemisphere mid-latitudes deviations up to 15–20% (0.35–0.40 in zonal mean  $\Omega_{\text{arag}}$ ) can be found, with maximum values from August to October and minimum values from February to April (Fig. 12). Largest seasonal amplitudes ( $\Delta\Omega_{\text{arag}}$  up to 0.7) can be found around 40° N in the western Pacific and Atlantic, as well as in the South Atlantic near the coast of Argentina. These regions correspond to regions of high primary productivity and large seasonality in production. Surface  $\Omega_{\text{arag}}$  is increased at the end and after the high production phase, when nutrient and DIC concentrations are reduced.

The seasonal variability in the Arctic Ocean is out of phase with the general Northern Hemisphere seasonal signal and shows the same seasonal pattern as modeled for the Southern Hemisphere. Melting of Arctic sea ice decreases the supersaturation over the summer. The variability in sea ice cover and the associated freshwater input lead to a strong seasonal variability of alkalinity (up to  $\pm 4\%$ ) and DIC, which outbalances the relatively small variability in temperature. Overall, seasonal variability in  $\Omega_{\text{arag}}$  is small ( $< 0.1$ ) compared to the simulated anthropogenic changes in the Arctic.

The modeled large scale amplitudes and patterns of the seasonal variability do not show significant changes between 1820 and 2100 in the transient simulation under the A2 scenario in most regions. In the Southern Ocean (south of 40° S) and around 50° N and 75° N the seasonal amplitude tends to be slightly reduced (Fig. 12). However, as the annual mean  $\Omega_{\text{arag}}$  decreases with increasing  $\text{CO}_2$ , the relative seasonal variability increases from about +10%/–5% at 40° N in 1820 to about +25%/–15% in 2100. Finally, we note that the interannual variability in zonal mean  $\Omega$  is small (less than  $\pm 0.1$  in surface  $\Omega_{\text{arag}}$ ) both compared to the anthropogenic change as well as compared to zonal variability (Fig. 12). Because calcification is linearly coupled with production in the CSM1.4, interannual variability that could be caused by variations in the  $\text{CaCO}_3$ /organic carbon formation and remineralization rates is not captured by the model.

## 4 Discussion and conclusions

The changes in the acid-base state of ocean waters in response to human induced carbon emissions have been analyzed with the NCAR CSM1.4-carbon coupled climate-carbon cycle model for two transient greenhouse gas emission scenarios. The model is able to reproduce the large-scale, contemporary distribution of carbonate and the saturation state with respect to aragonite reasonably well. A number of caveats are nevertheless worth mentioning. Results simulated for the North Pacific should be viewed with caution as the model overestimates saturation in this basin. The NCAR CSM1.4 model has a climate sensitivity near the lower bound of current estimates and simulated magnitudes of climate feedbacks on changes in saturation and pH might be somewhat low. However, these feedbacks play a minor role in most regions. Further, the model does not distinguish between different mineral phases (calcite, aragonite, high-Mg calcite) of calcium carbonate (Gangstø et al., 2008) and therefore has a single length-scale for calcium carbonate remineralization independent of seawater saturation state. The model does not include an active sediment reservoir or sediment burial and further research is needed to quantify the magnitude of a potential buffering of the simulated changes in saturation by re-dissolution of calcium carbonate from ocean sediments. Langdon (2002) suggests that dissolution rates on reefs, banks, and tropical shelves will not be able to buffer the  $[\text{CO}_3^{2-}]$  decrease over the next 50–100 years.

Our study with a coupled climate-carbon cycle model supports the important finding of earlier studies that the acid-base state of the global ocean is currently undergoing large and rapid changes (Orr et al., 2005; Caldeira and Wickett, 2003) in response to anthropogenic carbon emissions with potentially large negative socio-economic impacts (Hoegh-Guldberg, 2005). The saturation state with respect to aragonite, calcite, or high-magnesium calcite is decreasing worldwide, causing a gradual worsening of living conditions for corals, pteropods, foraminiferas and coccolitophorides and potentially also for non-calcifying marine organisms (e.g. Fabry et al., 2008). Orr et al. (2005) have pointed out that the Southern Ocean will become undersaturated for aragonite if

**BGD**

5, 4353–4393, 2008

### Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



atmospheric CO<sub>2</sub> increases beyond 560 ppm.

We highlight that the Arctic Ocean surface becomes undersaturated with respect to aragonite at even lower CO<sub>2</sub> concentration. The combination of observation-based estimates of  $\Omega_{\text{arag}}$  in the Arctic (Jutterström and Anderson, 2005) with NCAR CSM1.4-carbon model projection indicates that 10% of the surface water along the investigated Arctic transect will become undersaturated for at least one month of the year when atmospheric CO<sub>2</sub> exceeds 409 ppm. This concentration is only 25 ppm higher than today's value and projected to be reached within the next decade for all SRES scenarios (Platner et al., 2008; Meehl et al., 2007) and within the next two decades for the recent set of multi-gas mitigation scenarios (Van Vuuren et al., 2008; Strassmann et al., 2008). The entire water column in the Arctic is projected to become undersaturated within this century if anthropogenic carbon emissions continue to grow along the SRES A2 business-as-usual trajectory. Even under the relatively low-CO<sub>2</sub> scenario B1, more than 50% of the Arctic surface waters are projected to become undersaturated. These results are consistent with the study by (Orr et al., 2008) who analyzed Arctic acidification based on results from different AOGCMS, including the NCAR CSM1.4-carbon.

Atmospheric CO<sub>2</sub> will very likely exceed 400 ppm in the next one or two decades given current carbon emission trends and the inertia of the energy and the whole socio-economic system. The question is thus not whether undersaturation will occur in the Arctic, but how big its magnitude will be, how large an area will be affected by undersaturation, and over how many months of the year undersaturation will be prevalent. Our findings, combined with the emerging evidence on the impacts of aragonite undersaturation on marine organisms imply that not only radiative forcing or temperature, but also atmospheric CO<sub>2</sub> concentration should be included as a target in climate policy analyses and in the development of mitigation scenarios with integrated assessment models. Our subjective assessment is that atmospheric CO<sub>2</sub> should not exceed 450 ppm in order to avoid the risk of large changes in marine ecosystems.

The loss in volume of water with saturation has been quantified. We consider five saturation regimes for ease of discussion. Water with an aragonite saturation of more

**BGD**

5, 4353–4393, 2008

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean  
acidification**

M. Steinacher et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

than 300%, considered suitable for coral growth, are projected to vanish completely in the second half of the century for the A2 scenario. Water with a saturation of more than 400%, considered optimal for coral growth, covered about 16% of the surface at preindustrial time and are gone in the next few years. Overall, the volume of water supersaturated with respect to aragonite decreases strongly in the A2 and B1 scenarios considered here, with potentially adverse effect also on species living on the ocean floor and in the deep (Guinotte et al., 2006).

Our analysis shows that in addition to the uptake of anthropogenic carbon by air-sea gas exchange, freshwater input from increased precipitation and ice melt contributes to the decrease in Arctic saturation state, while an increase in marine biological production slightly counteracts the decline. Additional freshwater input from land ice melt has not been considered here and is expected to slightly lower projected saturation states (Orr et al., 2008). Seasonal variability in saturation is significant in mid-latitudes regions, but small at high latitudes. The anthropogenic changes in saturation are superimposed on the seasonal variations. We did not find substantial changes in seasonal variability over this century in the NCAR model. Interannual variability in saturation is found to be small. Work at time series station shows that the anthropogenic signal is clearly distinguishable from seasonal and interannual variability (Bates, 2001; Brix et al., 2004; Kleypas et al., 2006).

In conclusion, human activities are perturbing the ocean and the habitats for marine organisms. The results of this study and of Feely et al. (2008) for the coastal North Pacific and Orr et al. (2008) for the Arctic show that undersaturation of surface waters with respect to aragonite is likely to become reality in a few years only.

*Acknowledgements.* This work was funded by the European Union projects CARBOOCEAN (511176-2) and EUROCEANS (511106-2) and is a contribution to the “European Project on Ocean Acidification” (EPOCA) which received funding from the European Community’s Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 211384. Additional support was received from the Swiss National Science Foundation and SCD acknowledges support from the U.S. National Science Foundation (NSF) grant ATM-0628582. Simulations were carried out at the Swiss National Supercomputing Centre (CSCS) in Manno, Switzerland.

## References

- Ammann, C. M., Joos, F., Schimel, D. S., Otto-Bliesner, B. L., and Tomas, R. A.: Solar influence on climate during the past millennium: Results from transient simulations with the NCAR Climate System Model, *P. Natl. Acad. Sci. USA*, 104, 3713–3718, doi:10.1073/pnas.0605064, 2007. 4360
- Anderson, L. A. and Sarmiento, J. L.: Redfield ratios of remineralization determined by nutrient data analysis, *Global Biogeochem. Cy.*, 8, 65–80, 1994. 4359
- Bates, N. R.: Interannual variability of oceanic CO<sub>2</sub> and biogeochemical properties in the Western North Atlantic subtropical gyre, *Deep-Sea Res. Pt. II*, 48, 1507–1528, 2001. 4374
- Bettge, T. W., Weatherly, J. W., Washington, W. M., Pollard, D., Briegleb, B. P., and Strand, W. G.: The CSM Sea Ice Model, NCAR Tech. Note NCAR/TN-425+STR, National Center for Atmospheric Research, Boulder, Colorado, USA, 1996. 4359
- Bonan, G. B.: The NCAR Land Surface Model (LSM version 1.0) coupled to the NCAR Community Climate Model, NCAR Tech. Note NCAR/TN-429+STR, National Center for Atmospheric Research, Boulder, Colorado, USA, 1996. 4359
- Boville, B. A. and Gent, P. R.: The NCAR Climate System Model, version one, *J. Climate*, 11, 1115–1130, 1998. 4359
- Brix, H., Gruber, N., and Keeling, C. D.: Interannual variability of the upper ocean carbon cycle at station ALOHA near Hawaii, *Global Biogeochem. Cy.*, 18, GB4019, doi:10.1029/2004GB002245, 2004. 4374
- Caldeira, K. and Wickett, M. E.: Anthropogenic carbon and ocean pH, *Nature*, 425, p. 365, 2003. 4354, 4356, 4358, 4372
- Cao, L., Caldeira, K., and Jain, A. K.: Effects of carbon dioxide and climate change on ocean acidification and carbonate mineral saturation, *Geophys. Res. Lett.*, 34, L05607, doi:10.1029/2006GL028605, 2007. 4358, 4363, 4370
- Conkright, M. E., Antonov, J. I., Boyer, T. P., Locarnini, R. A., O'Brien, T. D., Stephens, C., and Garcia, H. E.: *World Ocean Atlas 2001*, vol. 49-52 of NOAA Atlas NEDIS, US Govt. Printing Office, Washington, D.C., 2002. 4360
- Dickson, A.: *Handbook of methods of the analysis of the various parameters of the carbon dioxide system in seawater*, US Dep. of Energy, Washington, D.C., 2002. 4361
- Doney, S., Lindsay, K., Fung, I., and John, J.: Natural variability in a stable, 1000-yr global coupled climate-carbon cycle simulation, *J. Climate*, 19, 3033–3054, 2006. 4359, 4360

**BGD**

5, 4353–4393, 2008

---

### Imminent ocean acidification

M. Steinacher et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Fabry, V. J.: Marine Calcifiers in a High-CO<sub>2</sub> Ocean, *Science*, 320, 1020–1022, 2008. 4357
- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES J. Mar. Sci.*, 65, 414–432, 2008. 4355, 4356, 4372
- 5 Feely, R. A., Sabine, C. L., Kitack, L., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of Anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> System in the Oceans, *Science*, 305, 362–366, 2004. 4356, 4366
- Feely, R. A., Sabine, C. L., Hernandez-Ayon, J. M., Ianson, D., and Hales, B.: Evidence for Upwelling of Corrosive "Acidified" Water onto the Continental Shelf, *Science*, 320, 1490–1492, doi:10.1126/science.1155676, 2008. 4354, 4374
- 10 Fine, M. and Tchernov, D.: Scleractinian Coral Species Survive and Recover from Decalcification, *Science*, 315, p. 1811, 2007. 4356
- Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Dorland, R. V.: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, chap. Changes in Atmospheric Constituents and in Radiative Forcing*, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 129–234, 2007. 4360
- 15 to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, chap. Changes in Atmospheric Constituents and in Radiative Forcing, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 129–234, 2007. 4360
- Friedlingstein, P., Cox, P., Betts, R., Bopp, L., Von Bloh, W., Brovkin, V., Cadule, P., Doney, S., Eby, M., Fung, I., Bala, G., John, J., Jones, C., Joos, F., Kato, T., Kawamiya, M., Knorr, W., Lindsay, K., Matthews, H. D., Raddatz, T., Rayner, P., Reick, C., Roeckner, E., Schnitzler, K. G., Schnur, R., Strassmann, K., Weaver, A. J., Yoshikawa, C., and Zeng, N.: Climate-Carbon Cycle Feedback Analysis: Results from the C<sup>4</sup>MIP Model Intercomparison: Evolution of carbon sinks in a changing climate, *J. Climate*, 19, 3337–3353, 2006. 4363
- 20 Frölicher, T. L., Joos, F., Plattner, G. K., Steinacher, M., and Doney, S. C.: Natural variability and anthropogenic trends in oceanic oxygen in a coupled carbon cycle-climate model ensemble, *Global Biogeochem. Cy.*, in revision, 2008. 4360, 4363
- Frölicher, T. L., Joos, F., Plattner, G. K., Steinacher, M., and Doney, S. C.: Natural variability and anthropogenic trends in oceanic oxygen in a coupled carbon cycle-climate model ensemble, *Global Biogeochem. Cy.*, in revision, 2008. 4360, 4363
- 25 Fung, I., Doney, S., Lindsay, K., and John, J.: Evolution of carbon sinks in a changing climate, *P. Natl. Acad. Sci. USA*, 102, 11 201–11 206, 2005. 4359, 4363
- Gangstø, R., Gehlen, M., Schneider, B., Bopp, L., Aumont, O., and Joos, F.: Modeling the marine aragonite cycle: changes under rising carbon dioxide and its role in shallow water CaCO<sub>3</sub> dissolution, *Biogeosciences*, 5, 1057–1072, 2008, <http://www.biogeosciences.net/5/1057/2008/>. 4355, 4358, 4372
- 30 Gattuso, J.-P., Frankignoulle, M., Bourge, I., Romaine, S., and Buddemeier, R. W.: Effect of

**BGD**

5, 4353–4393, 2008

---

**Imminent ocean acidification**M. Steinacher et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calcium carbonate saturation of seawater on coral calcification, *Global Planet. Change*, 18, 37–46, 1998. 4357

Gent, P. R., Bryan, F. O., Danabasoglu, G., Doney, S. C., Holland, W. R., Large, W. G., and McWilliams, J. C.: The NCAR Climate System Model global ocean component, *J. Climate*, 11, 1287–1306, 1998. 4359

German Advisory Council on Global Change: *The Future Oceans – Warming up, Rising High, Turning Sour*, WGBU Special Report, German Advisory Council on Global Change (WBGU), Berlin, Germany, 2006. 4354

Guinotte, J., Orr, J., Cairns, S., Freiwald, A., Morgan, L., and George, R.: Will human-induced changes in seawater chemistry alter the distribution of deep-sea scleractinian corals?, *Front. Ecol. Environ.*, 4, 141–146, 2006. 4374

Heinze, C.: Simulating oceanic CaCO<sub>3</sub> export production in the greenhouse, *Geophys. Res. Lett.*, 31, L16308, doi:10.1029/2004GL020613, 2004. 4357

Hoegh-Guldberg, O.: Low coral cover in a high-CO<sub>2</sub> world, *J. Geophys. Res.*, 110, C09S06, doi:10.1029/2004JC002528, 2005. 4355, 4357, 4372

Holland, M. M., Bitz, C. M., and Tremblay, B.: Future abrupt reductions in the summer Arctic sea ice, *Geophys. Res. Lett.*, 33, L23503, doi:10.1029/2006GL028024, 2006. 4355, 4369

Iglesias-Rodriguez, M. D., Halloran, P. R., Rickaby, R. E. M., Hall, I. R., Colmenero-Hidalgo, E., Gittins, J. R., Green, D. R. H., Tyrrell, T., Gibbs, S. J., von Dassow, P., Rehm, E., Armbrust, E. V., and Boessenkool, K. P.: Phytoplankton Calcification in a High-CO<sub>2</sub> World, *Science*, 320, 336–340, doi:10.1126/science.1154122, 2008. 4357

Joos, F. and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20 000 years, *P. Natl. Acad. Sci. USA*, 105, 1425–1430, 2008. 4355

Jutterström, S. and Anderson, L. G.: The saturation of calcite and aragonite in the Arctic Ocean, *Mar. Chem.*, 94, 101–110, 2005. 4358, 4361, 4373

Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J., Feely, R. A., Millero, F., Mordy, C., and Peng, T.-H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP), *Global Biogeochem. Cy.*, 18, GB4031, doi:10.1029/2004GB002247, 2004. 4360

Kiehl, J. T., Hack, J. J., Bonan, G. B., Boville, B. A., Williamson, D. L., and Rasch, P. J.: The National Center for Atmospheric Research Community Climate Model, *J. Climate*, 11, 1151–1178, 1998. 4359

Kleypas, J. A., Buddemeier, R. W., Archer, D., Gattuso, J.-P., Langdon, C., and Opdyke, B. N.:

**BGD**

5, 4353–4393, 2008

---

## Imminent ocean acidification

M. Steinacher et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Geochemical Consequences of Increased Atmospheric Carbon Dioxide on Coral Reefs, *Science*, 284, 118–120, 1999. 4357, 4358

Kleyapas, J. A., Feely, R. A., Fabry, V. J., Langdon, C., Sabine, C. L., and Robbins, L. L.: Impacts of Ocean Acidification on Coral Reefs and Other Marine Calcifiers: A Guide for Future Research, report of a workshop held 18–20 April 2005, St. Petersburg, FL, sponsored by NSF, NOAA, and the US Geological Survey, 2006. 4356, 4357, 4365, 4366, 4374

Langdon, C.: Review of Experimental Evidence for Effects of CO<sub>2</sub> on Calcification of Reef builders, *Proceedings 9th International Coral Reef Symposium*, 2, 1091–1098, 2002. 4355, 4372

Langdon, C. and Atkinson, M. J.: Effect of elevated pCO<sub>2</sub> on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment, *J. Geophys. Res.*, 110, L05607, doi:10.1029/2004JC002576, 2005. 4356, 4357, 4365

Lueker, T. J., Dickson, A. G., and D., K. C.: Ocean pCO<sub>2</sub> calculated from dissolved inorganic carbon, alkalinity, and equations for K-1 and K-2: validation based on laboratory measurements of CO<sub>2</sub> in gas and seawater at equilibrium, *Mar. Chem.*, 70, 105–119, 2000. 4361

Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650 000–800 000 years before present, *Science*, 453, 379–382, 2008. 4354

McNeil, B. I. and Matear, R. J.: Climate change feedbacks on future oceanic acidification, *Tellus B*, 59, 191–198, 2007. 4355, 4358, 4370

Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.-C.: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, chap. Global Climate Projections, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 747–846, 2007. 4373

Millero, F. J.: Thermodynamics of the carbon dioxide system in the oceans, *Geochim. Cosmochim. Ac.*, 59, 661–677, 1995. 4361

Najjar, R. G., Jin, X., Louanchi, F., Aumont, O., Caldeira, K., Doney, S. C., Dutay, J.-C., Follows, M., Gruber, N., Joos, F., Lindsay, K., Maier-Reimer, E., Matear, R. J., Matsumoto, K., Mouchet, A., Orr, J. C., Sarmiento, J. L., Schlitzer, R., Weirig, M. F., Yamanaka, Y., and

**BGD**

5, 4353–4393, 2008

---

## Imminent ocean acidification

M. Steinacher et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean  
acidification**

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Yool, A.: Impact of circulation on export production, dissolved organic matter, and dissolved oxygen in the ocean: Results from Phase II of the Ocean Carbon-cycle Model Intercomparison Project (OCMIP-2), *Global Biogeochem. Cy.*, 21, GB3007, doi:10.1029/2006GB002857, 2007. 4359
- 5 Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005. 4354, 4355, 4356, 4357, 4358, 4364, 4366, 4372
- 10 Orr, J. C., Jutterström, S., Bopp, L., Anderson, L. G., Fabry, V. J., Frölicher, T. L., Jones, P., Joos, F., Maier-Reimer, E., Segschneider, J., Steinacher, M., and Swingedouw, D: Arctic ocean acidification, *Nature*, submitted, 2008. 4361, 4373, 4374
- Plattner, G.-K., Knutti, R., Joos, F., Stocker, T. F., von Bloh, W., Brovkin, V., Cameron, D., Driesschaert, E., Dutkiewicz, S., Eby, M., Edwards, N. R., Fichefet, T., Hargreaves, J. C., Jones, C. D., Loutre, M. F., Matthews, H. D., Mouchet, A., Müller, S. A., Nawrath, S., Price, A., Sokolov, A., Strassmann, K. M., and Weaver, A. J.: Long-term climate commitments projected with climate – carbon cycle models, *J. Climate*, 21, 2721–2751, doi:10.1175/2007JCLI1905.1, 2008. 4373
- 20 Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Kheshgi, H. S., Le Quere, C., Scholes, R. J., and Wallace, D. W. R.: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, chap. *The Carbon Cycle and Atmospheric Carbon Dioxide*, pp. 183–237, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2001. 4355
- 25 Randerson, J. T., Thompson, M. V., Conway, T. J., Fung, I. Y., and Field, C. B.: The contribution of terrestrial sources and sinks to trends in the seasonal cycle of atmospheric carbon dioxide, *Global Biogeochem. Cy.*, 11, 535–560, 1997. 4359
- Royal Society: *Ocean acidification due to increasing atmospheric carbon dioxide*, Policy Document 12/05, Royal Society, London, 2005. 4354
- 30 Schlitzer, R.: *Ocean Data View (Software)*, <http://odv.awi.de>, 2007.
- Schneider, K. and Erez, J.: The effect of carbonate chemistry on calcification and photosynthesis in the hermatypic coral *Acropora eurystroma*, *Limnol. Oceanogr.*, 51, 1284–1293, 2006.

Siegenthaler, U., Stocker, T. F., Monnin, E., Lüthi, D., Schwander, J., Stauffer, B., Raynaud, D., Barnola, J.-M., Fischer, H., Masson-Delmotte, V., and Jouzel, J.: Stable carbon cycle-climate relationship during the late Pleistocene, *Science*, 310, 1313–1317, doi:10.1126/science.1120130, 2005. 4355

Steinacher, M.: Ocean acidification and changes in marine productivity in simulations with the fully coupled 3-D climate model CSM1.4-carbon, Master's thesis, University of Bern, Bern, Switzerland, 2007. 4355, 4369

Strassmann, K. M., Joos, F., and Plattner, G.-K.: CO<sub>2</sub> and non-CO<sub>2</sub> radiative forcing agents in 21st century climate change mitigation scenarios, *Clim. Dynam.*, submitted, 2008. 4355, 4373

Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34, L09501, doi:10.1029/2007GL029703, 2007. 4355

Thomas, H., Prowe, A. E. F., van Heuven, S., Bozec, Y., de Baar, H., Schiettecatte, L.-S., Suykens, K., Kon, M., Borges, A. V., Lima, I. D., and Doney, S. C.: Rapid decline of the CO<sub>2</sub> buffering capacity in the North Sea and implications for the North Atlantic Ocean, *Global Biogeochem. Cy.*, 21, GB4001, doi:10.1029/2006GB002825, 2007. 4364

Van Vuuren, D. P., Meinshausen, M., Plattner, G.-K., Joos, F., Strassmann, K. M., Smith, S. J., Wigley, T. M. L., Raper, S. C. B., Riahi, K., de la Chesnaye, F., den Elzen, M., Fujino, J., Jiang, K., Nakicenovic, N., Paltsev, S., and Reilly, J. M.: Temperature increase of 21st century mitigation scenarios, *Proc. Nat. Acad. Sci. USA*, 105, 15 258–15 262, doi:10.1073/pnas.0711129105, 2008. 4355, 4373

Wang, Y. M., Lean, J. L., and Sheeley, N. R.: Modeling the sun's magnetic field and irradiance since 1713, *Astrophys. J.*, 625, 522–538, 2005. 4360

Weatherly, J. W., Briegleb, B. P., Large, W. G., and Maslanik, J. A.: Sea Ice and Polar Climate in the NCAR CSM, *J. Climate*, 11, 1472–1486, 1998. 4359, 4370

## Imminent ocean acidification

M. Steinacher et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Imminent ocean acidification**

M. Steinacher et al.

**Table 1.** Different levels of aragonite undersaturation ( $\Omega_{\text{arag}} < 1$ ), atmospheric CO<sub>2</sub> concentration and year when these levels are reached in the Arctic for the A2 and B1 scenario. Levels are defined by the area fraction occupied by undersaturated surface waters in at least one month of the year, in the annual mean, and throughout the year.

Level (area fraction and seasonal occurrence)	CO <sub>2</sub> (ppm)	Year in A2	Year in B1
10%, ≥1 month	409	2016	2018
10%, annual mean	428	2023	2025
10%, year-round	450	2029	2034
50%, ≥1 month	520	2047	2068
50%, annual mean	534	2050	2085
50%, year-round	552	2054	–
100%, ≥1 month	634	2070	–
100%, annual mean	634	2070	–
100%, year-round	694	2080	–
Entire water column	765	2090	–

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Imminent ocean acidification

M. Steinacher et al.



**Fig. 1.** Prescribed fossil fuel and land use CO<sub>2</sub> emissions **(a)** based on historical data (1820–2000, black) and the SRES-A2 (red) and B1 (green) scenarios (2000–2100). Simulated global annual mean atmospheric CO<sub>2</sub> **(b)**, sea surface temperature (SST) **(c)**, dissolved inorganic carbon (DIC) **(d)**, aragonite saturation ( $\Omega_{\text{arag}}$ ) **(e)**, and pH **(f)** at the ocean surface.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Imminent ocean acidification

M. Steinacher et al.



**Fig. 2.** Taylor diagram conveying information about the pattern similarity between simulated and observation-based (GLODAP, WOA01) results for  $\Omega_{arag}$  (squares) and  $[CO_3^{2-}]$  (crosses). The observed field is represented by a point at unit distance from the origin along the abscissa. The distance from this point to the other points representing simulated fields, indicates the normalized r.m.s error. The simulated fields are averaged over the model years 1990–1999 and the results are shown for the entire global ocean (black), the top 200 m of the global ocean (blue), and the three major basins Atlantic (green), Pacific (red), and Indian Ocean (magenta), all of which include the respective part of the Southern Ocean.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Imminent ocean  
acidification

M. Steinacher et al.




**Fig. 3.** Meridional sections through the Pacific (left) and the Atlantic (right) showing zonal mean of observed (top) and modeled (middle) supersaturation with respect to aragonite ( $\Delta[\text{CO}_3^{2-}]$ ) in  $\mu\text{mol/l}$ . The saturation horizon is indicated by the thick line at  $\Delta[\text{CO}_3^{2-}] = 0$ . The bottom row shows the differences between model and measurements (model-observations). Simulated fields are averaged over the model years 1990–1999.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Imminent ocean acidification

M. Steinacher et al.



**Fig. 4.** (a) Observation-based present day aragonite saturation ( $\Omega_{\text{arag}}$ ) in the Arctic Ocean based on measurements from the ODEN-91, AOS-94, and ARCSYS-96 cruises during summer. The transect starts near the Bering Strait (0 km), leads then across the North Pole (2500 km) to Spitsbergen (3750 km), and from there to the Laptev Sea (8600 km), crossing the Eurasian Basin three times. (b) Difference between the modeled and the observation-based  $\Omega_{\text{arag}}$  for the same transect and time. (c) Model-only projection of summer  $\Omega_{\text{arag}}$  by 2099 under the A2 scenario, and (d) from combining modeled changes in  $\Omega_{\text{arag}}$  and observation-based  $\Omega_{\text{arag}}$  shown in (a).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Imminent ocean acidification

M. Steinacher et al.




**Fig. 5.** Simulated aragonite saturation at the surface by the years 1820 (a), 2000 (b), and 2100 (c, d) for SRES scenarios A2 and B1. The largest changes in  $\Omega_{\text{arag}}$  (e, f) are projected for the subtropics.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Imminent ocean acidification

M. Steinacher et al.



**Fig. 6.** Projected evolution of the **(a)** annual-mean and of the **(b)** lowest monthly mean zonally-averaged aragonite saturation  $\Omega_{\text{arag}}$  for the SRES A2 scenario (model only). The evolution is plotted as a function of time and of the annual-mean atmospheric  $\text{CO}_2$  mixing ratio at the ocean surface.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Imminent ocean  
acidification

M. Steinacher et al.



**Fig. 7.** Time-depth diagrams of simulated annual mean aragonite supersaturation  $\Delta[\text{CO}_3^{2-}]$  ( $\mu\text{mol/l}$ ) for SRES A2; **(a)** global average, **(b)** in the Arctic Ocean, and **(c)** in the Southern Ocean.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






**Fig. 8.** Simulated global changes in the volume of supersaturated waters with respect to aragonite under the SRES A2. Panel (a) shows changes in the entire ocean volume, whereas panel (b) shows only changes for the top 200 m. Waters that are more than four times supersaturated (orange) are projected to vanish by the year 2010, and those that are 3 to 4 times supersaturated (gray) by the year 2070. The volumes of 2–3 and 1–2 times supersaturated waters (blue and red) decrease to 25% and 70% of the preindustrial values by 2100. Thus the volume of undersaturated waters (green) increases by about 30% (or 17% of the total ocean volume) by 2100. The dashed lines indicate the volume changes under the SRES B1 scenario. The crosses denote the respective values derived from the GLODAP/WOA01 data set for 1995 and the estimated preindustrial values (1830) obtained by subtracting estimated anthropogenic  $\text{CO}_2$  from DIC concentrations.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Imminent ocean  
acidification

M. Steinacher et al.




**Fig. 9.** Differences in projected surface  $\Omega_{\text{arag}}$  **(a)** and pH **(b)** between the SRES A2 simulation with and without warming, averaged over the decade 2090–2099. Positive values indicate enhanced acidification by climate change, negative values indicate reduced acidification.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Imminent ocean acidification

M. Steinacher et al.



**Fig. 10.** Simulated annual mean alkalinity and DIC surface concentrations from 1820 to 2100 in the Arctic Ocean **(a)** and between 50°N and 70°N **(b)**. Without warming (blue) both regions show similar changes. Including climate change (red) changes alkalinity and DIC in the North Pacific and North Atlantic only slightly and the difference Alk-DIC, which is approx. proportional to  $[\text{CO}_3^{2-}]$ , is almost constant. This is representative for all regions, except the Arctic, where the alkalinity is strongly reduced and the difference Alk-DIC is virtually zero by the end of the 21st century.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





**Fig. 11.** Projected changes in the annual mean difference Alk-DIC, which is approximately proportional to  $[\text{CO}_3^{2-}]$ , from 1820 to 2100 under the A2 scenario in the top 200 m of the Arctic Ocean. The total change (black) is composed of changes in the net  $\text{CO}_2$  uptake by gas-exchange with the atmosphere (red), changes in the freshwater flux at the ocean surface (blue), as well as changes in biology (green) and in the export of excess carbon by surface-to-deep and lateral exchange (orange). The difference between the simulation without warming (hatched) and the standard global warming simulation (solid) quantifies the impact of climate change on the individual components and on the total amplification of the carbonate ion reduction. The inset relates the changes in the difference Alk-DIC to the changes in  $[\text{CO}_3^{2-}]$  for all grid boxes that contribute to the Arctic budget and for the warming (red) and no-warming (blue) simulations. The strong correlation indicates that the difference Alk-DIC is a reasonable proxy for  $[\text{CO}_3^{2-}]$  in the Arctic surface ocean.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Fig. 12.** Simulated seasonal variability of zonal mean surface  $\Omega_{\text{arag}}$ . Average deviation from the annual mean for the decade 1990–1999, in absolute values (a) and relative to the annual mean  $\Omega_{\text{arag}}$  (b, in percent). Maps of maximum positive (c) and negative (d) anomalies in surface  $\Omega_{\text{arag}}$  for the decade 1990–1999 (maximum deviation from the decadal mean at each grid point and for every month). Average deviation from the annual mean in March/April (e) and September/October (f) for the decades 1990–1999 (black line) and 2090–2099 (red line) under SRES A2. The shaded areas indicate the interannual variability (green,  $\pm\sigma$ ) and the variability across different longitudes (grey,  $\pm\sigma$ ) for the decade 1990–1999.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I ◀](#)
[▶ I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
