

Interactive comment on “Impact of anthropogenic ocean acidification on thermal tolerance of the spider crab *Hyas araneus*” by K. Walther et al.

K. Walther et al.

kathleen.walther@awi.de

Received and published: 17 August 2009

Author comment to I. J. McGaw – Referee 1

We appreciate the constructive comments by I. J. McGaw In the following we answered the questions and tried to develop a hypothesis to explain the observed results.

1. In our opinion the standard unit for CO₂ widely used in the community is ppm, the community, not us on our own would have to consider changing it, e.g. to SI units.
2. CO₂ penetrates the surface ocean level by air-to-sea equilibration. The distribution of CO₂ is occurs by ocean circulation (Orr et. al, 2001). In the German Bight the water depth is less than 100m (Pohlmann, 1996), in this case the whole water body will be equilibrated with CO₂.

3. *Hyas araneus* live on hard, stony, sandy and soft bottoms from < 1 down to 360 m, most commonly at depths less than 50 m. Males may reach a carapace length of 105 mm (Christiansen, 1969). We add this information to the manuscript. To our knowledge this species is not used commercially.

4. We prevented haemolymph clotting around the oxygen probes by rinsing the probes with a heparin solution. We added this information to the methods section. We calculated heart rates continuously at each temperature step independent from pauses of heart rate, which were not observed for significant time during the incubations.

5. We used 25°C as the upper temperature because the original aim of our study was to show effects of CO₂ on pejus limits. In this case it was not our aim to expose the crabs at temperatures beyond their critical temperature. However, the experiments showed that the critical temperature is largely affected by CO₂. In future studies it is necessary to quantify the critical temperature under normocapnic conditions.

6. We changed this section accordingly.

7. We eliminated these sentences.

8. We eliminated the repeated mentioning of results in the discussion part.

We developed a hypothesis why increased CO₂ affects heart rates and PO₂ levels: Short-term exposure to acidification results in enhanced ventilation and an accompanied rise in heart rate which supports CO₂ release and alleviation of pH disturbances (cf. Pörtner et al., 2005). The additional decrement in heart rate at low temperatures was possibly caused by the accumulation of adenosine under CO₂ exposure. Adenosine was found to depress ventilation rate in *Sipunculus nudus* (Reipschläger et al., 1997). In crustaceans, adenosine also depresses spontaneous activity and the responsiveness of interneurons to electrical and chemical stimuli in the brain (Derby et al., 1987) and elicits bradycardia (Brevard et al., 2003). In contrast, adenosine can display a stimulatory effect on heart rate, haemolymph flow and scaphognathite fre-

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

quency (Maurer et al., 2008; Stegen and Grieshaber, 2001). This apparent discrepancy resembles the contrasting CO₂ effects at low and at high temperatures. Such stimulatory effect might in fact be involved in the increase in heart rate with rising CO₂ concentrations in the warmth (Fig. 5). Further experiments are required to test these hypotheses.

9. Yes the North Sea around Helgoland showed an increase in temperature during the last 40 years by 1,1 °C to a mean temperature of 18 °C and maxima of about 20°C (Wilthire and Manly, 2004). Due to rising temperatures in the North Sea the projected critical temperatures of *H. araneus* can be reached in the near future.

10. We demonstrated the effect of CO₂ on the acute thermal tolerance window. During 24h of incubation the pH level of the extracellular space would reach a new level. This is the same during exposure to longer periods (days). It is conceivable that the crabs fully compensate their extracellular pH close to the starting point when exposed over a longer time (months).

11. We haven't done thermal preference experiments yet. To answer the question longterm exposure to CO₂ is necessary. This is a nice suggestion for future studies.

References:

Brevard, M. E., Duong, T. Q., King, J. A. and Ferris, C. F.: Changes in MRI signal intensity during hypercapnic challenge under conscious and anesthetized conditions, *Magnetic Resonance Imaging*, 21, 995-1001, 2003.

Christiansen, M. E. (Ed.): *Crustacea Decapoda Brachyura*, in: *Marine Invertebrates of Scandinavia*, No. 2, Universitetsforlaget, Oslo, 1969.

Derby, C. D., Ache, B. W. and Carr, W. E. S.: Purinergic modulation in the brain of the spiny lobster, *Brain Research*, 421, 57-64, 1987.

Maurer, G., Wilkens, J. L. and Grieshaber, M. K.: Modulatory effects of adenosine and adenine nucleotides on different heart preparations of the American lobster, *Homarus*

americanus, J. Exp. Biol., 211, 661-670, 2008.

Orr, J.C., Maier-Reimer, E., Mikolajewicz, U., Monfray, P., Sarmiento, J.L., Toggweiler, J.R., Taylor, N.K., Palmer, J., Gruber, N., Sabine, C.L., Le Quéré, C., Key, R.M. and Boutin, J.: Estimates of anthropogenic carbon uptake from four three-dimensional global ocean models. *Global Biogeochemical Cycles*, 15, 1, 43-60, 2001.

Pohlmann, T.: Predicting the thermocline in a circulation model of the North Sea – Part I: model description, calibration and verification. *Continental Shelf Research*, 16, 7, 131-146, 1996.

Pörtner, H. O., Langenbuch, M. and Michaelidis, B.: Synergistic effects of temperature extremes, hypoxia, and increases in CO₂ on marine animals: From Earth history to global change, *J. Geophys. Res.*, 110, C09S10, doi:10.1029/2004JC002561, 2005. Reipschläger, A., Nilsson, G. E. and Pörtner, H. O.: A role for adenosine in metabolic depression in the marine invertebrate *Sipunculus nudus*, *American Journal of physiology-regulatory integrative and comparative physiology*, 272, R350-R356, 1997.

Stegen, E. and Grieshaber, M. K.: Adenosine increases ventilation rate, cardiac performance and haemolymph velocity in the American lobster *Homarus americanus*, *J. Exp. Biol.*, 204, 947-957, 2001.

Wiltshire, K. H. and Manly, B. F. J.: The warming trend at Helgoland Roads, North Sea: phytoplankton response, *Helgoland Mar. Res.*, 58, 269-273, 2004.

Interactive comment on Biogeosciences Discuss., 6, 2837, 2009.

BGD

6, C1546–C1549, 2009

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper