

Abstract

Modeling fire as an integral part of an Earth system model (ESM) is vital for quantifying and understanding fire-climate-vegetation interactions on a global scale and from an Earth system perspective. In this study, we introduce to the Community Earth System Model (CESM) the new global fire parameterization proposed by Li et al. (2012), now with a more realistic representation of the anthropogenic impacts on fires, with a parameterization of peat fires, and with other minor modifications. The improved representation of the anthropogenic dimension includes the first attempt to parameterize agricultural fires, the economic influence on fire occurrence, and the socioeconomic influence on fire spread in a global fire model; also an alternative scheme for deforestation fires.

The global fire parameterization has been tested in CESM1's land component model CLM4 in a 1850–2004 transient simulation, and evaluated against the satellite-based Global Fire Emission Database version 3 (GFED3) for 1997–2004. The simulated 1997–2004 average global totals for the burned area and fire carbon emissions in the new fire scheme are 338 Mha yr^{-1} and 2.1 Pg C yr^{-1} . Its simulations on multi-year average burned area, fire seasonality, fire interannual variability, and fire carbon emissions are reasonable, and show better agreement with GFED3 than the current fire scheme in CESM1 and modified CTEM-FIRE. Moreover, the new fire scheme also estimates the contributions of global fire carbon emissions from different sources. During 1997–2004, the contributions are 8% from agricultural biomass burning, 27% from tropical deforestation and degradation fires, 5% from global peat fires (3.7% from tropical peat fires), and 60% from other fires, which are close to previous assessments based on satellite data, government statistics, or other information sources. In addition, we investigate the importance of direct anthropogenic influence (anthropogenic ignitions and fire suppression) on global fire regimes during 1850–2004, using CESM1 with the new fire scheme. Results show that the direct anthropogenic impact is the

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



main factor driving the trends of global burned area in the whole period and fire carbon emissions only before ~ 1870.

1 Introduction

Fire is an important Earth system process on a global scale (Bowman et al., 2009). It depends on vegetation characteristics, climate, and human activities, and generates feedbacks by affecting biogeochemical cycles, vegetation composition and structure, land-atmosphere water and heat exchanges, atmospheric chemistry and composition, and human health and property (Bond et al., 2004; IPCC, 2007; Cochrane and Ryan, 2009; Bond-Lamberty et al., 2009; van der Werf et al., 2010; Bowman et al., 2011).

To quantify the role of fire in the Earth system on a global scale from an Earth system perspective and project global change with variable fire regimes, it is vital to model fire as an integral part of an Earth system model (ESM). An ESM with a representation of the anthropogenic impact on fires may also help identify appropriate and sustainable strategies for fire management.

Existing global fire parameterizations suitable for ESMs aim to best match the observed fire regimes for the contemporary time period, given that only the contemporary global fire product is available with a sufficient quality to be used as evaluation data (Kloster et al., 2010; Prentice et al., 2011). These global fire parameterizations are generally large-scale (10^3 – 10^5 km²), and simplify the real world at various levels to improve the efficiency of computations. MC-FIRE is the first fire parameterization used in a Dynamic Global Vegetation Model (DGVM, a type of ecosystem model in ESM) (Lenihan et al., 1998). Then, Thonicke et al. (2001) proposed the most widely used fire scheme, Glob-FIRM. Arora and Boer (2005) introduced CTEM-FIRE in CTEM-DGVM, which was a process-based fire model of intermediate complexity as Glob-FIRM but considering the effect of wind speed on fire spread and certain anthropogenic effects on fire regimes. N. M. Mahowald (personal communication, 2012) modified Glob-FIRM by translating the original annual to a sub-daily time step to simulate fire seasonality

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that the simulated global total amount and spatial distribution of burned area and fire emissions were broadly consistent with the satellite-based GFED3 fire product.

Global fire parameterizations need further improvements, especially with respect to the representation of the human dimension of fire regimes. First, fires are used worldwide to clear agricultural residue, fertilize the soil, and eliminate pests (Chidumayo, 1987; Le Page et al., 2010a). Agricultural fires account for 8–11 % of the 2001–2003 global annual fire counts (Korontzi et al., 2006) and significantly change fire seasonality in many regions (Le Page et al., 2010a; Magi et al., 2012). These fires emit large amounts of hydrocarbons, nitrogen oxides, carbon monoxide, which are the precursors of tropospheric ozone and influence the chemistry of the OH radical (Logan et al., 1981; Logan, 1985), and smoke aerosols with significant consequences on human health (World Health Organization (WHO), 2000). Earlier global fire models assumed that there were no fires in cropland (Arora and Boer, 2005; Kloster et al., 2010; Thonicke et al., 2010; Prentice et al., 2011), or neglected any differences from non-agricultural fires, e.g. the current fire model in CESM1 (N. M. Mahowald, personal communication, 2012; Oleson et al., 2010). Second, deforestation fires and peat fires are important sources of fire carbon emissions (van der Werf et al., 2010). Of all the previous global fire models, only the modified CTEM-FIRE (Kloster et al., 2010) included deforestation fires based on land use change data. In Kloster et al. (2010), the fires in a grid cell were simulated as the sum of deforestation fires and fires due to natural and anthropogenic (intentional and accidental) ignitions. So far, no global fire model includes peat fires. The ESMs that do not include peat fires cannot simulate some large fires and associated extreme air pollution events, such as the famous 1997 Southeast Asian Haze that lasted several months and was caused by tropical peat fires and deforestation fires in Indonesia (http://en.wikipedia.org/wiki/1997_Southeast_Asian_haze), or the 2010 Russian Haze due to boreal peat fires and forest fires (http://en.wikipedia.org/wiki/2010_Russian_wildfires). Third, only the effect of population density on fire occurrence has been considered in earlier studies (Pechony and Shindell, 2009; Kloster et al., 2010; Thonicke et al., 2010; Li et al., 2012). Other anthropogenic impacts on

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



fires, for example, the fire management and fire fighting capability related to countries' socioeconomic situations (Chuvieco et al., 2008; Aldersley et al., 2011), have yet to be considered.

The aim of the present study is to introduce an expanded version of the Li et al. (2012) fire parameterization to the CESM. The expanded version includes a more realistic representation of anthropogenic impacts on fires, the parameterization of peat fires, and other minor modifications. The more realistic representation of anthropogenic impacts includes the first attempt to parameterize agricultural fires, the economic influence on fire occurrence, and the socioeconomic influence on fire spread in a global fire model; also an alternative scheme of deforestation fires. Its global performance in CESM1 is evaluated against the satellite-based Global Fire Emission Database version 3 (GFED3) (Giglio et al., 2010; van der Werf et al., 2010), and compared with the current fire parameterization implemented in CESM1 (N. M. Mahowald, personal communication, 2012; Oleson et al., 2010) and the modified CTEM-FIRE (Kloster et al., 2010). Methodologically, compared with the two fire schemes, the new fire scheme of CESM has not only the advantages of the Li et al. (2012)'s fire scheme (see the Para 3 in Sect. 1) but also the Li et al. (2012)'s future improvements listed above.

In this paper, Sect. 2 presents the data used to calibrate, drive and evaluate the model. Section 3 describes the new fire parameterization scheme. Section 4 introduces its application in CESM1 and simulations. Section 5 evaluates the global fire simulations. Based on the new fire parameterization, Sect. 6 investigates the importance of direct anthropogenic influence on the long-term trend of global fire regimes since the middle of the 19th century. Conclusions and future development appear in Sect. 7.

2 Data

Table 1 lists data used in the present study. The 1997–2009 monthly burned area and fire carbon emissions data at a 0.5° spatial resolution are provided by GFED3 (Giglio

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2010; van der Werf et al., 2010). The GFED3 burned area is a mixture of observations and satellite-based estimates, which are generated from 500-m MODIS burned area maps (MCD64A1), active fire detections from multiple satellites, local regression, and regional regression trees (Giglio et al., 2010; L. Giglio, personal communication, 2012). The GFED3 fire emissions data are the output of a revised CASA biogeochemical model driven by the GFED3 burned area, MODIS vegetation and land data, active fire detections from multiple satellites, weather observations, MODIS photosynthetically active radiation, and AVHRR NDVI data (van der Werf et al., 2010). The GFED3 fire product represents the most comprehensive attempt to date to derive the burned area and fire emissions from remote sensing data, and it is suitable for calibrating functions and parameters as well as for evaluating present-day simulations of global fire models (van der Werf et al., 2010; Prentice et al., 2011). The MODIS 0.5° global monthly fire count product for 2001–2010 is obtained from ftp://fuoco.geog.umd.edu (Giglio et al., 2006).

The 1948–2004 3-h surface air temperature, wind speed, specific humidity, and air pressure, and 6-h downward solar radiation data and precipitation at a T62 (~ 1.875°) spatial resolution are the atmospheric data that drive the CLM4 (Qian et al., 2006). The 1948–2004 6-h reanalysis relative humidity data at a 2.5° spatial resolution are derived from the National Centers for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) (Kalnay et al., 1996), which are bias-corrected using the 10' monthly Climate Research Unit (CRU) climatological data (New et al., 1999, 2000) according to the method of Qian et al. (2006). NASA LIS/OTD grid product v2.2 (http://ghrc.msfc.nasa.gov) provides the 2.5° daily lightning time series from May 1995 to December 2004 (4260 × 144 × 73) and 2-h climatological lightning data (365 × 12 × 144 × 73).

The population density data for 1990, 1995, 2000, and 2005 at a 0.5° spatial resolution are provided by the Gridded Population of the World version 3 (GPWv3) (CIESIN, 2005). The 5 min decadal population density data for 1850–1980 are from the Database of the Global Environment version 3.1 (HYDEv3.1) (Klein Goldewijk

et al., 2010). Gross Domestic Production (GDP) per capita in 2000 and at a 0.5° spatial resolution is from Van Vuuren et al. (2006), which is the base-year GDP data for IPCC-SRES and derived from country-level World Bank's World Development Indicators (WDI) measured in constant 1995 US\$ (World Bank, 2004) and the UN Statistics Database (UNSTAT, 2005). The peatland area fraction at a 0.5° spatial resolution is derived from three vector datasets: peatland data in Indonesia and Malaysian Borneo (Olson et al., 2001); peatland data in Canada (Tarnocai et al., 2011); and bog, fen and mire data in boreal regions (north of 45° N) outside Canada provided by the Global Lakes and Wetlands Database (GLWD) (Lehner and Döll, 2004). The climatological peak month data for agricultural waste burning at a 0.5° resolution (van der Werf et al., 2010) are used to estimate the timing of agricultural fires. The annual 0.47° × 0.63° land use and land cover change (LULCC) data for 1850–2005 and the present-day land cover data are from the CLM4 land surface data in CESM1 (Lawrence and Chase, 2007, 2010; Oleson et al., 2010). The LULCC data is based on Version 1 of the Land-Use History A product (LUHa.v1) (Hurtt et al., 2006), while the present-day data is based on the MODIS land surface data product (Justice et al., 2002; Hansen et al., 2003) and the cropping dataset of Ramankutty et al. (2008). The CLM represents vegetation as plant functional types (PFTs, Table 2) instead of species to generalize plant function on a global scale (Bonan et al., 2002). Thus, the land type data from various sources are translated into the PFT levels in the CLM4 land surface data (Oleson et al., 2010).

3 Fire parameterization

The new fire parameterization for CESM contains four components: agricultural fires in cropland (region A), deforestation fires in the tropical closed forests (region B), non-peat fires outside regions A and B (region C), and peat fires in all three regions (Fig. 2). In this fire parameterization, burned area is mainly affected by climate and weather conditions, vegetation composition and structure, and human activities. After burned

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



area is calculated, we estimate the fire impact, including biomass and peat burning, and vegetation mortality in the post-fire region; adjustment of the carbon and nitrogen (C/N) pools; and trace gas and aerosol emissions due to fuel burning.

This fire parameterization is a further development of the work presented in detail by Li et al. (2012). The Li et al. (2012) parameterization (Fig. 1) is used here in region C with modifications, mainly from adding parameterizations about the economic influence in the fire occurrence component and the socioeconomic impact in the fire spread component. The current paper only describes the further development of the Li et al. (2012) scheme, i.e. the parameterization schemes of agricultural fires, deforestation fires, peat fires, economic impact on fire occurrence, socioeconomic impact on fire spread, and some minor modifications. Table A1 lists all variables and parameters used in the present study and their meanings and units.

3.1 Agricultural fires

Korontzi et al. (2006) used the MODIS active fire product to describe global patterns of agricultural fires and indicated that the fuel load could affect the interannual variability of agricultural fires. Moreover, as an anthropogenic activity, agricultural fires are greatly affected by socioeconomic conditions (Chuvienco et al., 2008; Bowman et al., 2011), and generally occur after harvesting or before planting, broadly different from the fire seasonality of non-agricultural fires (Le Page et al., 2010a; Magi et al., 2012). Thus, the burned area of cropland (km^2 (time step) $^{-1}$) is taken as:

$$A_b = a_1 f_b f_{se} f_t f_{crop} A_g, \quad (1)$$

where constant a_1 ((time step) $^{-1}$) is estimated using an inverse method (see Sect. 4); f_b is the fuel availability factor; f_{se} represents the socioeconomic effect on fires; f_t determines the seasonality of agricultural fires; f_{crop} is the fractional coverage of cropland; and A_g is the area of the grid cell (km^2).

We assume that the fuel-load requirement of fires is universal, so f_b is set the same as that used for non-cropland regions in Li et al. (2012). The socioeconomic factor f_{se}

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is given as follows:

$$f_{se} = f_d f_e. \quad (2)$$

Here

$$f_d = 0.04 + 0.96 \times \exp \left[-\pi \left(\frac{D_p}{350} \right)^{0.5} \right] \quad (3)$$

5 and

$$f_e = 0.01 + 0.99 \times \exp \left(-\pi \frac{\text{GDP}}{10} \right) \quad (4)$$

are the effects of population density D_p (personkm⁻²) and Gross Domestic Product GDP (k1995US\$capita⁻¹) on burned area, respectively, which are identified by maximizing the explained variability of the GFED3 burned area fraction with both socioeconomic indices in grid cells where the area fraction of cropland is > 50 % (Fig. 3). Equations (3) and (4) reflect that less populated and less developed regions are more likely to use fires as a cheap and effective means of removing agricultural waste. f_t is simply set to 1 for rainless time steps during 1 month after harvesting or before planting, or to 0 otherwise, which is supported by previous analyses of agricultural fire seasonality (Korontzi et al., 2006; Le Page et al., 2010a; Magi et al., 2012). The specific month for agriculture fires (month after harvesting or before planting) can be set for various crop types for ESMs that simulate the harvesting and planting date, or set as the climatological peak month for agricultural fires derived from van der Werf et al. (2010) as we do here.

20 In the post-fire region, carbon transfers from agricultural waste (litter) to atmosphere. The combustion completeness factor is set to 0.8, which is the same as that used for grass litter in Li et al. (2012). Emission factors of trace gases and aerosols are shown in Table 3, which are based on Andreae and Merlet (2001) and M. O. Andreae (personal communication, 2011).

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Deforestation fires

Of the existing global fire models, only modified CTEM-FIRE (Kloster et al., 2010) simulated deforestation fires, in which fires in a grid cell were set as the sum of deforestation fires and fires due to natural and (intentional and accidental) human ignitions. The deforestation fires were assumed to occur whenever the land type was converted (e.g. from tree, grass or shrub PFTs to bare soil, from crop PFTs to grass or tree PFTs) and the soil was dry. This parameterization may lead to double-counting and, therefore an overestimation of fire carbon emissions and especially the burned area due to intentional human ignitions in land-type conversion regions, given that deforestation fires are a type of fires due to intentional human ignitions and many land-type conversions do not involve deforestation.

Here we provide an alternative scheme to parameterize deforestation fires in global fire models. It focuses on deforestation fires in tropical closed forests. Tropical closed forests are defined as grid cells with tropical tree coverage > 60 % according to the FAO classification. Deforestation fires are defined as fires caused by deforestation (including escaped deforestation fires, termed degradation fires) outside cropland of these grid cells. To avoid overestimating fires due to anthropogenic intentional ignitions, the Li et al. (2012) parameterization about fires due to natural and anthropogenic (intentional and accidental) ignitions is not used in these grid cells. In this way, fires due to natural and anthropogenic accidental ignitions are neglected, which are rare in these regions due to high humidity environments (van der Werf, et al., 2009; Le Page et al., 2010b). Deforestation fires are controlled by the deforestation rate and the climate conditions (Mortan et al., 2008; van der Werf et al., 2009; Le Page et al., 2010b), so the burned area is estimated as:

$$A_b = bf_{lu}f_{cli,d}A_g \quad (5)$$

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where b ((time step)⁻¹) is a global constant; f_{lu} and $f_{cli,d}$ represent the effects of decreased coverage fraction of tree PFTs derived from land use data and climate conditions on the burned area, respectively, and vary between 0.0 and 1.0.

The constant b and f_{lu} are calibrated in the Amazon rainforest (tropical closed forests within 15.5° S–10.5° N, 30.5° W–91° W). There are two reasons for our region selection: (i) Amazon rainforest is the dominant tropical deforestation region (Hansen et al., 2008); (ii) the peat accumulation and cropland coverage are low in this region (GACGC, 2000; Hurtt et al., 2006), so the effects of peat fires and agricultural fires on the estimations of b and f_{lu} are small.

The constant $b = 0.03 \text{ d}^{-1}$ is based on Eq. (5) and the following three datasets: atmospheric observations (Qian et al., 2006), CLM4's land use and land cover change data (Oleson et al., 2010), and the GFED3 burned area (Giglio et al., 2009). The multi-year (1997–2004) averages of these data are regridded to the T62 resolution of the Qian et al. (2006) data.

f_{lu} is defined as

$$f_{lu} = \max(0.0005, 0.19D - 0.0009). \quad (6)$$

Equation (6) can explain 83 % of the spatial variability of the 1997–2004 average 0.5° GFED3 annual burned area fraction in grid cells with annual decreased tree coverage D (yr⁻¹) in the Amazon rainforest, and reflects that the burned area in tropical closed forests generally increases with greater losses of tree coverage (Fig. 4). D (yr⁻¹) is based on CLM4's land use and land cover change data.

Fires in tropical closed forests only occur during the dry season, generally 2 months after the wet season has ended (Schroeder et al., 2005). On an intraseasonal time scale, 10-day wet periods in the dry season can lead to a drop in the fire activity levels (Le Page et al., 2010b). To represent the impact of climate on fires, we use 60-day running mean of precipitation (P_{60d} , mm d⁻¹) to reflect the desiccation dynamics of these slashed trees during the dry season (Carvalho et al., 2001), and use the 10-day running mean of precipitation (P_{10d} , mm d⁻¹) to reflect the rapid variability of superficial

Improved global fire modeling in the Community Earth System Model

F. Li et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

moisture due to the daily weather (Uhl and Kauffman, 1990; Holdsworth and Uhl, 1997; Ray et al., 2005). We also assume that humans do not use fires to clear biomass in deforestation regions during the time step when precipitation (P , mm d^{-1}) is greater than the level of drizzle (0.25 mm d^{-1}). In total, we parameterize the effect of climate on deforestation fires as:

$$f_{\text{cli,d}} = \max \left[0, \min \left(1, \frac{b_2 - P_{60\text{d}}}{b_2} \right) \right]^{0.5} \max \left[0, \min \left(1, \frac{b_3 - P_{10\text{d}}}{b_3} \right) \right]^{0.5} \max \left[0, \min \left(1, \frac{0.25 - P}{0.25} \right) \right], \quad (7)$$

where b_2 (mm d^{-1}) and b_3 (mm d^{-1}) are the grid-cell dependent thresholds of $P_{60\text{d}}$ and $P_{10\text{d}}$, respectively. Le Page et al. (2010b) analyzed the relationship between large-scale deforestation fire counts and precipitation during 2003–2006 in Southern Amazonia where tropical evergreen trees (BET Tropical) are dominant. Figure 2 in Le Page et al. (2010b) showed that fires generally occurred if both $P_{60\text{d}}$ and $P_{10\text{d}}$ were less than about 4.0 mm d^{-1} , and fires occurred more frequently in a drier environment. Based on the 30-yr (1985–2004) precipitation data in Qian et al. (2006), the climatological precipitation of dry months ($P < 4.0 \text{ mm d}^{-1}$) in a year over tropical deciduous tree (BDT Tropical) dominated regions is 46% of that over BET Tropical dominated regions, so we set the PFT-dependent thresholds of $P_{60\text{d}}$ and $P_{10\text{d}}$ as 4.0 mm d^{-1} for BET Tropical and 1.8 mm d^{-1} ($= 4.0 \text{ mm d}^{-1} \times 46\%$) for BDT Tropical, and b_2 and b_3 are the average of thresholds of BET Tropical and BDT Tropical weighted by their coverage.

Tropical deforestation fires usually run out of the conversion regions due to weak incentives and/or low ability to control fires (Cochrane, 2003; Baker and Bunyavejchewin, 2009), so, unlike Kloster et al. (2010), we do not limit the post-fire areas due to deforestation in land-type conversion regions. In CLM4 (Oleson et al., 2010; P. E. Thornton, personal communication, 2011), the carbon and nitrogen loss in the tree-reduced region is distributed into a wood production pool with a residence time of 100 yr, a paper

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



production pool with a residence time of 10 yr, and a conversion flux released to the atmosphere immediately; and the fraction assigned to the conversion flux is set to 60 % for tropical trees following Houghton et al. (1983). In conversion regions, the maximum fire carbon emissions are assumed to be 80 % of the total conversion flux as Kloster et al. (2010). To reach the maximum fire carbon emissions in a conversion region requires burning this region about twice ($80 \% \times 60 \% \approx 0.3 + (1 - 0.3) \times 0.3$) when we set combustion completeness factor to 0.3 for stem (the mean of 0.2–0.4 used in van der Werf et al., 2010). Therefore, when the burned area calculated from Eq. (5) is no more than twice the tree-reduced area, we assume no escaped fires outside the land-type conversion region, and the fire-related fraction of the total conversion flux is estimated as $\frac{A_b/A_g}{2D}$. Otherwise, 80 % of the total conversion flux is assumed to be fire carbon emissions, and the biomass combustion and vegetation mortality outside the tree-reduced regions with an area fraction of $\frac{A_b}{A_g} - 2D$ are set the same as Li et al. (2012). Emission of trace gases and aerosols are estimated using the emission factors in Tables 3 and 4 of Li et al. (2012).

3.3 Peat fires

The burned area due to peat fires is given as:

$$A_b = cf_{\text{cli,p}}f_{\text{peat}}(1 - f_{\text{sat}})A_g, \quad (8)$$

where the constant c (time step)⁻¹ is derived using an inverse method (see Sect. 4), $f_{\text{cli,p}}$ represents the effect of climate on the burned area; f_{peat} is the fractional coverage of peatland in the grid cell; and f_{sat} is the fraction of the grid cell with a water table at the surface or higher.

The combustion of tropical peatland is strongly controlled by climate (Field et al., 2009; Page et al., 2009). Similar to deforestation fires, peat fires occur late in the dry season, which represents the long-term memory of peat soil to the wet-season climate (Page et al., 2009). Peat fires are ground fires. Different from deforestation fires, peat

fires may persist for a very long time and are not so sensitive to intraseasonal and real-time precipitation (Cochrane and Ryan, 2009). Accordingly, we set $f_{\text{cli,p}}$ as a function of long-term precipitation $P_{60\text{d}}$:

$$f_{\text{cli,p}} = \max \left[0, \min \left(1, \frac{4 - P_{60\text{d}}}{4} \right) \right]^2, \quad (9)$$

and neglect the effects of intraseasonal and real-time precipitation on peat fires. In southern Central Kalimantan, Borneo, Indonesia, Eq. (9) simulates an obvious peak in 1997 and a smaller one in 2002 during 1997–2004, consistent with the observed inter-annual variability of peat fires (Page et al., 2009); in 1997, Eq. (9) peaks in September/October, consistent with the observed seasonality of peat fires (Koe et al., 2010).

In boreal peatlands, fires generally do not occur in frozen peat, and drier and/or warmer conditions are likely to increase the risk of peat burning (Turetsky et al., 2004). Thus, we set $f_{\text{cli,p}}$ as

$$f_{\text{cli,p}} = \exp \left(-\pi \frac{\theta_{17\text{cm}}}{0.3} \right) \cdot \max \left[0, \min \left(1, \frac{T_{17\text{cm}} - T_f}{10} \right) \right], \quad (10)$$

where $\theta_{17\text{cm}}$ and $T_{17\text{cm}}$ are the soil wetness and soil temperature of the top 17 cm; $T_f = 273.15\text{ K}$ is the freezing temperature. 17 cm is the observed maximum burned depth of boreal bogs (Benscoter et al., 2011), and also close to the maximum burned depth for boreal organic soil (15 cm) used in van der Werf et al. (2010).

Peat fires lead to peat combustion and the combustion and mortality of vegetation in peatlands. For tropical peat fires, based on Page et al. (2002), about 18 % of the peat soil carbon is emitted with 1 % of peatland burned (ref: about 6 % of the peat carbon loss from store caused by 33.9 % of the peatland burned, $6\%/33.9\% = 18\%$). Correspondingly, carbon emissions due to peat combustion ($\text{gC m}^{-2} (\text{time step})^{-1}$) are set as the product of 0.18, burned area fraction of peat fire ($(\text{time step})^{-1}$), and soil organic carbon (gC m^{-2}). For boreal peat fires, the carbon emissions due to peat combustion

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are set as 2.2 kg C m^{-2} peat fire area, based on Turetsky and Wieder (2001). The emissions of trace gases and aerosols due to peat combustion are estimated based on the emission factors of peat fires from van der Werf et al. (2010) (Table 3). The biomass combustion and related emissions of trace gases and aerosols, and vegetation mortality in post-fire peatlands are set similar to those for tree, grass, shrub PFTs in Li et al. (2012).

3.4 Socioeconomic influence on fires

Humans influence fire counts not only by adding ignition sources (intentionally and accidentally), but also by suppressing both anthropogenic and natural fires. Earlier studies (Arora and Boer, 2005; Pechony and Shindell, 2009; Thonicke et al., 2010; Kloster et al., 2010; Li et al., 2012) only considered the impact of population density on fire occurrence. In reality, the economic conditions may also affect fire regimes (Chuvieco et al., 2008; Aldersley et al., 2011). In the present study, the parameterizations of economic impact on fire occurrence and socioeconomic impact on fire spread are added to Li et al. (2012) for non-peat fires outside cropland and tropical closed forests. For scarcely populated regions ($D_p \leq 0.1 \text{ person km}^{-2}$), e.g. Northern Australia and parts of the boreal forest, fires are usually regarded as posing a very low risk to human life and infrastructure, no matter how well developed the country, so the fire management is very limited (Mollicone et al., 2006; Gill et al., 2009; Le Page et al., 2010a). Thus, we assume no economic influence on fire occurring and socioeconomic influence on fire spreading in regions of $D_p \leq 0.1 \text{ person km}^{-2}$. In regions of $D_p > 0.1 \text{ person km}^{-2}$, we parameterize the socioeconomic influence on fires as follows.

For shrub and grass PFTs, the economic influence on fire occurrence is parameterized as

$$f_{\text{eo}} = 0.1 + 0.9 \times \exp \left[-\pi \left(\frac{\text{GDP}}{8} \right)^{0.5} \right], \quad (11)$$

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which captures 73 % of the observed MODIS fire counts with variable GDP in regions where shrub and grass PFTs are dominant (fractional coverage of shrub and grass PFTs > 50 %) (Fig. 5a). In regions outside tropical closed forests and dominated by trees (fractional coverage of tree PFTs > 50 %), we could not find a skillful continuous function to fit the economic impact. Therefore, we divide the MODIS fire counts into only two bins (GDP ≥ 20 k 1995 US\$ capita⁻¹, and GDP < 20 k 1995 US\$ capita⁻¹) and parameterize the economic influence on fire occurring for tree PFTs as

$$f_{eo} = \begin{cases} 0.39, & \text{GDP} \geq 20 \\ 1, & \text{GDP} < 20 \end{cases} \quad (12)$$

to reproduce that the MODIS fire counts in tree-dominated regions of GDP ≥ 20 k 1995 US\$ capita⁻¹ is 39 % of that in other tree-dominated regions (Fig. 5b). Equations (11) and (12) reflect that more developed countries are better at fire detection and fuel management and thus have superior fire suppression. Moreover, Fig. 5 shows the weaker impact of economic conditions on fires for tree PFTs compared with that for grass and shrub PFTs. It is probably because the tree canopy generally conceals surface fires (the most common fire type) better and the tree trunks make fuel management more difficult, which does not vary much with regional economic situations.

The fire fighting capacity, which depends on socioeconomic conditions, affects the fire spread area. Due to a lack of observations, we consider the socioeconomic impact on the average burned area of a fire rather than separate fire spread rate and fire duration. The socioeconomic effect on fire spread area is given as:

$$F_{se} = F_d F_e, \quad (13)$$

where F_d and F_e are effects of the demographic and economic conditions on the average spread area of a fire, and are derived in a similar way to Eqs. (3) and (4).

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For shrub and grass PFTs, the demographic impact is

$$F_d = 0.2 + 0.8 \times \exp \left[-\pi \left(\frac{D_p}{450} \right)^{0.5} \right], \quad (14)$$

and the economic impact factor is

$$F_e = 0.2 + 0.8 \times \exp \left(-\pi \frac{\text{GDP}}{7} \right) \quad (15)$$

5 (Fig. 6a, b). For tree PFTs outside tropical closed forests, the demographic and economic impact factors are given as

$$F_d = 0.4 + 0.6 \times \exp \left(-\pi \frac{D_p}{125} \right), \quad (16)$$

and

$$F_e = \begin{cases} 0.62, & \text{GDP} > 20 \\ 0.83, & 8 < \text{GDP} \leq 20 \\ 1, & \text{GDP} \leq 8 \end{cases} \quad (17)$$

10 (Fig. 7a, b). Equations (14)–(17) reflect that more developed and more densely populated regions have a higher fire fighting capability.

3.5 Other minor modifications

First, effect of temperature on fire occurrence is taken into account to improve the fire seasonality in boreal regions, which is given as:

$$15 \quad f_T = \max \left[0, \min \left(1, \frac{T + 10}{10} \right) \right]. \quad (18)$$

16770

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This is consistent with the fact that temperature is the main constraint for fire occurrence when temperature is below freezing, and fire generally does not occur when the temperature is less than -10°C (Chen and Chen, 2000; Aldersley et al., 2011). Thonicke et al. (2001) also used freezing temperature as the upper temperature threshold for fire occurrence. Second, due to introducing the effects of economic conditions and temperature on fire occurrence, constant a in Eq. (5) of Li et al. (2012), which is the number of potential ignition sources per person per month, is changed to 0.0035 (count person $^{-1}$ mon $^{-1}$) according to the method introduced in Appendix A of Li et al. (2012). After adding the influence of economic conditions and temperature on fire occurrence, the explained variance of observed fire counts is improved to 82 % from original 67 % for the sample used in Appendix A of Li et al. (2012). When the socioeconomic influence on the average spread area for a fire is introduced, the meaning of u_{max} in Eq. (14) of Li et al. (2012) is changed to the expected maximum average fire spread rate not only when wind speed and fuel wetness are optimal for fire spread but also when there is no anthropogenic suppression. Correspondingly, u_{max} increases to 0.47 ms^{-1} for grass PFTs, 0.43 ms^{-1} for shrub PFTs, 0.40 ms^{-1} for needleleaf tree PFTs, and 0.37 ms^{-1} for other tree PFTs.

4 Application and CESM1 simulations

CESM is a global coupled Earth system model hosted at the National Center for Atmospheric Research (NCAR). CESM version 1 (CESM1) and its precursor CCSMs are used for Intergovernmental Panel on Climate Change (IPCC, <http://www.ipcc.ch/>) global change research. CESM1 provides state-of-the-art computer simulations of the Earth's past, present, and future climate states and is a platform to quantify and understand various Earth system processes and interactions. It contains five separate models simultaneously simulating the Earth's atmosphere, ocean, land, land-ice, and sea-ice, plus a central coupling component. The active (dynamical) components (CAM5, POP2, CLM4, CISM, or CICE4) are generally fully prognostic, and are state-of-the-art

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



climate prediction and analysis tools. A detailed description of CESM1 and its code can be found on its homepage <http://www.cesm.ucar.edu/index.html>. Presentations in the CESM annual workshops (<http://www.cesm.ucar.edu/events/workshops.html>) and the CESM1 special issue in the Journal of Climate (<http://journals.ametsoc.org/page/CCSM4/CESM1>) showed that CESM1 can broadly reproduce the observed mean and variability of many primary variables.

In CESM1, the fire dynamics module operates within the CLM4 carbon/nitrogen biogeochemistry model. The CLM4 succeeds CLM3.5 with updates to the soil hydrology, soil thermodynamics, snow dynamics, albedo parameters, the land surface type dataset, the river transport models, capability to apply transient land use and land cover change (LULCC) including wood harvesting, a new urban canyon model, and several other minor modifications (Oleson et al., 2010; Lawrence et al., 2011). The modeling of C/N dynamics in CLM4 is based on the Biome-BGC model, which estimates the states and fluxes of carbon and nitrogen for vegetation, litter, and soil organic matter, and the associated exchange with the atmosphere (Thornton et al., 2007; P. E. Thornton, personal communication, 2011). To generalize plant function to the global scale, CLM4 represents vegetation as 17 plant functional types (PFTs): 8 tree PFTs, 3 shrub PFTs, 3 grass PFTs, 2 crop PFTs, and bare soil PFT (Table 2).

When the new fire model is applied in CESM1, parameters a_1 in Eq. (1) and c in Eq. (8) are estimated using an inverse method, similar to the estimation of the propensity of people to produce ignition events in SPITFIRE (Thonicke et al., 2010). Specifically, they are derived by matching simulations with the following references: the long-term average contribution of cropland fires was 4.7% of the total global burned area (van der Werf et al., 2010); about 2.4 Mha peatland was burned over Indonesia in 1997 (Page et al., 2002); the average burned area of peat fires in Western Canada (Northwest Territories, Alberta, British Columbia, Saskatchewan) was 0.2 Mha yr^{-1} for 1980–1999 (Turesky et al., 2004).

CESM1 supports various component sets. According to a suggestion in the CESM1 user guide, the transient component set I20TRCN (rather than component sets which

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

use full coupled CESM1) is used to evaluate the new fire parameterization. I20TRCN is a 20th century simulation (1850–2004) with the CLM4 driven with the atmospheric observations, transient CO₂ concentration, transient nitrogen and aerosol deposition, and transient land use and land cover changes (wood harvesting included). The atmospheric observations are obtained by cycling 25-yr (1948–1972) Qian et al. (2006) and bias-corrected relative humidity for 1850–1949 followed by the full time series for the years 1950–2004, and 3-h lightning data which are derived from NASA LIS/OTD 2-h climatological lightning data before May 1995 and NASA LIS/OTD daily lightning time series and the 2-h climatological lightning data from May 1995 to December 2004. Other input data include the annual 1850–2004 population density that is obtained by temporal linear interpolation of the data from HYDEv3.1 (prior to 1990) and GPWv3 (since 1990), the present-day GDP data and peat map. The simulations are run globally at a 1.9° × 2.5° spatial resolution with a half-hourly temporal resolution. The input data are regridded to match the requirements of the CESM1 run. In I20TRCN run, $a_1 = 0.5 \text{ h}^{-1}$, and $c = 8.2 \times 10^{-4} \text{ h}^{-1}$ for tropical peat fires and $c = 2.2 \times 10^{-5} \text{ h}^{-1}$ for boreal peat fires. Moreover, in the present study, the climatological peak month of agricultural fires from van der Werf et al. (2010) is used to specify the month when cropland fires may occur.

5 Evaluation of fire simulations

The CESM1 simulations with the fire parameterization introduced in Sect. 3 (Mod-new) are evaluated using the GFED3 fire product. Performance of the fire parameterizations are assessed by evaluating the simulations of burned area and fire carbon emissions (two primary fire variables in ESMs). Burned area not only determines the fire carbon emissions (an important item for the calculation of land-atmosphere net carbon exchange), but also influences other carbon/nitrogen fluxes and pools as well as the vegetation composition and structure. The evaluation period is 1997–2004, which is the common period between GFED3 and the simulations. In addition, Mod-new is

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compared against CESM1 simulations with its current fire parameterization (a modified version of Glob-FIRM) (Mod-old) and modified CTEM-FIRE (Mod-CTEM). The code and description for the current fire parameterization in CESM1 can be obtained from <http://www.cesm.ucar.edu/models/cesm1.0/>; while the modified CTEM-FIRE was described by Kloster et al. (2010) in detail.

5.1 Burned area

Figure 8a shows the GFED3 and simulated global total of annual burned area averaged over 1997–2004. The mean annual global burned area with the new fire module is 338 Mha yr^{-1} , close to the GFED3 (380 Mha yr^{-1}). Relative to GFED3, both Mod-old (167 Mha yr^{-1}) and Mod-CTEM (178 Mha yr^{-1}) tend to underestimate the global burned area by at least 50%. With the new fire scheme, the contributions of four components to the 1997–2004 average global total are: 4.7% from agricultural fires, 2.9% from tropical deforestation and degradation fires, 92% from non-peat fires outside cropland and tropical closed forests, and 0.4% from peat fires. The estimated burned area in cropland is 12.4% with the old fire scheme which does not distinguish between agricultural fires and non-agricultural fires; whereas the modified CTEM-FIRE assumes no fires in cropland. The old one does not model deforestation fires explicitly, and the modified CTEM-FIRE simulates the contribution of global deforestation fires as 6.3%. Both old fire scheme and modified CTEM-FIRE do not include peat fires.

Figure 8b shows the spatial dispersion/heterogeneity of the annual burned area fraction on the global land surface grids at 1.9° (lat) \times 2.5° (lon) spatial resolution, which is quantified using the coefficient of variation, CV (the ratio of the standard deviation to the mean). The CV for Mod-new is 3.6, which is closer to GFED3 (CV = 4.5) than Mod-old (CV = 2.7) and Mod-CTEM (CV = 2.3), although all three schemes underestimate the spatial heterogeneity of the annual burned area fraction.

Mod-new also reproduces the main features of the global spatial distribution of the annual burned area fraction (Fig. 9). It correctly captures the high burned area fraction in tropical savannas, the moderate fraction in Northern Eurasia, and the low fraction

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in deserts due to low fuel availability and in humid forests due to low fuel combustibility. Mod-new shows an improved simulation of the spatial pattern relative to the Mod-old and Mod-CTEM, especially in the tropics. The global spatial correlation between GFED3 and the simulations increases from $Cor = 0.23$ for Mod-old and $Cor = 0.44$ for Mod-new to $Cor = 0.68$ for Mod-new.

The African continent contains the majority of global burned area, contributing 68 % of global total amount for 1997–2004 based on GFED3 (134 $Mha\text{yr}^{-1}$ for Northern Hemisphere Africa and 124 $Mha\text{yr}^{-1}$ for Southern Hemisphere Africa). All of the simulations have a low bias over Africa, but Mod-new (120 $Mha\text{yr}^{-1}$, 84 $Mha\text{yr}^{-1}$) simulates the high burned area in Africa better than Mod-old (18 $Mha\text{yr}^{-1}$, 38 $Mha\text{yr}^{-1}$) and Mod-CTEM (24 $Mha\text{yr}^{-1}$, 45 $Mha\text{yr}^{-1}$) (Table 4). For Amazon tropical closed forests where deforestation and degradation fires are primary, the relative mean squared error (MSE divided by the variance of GFED3 annual burned area fraction) is 0.39 for Mod-new, 0.72 for Mod-old, and 2.72 for Mod-CTEM (Fig. 10), justifying the improved simulation of tropical deforestation fires using the scheme described in Sect. 3.2.

It is noted that in the high-latitudes of North America and Eastern Siberia, Mod-new does not substantially improve the fire simulation compared with the other two fire schemes. Mod-new underestimates the burned area as Mod-CTEM, whereas Mod-old generally overestimates the burned area (Fig. 9). Mod-new and Mod-CTEM underestimate the burned area in boreal forests (moisture-limited fire regime region) partly due to the wet simulation bias of land surface in CLM4 (Figs. 9, 10 in Bonan et al. (2011) showed that CLM4 clearly underestimated the latent heat flux of boreal forests in Eastern Siberia and North America, especially during the fire season (summer). The latent heat flux is directly proportional to the water flux output from the land to the atmosphere, so underestimation of the latent heat flux means overestimation of the water retained by the land). Underestimation of the biomass due to CLM4's cold and dry simulation bias of soil over boreal tundra (Lawrence et al., 2011) where fuel load is limited factor of fires in summer partly explains the underestimation of the burned area simulated by the Mod-new and Mod-CTEM (H. Lee, personal communication, 2011).

Improved global fire modeling in the Community Earth System Model

F. Li et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In terms of the fire seasonality, GFED3 and the simulations with all three fire parameterizations produce the same dominant pattern, in which the peak month for burned area varies between the dry season in the tropics and the warm season in extratropical regions (Fig. 11). The transition from tropical to extratropical patterns reflects the effects of precipitation seasonality reinforced by the fact that climatic conditions in high latitudes are generally too cold to burn in winter (Prentice et al., 2011). The introduction of the agricultural fire scheme allows Mod-new to simulate the peak month in Eastern China, Western Russia, and Northern India better than Mod-old and Mod-CTEM. For central and Eastern North America (crop-dominated region), Mod-old and Mod-CTEM overestimate the annual burned area and they also simulate August/September as the peak month rather than April which is the observed peak month. Moreover, Mod-new identifies the peak month of burned area accurately in tropical closed forests, while Mod-CTEM tends to delay the peak month by 1–2 months in the tropical closed forests of the Southern Hemisphere.

Next, we test the simulated global spatial pattern of fire interannual variability, by using the standard deviation of the annual burned area fraction (Fig. 12). The new fire parameterization and GFED3 show generally similar patterns, e.g. a high interannual variation in the tropical savannas, a medium variation in Northeastern China and Northwestern Eurasia, and a low variation in the deserts and humid forests. The global spatial correlation between GFED3 and the simulations increases from $Cor = 0.14$ for Mod-old and $Cor = 0.26$ for Mod-CTEM to $Cor = 0.48$ for Mod-new. Figure 13 shows interannual variability of the global burned area from GFED3 and simulations. Mod-new and Mod-CTEM can capture the peak in 1998 shown in GFED3, but they overestimate the decline from 1998 to 1999 as LPX-DGVM (Prentice et al., 2011). Mod-new also reproduces the year-to-year variation from 2000 to 2004. The temporal correlation between Mod-new and GFED3 is 0.69, which is higher than the Mod-old (0.08) and Mod-CTEM (0.25).

5.2 Fire carbon emissions

The 1997–2004 average global fire carbon emissions simulated by the three fire parameterizations are 2.1 PgCyr^{-1} for Mod-new, 2.0 PgCyr^{-1} for Mod-CTEM, and 2.3 PgCyr^{-1} for Mod-old, close to GFED3 (2.1 PgCyr^{-1}) (Fig. 14a). The old scheme and CTEM-FIRE simulate the global total of fire carbon emissions reasonably by arbitrarily setting high combustion completeness of woody biomass in post-fire regions, when their simulated global burned areas are less than half the observed. For tree PFTs, the old scheme sets combustion completeness factor to 0.88 for fine root and live coarse roots and 0.35 for dead coarse roots, which are inconsistent with field observations showing that tree roots in post-fire regions are generally little combusted for carbon emissions to atmosphere directly due to a lack of oxygen below ground (Zhou and Lu, 2009). Indeed, GFED3 (van der Werf et al., 2010) and the global fire models of Arora and Boer (2006), Kloster et al. (2010), Li et al. (2012), and the present study consider the root mortality and assume no combustion of roots. The modified CTEM-FIRE (Kloster et al., 2010) set combustion completeness factor as 0.5 and 0.5–0.6 for tree stems and course woody debris, respectively, higher than and on the upper side of combustion completeness factors used by GFED3 (0.2–0.4 for tree stems and 0.4–0.6 for course woody debris). As shown in Fig. 14b, the combustion completeness for Mod-old (13.8 TgCMha^{-1}) and Mod-CTEM (11.2 TgCMha^{-1}) are more than double that for GFED3 (5.5 TgCMha^{-1}), and higher than Mod-new (6.1 TgCMha^{-1}).

The new fire scheme also provides fire carbon emissions from different sources (Fig. 15). 1997–2004 average contributions of the four components to the global total are: 8% from agricultural fires, 27% from tropical deforestation and degradation fires, 60% from non-peat fires outside cropland and tropical closed forests, and 5% from peat fires (3.7% from tropical peat fires). The contribution of agricultural waste burning is within the range of earlier assessments: $\sim 16\%$ from Yevich and Logan (2003) based on government statistics, energy assessments from the World Bank, technical reports and discussions with experts in various research fields in 1985 and 1995; 3%

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from satellite-based estimates of van der Werf et al. (2010) for 2001–2009; and ~ 10 % based on MODIS agricultural fire counts (Korontzi et al., 2006), global fire carbon emissions, and assumption of equal emissions for each fire count for 2001–2003 (van der Werf et al., 2010). The contributions of tropical deforestation and degradation fires and peat fires are similar to the satellite-based estimates of van der Werf et al. (2010) for 2001–2009: 20 % from tropical deforestation and degradation fires and 3 % from tropical peat fires. In Mod-CTEM, the contribution of global deforestation fires is 11 %.

For global spatial patterns, the new fire module can reproduce the high carbon emissions in African and South American savannas, the moderate carbon emissions in Canada and around 50° N in Eurasia, and the low emissions in desert and frozen soil regions (Fig. 16). Mod-new simulates the emissions in boreal Asia and boreal North America better than Mod-CTEM, although both underestimate the amount of carbon emissions. Mod-old overestimates the number of grid cells with fire carbon emissions. The global spatial correlation between Mod-new and GFED3 is 0.52, higher than that for Mod-old (Cor = 0.39) and Mod-CTEM (Cor = 0.32). In addition, all simulations produce similar interannual variability of global fire carbon emissions to GFED3, i.e. peaks in 1997 and 1998 followed by a decline (Fig. 17). The correlation between the simulations and GFED3 are 0.9 for the Mod-new and Mod-CTEM and 0.86 for the old fire parameterization.

6 Direct anthropogenic influence on historical fire regimes

Concerns have grown about the relative importance of climate change and human influence in shaping historical fire regimes (Marlon et al., 2008; Lehsten et al., 2009; Pechony and Shindell, 2009, 2010; Kloster et al., 2010; Archibald et al., 2012). However, the direct anthropogenic impact (fire ignition and suppression) on global fire regime that varies with climate change remain unclear.

Here we report our results based on CESM1 with the new fire scheme. Simulations with (default) and without direct anthropogenic influence are used to quantify the direct

anthropogenic influence on fires. In the simulation without direct anthropogenic influence, the deforestation fires, crop fires, anthropogenic ignitions, anthropogenic suppression, and peat fires due to anthropogenic ignitions are turned off. The fraction of non-peat burned area due to anthropogenic ignitions in the default simulation is applied to separate the anthropogenic peat fires from the natural peat fires.

Figure 18 shows the long-term trends of global burned area and fire carbon emissions from the two simulations. In the default simulation, global burned area and fire carbon emissions peak at 1873 and 1871, respectively, and then abruptly decline. This agrees with the reconstructed global biomass burning based on sedimentary charcoal records by Marlon et al. (2008), and is supported by the analysis of Antarctic ice-core CO (concentration and isotopic ratios) records (Wang et al., 2010; Prentice, 2010). The reconstructed global biomass burning based on Antarctic ice-core CH₄ records (Ferretti et al., 2005) also showed a rise from 1800 to 1870, but noted a sharp rise after 1870 which was contrary to Marlon et al. (2008), Wang et al. (2010), and our results. Houwelling et al. (2008) questioned Ferretti et al. (2005)'s source partitioning of historical variation of CH₄ emissions. The results of Houwelling et al. (2008) showed that biomass burning was not the main contributor for a sharp upward trend of the CH₄ emissions after 1870. In addition, estimation in Kloster et al. (2010) based on the modified CTEM-FIRE showed a downward trend in fire carbon emissions between 1900 and 1960, which is slighter than the downward trend in Marlon et al. (2008) and our results; and a clear upward trend during the last three decades of the 20th century, which is not observed in Marlon et al. (2008) and our results. The difference is in part because the increasing anthropogenic suppression on fire spreading (accompanying the abrupt growth of population) was not taken into account in the fire model of Kloster et al. (2010) which included only the impact of population on fire occurring.

The simulation without direct anthropogenic influence shows a downward trend in global burned area without the peak at ~1870. After ~1870, though burned area declines in both simulations, the downward trend in the default simulation is much stronger. Linear trends after 1873 are $-4.3\%(10\text{yr})^{-1}$ and $-1.2\%(10\text{yr})^{-1}$ for

Improved global fire modeling in the Community Earth System Model

F. Li et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

simulations with and without direct anthropogenic influence, respectively. This suggests that direct anthropogenic influence is the main driver of long-term trends in global burned area since the middle of the 19th century, consistent with the conclusions of Pechony and Shindell (2010) based on the simulations of fire counts that direct anthropogenic influence on fires controlled the global fire regime since the Industrial Revolution.

For fire carbon emissions, the simulation without direct anthropogenic influence shows abrupt decline before 1871, in contrast to the default simulation, indicating that direct anthropogenic influence is the major driver before 1871. After 1871, both simulations present a downward trend. However, the difference in their trends is smaller than that for global burned area simulations. Linear trends after 1873 are $-3.6\%(10\text{yr})^{-1}$ and $-2.3\%(10\text{yr})^{-1}$ for simulations with and without direct anthropogenic influence, respectively, suggesting that direct anthropogenic influence on fires is not the main driver of long-term trend in fire carbon emissions after ~ 1870 . This is supported by the results of numerical experiments from Kloster et al. (2010) that downward trend of global fire carbon emissions from 1900 to 1960 was mainly caused by reduced fuels as a consequence of land use and wood harvesting (i.e. the indirect anthropogenic influence on fires).

7 Conclusions and future development

The new fire parameterization for the Community Earth system model (CESM) contains four components: agricultural fires, non-peat fires outside cropland and tropical closed forests, tropical deforestation fires, and peat fires. The process-based fire parameterization of intermediate complexity described in Li et al. (2012) is revised by mainly adding economic impact on fire occurrence and socioeconomic impact on fire spread, for use by the third component. The new fire scheme has better structure, parameter estimation and mathematical derivation than the current fire scheme in CESM1 (a modified version of Glob-FIRM) and modified CTEM-FIRE (Li et al., 2012; Para. 3

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in Sect. 1). Moreover, introducing our parameterization for agricultural fires, tropical deforestation fires, peat fires, and the socioeconomic impact on fires into ESMs allows more realistic representation of fires and the simulation of some extreme air pollution events due to fires.

5 The new fire scheme is tested in a 1850–2004 transient simulation with the CESM1’s land component CLM4, and evaluated against the latest satellite-based GFED3 fire product for 1997–2004. Results show that the new fire scheme reasonably simulates the multi-year burned area, fire seasonality, fire interannual variability, and fire carbon emissions; and performs better than the current fire scheme in CESM1 (a modified
10 version of commonly-used Glob-FIRM) and modified CTEM-FIRE.

According to our fire scheme, for 1997–2004, the agricultural biomass burning accounts for 8% of global fire carbon emissions; the tropical deforestation and degradation fires and global peat fires account for only 2.9% and 0.4% of burned area, but 27% and 5% of fire carbon emissions; other fires contributes 60% of global fire carbon
15 emissions. The contributions of fire carbon emissions from various sources are close to previous assessments based on satellite data, government statistics, and other information.

The trends of global fire regimes since the middle 19th century simulated with the new fire parameterization are broadly consistent with earlier reconstructions based on sedimentary charcoal records (Marlon et al., 2008) and Antarctic ice-core CO records
20 (Wang et al., 2010; Prentice, 2010). Based on the new fire scheme, we investigate the direct anthropogenic impact (fire ignition and suppression) on global burned area and fire carbon emissions, by comparing 1850–2004 simulations with and without direct anthropogenic impact. Results show that the sharp increase trend of global burned
25 area before ~ 1870 followed by a clear downward trend are mainly caused by the direct anthropogenic impact. However, for fire carbon emissions, only the sharp upward trend before ~ 1870 is overwhelmingly controlled by the direct anthropogenic impact.

The present study contributes many potential applications. First, one major target of fire sciences is to quantify the role of fire in the Earth system on a global scale

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

part because the GDP data does not reflect the much higher economic level in East China than the country average level. Second, many peatlands in Siberia and peatland in Alaska are classified to other wetland type in GLWD (Lehner and Döll, 2004, a source of our peat map) when compared with Yu et al. (2010). The missing peatland areas may contribute to the underestimation of burned area and fire carbon emissions in these regions (Figs. 9, 16). Also, previous studies focus on the peat management and peat fires in regions where fire regimes are moisture-limited (i.e. the forests over Canada and Indonesia). The mechanisms of peat fires in boreal tundra are unknown. Third, the new fire module for CESM just parameterizes the deforestation and degradation fires in the tropical closed forests. Mechanism analyses and observations about deforestation and degradation fires in other regions will improve our understanding and help to quantify them in the future. In addition, Randerson et al. (2012) estimate the global contribution of small fires and point out that GFED3 may underestimate burned area and fire carbon emissions by missing small fires. This means that Eqs. (3)–(6) and (14)–(17) could be recalibrated (based on methods in Sect. 3) using the Randerson et al. (2012) data rather than GFED3 and results could be compared.

Acknowledgements. This study is co-supported by the State Key Project for Basic Research Program of China (973) under Grant No. 2010CB951801, the Strategic Priority Research Program of Chinese Academy of China under Grant No. XDA05110201, and the Key Program of National Natural Science Foundation under Grant No. 40830103. We are grateful to X.-D. Zeng and Q.-C. Zeng from Institute of Atmospheric Physics, Chinese Academy of Sciences, L. Giglio from NASA Goddard Space Flight Center, J. T. Randerson from University of California, H. Lee from NCAR, Z. Subin from Princeton University, N. M. Mahowald from Cornell University, and S. Kloster from Max Planck Institute for Meteorology for helpful suggestions and discussions.

References

- Aldersley, A., Murray, S. J., and Cornell, S. E.: Global and regional analysis of climate and human drivers of wildfire, *Sci. Total Environ.*, 409, 3472–3481, 2011.
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, 2001.
- Archibald, S., Staver, C., and Levin, S. A.: Evolution of human-driven fire regimes in Africa, *P. Natl. Acad. Sci.*, 109, 847–852, 2012.
- Arora, V. K. and Boer, G. J.: Fire as an interactive component of dynamic vegetation models, *J. Geophys. Res.*, 110, G02008, doi:10.1029/2005JG000042, 2005.
- Baker, P. J. and Bunyavejchewin, S.: Fire behavior and fire effects across the forest landscape mosaics of continental Southeast Asia, in: *Tropical Fire Ecology: Climate Change, Land Use and Ecosystem Dynamics*, edited by: Cochrane, M. A., Springer-Praxis, Heidelberg, Germany, 311–334, 2009.
- Benscoter, B., Thompson, D., Waddington, J. M., Wotton, M., Flannigan, M., de Groot, W., and Turetsky, M. R.: Interactive effects of vegetation, soil moisture, and bulk density on depth of burning of thick organic soils, *Int. J. Wildland Fire*, 20, 418–429, 2011.
- Bonan, G. B., Levis, S., Kergoat, L., and Oleson, K. W.: Landscapes as patches of plant functional types: an integrating concept for climate and ecosystem models, *Global Biogeochem. Cy.*, 16, 1021, doi:10.1029/2000GB001360, 2002.
- Bond, W. J., Woodward, F., and Midgley, G. F.: The global distribution of ecosystems in a world without fire, *New Phytol.*, 165, 525–538, doi:10.1111/j.1469-8137.2004.01252.x, 2004.
- Bond-Lamberty, B., Peckham, S. D., Gower, S. T., and Ewers, B. E.: Effects of fire on regional evapotranspiration in the Central Canadian boreal forest, *Glob. Change Biol.*, 15, 1242–1254, doi:10.1111/j.1365-2486.2008.01776.x, 2009.
- Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Mortiz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth system, *Science*, 324, 480–484, doi:10.1126/science.1163886, 2009.
- Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Cochrane, M. A., D'Antonio, C. M., DeFries, R., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Mack, M., Moritz, M. A., Pyne, S., Roos, C. I., Scott, A. C., Sodhi, N. S., and Swetnam, T. W.: The human dimension of

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

fire regimes on Earth, *J. Biogeogr.*, 38, 2223–2236, doi:10.1111/j.1365-2699.2011.02595.x, 2011.

Carvalho, J. A., Costa, F. S., Veras, C. A. G., Sandberg, D. V., Alvarado, E. C., Gielow, R., Serra, A. M., and Santos, J. C.: Biomass fire consumption and carbon release rates of rain-forestclearing experiments conducted in Northern Mato Grosso, Brazil, *J. Geophys. Res.*, 106, 17877–17887, 2001.

Chen, T.-X. and Chen W.-B.: The relationship between forest fire and meteorological elements and the forest of forest fire grades and forest fire's defence, *Meteo. J. Henan*, 4, 25–26, 2000.

Chen, Y., Randerson, J. T., Morton, D. C., DeFries, R. S., Collatz G. J., Kasibhatla, P. S., Giglio, L., Jin, Y., and Marlier, M. E.: Forecasting fire season severity in South America using sea surface temperature anomalies, *Science*, 334, 787–791, 2011.

Chidumayo, E. N.: A shifting cultivation land use system under population pressure in Zambia, *Agroforest. Syst.*, 5, 15–25, 1987.

Chuvieco, E., Giglio, L., and Justice, C.: Global characterization of fire activity: toward defining fire regimes from Earth observation data, *Glob. Change Biol.*, 14, 1488–1502, doi:10.1111/j.1365-2486.2008.01585.x, 2008.

CIESIN: Gridded population of the world version 3 (GPWv3): Population density grids, Technical report, Socioeconomic Data and Applications Center (SEDAC), Columbia University, Palisades, New York, USA, 2005.

Cochrane, M. A.: Fire science for rainforests, *Nature*, 421, 913–919, 2003.

Cochrane, M. A. and Ryan, K. C.: Fire and fire ecology: Concepts and principles, in: *Tropical Fire Ecology: Climate Change, Land Use and Ecosystem Dynamics*, edited by Cochrane M. A., Springer-PRAXIS, Heidelberg, Germany and Chichester, UK, 25–62, 2009.

Ferretti, D. F., Miller, J. B., White, J. W. C., Etheridge, D. M., Lassey, K. R., Lowe, D. C., MacFarling Meure, C. M., Dreier, M. F., Trudinger, C. M., van Ommen, T. D., and Langenfelds, R. L.: Unexpected changes to the global methane budget over the past 2000 years, *Science*, 309, 1714–1717, 2005.

Field, R. D., van der Werf, G. R., and Shen, S. S. P.: Human amplification of drought-induced biomass burning in Indonesia since 1960, *Natl. Geosci.*, 2, 185–188, doi:10.1038/NGEO443, 2009.

GACGC: World in transition: strategies for managing global environmental risks, German Advisory Council on Global Change, Annual Report 1998, Berlin, 2000.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Justice, C. O., Townshend, J. R. G., Vermote, E. F., Masuoka, E., Wolfe, R. E., Saleous, N., Roy, D. P., and Morisette, J. T.: An overview of MODIS Land data processing and product status, *Rem. Sens. Environ.*, 83, 3–15, 2002.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.

Klein Goldewijk, K., Beusen, A., and Janssen, P.: Long term dynamic modeling of global population and built-up area in a spatially explicit way, *HYDE 3.1, Holocene*, 20, 565–573, 2010.

Kloster, S., Mahowald, N. M., Randerson, J. T., Thornton, P. E., Hoffman, F. M., Levis, S., Lawrence, P. J., Feddesma, J. J., Oleson, K. W., and Lawrence, D. M.: Fire dynamics during the 20th century simulated by the Community Land Model, *Biogeosciences*, 7, 1877–1902, doi:10.5194/bg-7-1877-2010, 2010.

Korontzi, S., McCarty, J., Loboda, T., Kumar, S., and Justice, C.: Global distribution of agricultural fire in croplands from 3 years of Moderate Resolution Imaging Spectroradiometer (MODIS) data, *Global Biogeochem. Cy.*, 20, GB2021, doi:10.1029/2005GB002529, 2006.

Lawrence, P. J. and Chase, T. N.: Representing a new MODIS consistent land surface in the Community Land Model (CLM 3.0), *J. Geophys. Res.*, 112, G01023, doi:10.1029/2006JG000168, 2007.

Lawrence, P. J. and Chase, T. N.: Investigating the climate impacts of global land cover change in the community climate system model, *Int. J. Climatol.*, 30, 2066–2087, 2010.

Lawrence, D., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z. L., Levis, S., Skaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the community land model, *J. Adv. Model. Earth Syst.*, 3, M03001, doi:10.1029/2011MS000045, 2011.

Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296, 1–22, 2004.

Lehsten, V., Tansey, K., Balzter, H., Thonicke, K., Spessa, A., Weber, U., Smith, B., and Arneeth, A.: Estimating carbon emissions from African wildfires, *Biogeosciences*, 6, 349–360, doi:10.5194/bg-6-349-2009, 2009.

Lenihan, J. M. and Neilson, R. P.: Simulating broad-scale fire severity in a dynamic global vegetation model, *Northwest Sci.*, 72, 91–103, 1998.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Le Page, Y., Oom, D., Silva, J. M. N., Jönsson, P., and Pereira, J. M. C.: Seasonality of vegetation fires as modified by human action: observing the deviation from eco-climatic fire regimes, *Global Ecol. Biogeogr.*, 19, 575–588, 2010a.
- Le Page, Y., van der Werf, G. R., Morton, D. C., and Pereira, J. M. C.: Modeling fire-driven deforestation potential in Amazonia under current and projected climate conditions, *J. Geophys. Res.-Biogeosciences*, 115, G03012, doi:10.1029/2009JG001190, 2010b.
- Levis, S., Bonan, G. B., Vertenstein, M., and Oleson, K. W.: The Community Land Model's dynamic global vegetation model (CLM-DGVM): Technical description and user's guide, NCAR Tech. Note TN-459_IA, Terrestrial Sciences Section, Boulder, Colorado, 2004.
- Li, F., Zeng, X. D., and Levis, S.: A process-based fire parameterization of intermediate complexity in a Dynamic Global Vegetation Model, *Biogeosciences*, 9, 2761–2780, doi:10.5194/bg-9-2761-2012, 2012.
- Logan, J. A.: Tropospheric ozone: seasonal behavior, trends and anthropogenic influence, *J. Geophys. Res.*, 90, 10463–10482, 1985.
- Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: a global perspective, *J. Geophys. Res.*, 86, 7210–7254, 1981.
- Luiz, E. O., Aragão, C., and Shimabukuo, Y. E.: The incidence of fire in Amazonian forests with implications for REDD, *Science* 328, 1275–1278, 2010.
- Magi, B. I., Rabin, S., Shevliakova, E., and Pacala, S.: Separating agricultural and non-agricultural fire seasonality at regional scales, *Biogeosciences*, 9, 3003–3012, doi:10.5194/bg-9-3003-2012, 2012.
- Marlon, J. R., Bartlein, P. J., Carcaillet, C., Gavin, D. G., Higuera, P. E., Joos, F., Power, M. J., and Prentice, I. C.: Climate and human influences on global biomass burning over the past two millennia, *Nat. Geosci.*, 1, 697–702, doi:10.1038/ngeo313, 2008.
- Mollicone, D., Eva, H. D., and Achard, F.: Human impact on “wild” fires in boreal Eurasian forests, *Nature*, 440, 436–437, 2006.
- Morton, D. C., DeFries, R. S., Randerson, J. T., Giglio, L., Schroeder, W., and van der Werf, G. R.: Agricultural intensification increases deforestation fire activity in Amazonia, *Glob. Change Biol.*, 14, 2262–2275, doi:10.1111/j.1365-2486.2008.01652.x, 2008.
- New, M., Hulme, M., and Jones, P.: Representing twentieth century space–time climate variability, Pt. I: development of a 1961–90 mean monthly terrestrial climatology, *J. Climate*, 12, 829–856, 1999.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

New, M., Hulme, M., and Jones, P.: Representing twentieth-century space–time climate variability, Pt. II: development of 1901–96 monthly grids of terrestrial surface climate, *J. Climate*, 13, 2217–2238, 2000.

Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C. L., Hoffman, F., Lamarque, J. F., Mahowald, N. M., Niu, G. Y., Qian, T., Rander-
son, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang, Z. L., Zeng, X. D., and Zeng, X.: Technical description of version 4.0 of the community land model, NCAR Tech. Note NCAR/TN-478+STR, Terrestrial Sciences Section, Boulder, Colorado, 2010.

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Under-
wood, E. C., D’Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Heda, P., and Kassem, K. R.: Ter-
restrial ecoregions of the world a new map of life on earth, *Bioscience*, 51, 933–938, 2001.

Ottmar R. D., Sandberg, D. V., Riccardi, C. L., and Prichard, S. J.: An overview of the fuel char-
acteristic classification system – quantifying, classifying, and creating fuelbeds for resource
planning, *Can. J. Forest Res.*, 37, 2383–2393, 2007.

Page, S. E., Siegert, F., Rieley, J. O., Boehm, H.-D. V., Jaya, A., and Limin, S.: The amount of
carbon released from peat and forest fires in Indonesia in 1997, *Nature*, 420, 61–65, 2002.

Page, S. E., Hoscilo, A., Langner, A., Tansey, K. J., Siegert, F., Limin, S., and Rieley, J. O.:
Tropical peatland fires in Southeast Asia, in: *Tropical Fire Ecology: Climate Change, Land
Use and Ecosystem Dynamics*, edited by: Cochrane, M. A., Springer-Praxis, Heidelberg,
Germany, 263–283, 2009.

Pechony, O. and Shindell, D. T.: Fire parameterization on a global scale, *J. Geophys. Res.*, 114,
D16115, doi:10.1029/2009JD011927, 2009.

Pechony, O. and Shindell, D. T.: Driving forces of global wildfires over the past
millennium and the forthcoming century, *P. Natl. Acad. Sci.*, 107, 19167–19170,
doi:10.1073/pnas.1003669107, 2010.

Prentice, I. C.: The burning issue, *Sciences*, 330, 1636–1637, 2010.

Prentice, I. C., Kelley, D. I., Foster, P. N., Friedlingstein, P., Harrison, S. P., and Bartlein, P. J.:
Modeling fire and the terrestrial carbon balance, *Global Biogeochem. Cy.*, 25, GB3005,
doi:10.1029/2010GB003906, 2011.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Qian, T., Dai, A., Trenberth, K. E., and Oleson, K. W.: Simulation of global land surface conditions from 1948 to 2004, Pt I: forcing data and evaluations, *J. Hydrometeorol.*, 7, 953–975, 2006.

Ramankutty, N., Evan, A. T., Monfreda, C., and Foley, J. A.: Farming the planet: 1. geographic distribution of global agricultural lands in the year 2000, *Global Biogeochem. Cy.*, 22, GB1003, doi:10.1029/2007GB002952, 2008.

Randerson, J. T., Chen, Y., van der Werf, G. R., Rogers, B. M., and Morton D.: Small fire contributions to global burned area and biomass burning emissions, *J. Geophys. Res.*, doi:10.1029/2012JG002128, in press, 2012.

Schroeder, W., Morisette, J. T., Csiszar, I., Giglio, L., Morton, D., and Justice, C. O.: Characterizing vegetation fire dynamics in Brazil through multisatellite data: common trends and practical issues, *Earth Interact.*, 9, 1–26, 2005.

Tarnocai, C., Kettles, I. M., and Lacelle, B.: Peatlands of Canada, Geological Survey of Canada, Open File 6561, CD-ROM, doi:10.495/288786, 2011.

Thonicke, K., Venevsky, S., Sitch, S., and Cramer, W.: The role of fire disturbance for global vegetation dynamics: coupling fire into a dynamic global vegetation model, *Global Ecol. Biogeogr.*, 10, 661–677, 2001.

Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L., and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model, *Biogeosciences*, 7, 1991–2011, doi:10.5194/bg-7-1991-2010, 2010.

Thornton, P. E., Lamarque, J., Rosenbloom, N., and Mahowald, N. M.: Influence of carbon-nitrogen cycle coupling on land model response to CO₂ fertilization and climate variability, *Global Biogeochem. Cy.*, 21, GB4018, doi:10.1029/2006GB002868, 2007.

Turetsky, M. R., Wieder, R. K., Halsey, L. A., and Vitt, D. H.: Current disturbance and the diminishing peatland carbon sink, *Geophys. Res. Lett.*, 29, 1526, doi:10.1029/2001GL014000, 2002.

Turetsky, M. R., Amiro, B. D., Bosch, E., and Bhatti, J. S.: Historical burn area in Western Canadian peatlands and its relationship to fire weather indices, *Global Biogeochem. Cy.*, 18, GB4014, doi:10.1029/2004GB002222, 2004.

Uhl, C. and Kauffman, J. B.: Deforestation, fire susceptibility, and potential responses to fire in the Eastern Amazon, *Ecology*, 71, 437–449, 1990.

UNSTAT: National Accounts Main Aggregates Database, United Nations Statistics Division, 2005.

van der Werf, G. R., Morton, D. C., DeFries, R. S., Giglio, L., Randerson, J. T., Collatz, G. J., and Kasibhatla, P. S.: Estimates of fire emissions from an active deforestation region in the southern Amazon based on satellite data and biogeochemical modelling, *Biogeosciences*, 6, 235–249, doi:10.5194/bg-6-235-2009, 2009.

van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Van Vuuren, D. P., Lucas, P. S., and Hilderink, H. B. M., Downscaling drivers of global environmental change: enabling use of global SRES scenarios at the national and grid levels, Report 550025001, Netherlands Environmental Assessment Agency, 45 pp., 2006.

Wang, Z., Chappellaz, J., Park, K., and Mark, J. E.: Large variations in Southern Hemisphere biomass burning during the last 650 years, *Science*, 330, 1663–1666, 2010.

World Bank: World development indicators 2004, Oxford University Press, New York, 416 pp., 2004.

World Health Organization (WHO): Vegetation fires, Geneva, Switzerland, Fact Sheet 254, 2000.

Zeng, X. D.: Evaluating the dependence of vegetation on climate in an improved dynamic global vegetation model, *Adv. Atmos. Sci.*, 27, 977–991, doi:10.1007/s00376-009-9186-0, 2010.

Zeng, X. D., Zeng, X., and Barlage, M.: Growing temperate shrubs over arid and semiarid regions in the NCAR dynamic global vegetation model (CLM-DGVM), *Global Biogeochem. Cy.*, 22, GB3003, doi:10.1029/2007GB003014, 2008.

Zhou, G.-S. and Lu, Q.: Meteorology and Fires in Forests and Grassland, China Meteorological Press, Beijing, 2009.

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Datasets used to drive CLM-DGVM and evaluate simulations.

Variables	Sources	Resolutions
Burned area Fire emissions	GFED3 (Giglio et al., 2010; van der Werf et al., 2010)	0.5°, monthly
Fire counts	MODIS (Giglio et al., 2006)	0.5°, monthly
Surface air temperature Wind speed Specific humidity Air pressure Precipitation Incoming solar radiation	Qian et al. (2006)	T62 (~ 1.875°), 3-hourly T62 (~ 1.875°), 6-hourly
Relative humidity	NCEP/NCAR (Kalnay et al., 1996); CRU (New et al., 1999, 2000)	2.5°, 6-hourly 10 min, monthly*
Lightning frequency	NASA LIS/OTD v2.2 (http://ghrc.msfc.nasa.gov)	2.5°, 2-hourly* 2.5°, daily
Population density	GPWv3 (CIESIN, 2005); HYDE v3.1 (Klein Goldewijk et al., 2010)	0.5°, 5-yr 5 min, 10-yr
Gross domestic product (GDP)	van Vuuren et al. (2006)	0.5°, in 2000
Peat area fraction	Olson et al. (2001); Tarnocai et al. (2011); GLWD (Lehner and Döll, 2004)	vector, for present day
Peak month of agricultural waste burning	van der Werf et al. (2010)	0.5°, for present day
Land use and land cover change (LULCC) Present-day land cover	CLM4 land surface data	0.47° × 0.63°, annual 0.47° × 0.63°, for present day

* Climatology data.

**Improved global fire
modeling in the
Community Earth
System Model**

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 2.** Plant functional types (PFTs) used in the present study.

PFT	Abbreviation
Trees	
Broadleaf Evergreen Tropical	BET Tropical
Broadleaf Deciduous Tropical	BDT Tropical
Broadleaf Evergreen Temperate	BET Temperate
Needleleaf Evergreen Temperate	NET Temperate
Broadleaf Deciduous Temperate	BDT Temperate
Needleleaf Evergreen Boreal	NET Boreal
Broadleaf Deciduous Boreal	BDT Boreal
Shrubs	
Broadleaf Deciduous Temperate	BDS Temperate
Broadleaf Deciduous Boreal	BDS Boreal
Grasses	
C4	–
C3 Non-arctic	–
C3 Arctic	–
Crop	
Crop 1	–
Crop 2	–

Table A1. List of model variables.

Variable	Description	Unit
a	Potential anthropogenic ignitions	count person ⁻¹ mon ⁻¹
A_b	Burned area	km ² (time step) ⁻¹
A_g	Area of grid cell	km ²
b_2	Threshold for P_{60d}	mm d ⁻¹
b_3	Threshold for P_{10d}	mm d ⁻¹
D_p	Population density	person km ⁻²
D	Decreased tree coverage	yr ⁻¹
f_b	Fuel availability factor	–
$f_{cli,d}$	Effect of climate on deforestation fires	–
$f_{cli,p}$	Effect of climate on peatland burned area	–
f_{crop}	Fractional coverage of cropland	–
f_d	Effects of population density and	–
f_e	GDP on agricultural burned area	–
f_{eo}	Effect of economic situation on fire occurrence	–
f_{lu}	Effect of decreases of tree coverage on burned area	–
f_{peat}	Coverage fraction of peatland	–
f_{sat}	Area fraction with water table higher/at the surface	–
f_{se}	Socioeconomic effect on burned area of agricultural fires	–
f_t	Factor to determine agricultural fire seasonality	–
f_T	Effect of temperature on fire occurrence	–
F_d	Effects of population density and	–
F_e	GDP on average spread area of a fire	–
F_{se}	Socioeconomic effect on fire spread area	–
D	Decreased tree coverage	yr ⁻¹
GDP	Gross Domestic Product	k 1995 US\$ capita ⁻¹
P_{60d}	60-day running mean of precipitation	mm d ⁻¹
P_{10d}	10-day running mean of precipitation	mm d ⁻¹
P	Precipitation	mm d ⁻¹
T_{17cm}	Soil temperature of the top 17 cm	K
T_f	Freezing temperature	K
T	Temperature	K
u_{max}	Maximum average fire spread rate	mm s ⁻¹
θ_{17cm}	Soil wetness of the top 17 cm	mm ³ mm ⁻³

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 1. Fire parameterization of Li et al. (2012). It contains three components: fire occurrence, fire spread, and fire impact.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 2. Structure of new fire parameterization. Fire scheme described in Li et al. (2012) is used in Reg C with modifications by mainly adding the economic influence in the fire occurrence component and the socioeconomic influence in the fire spread component.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
⏪	⏩
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 3. Effects of **(a)** demographic situation f_d , and **(b)** economic situation f_e on burned area fraction (lines). Population density (person km^{-2}) and Gross Domestic Product (GDP, $\text{k1995US\$ capita}^{-1}$) are used as socioeconomic indicators. In regions where area fraction of crop PFTs is larger than 50 %, we partition 1997–2004 average 0.5° GFED3 burned area fraction divided by f_e (i.e. $\frac{A_b}{A_g} \text{GFED3} / f_e$) over 25 person km^{-2} population density bins, and GFED3 burned area fraction divided by f_d (i.e. $\frac{A_b}{A_g} \text{GFED3} / f_d$) over 1 $\text{k1995US\$ capita}^{-1}$ GDP bins. Black circles indicate the (normalized) average in bins with sample size > 5 . Goodness of fit are $R^2 = 0.46$ ($n = 42$) in **(a)** and $R^2 = 0.70$ ($n = 13$) in **(b)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Effects of decreases of tree coverage on annual burned area, f_{lu} (Line). 0.5° GFED3 burned area fraction in Amazon rainforest (tropical closed forest in 15.25° S– 10.25° N, 30.75° – 90.75° W) are partitioned over $0.01\% \text{ yr}^{-1}$ deforestation rate bins. Black circles indicate the average burned area fraction in bins with sample size > 5 . Goodness of fit is 0.83 ($n = 19$).

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper




Fig. 5. Effect of economic situation on fire occurrence, f_{eo} , for **(a)** grass and shrub PFTs and **(b)** tree PFTs (lines). In shrubs and grasses dominated regions (fractional coverage of shrub and grass PFTs > 50%), 2001–2004 average 0.5° MODIS fire counts are partitioned over 1 k 1995US\$capita⁻¹ GDP bins. Black circles in **(a)** indicate the (normalized) average fire counts in bins with sample size > 5. Goodness of fit are $R^2 = 0.73$ ($n = 14$). f_{eo} in **(b)** reproduces that MODIS fire counts in tree-dominated regions of GDP ≥ 20 k 1995US\$capita⁻¹ is 0.39 of that in other tree-dominated regions.

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 6. Effects of **(a)** demographic situation F_d and **(b)** economic situations F_e on average burned area of a fire (Lines). In shrub-and-grass-dominated regions, we partition average burned area of a fire (derived from 2001–2004 GFED3 burned area and MODIS fire counts at 0.5° spatial resolution) divided by F_e (i.e. $a_{\text{MODIS,GFED3}}/F_e$) over 25 person km^{-2} population density bins, and average burned area of a fire divided by F_d (i.e. $a_{\text{MODIS,GFED3}}/F_d$) over 1 k 1995 US\$ capita^{-1} GDP bins. Black circles indicate the (normalized) average in bins with sample size > 5 . Goodness of fit are $R^2 = 0.71$ ($n = 13$) in **(a)** and $R^2 = 0.81$ ($n = 11$) in **(b)**.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 7. Same as Fig. 6, but for tree-dominated regions outside tropical closed forests. Goodness of fit is $R^2 = 0.43$ ($n = 9$) in (a). F_e in (b) reproduces that $a_{\text{MODIS,GFED}}/F_d$ in tree-dominated regions of $\text{GDP} > 20 \text{ k1995US\$ capita}^{-1}$ and $\text{GDP} > 8 \text{ k1995US\$ capita}^{-1}$ is 0.62 and 0.83 of that in other tree-dominated regions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 8. 1997–2004 average **(a)** global annual burned area and **(b)** spatial dispersion of annual burned area fraction on global land surface grids quantified by coefficient of variation CV from GFED3 and CESM1 simulations with the fire parameterization introduced in Sect. 3 (Mod-new), CESM1 current fire parameterization (Mod-old) and modified CTEM-FIRE (Mod-CTEM).

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 9. Spatial distribution of annual burned area fraction averaged over 1997–2004 for GFED3 and CESM1 simulations with different fire parameterizations. The global spatial correlation (Cor) between GFED3 and simulation is also given.

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 10. Comparison of GFED3 and modeled 1997–2004 average annual burned area fraction over Amazon tropical closed forest. The black line depicts the 1 : 1 slope and represents the perfect simulation. Relative mean squared error (MSE divided by the variance of GFED3 annual burned area fraction) is 0.39 for Mod-new, 0.72 for Mod-old and 2.72 for Mod-CTEM.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 11. Same as Fig. 9, but for month of maximum burned area for those areas where burned area fraction > 0.1 %.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 12. Same as Fig. 9, but for standard deviation (Std) of annual burned area fraction which is used as a spatially-explicit measure of fire interannual variability.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 13. Annual global burned area normalized by the mean for 1997–2004 from GFED3 and CESM1 simulations with different fire parameterizations. The numbers in brackets denote temporal correlation between GFED3 and simulations.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 14. 1997–2004 average **(a)** global annual fire carbon emissions and **(b)** combustion completeness in global post-fire regions.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 15. Contribution of agricultural fires, tropical deforestation and degradation fires, non-peat fires outside cropland and tropical closed forests, and peat fires to 1997–2004 global fire carbon emissions from Mod-new.

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 16. Same as Fig. 9, but for annual fire carbon emissions.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 17. Same as Fig. 13, but for normalized global fire carbon emissions.

BGD

9, 16753–16814, 2012

Improved global fire modeling in the Community Earth System Model

F. Li et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Improved global fire modeling in the Community Earth System Model

F. Li et al.




Fig. 18. Historical variation of **(a)** global burned area and **(b)** fire carbon emissions in simulations with and without direct anthropogenic influence. 25-yr low-pass filter is used.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)