

Dear editor,

We would like to thank both referees for their constructive and helpful comments on our manuscript. We detail below point by point how we plan to revise the article for publication in Biogeosciences.

Anonymous Referee #1

My main concern is about the utilization of the steady state model ('Open') to describe a system where there is a seasonal nearly-complete DSi consumption (Kerguelen Plateau). This model assumes steady state conditions in the mixed layer and, therefore, no changes in concentrations with time.

It is correct that the steady state model when it reproduces a flow-through reactor assumes a continuous supply of substrate and thus no change in concentration with time. Theoretically, a series of sequential steady state models would reproduce finally the isotope dynamics of a closed system and allow consumption of substrate (Fry, 2006). It appears that in some oceanic Si or N isotopic systems, the steady state reproduces better the data (e.g. Sigman et al. 1999; Cardinal et al. 2005) than closed Rayleigh model. More particularly, in the case of the Kerguelen Plateau, Fripiat et al. (2011) have already shown that the steady state model better describes the seasonality of the silicon isotopes in the ML and it seems that our data confirm this observation. In their study, authors have suggested that there was a significant ventilation of the ML above the Plateau which supply DSi with low $\delta^{30}\text{Si}$ value into the system. We acknowledge that it is conceptually hard to reconcile a seasonal decrease of concentration and increase of isotopic composition with the steady state but so far, no better model has been shown. In the revised version, we will make clearer this apparent contradiction and raises that we can consider each steady-state observed in the area as a snapshot highlighting a Si supply from below and that the steady state must be broken and formed again when surface concentration is decreasing (i.e. when supply does not exactly compensate uptake)

Authors also look at figure 4 to infer which models fit the most with the observations. I'm not sure that the level of precision is sufficient to distinguish between models. Especially if we take into account the precision on both the fractionation factor and the Si-source: on figure 4, the former is shown only for the steady state model and the latter is not taking into account.

We didn't add the precision on the fractionation factor for the Rayleigh model to avoid overloading the figure but we will add it on fig.4 as for the steady state model. When adding this uncertainty, the productive stations above the plateau (A3-2, E5 and K1) remain clearly out of the Rayleigh fractionation trend. The isotopic composition and the DSi concentration of the Si-source are much better constrained ($\pm 2.2 \mu\text{mol L}^{-1}$ and $\pm 0.03 \text{‰}$, respectively) in this study and do not change the fact that Rayleigh model – contrary to steady state – cannot explain the measured isotopic values especially at high consumption in the late summer. We will mention this in the text of the revised version.

In late summer, the authors also infer that the biogenic silica pool is a mixture between new (high- $\delta^{30}\text{Si}$) and old (low- $\delta^{30}\text{Si}$) biogenic silica (from figure 4). But late summer DSi $\delta^{30}\text{Si}$ is low (figure 4),

and actually not significantly different than biogenic silica $\delta^{30}\text{Si}$ (error bars in figure 4). How is it possible to produce high- $\delta^{30}\text{Si}$ biogenic silica from a low- $\delta^{30}\text{Si}$ pool? I agree that biogenic silica $\delta^{30}\text{Si}$ could be explained by being a mixture between instantaneous and accumulated products in the Rayleigh model ('closed'). But in regard of the figure 4, assuming a constant fractionation factor, such high- $\delta^{30}\text{Si}$ biogenic silica requires a DSi pool being significantly higher than described with the steady state model at any time, or than the observations.

The isotopic dynamics in this region is complex regarding the various processes that can affect them. Several explanations are possible to describe what we observed, but it is difficult to discriminate which one is the more relevant:

- The fractionation factor of diatoms ($^{30}\epsilon$) may vary over the season. Since there is no study yet that focuses on $^{30}\epsilon$ seasonal variations (only one study has highlighted interspecific – but not seasonal – variations, see Sutton et al., 2013), we have considered it as constant, but we are aware that this may not reflect the reality in the ocean and that the constancy of the fractionation factor can be challenged.

- In the ocean, the distinction between steady state and Rayleigh models can be tricky. Indeed, many situations, such as the Kerguelen region, can combine or switch between characteristics of both closed and open system dynamics resulting in changes of the apparent fractionation factor ($\Delta^{30}\text{Si} = \delta^{30}\text{Si}_{\text{DSi}} - \delta^{30}\text{Si}_{\text{BSi}}$). As already proposed by Fripiat et al. (2012) we suggest that these variations are mainly controlled by the Si uptake:Si supply ratio of the system. When Si uptake:Si supply ratio is high (during the bloom period), the system roughly follows a Rayleigh fractionation model. However, when the DSi pool limits Si uptake (at the end of the productive period), the Si uptake:Si supply ratio decreases and the supply of light DSi into the ML by vertical mixing decreases significantly the $\delta^{30}\text{Si}_{\text{DSi}}$. The $\delta^{30}\text{Si}_{\text{BSi}}$ is affected differently since the BSi pool is composed by a mixing between newly formed (and light) diatoms and isotopically heavy diatoms that have been produced previously in the ML. Such process would explain why we observe late summer high $\delta^{30}\text{Si}_{\text{BSi}}$ associated to low $\delta^{30}\text{Si}$ of the DSi pool.

- Other processes such as mixing or BSi dissolution with or without isotopic fractionation, can affect differently the different pools and would lead to variations of the apparent fractionation factor. Mixing brings light DSi into the ML and decreases instantaneously the $\delta^{30}\text{Si}$ of the DSi pool while the dissolution of biogenic silica may affect mainly light and dead diatoms that were produced in the beginning of the bloom resulting in a progressive increase of the $\delta^{30}\text{Si}$ of the BSi pool. Such effects are temporally decoupled and would result in a decoupling of the information recorded in the dissolved and particulate phase (see also our reply to main concern of reviewer 2).

Hence even if these theoretical models seem to be appropriate for interpreting paleoceanographic isotope records that integrate longer temporal scale, it appears that they have limitations when applied to the modern ocean at shorter and seasonal scales.

We agree that this part of the discussion is confusing. We will rearrange the text to make it clearer.

Minor comments:

Pages 8-9 Lines 32-1: What is real WW Si properties? As written, it is not clear.

In this sentence, we will change “real WW Si-properties” by “specific Si concentration and isotopic composition of the WW”.

Page 10 Lines 28-29: Not sure to understand why the location of the meander being upstream of the Kerguelen Plateau is indicative of a delay in the initiation of the bloom.

As shown in Fig.1, and explained in the text, the meander is located downstream of the Plateau. Water comes from the South (see Park et al., 2014 and Sanial et al., 2015) and undergoes a progressive phytoplankton bloom and consumption of nutrients.

Page 11 Lines 9-11: Such accumulation is not seen on BSi concentration profiles? In addition, an accumulation should imply low biogenic silica $\delta^{30}\text{Si}$, no?

Yes, we realize that the interpretation here is inadequate. Actually we do not see subsurface accumulation of heavy BSi here, but a production of light BSi in surface. This lower surface BSi isotope composition could result from the supply of light DSi into the ML by sporadic mixing events. The DSi stock in the ML would have a lower isotopic signature than previously and diatoms would produce lighter BSi at the top of the ML. We will change our discussion (from line 6 to line 18) according to this.

Page 11 Line 30: The authors need to define better the concept behind "new silicic acid" and its relationship with silicon isotopes. As written, it is not clear.

We will remove “new” in this sentence as it can generate confusions with other parts of the discussion.

Page 12 Lines 1-2: the northern Kerguelen shelf...(to be added)...or from the mixing with waters north of the polar front bearing higher DSi $\delta^{30}\text{Si}$ due to the progressive export of low BSi $\delta^{30}\text{Si}$ along the meridional overturning circulation path. From my point of view (given the circumpolar boundary between APF and PFZ), this should be the dominant driver of such increase in subsurface DSi $\delta^{30}\text{Si}$.

We agree, this is what we wanted to say. We will replace our sentence by the referee’s one and add “including from coastal waters”.

Page 12 Line 18: HNLC area in KEOPS 1&2 are located in different areas, East and West of the Kerguelen Plateau. Are they really comparable? It should be discussed somewhere when you highlight the differences and make some hypotheses on the water sources. For example, HNLC WW during KEOPS 1 appear to be actually more representative (from their geographical locations) of this expected source of waters originating from the South. But I agree that measurements show the opposite, as being more similar to the HNLC area west of the Kerguelen Plateau. So, I do not really understand how the authors (latter in this paragraph) infer a source for the Kerguelen Plateau WW coming from the South based on these observations.

We agree that HNLC area in KEOPS 1 and 2 cannot be compared and be used as water sources for the model. We discuss that later (line 23-24 p12). Actually we did not choose a source coming from the South based on the observations. We have defined our source as the WW above the Plateau and discuss that this water mass likely originates from the South (based on hydrological and other biogeochemical observations, see Park et al., 2014 and Sanial et al., 2015). The Si isotopic composition and concentration of this water mass source in the HNLC WW might have been modified during its transport in subsurface from HNLC to the plateau (e.g. by sporadic mixing as seen in the region) and before being

supplied to the ML of the Plateau. This could explain why the WW HNLC seems to be inadequate to characterize Si concentration and isotopic composition of the Si supplied in the ML over the Plateau. In the revised version, we'll more clearly explain this.

Figure 3: DSi concentration between KEOPS1 & 2 would be useful here.

We will add DSi concentrations in this figure as suggested.

Page 13 Line 4: In the next paragraph, the authors indicate that Closset et al. (2014) did not report measurements in the deep silica maximum. There are other studies reporting direct measurements of high D/P in deep silica maximum of the Southern Ocean.

Our wording was not clear in the previous version. Actually Closset et al. (2014) did measure BSi production but they are close to 0 in the deep silica maximum, and thus high D/P ratios. High D/P ratios at depth have also been measured by Fripiat et al. (2011) in the PFZ of the Australian sector of the Southern Ocean. We will clarify this and add this reference to the text.

Page 14 Line 11: But not for biogenic silica $\delta^{30}\text{Si}$, following more closely the Rayleigh trends.

We are sorry, but we did not understand this comment. All the delta values discussed in this section 3.3.2 on WW correspond to DSi and not BSi. The $\delta^{30}\text{Si}$ of BSi that follow more closely a Rayleigh trend, corresponds to the ML pool (and not BSi in the WW).

Figure 4: Why the error propagation for the Rayleigh model is not shown? If presented, I expect that most of the DSi $\delta^{30}\text{Si}$ would still fit with the Rayleigh model. Especially if you take also into account the uncertainties related to the initial Si-source.

As previously discussed, we have excluded the precision on the fractionation factor for the Rayleigh model to avoid overloading the figure. When adding this uncertainty, the productive stations above the plateau (A3-2, E5 and K1) remain clearly out of the Rayleigh fractionation trend.

Page 14 Lines 7-22: I like this paragraph. Maybe it would be worth to say that such process is analog of what is observed for DSi $\delta^{30}\text{Si}$ across the ACC meridional overturning circulation.

We agree and we will mention this point in this paragraph.

Page 14 Lines 32-34: The BSi accumulation/export ratio appears to be the main driver. This should be said more strongly.

Yes, we will underline more this point.

Page 15 line 16: The authors suggest that late summer newly-formed biogenic silica presents high $\delta^{30}\text{Si}$, but the residual DSi $\delta^{30}\text{Si}$ is low (Figure 4). As being the Si-source, how can you produce high- $\delta^{30}\text{Si}$ biogenic silica from this DSi pool (assuming constant fractionation factor)?

Please see the above answer (first page of this document). We will change the text according to this explanation and provide a clearer discussion.

Page 15 Lines 27-29: Why low Si-uptake/supply ratios and a combination of open and closed system (also implying variable Si-uptake/supply ratios) are different approaches? As defined, the open model

assumes that Si-supply equals the sum of BSi accumulation/export and residual DSi (implying also low or close to unity Si-uptake/supply ratio, depending of the relative Si-utilization).

Yes, variations of Si uptake:Si ratio and the combination of open and closed model are not different approaches but are closely linked together. We will correct this part of the text.

Page 15 Lines 29-31: Why the export of biogenic silica will decrease $\Delta^{30}\text{Si}$? It should be the opposite (as discussed previously). As written it is not clear.

We have realized that this part of the discussion is not clear (as raised by the reviewer). We will remove line 27 to 31 and replace it by "Our results point out that an alternative approach to change the $\Delta^{30}\text{Si}$ could be obtained when the BSi pool in the system switches alternatively between the Rayleigh instantaneous or accumulated product depending on the Si production:Si export ratio."

Page 15: Lines 31-35: Same comment than above. The authors suggest that late summer newly-formed biogenic silica presents high $\delta^{30}\text{Si}$, but the residual DSi $\delta^{30}\text{Si}$ is low (Figure 4).

Please see the above answer (first page of this document). We will change the text according to this explanation and provide a clearer discussion.

Page 16 Lines 19-20: Dissolution occurs in subsurface (Nelson et al., 2002, DSR). This process can therefore also explain the observed trends. The authors should be more caution before completely ruling out the dissolution isotope effect.

We agree, dissolution can occur in subsurface. But during KEOPS-2, dissolution rates measured from 0 to 80 m were very low and since we were at the beginning of the productive period, we can expect that they remain low in deep waters. Moreover, even if we can expect some dissolution of biogenic silica in the water column, some recent studies have shown that isotopic fractionation during dissolution did not affect the BSi $\delta^{30}\text{Si}$ along the water column (Fripiat et al., 2012; Closset et al., 2015) and may not occur (Wetzel et al., 2014). This is why we have chosen to rule out the dissolution isotope effect.

Page 18 Lines 27-28: Why here the steady state model assumes, as it should be, no variations in concentrations with time, but it has been previously used to describe a situation over the Kerguelen Plateau exhibiting large variations in DSi concentration?

The steady state model has been previously used to describe the system over the Kerguelen Plateau by Fripiat et al. (2011). In their study, authors have suggested that there was a significant ventilation of the ML above the Plateau which supply DSi with low $\delta^{30}\text{Si}$ value into the system even if they have observed a consumption of DSi in surface waters. In our study, the steady state model assumes no variations in concentrations with time and this is exactly what we have observed in spring in the Meander. Vertical and/or horizontal mixings supplied DSi in the ML that allow the system to behave following a steady state model.

Reviewer #2

General comment: Manuscript organization

Currently, the manuscript requires very careful reading (and re-reading) in order to understand the authors' argumentation and get a sense of the various settings. I think the authors would do well to try to make the data more accessible to readers unfamiliar with the details of KEOPS-2. A few suggestions:

- I would spend some time at the beginning of the Discussion (rather than in the Methods section) to introduce the groups of stations and their oceanographic settings, and make sure to refer to these settings consistently (currently, the text sometimes refers just to station numbers, sometimes to stations "north of the PF" even though this has not been referred to before, etc.). Such clear and consistent nomenclature is vital for the reader to be able to follow the authors' reasoning easily.

Ok, we will add a paragraph and a table at the beginning of the Discussion to introduce the stations and describe briefly their oceanographic settings and make their designation more consistently in the text.

- The groupings used in the text could be shown schematically on Fig. 1, i.e. show the "HNLC" region, the "Plateau" region and the "Meander" region in different shadings. Also, Fig. 1 would benefit from a more schematic representation of the flow, especially together with frontal positions.

Ok, we will simplify the figure and make the distinction between "HNLC", "Plateau", "Meander" and "North of the Front" stations in Fig. 1. Note that in order to be consistent with the many other articles from the same KEOPS-2 cruise (including a special volume in Biogeosciences) we need to keep the same labels for the stations.

- The timing of sampling is key to the authors' seasonal interpretation of the data but is not referred to very clearly: e.g. given their similarities in WW Si properties and use as the seasonal Si source, it is very important to know that TNS06 and A3-1 were sampled just 2 days apart. But this information is only visible in the supplementary information.

We will include this information in the table that will describe the stations (please see previous comment).

Major comment 1: Interpretation of isotope systematics

In Section 3.3.3, the authors discuss the mixed layer isotope systematics of the plateau and meander regions on the basis of the Rayleigh and steady-state models. Clearly, the isotopic behaviour in this region is complex, and the authors' discussion does justice to this, taking into consideration the various processes that may be affecting the isotope dynamics. However, I have a fundamental problem with the interpretation that BSi follows Rayleigh systematics whilst DSi does not – this simply cannot work. The evolution of the Rayleigh product (instantaneous or accumulated) is intimately tied to the evolution of the DSi pool – i.e. the product cannot follow the Rayleigh curve unless the DSi pool also experiences the strong fractionation associated with closed-system systematics. Thus whilst there is clearly complexity in the isotopic dynamics here, the authors' interpretation of "decoupled" systematics (Rayleigh for BSi, steady-state for DSi) cannot be correct. Could the results instead be due to a difference in the timescales over which the two sampled pools are integrating? Regardless of the specific reason, the author's analysis shows that whilst the model framework is useful for interpreting

isotopic systematics, it also has limitations when applied to the real ocean, and I would recommend a more careful interpretation that is less strictly dependent on conformance to idealised model curves.

We totally agree and we understand it was unclear from our previous version. Indeed the most likely explanation for D_{Si} pool to follow steady-state and B_{Si} to follow Rayleigh is the different timescales they integrate as suggested by the reviewer. This as already raised previously by other authors (e.g. Cardinal et al. 2007; Fripiat et al., 2011). As already suggested by reviewer #1, we will rearrange the text to make it less confusing for the reader. Please see previous comment on the fractionation factor (1st page of this document) where we give a potential explanation of this apparent decoupling between the particulate and dissolved pool.

Also, there is an inconsistency between the text and the caption of Fig. 4 regarding the Si source considered for the analysis. The text (L28 on p12) mentions “averaged Plateau-WW”, whilst the figure caption mentions “TNS6-WW”.

We will correct the “TNS6-WW” by “averaged Plateau-WW”.

Major comment 2: Error in plotting $\delta^{30}\text{Si}/\text{BSi}$ data

Based on the data presented in the supplementary material, the depth axis of Fig. 3h appears to be wrong. This does not allow a direct comparison of B_{Si} concentration and isotope composition for the meanders stations and makes it hard to assess the author’s interpretation of these data in Section 3.2.3, since it is not clear whether their interpretation is based on the faulty figure or not. This must be corrected. Regardless of this error, the authors’ argument for accumulation in the deeper ML or just below does not seem to be borne out by the data, given that B_{Si} concentrations decrease strongly below 100m. Given the many references to WML and ML in the discussion here and elsewhere, I think it would be useful to show physical parameters (or at least MLD) for the stations. Barring that, it is difficult for the reader to follow the authors’ reasoning.

We thank reviewer 2 for his careful look at our data. We apologise for this error which was on the depth axis on Fig. 3h. As suggested by reviewer #1, we will also change the discussion regarding the B_{Si} accumulation in this part of the text and we will add the MLD in each panels of Fig. 3.

Major comment 3: Calculation of quantitative results

In Sections 3.3.1 and 3.3.4 respectively, the authors calculate seasonal net B_{Si} production and estimate surface D_{Si} concentrations from isotope data, without giving details of the equations used (Section 3.3.1) or the assumptions made (Section 3.3.4) in order to achieve the results. Whilst I appreciate that the authors have done this before in other publications, it would be much better if their methods used for this study were documented here as well.

In Section 3.3.1 we used the same equations as those used in Fripiat et al., (2011). We will add the equations in the text or in the supp. material.

In Section 3.3.4, as written p16 L31 or p17 L31, we calculate the net BSi production in early and late summer assuming that the system is described by Rayleigh equations. The assumptions behind that are the following:

- The fractionation factor $^{30}\epsilon$ is constant.*
- There is no significant BSi export during this period so the BSi export:BSi production ratio is very low.*
- There is no significant BSi supply during this period so the Si supply:Si uptake ratio is very low.*

For example, in early summer, these two last assumptions are very consistent with the situation when the bloom has just started, when diatoms accumulate in the ML and has not consumed all the DSi stock.

In Mid-summer, the sharp decrease observed in our data can be associated to a mixing event. As specify p17 L11, we assume that the system is describe by steady state equations. The assumption behind that are the following:

- The fractionation factor $^{30}\epsilon$ is constant.*
- The DSi supplied into the system has the same isotopic signature as the source. The DSi source for the Plateau stations is the WW. Since the mixing event bring Si into the ML from deeper water, this assumption is clearly realistic.*
- All the BSi and the remaining DSi should be removed from the system. We cannot verify these assumptions but we can expect that a large part of the BSi was exported out of the ML Mid-summer. This assumption might be supported by the deep silica maximum observed late summer below the ML (150 m) by Mosseri et al. (2008) at the same station.*

We will list these assumptions in the revised text.

Minor comments

- Meander WW evolution: In Section 3.3.2, the authors discuss the evolution of WW $\delta^{30}\text{Si}$ in the meander region. Whilst it is possible that diapycnal interaction is responsible for this evolution as the authors argue, clearly the possibility of lateral interactions across the PF must be considered as well. After all, the intense mesoscale activity at the front acts strongly to stir/exchange tracers laterally/isopycnally as well as mixing them diapycnally (e.g. Dufour et al. 2015, J. Phys. Oc., 10.1175/JPO-D-14-0240.1) . As the authors argue for the ML in Section 3.2.3, lateral mixing of Sipoor PFZ subsurface waters with WW could thus well produce the evolution observed. In Fig. 5, it would be interesting to compare the evolution expected from such lateral subsurface mixing with that expected from diapycnal mixing.

The diapycnal mixing is sufficient to fully explain the Si characteristics of WW at stations E1 and E2 but cannot explain what we observed at stations E3, E4E and E5. We agree that a combination of vertical mixing and lateral advection is likely the reason of these observations and we will discuss this argument at the end of this paragraph with the reference to Dufour et al. (2015). However we will probably not change Fig. 5 as, considering our dataset, it is much more difficult to find the appropriate end-members for diapycnal advection than for vertical mixing and we think that we may not have enough measurements to do that properly (we would need more stations North of the polar front).

- Strange HNLC station: In my opinion, not enough reference is made to the fact that the HNLC station has a unique isotopic behaviour – its $\delta^{30}\text{SiDSi}$ evolves to very heavy values whilst $\delta^{30}\text{SiBSi}$ is very light, suggesting a very different expression of fractionation here. The authors explanation of the $\delta^{30}\text{SiBSi}$ data from this station (L25-21, p9) is unclear and thus not convincing.

We are sorry, we realized that there was an error in fig.2b. The blue profile that evolves toward very heavy $\delta^{30}\text{Si}$ values corresponds to station F and not R. This isotopic behavior is well consistent with the low DSi concentrations measured at station F. We will correct that.

- Silicate pump: In Section 4 (L7-10, p19), I do not understand how the silicate pump could lead to low BSi production despite high primary productivity. Should it not be the other way around if silica is more efficiently exported than organic matter? As a mechanism of decoupling Si, N and C, I am surprised that no mention is made of the Fe-related plasticity in Si:N ratios (as shown by the canonical 1998 papers by Sunda and Huntsman or Takeda), which I think would explain direction of change better.

As discussed in the text, the different diatom communities living under different biogeochemical conditions lead to variations in the silicon pump efficiency. These communities may have different degrees of silicification, explaining how we can observe high primary production but only moderate BSi production. By measuring Si uptake rates during the KEOPS-2 cruise, Closset et al. (2014) have suggested that these differences may not be primarily controlled by iron limitation such as already shown in artificial iron enrichment experiments (Takeda, 1998 or Hutchins and Bruland, 1998). Moreover, other recent studies in the Southern Ocean have highlighted that diatom community composition could better explain differences in silicification than physiological response to iron enrichment (see Baines et al., 2010; Assmy et al., 2013).

Various minor comments:

- UCDW/LCDW identification (L3-5, p5): It is UCDW that is associated with the O₂ minimum, not LCDW (e.g. Talley 2013, Oceanography 26, 80-97)

We will correct that in the text.

- Sections 2.2 and 2.3 both have the same heading. These should be “Sample collection” and “Sample preparation” respectively. The sub-sections of Section 2.3 are numbered incorrectly. On L6, p6, could there be an error in the units given for NaOH concentration?

Yes, we will modify as: “2.2 Sample collection” and “2.3 Sample preparation” and change the numbers in section 2.3. The unit given for NaOH concentration is 0.2 mol L^{-1} and not $\mu\text{mol L}^{-1}$ as written in the text and will be corrected.

- Section 2.5: it would be better to refer to “full external” rather than “global” reproducibility.

Ok, we will change “global reproducibility” by “full external reproducibility”.

- Section 3.1: Cao et al. 2012 is not a Southern Ocean study.

We agree, but they have made a figure (Fig.6) that compiled all the $\delta^{30}\text{Si}$ ranges of values for the dissolved and particulate pools in the global ocean (included in the Southern Ocean). We will more clearly mention this reference for their compilation.

- Section 3.3 and Fig. 3: The comparison with late summer conditions during KEOPS-1 is useful, but Fig. 3 is confusing since it combines the seasonal signal (i.e. KEOPS-1 vs. KEOPS-2) with a regional/biogeochemical signal (i.e. HNLC versus Plateau). The figure should separate these regions into separate panels and also show concentration profiles to provide context for the isotope data.

As previously suggested, we will add concentrations profiles in Fig.3. But we will probably not separate the regions in different panels as we have used different colors to discriminate between the HNLC and Plateau profiles.

- Section 3.3.4: It would be good to mention once again that the sediment trap is located at Station A3. On L33, p16, I don't understand why the ML concentration estimated from sediment trap data is compared with mid-November surface concentrations at station E5, rather than with mid-November concentrations at the station where the sediment trap is located (A3-2). Concentrations here are about 2x higher than estimated from the sediment trap $\delta^{30}\text{Si}$ data. Also, on L29, p17, it should be mentioned that the high values seen in late summer cannot be explained by the steady-state model.

Yes, we will change the sentence P16 L33 by "The remaining DSi concentration in the ML would be $9.53 \mu\text{mol L}^{-1}$ in November 2011 which is half of the ML DSi concentration measured there mid-November but which is close to the $12 \mu\text{mol L}^{-1}$ measured at E5." We will also specify that the high values seen in late summer cannot be explained by the steady state model.

- Finally, although in general the manuscript's English is good, it would profit from being proof-read by a native English speaker in order to iron out small grammatical errors.

Ok, we will take into consideration this comment.

Moreover, even though none of the reviewers has raised concerns on the accuracy of the data, we would like to add a section in the revised version of the supplementary material. Indeed, there have been some concerns regarding Si isotopic offsets among labs on North Atlantic data (see Brzezinski and Jones, 2015) as well as more recently on the GEOTRACES intercalibration (Grasse et al., to be submitted). We will show in the revised version that we have compared the deep samples of our KEOPS-2 study using a Neptune⁺ MC-ICP-MS (Closset et al. 2015) and a chemical purification adapted from Hughes et al. (2011): cationic exchange + anion doping with data from KEOPS-1 (Fripiat et al. 2011) using a Nu Instrument MC-ICP-MS (Cardinal et al. 2003) and chemical purification adapted from De La Rocha et al. (1996). On a $\delta^{30}\text{Si}$ vs. $1/\text{Si}$ plot the two dataset show an offset of only 0.08 ‰ which is not analytically significant. We propose to modify the Supplementary Material Figure S4 to better highlight this comparison.

List of relevant changes in the revised manuscript

- * P7-8: We have removed the paragraph “KEOPS-2, cruise and hydrological settings” in the Material and Methods and we have combined it with “General considerations” in the Results and Discussion. We have detailed the different groups of stations in this paragraph.
- * We have added a table with station settings in the revised manuscript (Table 1).
- * P8: We have included a paragraph about the accuracy of our data and their comparison with previous $\delta^{30}\text{Si}$ values in this region, and we have modified the corresponding figure (Fig. S4).
- * P11: As suggested by the reviewers, we have changed our interpretation of low $\delta^{30}\text{Si}$ in surface compared to the subsurface. We have included a comparison with the meridional overturning circulation.
- * P12: we have detailed the reason why we used preferentially the steady-state model to calculate BSi production in the Kerguelen region.
- * P14: We have included the interpretation suggested by a reviewer about isopycnal mixings for explaining the winter water DSi and $\delta^{30}\text{Si}$ signatures.
- * P15: We have modified the paragraph about the decoupling between the dissolved and the particulate phase to make it clearer.
- * P17-18: As suggested by the reviewers, we have detailed the assumptions behind our estimations of BSi production and DSi consumption.
- * P20: We have added some references to support our conclusions>
- * Figures: We have simplified figure 1 and corrected figure 2. We have added 2 panels on figure 3 and included the uncertainties on figure 4.

Marked-up manuscript version (changes are in green color)

Unveiling the Si cycle using isotopes in an iron fertilized zone of the Southern Ocean: from mixed layer supply to export

Ivia Closset¹, Damien Cardinal¹, Mathieu Rembauville², François Thil³, Stéphane Blain²

¹Sorbonne Universités (UPMC, Univ Paris 06)-CNRS-IRD-MNHN, LOCEAN Laboratory, 4 place Jussieu, F-75005 Paris, France

²Sorbonne Universités (UPMC, Univ Paris 06)-CNRS, Laboratoire d’Océanographie Microbienne (LOMIC), Observatoire Océanologique, F-66650 Banyuls/mer, France

³Laboratoire des Sciences du Climat et de l’Environnement, CNRS 91190 Gif-sur-Yvette, France

Correspondence to: Ivia Closset (ivia.closset@locean-ipsl.upmc.fr)

Abstract. A massive diatom-bloom forms annually in the surface waters of the naturally Fe-fertilized Kerguelen Plateau (Southern Ocean). In this study, silicon isotopic signatures ($\delta^{30}\text{Si}$) of silicic acid (DSi) and suspended biogenic silica (BSi) were investigated through the whole water column with unprecedented spatial resolution, during the KEOPS-2 experiment (spring 2011). We used $\delta^{30}\text{Si}$ measurements to track the sources of silicon that fuelled the bloom, and investigated the seasonal evolution of the Si biogeochemical cycle in the iron fertilized area. We compared the results from stations with various degrees of iron enrichment and bloom conditions to an HNLC reference station. Dissolved and particulate $\delta^{30}\text{Si}$ signatures were highly variable in the upper 500 m, reflecting the effect of intense silicon utilization in spring, while they were quite homogeneous in deeper waters. The Si-isotopic and mass balance identified a unique Winter Water (WW) Si-source for the iron-fertilized area that originated from southeast of the Kerguelen Plateau and spread northward. When the WW reached a retroflexion of the Polar Front (PF), the $\delta^{30}\text{Si}$ composition of the silicic acid pool became progressively heavier. This would result from sequential diapycnal and isopycnal mixings between the initial WW and ML water masses, highlighting the strong circulation of surface waters that defined this zone. When comparing the results from the two KEOPS expeditions, the relationship between DSi depletion, BSi production and their isotopic composition appears decoupled in the iron fertilized area. This seasonal decoupling could help to explain the low apparent fractionation factor observed in the ML at the end of summer. Taking into account these considerations, we refined the seasonal net BSi production in the ML of the iron-fertilized area to $3.0 \pm 0.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$, that was exclusively sustained by surface water phytoplankton populations. These insights confirm that the isotopic composition of dissolved and particulate silicon is a promising tool to improve our understanding on the Si-biogeochemical cycle since the isotopic and mass balance allows resolution of processes in the Si cycle (i.e. uptake, dissolution, mixing).

1 Introduction

The Southern Ocean plays a crucial role in the regulation of the global climate as it contributes significantly to the world’s ocean primary production and represents a major carbon sink (Takahashi et al., 2009). In the Southern Ocean, diatoms, a phytoplankton group that produce an opaline cell wall called a frustule (amorphous $\text{SiO}_2 \cdot n\text{H}_2\text{O}$), are responsible for more than 75 % of the annual primary production. Diatoms are especially dominant in the area south of the Antarctic Polar Front (PF) where they form massive blooms (e.g. Brzezinski et al., 2001). These blooms are often associated with high export events when diatoms are removed from the mixed layer (ML) via aggregation, settling or grazing processes (Boyd & Trull, 2007; Buesseler 1998) and thus contribute to the extensive biogenic silica (hereafter referred to as BSi) deposition that characterizes the sediments of this region known as the “opal belt” (Tréguer & De la Rocha, 2013; Ragueneau et al., 2000).

Since diatoms preferentially take up light silicon isotopes (^{28}Si), their biological activity leaves a clear imprint on the isotopic composition of both silicic acid (H_4SiO_4 , hereafter referred to as DSi) and BSi, enriching the DSi pool with heavy ^{30}Si isotope (De la Rocha et al., 1997). While a recent in vitro study has pointed out that this preferential uptake may vary among diatom species (ranging from -0.54 to -2.09 ‰, Sutton et al., 2013), field studies have reported constant Si isotopic fractionation factor ($^{30}\epsilon$) for the Antarctic Circumpolar Current, ACC (e.g. De la Rocha et al., 2000; Cardinal et al., 2005), estimated to -1.2 ± 0.2 ‰ on average (Fripiat et al., 2011a). In the reverse reaction, it is not clear whether BSi dissolution fractionates silicon isotopes. Two studies based on laboratory experiments provide contradictory results. Wetzel et al., (2014) performed alkaline digestion of diatom opal in sediment cores and reported no preferential release of a given isotope. Conversely, Demarest et al. (2009) placed fresh siliceous particles in undersaturated seawater and reported a fractionation factor of -0.55 ‰ associated with the dissolution process. The fractionation of dissolution opposes the one occurring during Si uptake (BSi dissolution preferentially releases ^{28}Si) which would reduce the overall (or net) fractionation factor. Thus, information on the silicon isotopic composition ($\delta^{30}\text{Si}$) potentially enables to identify DSi sources, discriminate and quantify different processes such as Si-uptake, BSi dissolution or physical mixing (e.g. Fripiat et al., 2011a).

Diatoms generate the largest latitudinal gradient of DSi in the world in the Southern Ocean because of their absolute requirement for silicic acid, from replete conditions south of the PF, in the Antarctic Zone (AZ), to depleted conditions in the Sub-Antarctic Zone (SAZ) (e.g. Brzezinski et al., 2001; Sarmiento et al., 2004). This contrasts strongly with the High Nutrient Low Chlorophyll (HNLC) characteristics of the Southern Ocean which could be more specifically defined as a High Nitrate Low Silicon Low Chlorophyll area (HNLSiLC, Dugdale et al., 1995). Several studies based on in vitro and artificial Fe-enrichments have shown that Fe stress is partly responsible for the HNLSiLC status of the ACC (see review in De Baar et al., 2005; Boyd et al., 2007). In parallel, several field studies have investigated the effect of natural iron delivery on the biogeochemical cycles (e.g. Blain et al., 2007; Pollard et al. 2009).

The KEOPS project consisted of two expeditions (late-summer 2005 and early-spring 2011) conducted in a naturally iron-fertilized area in the vicinity of the Kerguelen Plateau where a massive phytoplankton bloom is observed annually (Mongin et al., 2008). The first KEOPS expedition (January-February 2005) has highlighted that this bloom is sustained by iron supply originating from iron-rich deep waters through winter mixing and vertical diffusion in summer (Blain et al., 2007). The second expedition also revealed that episodic deepening of the mixed layer (ML) could also contribute to the vertical supply of Fe (Bowie et al. 2015). Silicon isotopes studied during the first cruise highlighted important aspects of the Si biogeochemical cycle in both the HNLC and the fertilized area (Fripiat et al., 2011a). It was suggested that the decline of the bloom was strongly controlled by silicic acid and iron co-limitations (Mosseri et al., 2008). Results from KEOPS-2 (October-November 2011) have shown that the Kerguelen bloom was characterized by a complex and heterogeneous distribution of phytoplankton communities (e.g. Lasbleiz et al., 2014), constrained by a mosaic of biogeochemical conditions (Park et al., 2014).

Isotopic variations induced by biological Si-utilization in the Southern Ocean ML can be described in a first approximation using two different models for closed and open systems. In the closed system (also referred-to as Rayleigh model) the surface ocean is considered to have a finite pool of DSi and that is not replenished by any external sources. In this case, the reaction progresses in a sequential mode over time: the initial substrate (here silicic acid) is consumed in the ML, which exponentially increases the $\delta^{30}\text{Si}$ of the remaining substrate (Eq. 1). The isotopic composition of the short-term or instantaneous product (here exported BSi, Eq. 2) differs from the long-term product that accumulates (here BSi in the ML) which holds the same signature as the initial substrate when everything is consumed in the system (Eq. 3):

$$\delta^{30}\text{Si}_{sub} = \delta^{30}\text{Si}_{init} - {}^{30}\epsilon \ln(1 - f) \quad (1)$$

$$\delta^{30}\text{Si}_{inst} = \delta^{30}\text{Si}_{sub} + {}^{30}\epsilon \quad (2)$$

$$\delta^{30}\text{Si}_{acc} = \delta^{30}\text{Si}_{init} - {}^{30}\epsilon \left(\frac{f \times \ln f}{1-f} \right) \quad (3)$$

where ${}^{30}\epsilon$ is the isotopic fractionation factor of the reaction, f the fraction of the remaining substrate and the subscripts “sub”, “init”, “acc” and “inst” refer to the remaining substrate, the initial substrate, the accumulated product and the instantaneous product respectively.

A distinct system is an open, flow-through system (also referred to as steady-state model) where a continuous supply of substrate balances the export of product. In this model, the DSi supply equals the sum of the BSi produced and immediately exported and the residual DSi stock leaving the system. Only one product forms from substrate and both $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ signatures display linear changes (Eq. 4 and 5):

$$\delta^{30}\text{Si}_{sub} = \delta^{30}\text{Si}_{init} - {}^{30}\epsilon(1 - f) \quad (4)$$

$$\delta^{30}\text{Si}_{prod} = \delta^{30}\text{Si}_{init} + {}^{30}\epsilon \times f \quad (5)$$

In these two ideal situations, the $\Delta^{30}\text{Si}$, i.e. the difference between $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$, is considered as constant. However, in the ocean, it could actually be altered by processes such as water mixing and BSi dissolution that would affect the isotopic and mass balance (e.g. Demarest et al., 2009). It is likely that natural environments reflect a combination of these closed and open systems, with the real fractionation trend lying between the two curves and depending on the Si-uptake:Si-supply ratio occurring in the system (Fry 2006). This could be particularly the case in the Southern Ocean where large nutrient consumption was observed in regions characterized by intense mixing events that strongly deepen the ML and bring DSi in surface waters (e.g. Brzezinski et al., 2001; Nelson et al., 2001).

In this paper, we investigate the spatial and seasonal variability of silicon isotopic composition of both seawater ($\delta^{30}\text{Si}_{\text{DSi}}$) and siliceous particles ($\delta^{30}\text{Si}_{\text{BSi}}$) in deep and surface waters of the Kerguelen area during the spring period. Our specific objectives are the following:

- Describe the early spring spatial distribution of silicon isotopes across the range of productivity observed near the Kerguelen Islands.
- Discuss the potential role of iron in regulating the Si biogeochemical cycle and isotope dynamics in the Southern Ocean surface and subsurface waters.
- Refine the net BSi production estimated by Fripiat et al. (2011a) using the appropriate spring biogeochemical conditions (DSi concentration and $\delta^{30}\text{Si}$).
- Finally, identify and track the Si sources that fuel the phytoplankton bloom above the Plateau and characterize the temporal evolution of the silicon biogeochemical cycle by combining our KEOPS-2 results with KEOPS-1 data.

2 Material and Methods

2.1 Sample collection

Si isotopic compositions from a total of 224 seawater samples and 137 particulate samples are presented in this study (Table S2). The particulate samples included 91 samples collected in the ML using Niskin bottles and 46 samples coming from in situ pumps (ISP). At all stations, seawater and particles were collected using a CTD (Conductivity-Temperature-Depth) rosette equipped with 12 L Niskin bottles. Seawater (approximately 5 L) was immediately filtered through polycarbonate membranes (Nuclepore, 0.4 μm) using large volume filtration units. Filtered water samples were stored in the dark in acid-cleaned polypropylene bottles and membranes were dried overnight at 50 °C and stored in polycarbonate Petri dishes at room temperature. Since particle concentration decreases with depth, deep-water particles were collected using in-situ pumps at 7 stations by filtering 70 to 1800 L of seawater through hydrophilic polyestersulphone membranes (SUPOR, 0.8 μm). Approximately 1/8 of the SUPOR membranes were dedicated to silicon isotopic analysis and were dried overnight at 50 °C and stored in Petri dishes at room temperature.

Sinking particles were collected from a moored sediment trap (Technicap PPS3) that was deployed at station A3 over the central Kerguelen Plateau at 289 m (seafloor depth 527 m). The sediment trap carousel was composed of 12 sampling cups (250 mL) collecting sinking particles from the 21 October 2011 to 7 September 2012. Sampling intervals were programmed to be short (10-14 days) in spring and summer and longer (99 days) in autumn and winter. A description of the physical environment of the deployment together with the detailed methods for sediment trap sample processing are reported in Rembauville et al. (2015a). After the BSi extraction (Rembauville et al., 2015b), samples were purified and analysed for the Si isotopic composition as described below.

2.2 Sample preparation

2.2.1 Particles digestion and BSi analyses

The membranes (Nuclepore and SUPOR) were subjected to a wet-alkaline digestion (adapted from Ragueneau et al., 2005). BSi was dissolved in Teflon tubes using a 0.2 mol L⁻¹ NaOH solution (pH 13.3) at 100°C for 40 min followed by neutralization with HCl (1 mol L⁻¹). As this digestion can also solubilize

some lithogenic silica (mainly clay minerals), a second and identical digestion was applied to the membranes that were then analyzed for aluminum (Al, a tracer of lithogenic source) using an Inductively Coupled Plasma Mass Spectrometer (ICP-MS; detection limit = 3.18 ppb). Using the Si:Al ratio measured in the second digestion, the lithogenic silicon dissolved in the first digestion can be estimated (Ragueneau et al., 2005). Unfortunately, the polycarbonate membranes used for silicon isotopic analysis were contaminated by Al during filtrations onboard. However, SUPOR filters as well as Al concentrations estimated on other filters (Lasbleiz et al., 2014) revealed negligible lithogenic silicon in the first leachate (on average 1.26 % for SUPOR filters). Such a lithogenic contribution should not bias significantly our $\delta^{30}\text{Si}$ value. Indeed, using a light end-member $\delta^{30}\text{Si}_{\text{LSi}}$ of -2.3 ‰ reported in clays (Opfergelt & Delmelle, 2012) and our maximum $\delta^{30}\text{Si}_{\text{BSi}}$ (2.06 ‰ station R2, 2400m) as extreme end-members, we calculated a maximum interference in the isotopic signal of 0.05 ‰ which is similar to our analytical precision for $\delta^{30}\text{Si}$. BSi concentrations were determined with a colorimetric method according to Grasshoff et al. (1999) and by ICP-MS on the same samples used to measure Si-isotopic composition. Every ISP sample was analyzed in full duplicates (i.e. on a second fraction of the same membrane) with a pooled standard deviation of 5.0 ± 4.6 % (n = 58), which corresponds to our average reproducibility of BSi measurements, and which is slightly better than the uncertainty estimated for this method (10 %, Ragueneau et al., 2005). Moreover, independent measurements of BSi concentration performed with Niskin bottles at the same stations and depths during KEOPS-2 by Lasbleiz et al. (2014) were similar to our results, suggesting that the ISP method was robust.

2.2.2 Seawater preconcentration and DSi analyses

A two-step preconcentration procedure adapted from the MAGIC method (Karl & Tien, 1992; Reynolds et al., 2006) was performed on seawater samples to increase H_4SiO_4 concentration and reduce the anionic matrix that could interfere with Si during isotopic analysis (e.g. sulfates, SO_4^{2-} ; Hughes et al., 2011). DSi was co-precipitated in two steps with brucite ($\text{Mg}(\text{OH})_2$) by adding 2 % (v/v), following by 1 % (v/v) of 1M NaOH to the seawater sample. This solution was shaken and left for 1h and the precipitate was recovered by centrifugation and redissolved with 1M HCl. The supernatant was removed and Si recovery was monitored by checking systematically that no detectable amount of silicic acid remained in the supernatant after coprecipitation and centrifugation. DSi concentrations in seawater samples were determined with a colorimetric method (Grasshoff et al., 1999) on the same samples as for Si-isotopic composition. Average reproducibility of DSi measurements was 6.7 % (calculated from 98 in-house silicon solution analyses at the ± 1 sd level).

2.3 Purification

Separation of Si from other ions in the sample was achieved by passing the solution through a cation-exchange column (BioRad cation exchange resin DOWEX 50W-X12, 200 to 400 mesh, in H⁺ form) using the protocol described in Georg et al. (2006). After purification, systematic analysis of major elements (such as Mg, Ca, Na, Al) were performed by ICP-MS to ensure sample purity prior to isotopic analyses (Si/X weight ratio usually > 50). Si concentrations were also measured in the purified solutions to check for complete recovery. This purification step did not allow the complete removal of the anionic matrix, which consists primarily of Cl^- , SO_4^{2-} and to a lesser extent PO_4^{3-} and NO_3^- . Cl^- originating from seawater

can be neglected compared to Cl⁻ added as HCl to dissolve the brucite. Therefore the solutions dedicated to DSi isotopic measurements were analyzed by anionic chromatography to control the concentration of sulfates. Indeed, in these samples, SO₄²⁻ concentrations could induce a significant shift in isotopic measurements (see supplementary method and Van den Boorn et al. (2009) for rock digestion solutions). Thus, as proposed by Hughes et al. (2011), samples and standards used for DSi isotopic analyses were doped with sulfuric acid in large excess compared to the natural SO₄²⁻ concentrations in order to control this sulfate matrix effect.

2.4 Isotopic measurements

The purified and sulfate doped Si solutions were analyzed for isotopic measurements on a Thermo Neptune⁺ Multicollector Inductively Coupled Plasma Mass Spectrometer (MC-ICP-MS; LSCE, Gif-sur-Yvette) in dry plasma mode using Mg external doping to correct for the mass bias (Cardinal et al., 2003; Abraham et al., 2008). Samples were injected into the plasma with an Apex desolvating nebulization system connected with a PFA nebulizer (100 μL min⁻¹ uptake rate) and without additional gas. Silicon isotopic compositions (δ³⁰Si) were calculated as the permil deviation from the quartz standard NBS28 (RM8546). They were measured relative to an in-house standard Quartz Merck, which was not significantly different from NBS28 (Abraham et al., 2008), analyzed immediately after and before the sample and expressed as:

$$\delta^{30}\text{Si} (\text{‰}) = \left[\frac{(^{30}\text{Si}/^{28}\text{Si})_{\text{sample}}}{(^{30}\text{Si}/^{28}\text{Si})_{\text{standard}}} - 1 \right] \times 1000 \quad (6)$$

Blanks levels were below 1 % of the main signal and were subtracted from each sample and standard analyses. All measurements were carried out in a matrix composed of HNO₃ 0.5 mol L⁻¹, HCl 0.5 mol L⁻¹, H₂SO₄ 1 mmol L⁻¹ and medium resolution mode (M/ΔM > 6000) to optimize the separation of ³⁰Si peak and ¹⁴N¹⁶O interference and were performed on the interference-free left side of the peak (Abraham et al., 2008). δ²⁹Si and δ³⁰Si were compared to the mass dependent fractionation line (Fig. S3) and samples falling outside of its analytical error were excluded from final dataset. Typical analytical conditions are provided in table S3.

Numerous analyses of a secondary reference material such as Diatomite (δ³⁰Si = 1.26 ‰, Reynolds et al., 2007) generated over the entire procedure indicated an average precision and a long-term analytical reproducibility (24 months) on δ³⁰Si values of 1.28 ± 0.05 ‰ (1 sd, n = 128) and confirmed that no uncorrected isotopic bias occurred. All BSi samples and some DSi samples were fully replicated and measured on separate days (chemical preparation plus isotopic measurements). In these cases, the average full external reproducibility on full duplicates δ³⁰Si is 0.06 ‰ (1 sd, n= 108) and 0.04 ‰ (n= 78) for δ³⁰Si_{BSi} and δ³⁰Si_{DSi} respectively. Error bars shown in all figures correspond to the analytical reproducibility or to the full external reproducibility if greater than 0.05 ‰.

3 Results and Discussion

3.1 KEOPS-2 cruise, hydrological settings and general considerations

The KEOPS-2 expedition was conducted in the Kerguelen Plateau region (Indian sector of the Southern Ocean) from 10 October to 20 November 2011 (austral spring) on board of the R/V Marion Dufresne. The Plateau acts as a barrier to the ACC, 60 % of which passes north of the Kerguelen Islands, mostly associated with the SAF, while 40 % is transported across the southern part of the Plateau and forms the jet of the PF (Park et al., 2014). This PF current is then deflected to the north following the eastern escarpment of the Plateau and forms a permanent cyclonic meandering associated with strong mesoscale activity (Park et al., 2014; Zhou et al., 2014). The circulation above the central Kerguelen Plateau is relatively weak ($< 3\text{ cm s}^{-1}$, Park et al., 2008b), providing good conditions for elevated primary production (Mongin et al., 2008). These particular hydrographic features generate contrasted biogeochemical and physical environments where phytoplankton communities will respond differently to iron availability. During KEOPS-2, the vertical distribution of water masses was characteristic of the Antarctic zone (AZ) in the vicinity of the PF (Park et al., 2014). Except for the first visit at over the Plateau (A3) where they reached the surface, the remnant Winter Water (WW, generally from 100 to 400 m), was capped by a homogeneous, warm and fresh mixed layer (ML) or Antarctic Surface Water (AASW) induced by seasonal stratification. Below these subsurface waters, a subsurface temperature maximum between 400 and 1400 m associated to the Upper Circumpolar Deep Waters (UCDW), followed the Lower Circumpolar Deep Water (LCDW) in all of the out-Plateau stations. The deeper Antarctic Bottom Waters (AABW) were found only at station F, north of the PF. The cruise consisted in two transects north to south (TNS stations) and east to west (TEW stations) aimed at documenting the spatial extension of the bloom and its coastal-offshore gradient; and 11 long-term stations devoted to process studies (see Fig. 1 and Table 1):

- A HNLC reference station (R2) located upstream of the eastward ACC flow, south of the Kerguelen Islands, hereafter referred to as HNLC station.

- 2 visits at the KEOPS-1 Plateau bloom reference station (A3-1 and A3-2), and at a moderate productive station (E4W) located in the plume of chlorophyll north of A3, hereafter referred to as Plateau stations.

- An open ocean station (F) influenced by warmer Subantarctic Surface Waters, located north of the PF, hereafter referred to as Polar Front Zone.

- 6 stations (TNS6 and E1 to E5) constituting a pseudo-Lagrangian survey and located in the area of PF retroflexion characterized by strong mesoscale activity (Zhou et al., 2014) and hereafter referred to as Meander stations.

During KEOPS-2, $\delta^{30}\text{Si}_{\text{DSi}}$ displayed a clear inverse relationship with silicic acid concentration as commonly observed in the Southern Ocean (Fig. S4 and e.g. in Fripiat et al., 2012; De Souza et al., 2012). The water column profiles showed a general increase in DSi concentrations. $\delta^{30}\text{Si}_{\text{DSi}}$ became gradually lighter with depth. In the upper 500 m, $\delta^{30}\text{Si}_{\text{BSi}}$ values were systematically lighter than $\delta^{30}\text{Si}_{\text{DSi}}$ values in agreement with the preferential uptake of ^{28}Si by diatoms (De La Rocha et al., 1997). Below 500 m there was a slight increase of $\delta^{30}\text{Si}_{\text{BSi}}$ and a decrease of $\delta^{30}\text{Si}_{\text{DSi}}$ values coincident with increasing

DSi concentrations. In contrast to the ML, the $\delta^{30}\text{Si}_{\text{BSi}}$ below 500 m was systematically higher than $\delta^{30}\text{Si}_{\text{DSi}}$ (on average 1.74 ± 0.13 and 1.20 ± 0.17 ‰, respectively). This observation differs significantly from the one observed in the Atlantic sector in summer by Fripiat et al. (2012) where the isotopic signature of BSi exported to depth was directly comparable to the $\delta^{30}\text{Si}_{\text{BSi}}$ values of particles in the ML. The surface KEOPS-2 $\delta^{30}\text{Si}_{\text{BSi}}$ (ranging from 0.47 ‰ to 2.04 ‰) and $\delta^{30}\text{Si}_{\text{DSi}}$ (ranging from 1.96 ‰ to 2.79 ‰) encompassed nearly the full range of delta results reported from previous Southern Ocean studies (-0.7 ‰ to 2.8 ‰ for BSi and 0.5 ‰ to 4.4 ‰ for DSi, e.g. De Souza et al., 2012; Fripiat et al., 2012). Recently, there have been some concerns regarding Si isotopic offsets among laboratories on North Atlantic data (see Brzezinski & Jones, 2015) as well as on the GEOTRACES intercalibration (Grasse et al., submitted to Journal of Analytical Atomic Spectrometry). We have compared the deep KEOPS-2 samples, measured using a Neptune+ MC-ICP-MS (this study) after a chemical purification adapted from Hughes et al. (2011) with data from KEOPS-1, measured using a Nu Instrument MC-ICP-MS (Fripiat et al., 2011a) after a chemical purification adapted from De La Rocha et al. (1996). Despite the two different sample processing methods and instrumentation, the two dataset show an offset of only 0.08 ‰ (Fig. S4) which is not analytically significant and allow the direct comparison between KEOPS-1 and KEOPS-2 delta values.

In general, $\delta^{30}\text{Si}_{\text{DSi}}$ signatures in deep waters were homogeneous and did not show large changes with depth or between stations (on average 1.28 ± 0.08 ‰ and 1.05 ± 0.06 ‰ for UCDW and LCDW respectively). There was much higher variability in $\delta^{30}\text{Si}_{\text{DSi}}$ in the ML and WW between stations (1.99 ‰ to 2.53 ‰ and 1.46 ‰ to 2.03 ‰, respectively, Table S4). Note that the WW exhibited systematically the largest dispersion of delta values and were systematically associated to a strong isotopic gradient toward light $\delta^{30}\text{Si}_{\text{DSi}}$ with depth (not shown). Consequently, the mean and median values were probably not representative of the specific Si concentration and isotopic composition of the WW. Usually, the temperature-minimum layer (T-min between 1.5 and 2°C in the Kerguelen area; Park et al., 2014) is the traditional definition of the remnant surface winter water. However, in some stations, it was not possible to clearly determine T-min (e.g. R2 and E2) and/or there was a significant salinity gradient above the T-min depth (e.g. F). Trull et al. (2015) have proposed that a shallower depth based on a threshold increase in salinity of 0.05 (S-threshold depth) could better represent the WW characteristics. They ascribed undeterminable T-min to weaker winter mixing compared to the previous year and thus, the nutrient depletion between the T-min and S-threshold depths could not be associated to recent consumption. In the following we will use the S-threshold approach since it is the most appropriate reference to reflect the pre-bloom biogeochemical conditions.

3.2 Distribution of Si isotopes vs. source and supply of iron

3.2.1 The HNLC reference station

In the ML, the Plateau and HNLC areas exhibited significant different silicon isotopic composition of both seawater and particles, reflecting the different degrees of Si-utilization by diatoms. The HNLC reference station (R2) displayed low chlorophyll *a* and BSi and very low Si-uptake rates, consistent with its iron-depleted condition and the dominance of non-siliceous organisms (see details in Lasbleiz et al., 2014 and Closset et al., 2014). In surface waters of the HNLC area, the concentration of biogenic silica

was the lowest measured in the surface during KEOPS-2 ($0.30 \pm 0.03 \mu\text{mol L}^{-1}$) and its silicon isotopic composition was low ($0.73 \pm 0.04 \text{‰}$) and similar to A3-1, (Fig. 2e, 2f). This is typical of non-bloom conditions and in the same range as those measured in HNLC waters of the Southern Ocean (e.g. Fripiat et al., 2011a, 2011b; Mosseri et al., 2008). However, the HNLC station displayed unexpectedly low silicic acid concentration and heavy $\delta^{30}\text{Si}_{\text{DSi}}$ in the ML ($12.94 \pm 0.49 \mu\text{mol L}^{-1}$ and $2.21 \pm 0.06 \text{‰}$, respectively; Fig. 2a, 2b). This latter is significantly heavier than the Si-isotopic composition of the fertilized area measured few days before ($1.99 \pm 0.03 \text{‰}$, A3-1). As already proposed by Closset et al. (2014) and Lasbleiz et al. (2014), this suggests that a development of diatoms could have occurred before our sampling, consuming a fraction of the DSi standing stock and increasing the $\delta^{30}\text{Si}_{\text{DSi}}$ of surface waters. The same evidence of surface production has also been deduced by Dehairs et al. (2015), who observed a slight nitrate depletion and enrichment of $\delta^{15}\text{N-NO}_3^-$. The low iron concentration that prevails at this station in early spring might have been sufficient to trigger a short phytoplankton growth (dominated by nanophytoplankton; Lasbleiz et al., 2014) as soon as light conditions became favorable. Then, both the high silica dissolution to production ratio (D:P > 1) observed in the ML (Closset et al., 2014) and the high barium excess measured between 200 and 400 m (Jacquet et al., 2015) suggest that this material was exported and remineralized by the time we visited the station. This could be confirmed by the clear $\delta^{30}\text{Si}_{\text{BSi}}$ maximum observed between 100 and 200 m (Fig. 2f) that could result from the dissolution isotopically light diatoms initially produced from lighter DSi.

By contrast, all other KEOPS-2 stations outside of the HNLC were characterized by the development of large spring blooms that were not homogeneous in time and space depending on the degree, the mode and the timing of their iron fertilization (Bowie et al., 2015; Trull et al., 2015). These blooms were organized in three main clusters related to their different iron supplies and are discussed separately in the following.

3.2.2 The Kerguelen Plateau zone

This area was characterized by large and recurrent blooms located southeast of the islands, mainly above the Kerguelen Plateau and delimited northward by the Polar Front (Blain et al., 2001, 2007). During our first visit to A3 (A3-1), low $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ ($1.99 \pm 0.03 \text{‰}$ and $0.77 \pm 0.05 \text{‰}$, respectively, Fig. 2b, 2f) were measured in the ML indicating that biogeochemical conditions prevailing there were characteristics of a pre-bloom or early-bloom period. Indeed, low chlorophyll *a* and BSi concentrations were observed despite high nutrients standing stocks (H_4SiO_4 and NO_3^- , see Blain et al., 2015 for NO_3^-) and relatively high iron concentrations (Bowie et al., 2015).

The largest phytoplankton development was observed during the second visit to A3, where chlorophyll *a* and BSi concentrations increased more than two fold over one month. This growth was reflected by the significant increase of both $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ (from $1.99 \pm 0.03 \text{‰}$ to $2.10 \pm 0.05 \text{‰}$ and from $0.77 \pm 0.05 \text{‰}$ to 0.96 ± 0.08 , respectively, Fig. 2b, 2f). There, the supply of nutrients and iron to the ML coming from the WW both in winter and during the productive period allowed a spring biogenic silica

production comparable to the most productive regions in the world, such as upwelling systems or river plumes (up to $43.4 \pm 0.4 \text{ mmol m}^{-2} \text{ d}^{-1}$ for net opal-production integrated over the euphotic zone, Fripiat et al., 2011a; Closset et al., 2014).

3.2.3 The recirculation zone in the Polar Front Meander

The central part of the Meander was characterized by a complex and slowly flowing water circulation associated with low to moderate dissolved iron concentrations (Bowie et al., 2015; Qu  rou   et al., 2015). Here, surface waters displayed generally higher $\delta^{30}\text{Si}_{\text{DSi}}$ signatures and lower DSi concentrations compared to above the Plateau (TNS6, E1 to E5; Fig. 2d), supporting the idea of a northward surface circulation with a progressive consumption and enrichment in ^{30}Si of the dissolved pool. According to Park et al. (2014), this zone could correspond to the latest arrival of water originated from the shallow Plateau located south of these stations. This would explain the delay observed between the initiation of production here and the southernmost bloom located above the Plateau and in the PF plume (Fig. 1). Moreover, radium isotopes signature (Sanial et al., 2015) suggested that the southward transport of chemical elements (such as iron) across the PF could also occur and significantly fuel the phytoplankton bloom in this area, mixing the Si-poor PF ASW with heavy $\delta^{30}\text{Si}_{\text{DSi}}$ and Si-rich AASW with light $\delta^{30}\text{Si}_{\text{DSi}}$.

The moderate iron fertilization occurring in the meander should have increased both the BSi production rate and biomass, with larger increase at the two last visits (E4E and E5, Closset et al., 2014; Lasbleiz et al. 2014). However, no clear trend was identified in the ML values of $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ since these two parameters were not significantly different from TNS6 to E4E (Table S4). This observation is strengthened by the homogeneous $\delta^{30}\text{Si}_{\text{DSi}}$ profiles in all the meander's stations (Fig. 2d). TSN6 and E1 to E3 stations display relatively homogeneous BSi concentrations and isotopic compositions. Interestingly, stations E4E and E5 were characterized by a surface accumulation of significantly lighter BSi when compared to the subsurface (respectively 1.26 ‰ and 1.46 ‰ at E5). This lower BSi isotope composition may likely result from the supply of light DSi into the ML by episodic mixing events. This would cause the DSi stock in the ML to have a lower isotopic signature than previously (as we observed slightly in Fig. 2d), and diatoms would produce lighter BSi at the top of the ML.

3.2.4 The Polar Front Zone

A plume of chlorophyll *a* was present in the Polar Front Zone (station F, Fig. 1), extending eastward of the Kerguelen Plateau and showing strong mesoscale activity associated to the temporal and spatial variability of the Polar Front (Park et al., 2014). There, the productive waters were characterized by a shallow and relatively warm ML ($> 4^\circ\text{C}$) associated with moderate to high iron concentration (Qu  rou   et al., 2015; Bowie et al., 2015), providing favorable conditions for phytoplankton development. These waters were characterized by distinctly different phytoplankton communities (mostly small diatoms and nanoflagellates) than those found above the Plateau (Lasbleiz et al., 2014). The Si-isotopic composition of ML seawater and particles measured at F were the heaviest of KEOPS-2 stations coinciding with strong Si-depletion in surface water (Fig. 2b, 2f). These $\delta^{30}\text{Si}_{\text{DSi}}$ signatures fall in the

range of previous reported values in the PFZ ML, from 2.05 ± 0.03 ‰ in spring (Cardinal et al., 2005) to 2.77 ± 0.23 ‰ in late summer (Fripiat et al., 2011a) in the Australian and Atlantic sector of the Southern Ocean respectively. In the PFZ, heavier $\delta^{30}\text{Si}$ signatures can be attributed to a higher utilization of silicic acid by diatoms leading to high net BSi production (Closset et al., 2014). Note that higher $\delta^{30}\text{Si}_{\text{DSi}}$ values were also measured in the PFZ WW (on average 2.14 ± 0.08 ‰ for all stations located north of the PF, Fig. 2b). This confirms that the source of Si is different in the PFZ than in the AZ. These PFZ waters can be advected by mixing with waters north of the polar front, including coastal waters, bearing higher $\delta^{30}\text{Si}_{\text{DSi}}$ due to the progressive export of low $\delta^{30}\text{Si}_{\text{BSi}}$ along their eastward advection (Bowie et al., 2015; Park et al., 2014).

3.3 Spatio-temporal variability of Si mass and isotopic balance in the iron fertilized area

By comparing the present dataset with the one collected during KEOPS-1 (summer 2005) the seasonality of the Si isotopic composition can be described, helping to refine the Si biogeochemical cycle in this area (Fripiat et al., 2011a; De Brauwere et al., 2012). To this purpose, the short term temporal evolution of the bloom initiation (two visits at A3) was examined by several visits in the meander that constituted a pseudo-lagrangian time-series (covering 27 days, Table 1). The silicon isotopic properties measured in the ML during the austral spring (KEOPS-2) were significantly lighter than those measured by Fripiat et al. (2011a) at the end of summer (Fig. 3) as expected from the seasonal progression of the preferential light Si isotopes consumption (De la Rocha et al., 1997). Temporally, the bloom above the plateau usually peaks in late November and declines gradually until January as nutrients (mainly iron and silicic acid) became limiting (Blain et al., 2007; Mongin et al., 2008). Then, a second and smaller bloom could persist at steady state until May when light level started to be insufficient to maintain photosynthetic activity (Blain et al., 2013). The first visit to A3 revealed relatively high DSi concentrations and low BSi and chlorophyll *a* stocks in the ML and WW that can be associated to the initial conditions prevailing before the summer stratification (Park et al., 2014; Blain et al. 2015) whereas concentrations and $\delta^{30}\text{Si}$ measured at A3 during KEOPS-1 represent the conditions at the end of the season (Fig. 3). These initial conditions (31.55 ± 2.21 $\mu\text{mol L}^{-1}$ and 1.76 ± 0.03 ‰ for A3-1 WW DSi concentration and isotopic composition respectively) differ from the HNLC WW chosen by Fripiat et al. (2011a) as the ultimate Si-source fuelling the bloom above the Plateau (52.5 ± 3.3 $\mu\text{mol L}^{-1}$ and 1.5 ± 0.0 ‰ for DSi concentration and $\delta^{30}\text{Si}_{\text{DSi}}$, respectively), but were not significantly different from the A3 WW reported by Fripiat et al. (2011a). Indeed, ML $\delta^{30}\text{Si}_{\text{DSi}}$ in the early spring were clearly off the steady state fractionation trend (not shown) when using the same initial Si-pool conditions as in Fripiat et al. (2011a). Thus, it appears that the WW Si-pool isotopic and contents properties of HNLC KEOPS-1 or KEOPS-2 reference stations could not be ascribed as a common Si-source. In the meander, the initial conditions were represented by the Si-properties of TNS6 WW (31.3 $\mu\text{mol L}^{-1}$ and 1.71 ± 0.02 ‰ for DSi concentration and $\delta^{30}\text{Si}_{\text{DSi}}$, respectively) which were similar to those used for the Plateau. This strengthens the idea of a unique Si-source originated from the south and that flow northward above the shallow Kerguelen Plateau to reach finally the PF retroflexion area. Thus, using the mean ACC $^{30}\epsilon$ value of -1.2 ± 0.2 ‰ compiled by Fripiat et al. (2011a) and the averaged Plateau-WW as initial conditions, we will attempt to describe the seasonal dynamic of the Si-biogeochemical cycle in the fertilized area off Kerguelen Islands.

3.3.1 Mixed layer dissolved Si-pool

The seasonal evolution of $\delta^{30}\text{Si}_{\text{DSi}}$ in the fertilized area is best described by steady state fractionation equations (Fig. 4). The steady state model assumes a continuous supply of substrate and thus no change in concentration with time. It should only represent a final snapshot and cannot reflect any temporal evolution. However, when steady state equations are used in a sequential time step, they allow consumption of substrate (Fry, 2006). In the case of the Kerguelen Plateau, Fripiat et al. (2011a) have already shown that the steady state model better describes the seasonality of the silicon isotopes in the ML and it seems that our data confirm this observation. Indeed, in Fig. 4, most of the Plateau and Meander ML $\delta^{30}\text{Si}_{\text{DSi}}$ stations were clearly off the Rayleigh fractionation trend. In contrast all ML $\delta^{30}\text{Si}_{\text{DSi}}$ data align along a steady state trend with decreasing f (i.e. increasing Si utilization): A3-1, A3-2, E1 to E4E Meander stations, E5 and finally A3 KEOPS-1 representing end of growth season. These open system equations are thus appropriate to describe the Si-utilization that occurred above the Kerguelen Plateau where deep and regular mixing events are expected to supply nutrients in surface waters (Park et al., 2008b). Surprisingly, in this model, Meander stations exhibit higher isotopic signatures and higher DSi depletion compared to the productive Plateau stations A3-2 and E4W while they were sampled several days before these two stations. This situation was not due to a higher Si-uptake in the Meander ML as the net BSi production at A3-2 was twice that measured at E5 (Closset et al., 2014), but could be explained by a strong mixing event that occurred just before our second visit to A3. This vertical mixing was induced by strong temporary winds that supplied water with high silicic acid content and light $\delta^{30}\text{Si}$ to the euphotic zone. This would thus decrease the apparent Si utilization (f) and lighten the isotopic signature of seawater. Note that, as previously mentioned, no clear temporal evolution can be shown in the Meander ML $\delta^{30}\text{Si}_{\text{DSi}}$ signatures. The hydrological conditions that lead to ML instabilities and regular mixing of the water masses was likely the reason of this situation as we will discuss in the following.

Using KEOPS-1 HNLC WW characteristics ($52.5 \pm 3.3 \mu\text{mol Si L}^{-1}$ and $1.5 \pm 0.0 \text{‰}$) to represent the initial conditions of WW in the fertilized area and assuming that the system behaves following an open mode, Fripiat et al. (2011a) estimated a seasonal depletion in the ML at $5.0 \pm 0.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$. By identifying a more appropriate Si-source for the Plateau ML averaged Plateau-WW ($32.9 \pm 2.2 \mu\text{mol Si L}^{-1}$ and $1.71 \pm 0.03 \text{‰}$) and using the same equations, we can refine this calculation and reduce the seasonal net BSi production in the upper 100 m to $3.0 \pm 0.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$. This flux is 40 % lower than the previous estimate but corresponds well to the range of published net BSi production values for the AZ (2.4 to $3.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$; Pondaven et al., 2000; Nelson et al., 2002; Pollard et al., 2006) and still balance the total Si supply estimated for the Plateau ML ($4.0 \pm 0.7 \text{ mol Si m}^{-2} \text{ y}^{-1}$, Fripiat et al. (2011a)). The mean net BSi production occurring during the 27 days that separate the two samplings at A3 can be also estimated at $10.7 \text{ mmol m}^{-2} \text{ d}^{-1}$ using the steady state equations and is consistent with the $14.3 \text{ mmol m}^{-2} \text{ d}^{-1}$ computed for the same period by Closset et al. (2014) in their seasonal budget of Si cycle above the Plateau.

3.3.2 Winter Waters dissolved Si-pool

By comparing WW DSi-properties from KEOPS-1 HNLC and fertilized area ($52.5 \pm 3.3 \mu\text{mol Si L}^{-1}$ and $34.2 \pm 1.9 \mu\text{mol Si L}^{-1}$), Fripiat et al. (2011a) estimated a seasonal depletion in the 100 to 400 m layer of $5.5 \pm 0.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$ and ascribe it to a net BSi production by a subsurface diatom population since a deep BSi and chlorophyll maximum was observed here in January (Mosseri et al., 2008; Uitz et al., 2009). The WW characteristics measured during our first visit at A3 ($31.5 \pm 2.2 \mu\text{mol Si L}^{-1}$; $1.76 \pm 0.03 \text{ ‰}$) were similar to the one observed at the end of the productive season by Fripiat et al. (2011a) suggesting that, contrary to what was previously proposed, WW did not undergo seasonal Si uptake above the Plateau. This conclusion is not contradictory to the development of a subsurface diatom community that could partly explain the deep BSi accumulation observed here during KEOPS-1. Indeed, in such communities, the BSi production may likely be sustained by regenerated source of Si (high D:P ratio, see Closset et al., 2014 and Fripiat et al., 2011c at the same station and in the Australian Sector of the Southern Ocean respectively) or may benefit from high diffusive fluxes ($> 1 \text{ mmol m}^{-2} \text{ d}^{-1}$, Rembauville et al., 2016) that characterize these transition layers and would not finally consume the DSi standing stock.

In contrast to the ML, the WW $\delta^{30}\text{Si}_{\text{DSi}}$ signatures of the fertilized area (Table S4) display significant variations (from $1.71 \pm 0.02 \text{ ‰}$ at TNS6 to $2.21 \pm 0.02 \text{ ‰}$ at E4E). It seems unlikely that such shift of WW toward higher $\delta^{30}\text{Si}$ values could be explained by the progressive consumption of silicic acid by a secondary diatom community just below the ML. Indeed, although deep silica maximum are common features in the Southern Ocean (Parslow et al., 2001), the Si uptake rates measured below the euphotic layer during KEOPS-2 were very low (Closset et al., 2014). Since this area has strong mesoscale physical activity (Zhou et al., 2014; Park et al., 2014), an alternative process that could decrease the WW DSi pool and increase the WW $\delta^{30}\text{Si}_{\text{DSi}}$ may be the vertical mixing between initial WW and several sequential surface ML water masses (Fig. 5). Such a process is similar to what was observed for $\delta^{30}\text{Si}_{\text{DSi}}$ across the major ACC meridional overturning circulation and could result here from (i) the deepening of the MLD induced by wind stress and followed by the stratification of the water column; (ii) mesoscale activity that generates negative vertical velocities of water masses. In this case, the resulting water mass should lie on a theoretical mixing curve involving the TNS6 WW and different ML water masses with Si-properties located between the initial and final ML end-members (TNS6 ML and E5 ML). For example, E1 WW could be fully explained by the mixing of TNS6 WW and TNS6 ML, and E2 WW by the mixing between TNS6 WW and E1 ML etc. Although this hypothesis seems valuable for these two examples, the large variability of the Si isotopic properties of WW water masses prevents us from identifying, with sufficient precision, the exact end-members of these mixings. **Moreover, isopycnal mixings through mesoscale eddy activity could also explain this complexity. Indeed, the intense mesoscale activity associated with the meanders of the Polar Front and with major topographic obstacles such as the Kerguelen region, allows the transport of tracers and nutrients laterally through the PF (Dufour et al., 2015).** Consequently, only a range of potential mixing curves has been identified in Fig. 5, but all the Meander WW fall between these two extreme situations. Thus, when the water masses joined the meander, mixing between WW and ML as well as lateral advection led to an enrichment of the Si isotopic signature and a dilution of the DSi pool of this source.

3.3.3 Mixed layer particulate Si-pool

While the open system model seems appropriate to describe the evolution of DSi isotopic composition, it is arguable when considering the $\delta^{30}\text{Si}_{\text{BSi}}$, since this model conceptually assumes the lack of BSi accumulation (Fry, 2009). As shown in Fig. 4, the signature of opal during the two KEOPS expeditions did not fall on the steady-state fractionation trend for BSi but rather fell between the two product-curves involved in the closed-model. The Rayleigh fractionation equations describe two extreme ideal situations: all the BSi produced in the ML is immediately exported from the system (the so-called instantaneous BSi, Eq. 2), or all BSi accumulates in the ML (the so-called accumulated BSi, Eq. 3). During diatom bloom development and through the seasons in the Kerguelen iron fertilized area, the system shifted from one situation to another or remained between these two extremes depending on the BSi accumulation:BSi export ratio which is impacted by the bloom maturity and/or the hydrodynamical conditions of the surface waters. The instability of the ML depth at TNS6, “bloom-initiation” stations (E1, E2 and E3) and at E4E impacted the phytoplankton growth as discussed previously and may have led to important biomass export through detrainment. In these systems, diatoms did not accumulate in the ML but settled toward deeper layers, exporting carbon and BSi (Laurenceau-Cornec et al., 2015; Planchon et al., 2015). The Si-isotopic properties at these stations fell close to the instantaneous product curve of the Rayleigh fractionation law (Fig. 4). Considering the early stage of the season during KEOPS-2 expedition and since Si-uptake rates above the Plateau were among the highest reported so far for the Southern Ocean, it appears that BSi and C export were very low at A3 and E4W (Jacquet et al., 2015; Planchon et al., 2015; Rembauville et al., 2015a, 2015b). At A3 and E4W BSi accumulated in the ML, resulting in Si-isotopic properties that seemed to be better described by the accumulated product curve of the Rayleigh model. The last two stations (E5, KEOPS-2, and A3, KEOPS-1) lied between the two curves since they combined the two biogeochemical processes (BSi accumulation and export). Indeed, the high Si-uptake rates observed at E5 ($20.5 \pm 0.2 \text{ mmol m}^{-2} \text{ d}^{-1}$; Closset et al., 2014) associated with the strengthening of the summer stratification would progressively reduce vertical Si supply and would support higher phytoplankton retention in the ML. KEOPS-1 expedition occurred during the decaying phase of the bloom. Thus, the biological material present in the ML at the end of summer was likely composed of old and detrital diatoms that were partly exported in December (Rembauville et al., 2015b) and living cells that were produced from regenerated Si-sources (Closset et al., 2014). The $\delta^{30}\text{Si}_{\text{BSi}}$ observed in the late-summer surface waters would be a combination of old particles that remained in the ML, with a lighter isotopic signature, and of new and isotopically heavier diatoms that would increase the delta value. This situation is consistent with the strong seasonality of primary and export production that characterizes the Southern Ocean and leads to a temporal decoupling between these two processes in the ML (see e.g. in Rembauville et al., 2015a).

The contrasted seasonal evolution of $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ (following an open or a closed system respectively) is similar to BSi and DSi isotopic offsets previously reported (e.g. Varela et al. 2004; Fripiat et al., 2012) leading to spatio-temporal variations of the ML $\Delta^{30}\text{Si}$ ($\Delta^{30}\text{Si} = \delta^{30}\text{Si}_{\text{BSi}} - \delta^{30}\text{Si}_{\text{DSi}}$), with low $\Delta^{30}\text{Si}$ at the end of summer when the ML silicic acid pool was highly depleted. Fripiat et al. (2012) attributed such offsets to modifications of the Si-uptake:Si-supply ratio in surface waters due to spatial variability of the biogeochemical conditions among the different zones of the ACC (AZ vs. PFZ). **Our results point out an alternative: the change in $\Delta^{30}\text{Si}$ can be obtained when the BSi pool in the system switches between the Rayleigh instantaneous or accumulated product depending on the BSi**

accumulation:BSi export ratio. Indeed, the processes that can affect the isotopic dynamics in this region are complex.

Thus, several explanations are possible to describe the variations of $\Delta^{30}\text{Si}$ we observed (Fig. 4), however it is difficult to discriminate which one is the most relevant:

- The fractionation factor of diatoms ($^{30}\epsilon$) may vary over the season. Since there is no study that so far has focused on $^{30}\epsilon$ seasonal variations, we have considered the fractionation factor as constant. Note that this may not reflect the reality in the ocean and that the constancy of the fractionation factor can be discussed. Moreover one in vitro study has reported inter-specific variations of fractionation factor (Sutton et al. 2013). This could yield to seasonal variation of the mean $^{30}\epsilon$ due to the succession of diatoms species during the bloom (Lasbleiz et al. 2014).
- In the ocean, the distinction between steady state and Rayleigh models can be tricky. Many situations, such as the Kerguelen bloom, can switch between characteristics of both closed and open system dynamics resulting in changes of the apparent fractionation factor ($\Delta^{30}\text{Si}$). As already proposed by Fripiat et al. (2012), we suggest that these variations are mainly controlled by the Si-uptake:Si-supply ratio of the system. When Si-uptake:Si-supply ratio is high (during the bloom period), the system roughly follows a Rayleigh fractionation model. However, when the DSi pool limits Si uptake (at the end of the productive period), the Si-uptake:Si-supply ratio decreases and the supply of isotopically light DSi into the ML by vertical mixing significantly decreases the $\delta^{30}\text{Si}_{\text{DSi}}$. The $\delta^{30}\text{Si}_{\text{BSi}}$ is affected differently since the BSi pool is composed by a mix between newly formed (and light) diatoms and isotopically heavy diatoms that have been produced previously in the ML. Such process would explain why we observe late summer high $\delta^{30}\text{Si}_{\text{BSi}}$ associated with low $\delta^{30}\text{Si}$ of the DSi pool.
- Other isotope effects, such as mixing or BSi dissolution, can affect Si pools differently and would lead to variations of the apparent fractionation factor. Mixings brings isotopically light DSi into the ML and instantaneously decreases the $\delta^{30}\text{Si}$ of the DSi pool while the dissolution of biogenic silica may affect mainly dead and light diatoms that were produced in the beginning of the bloom, resulting in a progressive increase of the $\delta^{30}\text{Si}$ of the BSi pool. Such effects are temporally decoupled and would result in a decoupling of the information recorded in the dissolved and particulate phase.

To conclude, while the closed and open theoretical models seem appropriate for interpreting paleoceanographic isotope records that integrate over longer temporal scales, they have limitations when applied to the modern ocean at shorter and seasonal scales.

3.3.4 Deep particulate Si fluxes

BSi export flux collected in the sediment trap located above the Plateau shows two summer maxima ($> 2 \text{ mmol m}^{-2} \text{ d}^{-1}$ in early December 2011 and early January 2012) separated by a period of reduced flux (Fig. 6). In contrast, export during austral winter exhibited a long period of very low particle flux ($< 0.2 \text{ mmol m}^{-2} \text{ d}^{-1}$, from March to September 2012). These seasonal variations are associated with the evolution of surface chlorophyll *a* concentrations with a delay of approximately one month, and were described in detail in Rembauville et al. (2015a). This sediment trap dataset allows us to investigate the variations of exported opal $\delta^{30}\text{Si}$ and subsequently the mechanisms governing the seasonal variations of Si stocks and fluxes in the ML. Recent studies have suggested that the seasonal variations of the isotopic composition of exported BSi may reflect the evolution of $\delta^{30}\text{Si}_{\text{BSi}}$ in the ML (Varela et al., 2004; Closset et al., 2015). In Fig. 6, we can identify five key periods in the seasonal Si biogeochemical cycle above the Kerguelen Plateau:

(i) Early spring: The relatively low $\delta^{30}\text{Si}$ (1.34 ‰ in October to 1.23 ‰ in December) did not directly relate to the isotopic signature of diatoms in the ML at the same station for the same period (0.79 ± 0.1 ‰ late October, and 1.00 ± 0.09 ‰ mid-November) but did correspond to the signature of particles collected between 200 m and 450 m (1.21 ± 0.1 ‰ late October, A3-1, and 1.34 ± 0.07 ‰ mid-November, A3-2; Fig. 2f). Although Demarest et al. (2009) estimated a fractionation factor of -0.55 ‰ during the dissolution of BSi, it seems unlikely that this process would be responsible for the heavy isotopic signature of the deep particles since dissolution rates are low during early spring (the dissolution to production ratio measured in the ML during KEOPS-2 was very low above the Plateau, less than 0.1, Closset et al., 2014). Moreover, recent studies have pointed out that the isotopic signature of particles is well conserved through the water column and can be closely related to the DSi consumption in surface waters (Fripiat et al., 2012; Closset et al., 2015). Therefore, assuming that the $\delta^{30}\text{Si}$ of exported BSi followed either the accumulated Rayleigh or a steady state equations, the 1.23 ‰ signature measured during the first export event corresponds to a DSi consumption ranging between 45 % and 60 % of the winter DSi standing stock in the ML, respectively, and results in a DSi concentration remaining in the ML ranging between $18 \mu\text{mol L}^{-1}$ (Rayleigh) and $13 \mu\text{mol L}^{-1}$ (steady-state) in November 2011 (one month delay between when the signal was produced in ML and recorded in the trap). The remaining DSi concentration calculated from Rayleigh accumulated product equation is consistent with the DSi concentration measured in the ML during the second visit to A3 in November ($19.7 \pm 0.3 \mu\text{mol L}^{-1}$, Fig. 2a) and confirms that, in early spring, the BSi production in the ML follows more closely to the closed system mode.

(ii) Early summer: During December, the $\delta^{30}\text{Si}$ of settling particles increased from 1.23 to 1.99 ‰ as a result of diatom activity in the ML. These high values cannot be explained by the steady state model. Applying the Rayleigh equations, we estimate a consumption of 71 % of the winter DSi standing stock.

The assumptions behind this estimation are the following:

- The fractionation factor $^{30}\epsilon$ is constant.
- There is no significant BSi export during this period so the BSi accumulation:BSi export ratio is very high.

- There is no significant DSi supply during this period so the Si-supply:Si-uptake ratio is very low.

These two last assumptions are consistent with the situation encountered early summer, i.e. when the bloom has just started and when diatoms accumulate in the ML and have not consumed all the DSi stock. The remaining DSi concentration in the ML would be $9.53 \mu\text{mol L}^{-1}$ in November 2011 which is half of concentration measured at A3 mid-November but which is close to the $12 \mu\text{mol L}^{-1}$ measured mid-November at E5. Such an increase between the two BSi delta values corresponds to a net BSi production of $22.79 \text{ mmol m}^{-2} \text{ d}^{-1}$ (integrated over a 80 m ML as in Closset et al., 2014). This production is half the production measured at A3-2 during 24h incubations ($46.8 \text{ mmol m}^{-2} \text{ d}^{-1}$ Closset et al., 2014). The difference can be explained by the different integration time associated with the two methods: the $\delta^{30}\text{Si}$ of exported BSi integrates one month, while a daily incubation only represents a snapshot of BSi production. The high production rate measured from 24h incubation may not have been maintained at this rate for the whole December. Additionally, DSi supply also occurs in summer (Fripiat et al., 2011a; Closset et al. 2014), a process that is not included in Rayleigh equations but causes an underestimate BSi production.

(iii) Mid-summer. The sharp decrease of $\delta^{30}\text{Si}$ of settling particles observed in January 2012 (from 1.99 to 1.72 ‰) can be associated with a strong vertical mixing event that brings new and isotopically light DSi in the ML. This phenomenon has already been observed in open ocean sediment traps time-series in the Antarctic Zone (Varela et al., 2004; Closset et al., 2015). Assuming that at this time of vertical mixing, the system behaves following an open mode (Fig. 4), this 0.27 ‰ $\delta^{30}\text{Si}_{\text{BSi}}$ decrease would correspond to the same change in $\delta^{30}\text{Si}_{\text{DSi}}$ and to an increase of $7.56 \mu\text{mol L}^{-1}$ of the DSi concentration in the ML (1.99 and 1.72 ‰ correspond to a DSi concentration of 32.54 and $24.98 \mu\text{mol L}^{-1}$, respectively). The assumptions behind this calculation are the following:

- The fractionation factor $^{30}\epsilon$ is constant.
- The DSi supplied into the system has the same isotopic signature as the source. This assumption is realistic since the mixing event brings DSi from WW (source) into the ML.
- All the BSi and the remaining DSi should be removed from the system. We cannot verify this last assumption but we can expect that a large fraction of the BSi was exported out of the ML mid-summer. This might be supported by the deep silica maximum observed late summer below the ML (150 m) by Mosseri et al. (2008) at the same station.

The subsequent DSi stock in the ML would be $17.06 \mu\text{mol L}^{-1}$ in December 2011. Using a simplistic approach (see supplementary method) we estimate that such Si supply would require a ML deepening of ca. 37 m which is consistent with the order of magnitude of ML variation in this area (Park et al., 2008b). High wind events could induce vertical mixing of the upper ocean, entraining cold water into the ML and bringing DSi into the euphotic zone. Integrating the DSi increase of $7.56 \mu\text{mol L}^{-1}$ over this new ML (80 m + 37 m) suggests a supply of 884 mmol m^{-2} . Such a mixing event would represent approximately 74 % of the total summer supply (1200 mmol m^{-2} ; Closset et al., 2014) and would allow the second bloom to appear as observed in January 2012 (Rembauville et al., 2015a). Sporadic upwelling through Ekman pumping above the Plateau (see Gille et al., 2014) could also bring DSi into the ML, but considering its low intensity, it is unlikely that it could sustain the needed flux of DSi. Note that if the closed model is considered, the $\delta^{30}\text{Si}_{\text{BSi}}$ signal would correspond to a consumption of 64 %

of the winter DSi stock and thus an increase of the DSi concentration of $2.3 \mu\text{mol L}^{-1}$ in the ML. This value would imply a deepening of the ML by only 8.75 m and a summer supply of 204 mmol m^{-2} which does not correspond to the one estimated by Closset et al. (2014). This highlights, as proposed previously, that BSi isotopic and mass balance can follow either a closed or an open mode depending on physical and biogeochemical conditions which vary during the growing season.

(iv) Late summer: The $\delta^{30}\text{Si}$ increased from 1.72 to 2.54 ‰ due to a second episode of enhanced biological activity in the ML. Using the Rayleigh equations (Fig. 4), this highest value corresponds to a consumption of 82 % of the winter DSi standing stock which results in a ML DSi concentration of $6.05 \mu\text{mol L}^{-1}$ and thus a consumption of $10.87 \mu\text{mol L}^{-1}$ of the DSi concentration in the ML between December 2011 and the end of January 2012 (52 days). Integrated over a 80 m ML, this value corresponds to a net BSi production of $16 \text{ mmol m}^{-2} \text{ d}^{-1}$. This production is consistent with the mean net BSi production of $10 \text{ mmol m}^{-2} \text{ d}^{-1}$ estimated for the same period by Closset et al. (2014) which decreases progressively to values lower than $1 \text{ mmol m}^{-2} \text{ d}^{-1}$.

(v) Winter: From March to September, the $\delta^{30}\text{Si}$ of settling particles remained relatively heavy but decreased gradually toward lower values as winter deep convection progressively took place above the Plateau (Blain et al., 2013).

4 Conclusions

The spatial distribution of $\delta^{30}\text{Si}$ of seawater and particulate matter in the ML was strongly impacted by the complex structure of water masses generated by the interaction between the PF and the bathymetry. In contrast, in deeper layers ($> 500 \text{ m}$), $\delta^{30}\text{Si}_{\text{DSi}}$ were remarkably homogeneous (on average $1.28 \pm 0.08 \text{ ‰}$ and $1.05 \pm 0.06 \text{ ‰}$ for UCDW and LCDW respectively) and deep $\delta^{30}\text{Si}_{\text{BSi}}$ exhibited constant values in all out-plateau KEOPS-2 stations (on average $1.74 \pm 0.13 \text{ ‰}$), suggesting that dissolution of opal did not have any significant isotopic effect during this early season.

The measured silicon properties of the ML can be considered as representative of the bloom initiation as also supported by companion studies from the same cruise (Blain et al., 2015; Cavagna et al., 2015; Closset et al., 2014; Lasbleiz et al., 2014). During this period, the Kerguelen area was characterized by a mosaic of biogeochemical environments (Trull et al., 2015). The HNLC area, strongly iron-limited station R2, exhibited very low biomass, low BSi-production and isotopically light BSi. However, moderate production in agreement with other proxies (Closset et al., 2014; Dehairs et al., 2015; Jacquet et al., 2015) could explain the relatively high $\delta^{30}\text{Si}_{\text{DSi}}$ values measured at this station, precluding its application as non-productive reference station for silicon biogeochemical cycle. The iron flux in the ML above the Plateau strongly stimulated diatom production, increasing significantly its $\delta^{30}\text{Si}_{\text{BSi}}$ (from $0.77 \pm 0.05 \text{ ‰}$ to $0.96 \pm 0.08 \text{ ‰}$ over 27 days during KEOPS-2) and simultaneously enriching the ML in ^{30}Si (up to $2.10 \pm 0.05 \text{ ‰}$). North of the PF, stocks and Si-isotopic composition of seawater and particles

were respectively lower and heavier than above the Plateau in accordance with its PFZ characteristics as already reported by Cardinal et al. (2007, 2005), Varela et al. (2004) and Fripiat et al. (2011b) in other sectors of the Southern Ocean. The situation was different in the Meander since these stations received only moderate and sporadic iron supplies. There, non-optimal light-mixing regime and nutrients availability (including iron) delayed the bloom development and lead the system to behave near steady-state with low DSi utilization. Indeed, both Si isotopic signatures and mass balance did not evolve significantly in the Meander ML.

Our Si-isotopic data allow identification of the initial conditions before diatom growth, Si sources, and the connections between different water masses through local circulation. The same Si source (WW Plateau and not WW HNLC as initially thought) can be used for both the Plateau and Meander stations, suggesting that water masses and the ML bloom originated from the southeast part of the Kerguelen Plateau and spread northward. The Si-properties of this source ($32.9 \pm 2.2 \mu\text{mol Si L}^{-1}$ and $1.71 \pm 0.03 \text{‰}$) allow us to refine the seasonal net BSi production at $3.0 \pm 0.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$ above the Plateau (instead of $10.5 \pm 1.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$ estimated by Fripiat et al., 2011a), which is more consistent with the published values for the AZ (2.4 to $3.3 \text{ mol Si m}^{-2} \text{ y}^{-1}$; e.g. Pondaven et al., 2000; Nelson et al., 2002). This suggests that, even if the iron-fertilization in some regions of the Southern Ocean stimulates the uptake of C and N and the production of organic matter compared to non-fertilized area, it does not necessary enhance consumption of silicic acid and production of BSi. This observation can be explained by the decoupling between the biogeochemical cycles of Si, C and N of diatom communities living under different biogeochemical conditions. These distinctive local communities may induce variations of the efficiency of the silicon pump (Dugdale et al., 1995) that preferentially recycles organic matter over biogenic silica in surface waters and thus lead to moderate net BSi production despite high primary production. *Indeed, other recent studies in the Southern Ocean have highlighted that diatom community composition could better explain differences in silicification than physiological response to iron enrichment (see Baines et al., 2010; Assmy et al., 2013).* Moreover, the isotope characteristics of the WW was similar through the whole season meaning there was no net BSi production above the Plateau. In contrast, in the Meander the variation of the WW Si characteristics (concentration and isotopic composition) can be explained by successive mixings between ML and WW or by lateral advection.

In the naturally Fe-fertilized area of Kerguelen, we have shown that the dissolved and particulate Si-pools were temporally decoupled. In this highly productive area, strong activity that promoted vertical and lateral nutrient exchanges, drove the evolution of $\delta^{30}\text{Si}_{\text{DSi}}$ to follow the steady state equations, while the $\delta^{30}\text{Si}_{\text{BSi}}$ seemed to follow more closely the Rayleigh equations. Depending on the regime of BSi export (e.g. BSi accumulation in the ML in early spring vs. massive export event at the end of the productive season), the ML $\delta^{30}\text{Si}_{\text{BSi}}$ can be described by the accumulation or instantaneous product, or could lie between these two ideal situations. If confirmed in other productive regions of the Southern Ocean, this observation could have great implications for paleoceanographic studies. Indeed, the $\delta^{30}\text{Si}_{\text{BSi}}$ of sediments is currently used as a proxy for past reconstructions of surface Si-utilization (e.g. De la Rocha et al., 1998; Ehlert et al., 2013). In these studies, the isotopic signature of sedimentary opal is used to calculate the $\delta^{30}\text{Si}_{\text{DSi}}$ and the extent of DSi consumption in surface waters using either

the Rayleigh's or steady state equations without taking into account the possible decoupling between $\delta^{30}\text{Si}_{\text{DSi}}$ and $\delta^{30}\text{Si}_{\text{BSi}}$ and its seasonal variability.

Finally, we have identified and quantified the processes that control the biogeochemical cycle of silicon in the ML over nearly one complete year. We show that the $\delta^{30}\text{Si}$ of settling diatoms collected in sediment traps is a powerful proxy to quantify Si fluxes in the ML, such as silicic acid consumption in spring and summer or Si supply to ML during mixing events.

Acknowledgements. The authors would like to thank B. Quéguiner as the KEOPS-2 chief scientist, the captain and the crew of the R/V Marion-Dufresne II for assistance on board and Heather McNair for improvement of English. We are grateful to G. de Souza and one anonymous reviewer for their constructive comments that helped us to improve the manuscript. The research leading to these results on silicon has been funded by the European Union Seventh Framework Programme under grant agreement n°294146 (MuSiCC Marie Curie CIG). KEOPS-2 was supported by the French Research program of INSU-CNRS LEFE-CYBER ('Les enveloppes fluides et l'environnement' – 'Cycles biogéochimiques, environnement et ressources'), the French ANR ('Agence Nationale de la Recherche', SIMI-6 program), the French CNES ('Centre National d'Etudes Spatiales') and the French Polar Institute IPEV (Institut Polaire Paul-Emile Victor).

References

Abraham, K., Opfergelt, S., Fripiat, F., Cavagna, A.-J., de Jong, J. T. M., Foley, S. F., André, L., Cardinal, D.: $\delta^{30}\text{Si}$ and $\delta^{29}\text{Si}$ Determination on USGS BHVO-1 and BHVO-2 Reference Materials with a New Configuration on a Nu Plasma Multi-Collector ICP-MS, *Geostandards and Geoanalytical Research*, 32(2), 193-202, 2008.

Blain, S., Tréguer, P., Belviso, S., Bucciarelli, E., Denis, M., Desabre, S., Fiala, M., Martin Jézéquel, V., Le Fèvre, J., Mayzaud, P., Marty, J.-C., Razouls, S.: A biogeochemical study of the island mass effect in the context of the iron hypothesis: Kerguelen Islands, Southern Ocean, *Deep-Sea Research I*, 48, 163-187, 2001.

Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillermin, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., Wagener, T.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, *Nature*, 446, 1070-1075, doi:10.1038/nature05700, 2007.

Blain, S., Renaut, S., Xing, X., Claustre, H., Guinet, C.: Instrumented elephant seals reveal the seasonality in chlorophyll and light-mixing regime in the iron-fertilized Southern Ocean, *Geophysical Research Letters*, 40, doi:10.1002/2013GL058065, 2013.

Blain, S., Capparos, J., Guéneuguès, A., Obernosterer, I., Oriol, L.: Distributions and stoichiometry of dissolved nitrogen and phosphorus in the iron fertilized region near Kerguelen (Southern Ocean), *Biogeosciences*, 12, 623-635, doi:105194/bg-12-623-2015, 2015.

Bowie, A. R., Van der Merwe, P., Quéroùé, F., Trull, T., Fourquez, M., Planchon, F., Sarthou, G., Chever, F., Townsend, A. T., Obernosterer, I., Sallée, J.-B., Blain, S.: Iron budgets for three distinct biogeochemical sites around the Kerguelen archipelago (Southern Ocean) during the natural fertilization experiment KEOPS-2. *Biogeosciences*. 12, 4421-4445, doi:10.5194/bg-12-4421-2015, 2015.

Boyd, P. W and Trull, T. W.: Understanding the export of biogenic particles in oceanic waters: Is there consensus?, *Progress in Oceanography*, 72, 276-312, 2007.

Brzezinski, M. A., Nelson, D. M., Franck, V. M., Sigmon, D. E.: Silicon dynamics within an intense open-ocean diatom bloom in the Pacific sector of the Southern Ocean, *Deep-sea research II*, 48, 3997-4018, 2001.

Brzezinski, M. A., Pride, C. J., Frank, V. M.: A switch from Si(OH)_4 to NO_3^- depletion in the glacial Southern Ocean, *Geophysical research letters*, 29(12), doi:10.1029/2001GL014349, 2002.

Bucciarelli, E., Pondaven, P., Sarthou, G.: Effects of an iron-light co-limitation on the elemental composition (Si, C, N) of the marine diatoms *Thalassiosira oceanica* and *Ditylum brightwellii*, *Biogeosciences*, 7, 657-669, 2010.

Buesseler, K. O.: The decoupling of production and particulate export in the surface ocean, *Global Biogeochemical Cycles*, 12(2), 297-310, 1998.

Cao, Z., Frank, M., Dai, M., Grasse, P., Ehlert, C.: Silicon isotope constraints on sources and utilization of silicic acid in the northern South China Sea, *Geochimica et Cosmochimica Acta*, 97, 88-104, 2012.

Cardinal, D., Alleman, L. Y., De Jong, J., Ziegler, K., André, L.: Isotopic composition of silicon measured by multicollector plasma source mass spectrometry in dry plasma mode, *Journal of Analytical Atomic Spectrometry*, 18, 213-218, 2003.

Cardinal, D., Alleman, L. Y., Dehairs, F., Savoye, N., Trull, T. W., André, L.: Relevance of silicon isotopes to Si-nutrient utilization and Si-source assessment in Antarctic waters, *Global biogeochemical cycles*, 19(GB2007), doi:10.1029/2004GB002364, 2005.

Cardinal, D., Savoye, N., Trull, T. W., Dehairs, F., Kopczynska, E. E., Fripiat, F., Tison, J.-L., André, L.: Silicon isotopes in spring Southern Ocean diatoms: Large zonal changes despite homogeneity among size fractions, *Marine Chemistry*, 106, 46-62, 2007.

Cavagna, A.-J., Fripiat, F., Dehairs, F., Wolf-Gladrow, D., Cisewski, B., Savoye, N., André, L., Cardinal, D.: Silicon uptake and supply during a Southern Ocean iron fertilization experiment (EIFEX) tracked by Si isotopes, *Limnology and Oceanography*, 56(1), 147-160, doi:10.4319/lo.2011.56.1.0147, 2011.

Cavagna, A.-J., Fripiat, F., Elskens, M., Dehairs, F., Mangion, P., Chirurgien, L., Closset, I., Lasleiz, M., Flores-Leiva, L., Cardinal, D., Leblanc, K., Fernandez, C., Lefèvre, D., Oriol, L., Blain, S., Quéguiner, B.: Production regime and associated N cycling in the vicinity of Kerguelen Island, Southern Ocean, *Biogeosciences*, 12, 6515-6528, doi:10.5194/bg-12-6515-2015, 2015.

Closset, I., Lasleiz, M., Leblanc, K., Quéguiner, B., Cavagna, A.-J., Elskens, M., Navez, J., Cardinal, D.: Seasonal evolution of net and regenerated silica production around a natural Fe-fertilized area in the Southern Ocean estimated with Si isotopic approaches, *Biogeosciences*, 11, 5827-5846, doi:10.5194/bg-11-5827-2014, 2014.

Closset, I., Cardinal, D., Bray, S., Thil, F., Djoureaev, I., Rigual-Hernandez, A. S., Trull, T.: Seasonal variations, origin and fate of settling diatoms in the Southern Ocean tracked by silicon isotope records in deep sediment traps, *Global Biogeochemical Cycles*, 29, 1495-1510, doi:10.1002/2015GB005180, 2015.

Coffineau, N., De la Rocha, C. L., Pondaven, P.: Exploring interacting influences on the silicon isotopic composition of the surface ocean: a case study from the Kerguelen Plateau, *Biogeosciences*, 11, 1371-1391, 2014.

De Baar, H. J. W., Boys, P. W., Coale, K. H., Landry, M. R., Tsuda, A., Assmy, P., Bakker, D. C. E., Bozec, Y., Barber, R. T., Brzezinski, M. A., Buesseler, K. O., Boyé, M., Croot, P. L., Gervais, F., Gorbunov, M. Y., Harrison, P. J., Hiscock, W. T., Laan, P., Lancelot, C., Law, C. S., Lvasseur, M., Marchetti, A., Millero, F. J., Nishioka, J., Nojiri, Y., van Oijen, T., Riebesell, U., Rijkenberg, M. J. A., Saito, H., Takeda, S., Timmermans, K. R., Veldhuis, M. J. W., Waite, A. M., Wong, C.-S.: Synthesis of iron fertilization experiments: From the Iron Age in the Age of Enlightenment, *Journal of Geophysical Research*, 110, C09S16, doi:10.1029/2004JC002601, 2005.

De Brauwere, A., Fripiat, F., Cardinal, D., Cavagna, A.-J., De Ridder, F., André, L., Elskens, M.: Isotopic model of oceanic silicon cycling: The Kerguelen Plateau case study, *Deep-Sea Research I*, 70, 42-59, 2012.

Dehairs, F., Fripiat, F., Cavagna, A.-J., Trull, T. W., Fernandez, C., Davies, D., Roukaerts, A., Fonseca Batista, D., Planchon, F., Elskens, M.: Nitrogen cycling in the Southern Ocean Kerguelen Plateau area: evidence for significant surface nitrification from nitrate isotopic compositions, *Biogeosciences*, 15, 1459-1482, doi:10.5194/bg-12-1459-2015, 2015.

De La Rocha, C.L., Brzezinski, M. A., DeNiro, M. J.: Purification, Recovery, and Laser-Driven Fluorination of Silicon from Dissolved and Particulate Silica for the Measurement of Natural Stable Isotope Abundances, *Analytical Chemistry*, 68(21), 3746-3750, 1996.

De La Rocha, C.L., Brzezinski, M. A., DeNiro, M. J.: Fractionation of silicon isotopes by marine diatoms during biogenic silica formation, *Geochimica et Cosmochimica Acta*, 61(23), 5051-5056, 1997.

De La Rocha, C.L., Brzezinski, M. A., DeNiro, M. J., Shemesh, A.: Silicon-isotope composition of diatoms as an indicator of past oceanic change, *Nature*, 395, 680-683, 1998.

De La Rocha, C.L., Hutchins, D. A., Brzezinski, M. A., Zhang, Y.: Effects of iron and zinc deficiency on elemental composition and silica production by diatoms, *Marine Ecology Progress Series*, 195,71-79, 2000.

De Souza, G. F., Reynolds, B. C., Rickli, J., Frank, M., Saito, M. A., Gerringa, L. J. A., Bourdon, B.: Southern Ocean control of silicon stable isotope distribution in the deep Atlantic Ocean, *Global Biogeochemical Cycles*, 26, GB2035, doi:10.1029/2011GB004141, 2012.

Demarest, M.S., Brzezinski, M. A., Beucher, C. P.: Fractionation of silicon isotopes during biogenic silica dissolution, *Geochimica et Cosmochimica Acta*, 73, 5572-5583, doi:10.1016/j.gca.2009.06.019, 2009.

Dufour, C.O., Griffies, S.M., De Souza, G.F., Frenger, I., Morrison, A.K., Palter, J.B., Sarmiento, J.L., Galbraith, E.D., Dunne, J.P., Anderson, W.G., Salter, R.D.: Role of Mesoscale Eddies in Cross-Frontal Transport of Heat and Biogeochemical Tracers in the Southern Ocean, *Journal of Physical Oceanography*, doi:10.1175/JPO-D-14-0240.1, 2015.

Dugdale, R.C., Wilkerson, F. P., Minas, H. J.: The role of a silicate pump in driving new production, *Deep-Sea Research I*, 42(5), 697-719, 1995.

Ehlert, C., Grasse, P., Frank, M.: Changes in silicate utilization and upwelling intensity off Peru since the Last Glacial Maximum – insights from silicon and neodymium isotopes, *Quaternary Science Reviews*, 72, 18-35, 2013.

Etourneau, J., Ehlert, C., Frank, M., Martinez, P., Scheider, R.: Contribution of changes in opal productivity and nutrient distribution in the coastal upwelling systems to Late Pliocene/Early Pleistocene climate cooling, *Climate of the Past*, 8, 1435-1445, 2012.

Frank, V. M., Brzezinski, M. A., Coale, K. H., Nelson, D. M.: Iron and silicic acid concentrations regulate Si uptake north and south of the Polar Frontal Zone in the Pacific Sector of the Southern Ocean, *Deep-Sea Research II*, 47,3315-3338, 2000.

Fripiat, F., Cavagna, A.-J., Savoye, N., Dehairs, F., André, L., Cardinal, D.: Isotopic constraints on the Si-biogeochemical cycle of the Antarctic Zone in the Kerguelen area (KEOPS), *Marine Chemistry*, 123, 11-22, doi:10.1016/j.marchem.2010.08.005, 2011a.

Fripiat, F., Cavagna, A.-J., Dehairs, F., Speich, S., André, L., Cardinal, D.: Silicon pool dynamics and biogenic silica export in the Southern Ocean inferred from Si-isotopes, *Ocean Science*, 7, 533-547, doi:10.5194/os-7-533-2011, 2011b.

Fripiat, F., Leblanc K., Elskens M., Cavagna A.-J., Armand L., André L., Dehairs F., Cardinal D.: Efficient silicon recycling in summer in both the Polar Frontal and Subantarctic Zones of the Southern Ocean, *Marine Ecology Progress Series*, 435, 47-61, doi:10.3354/meps09237, 2011c.

Fripiat, F., Cavagna, A.-J., Dehairs, F., de Brauwere, A., André, L., Cardinal, D.: Processes controlling the Si-isotopic composition in the Southern Ocean and application for paleoceanography, *Biogeosciences*, 9, 2443-2457, doi:10.5194/bg-9-2443-2012, 2012.

Fry, B.: *Stable Isotope Ecology*, Springer, New York, pp 308, 2006

Georg, R. B., Reynolds, B. C., Frank, M., Halliday, A. N.: New sample preparation techniques for the determination of Si isotopic compositions using MC-ICPMS, *Chemical Geology*, 235, 95-104, 2006.

Gille, S. T., Carranza, M. M., Cambra, R., Morrow, R.: Wind-induced upwelling in the Kerguelen Plateau region, *Biogeosciences*, 11, 6389-6400, doi:10.5194/bg-11-6389-2014, 2014.

Grasshoff, K., Kremling, K., Ehrhardt, M.: *Methods of Seawater Analysis* 3rd edition completely Revised and Extended edition. Wiley-VCH. ISBN 3-527-29589-5, 1999.

Hughes, H. J., Delvigne, C., Korntheuer, M., De Jong, J., André, L., Cardinal, D.: Controlling the mass bias introduced by anionic and organic matrices in silicon isotopic measurements, *Journal of Analytical Atomic Spectrometry*, 26, 1892-1896, 2011.

Hutchins, D. A., Bruland, K. W.: Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime, *Nature*, 393, 561-564, 1998.

Jacquet, S. H. M., Dehairs, F., Cavagna, A.-J., Planchon, F., Monin, L., André, L., Closset, I., Cardinal, D.: Early season mesopelagic carbon remineralization and transfer efficiency in the naturally iron-fertilized Kerguelen area, *Biogeosciences*, 12, 1713-1731, doi:10.5194/bg-12-1713-2015, 2015.

Karl, D. M. and Tien, G.: MAGIC: A sensitive and precise method for measuring dissolved phosphorus in aquatic environments, *Limnology and Oceanography*, 37(1), 105-116, 1992.

Laurenceau-Cornec, E. C., Trull, T. W., Davies, D. M., Bray, S. G., Doran, J., Planchon, F., Carlotti, F., Jouandet, M.-P., Cavagna, A.-J., Waite, A. M., Blain, S.: The relative importance phytoplankton aggregates and zooplankton fecal pellets to carbon export: insights from free-drifting sediment trap deployments in naturally iron-fertilised waters near the Kerguelen plateau, *Biogeosciences*, 12, 1007-1027, doi:10.5195/bg-12-1007-2015, 2015.

Lasbleiz, M., Leblanc, K., Blain, S., Ras, J., Cornet-Barthaux, V., Hélias Nunige, S., Quéguiner, B.: Pigments, elemental composition (C, N, P and Si), and stoichiometry of particulate matter in the naturally iron fertilized region of Kerguelen in the Southern Ocean, *Biogeosciences*, 11, 5931-5955, doi:10.5194/bg-11-5931-2014, 2014.

Leynaert, A., Bucciarelli, E., Claquin, P., Dugdale, R. C., Martin-Jézéquel, V., Pondaven, P., Raguegeau, O.: Effect of iron deficiency on diatom cell size and silicic acid uptake kinetics, *Limnology and Oceanography*, 49(4), 1134-1143, 2004.

Matsumoto, K., Sarmiento, J. L.: A corollary to the silicic acid leakage hypothesis, *Paleoceanography*, 23, PA2203, doi:10.1029/2007PA001515, 2008.

Matsumoto, K., Sarmiento, J. L., Brzezinski, M. A.: Silicic acid leakage from the Southern Ocean: A possible explanation for glacial atmospheric pCO₂, *Global Biogeochemical Cycles*, 16(3), doi:10.1029/2001GB001442, 2002.

Mongin, M., Molina, E. , Trull, T. W.: Seasonality and scale of the Kerguelen plateau phytoplankton bloom: A remote sensing and modeling analysis of the influence of natural iron fertilization in the Southern Ocean, *Deep-Sea Research II*, 55, 880-892, 2008.

Mosseri, J., Quéguiner, B., Armand, L., Cornet-Barthaux, V.: Impact of iron on silicon utilization by diatoms in the Southern Ocean: A case study of Si/N cycle decoupling in a naturally iron-enriched area, *Deep-sea research II*, 55, 801-819, 2008.

Nelson, D.M., Brzezinski, M. A., Sigmon, D. E., Franck, V. M.: A seasonal progression of Si limitation in the Pacific sector of the Southern Ocean, *Deep-sea research II*, 48, 3973-3995, 2001.

Nelson, D.M., Anderson, R. F., Barber, R. T., Brzezinski, M. A., Buesseler, K. O., Chase, Z., Collier, R. W., Dickson, M.-L., François, R., Hiscock, M. R., Honjo, S., Marra, J., Martin, W. R., Sambrotto, R. N., Sayles, F. L., Sigmon, D. E.: Vertical budgets for organic carbon and biogenic silica in the Pacific sector of the Southern Ocean, 1996-1998, *Deep-Sea Research II*, 49, 1645-1674, 2002.

Opfergelt, S., Delmelle, P.: silicon isotopes and continental weathering processes: Assessing controls on Si transfer to the ocean, *Comptes Rendus Geoscience*, 344, 723-738, doi:10.1016/j.crte.2012.09.006, 2012.

Park, Y.-H., Roquet, F., Durand, I., Fuda, J.-L.: Large-scale circulation over and around the Northern Kerguelen Plateau, *Deep-Sea Research II*, 55, 566-581, 2008a.

Park, Y.-H., Fuda, J.-L., Durand, I., Naveira Garabato, A. C.: Internal tides and vertical mixing over the Kerguelen Plateau, *Deep-Sea Research II*, 55, 582-593, 2008b.

Park, Y.-H., Durand, I., Kestenare, E., Rougier, G., Zhou, M., d'Ovidio, F., Cotté, C., Lee, J.-H.: Polar Front around the Kerguelen Islands: An up-to-date determination and associated circulation of surface/subsurface water *Journal of Geophysical Research: Oceans*, 119, doi:10.1002/2014JC010061, 2014.

Parslow, J. S., Boyd, P. W., Rintoul, S. R., Griffiths, F. B.: A persistent subsurface chlorophyll maximum in the Interpolar Frontal Zone south of Australia: Seasonal progression and implications for phytoplankton-light-nutrient interactions, *Journal of geophysical research*, 106(C12), 31543-31557, 2001.

Planchon, F., Ballas, D., Cavagna, A.-J., Bowie, A. R., Davies, D., Trull, T., Laurenceau-Cornec, E., Van der Merwe, P., Dehairs, F.: Carbon export in the naturally iron-fertilized Kerguelen area of the Southern Ocean based on the ²³⁴Th approach, *Biogeosciences*, 12, 3831-3848, doi:10.5195/bg-12-3831-2015, 2015.

Pollard, R. T., Lucas, M. I., Read, J. F.: Physical controls on biogeochemical zonation in the Southern Ocean, *Deep-Sea Research II*, 49, 3289-3305, 2002.

Pollard, R. T., Tréguer, P., Read, J.: Quantifying nutrient supply to the Southern Ocean, *Journal of Geophysical Research*, 111, C05011, doi:10.1029/2005JC003076, 2006.

Pollard, R. T., Salter, I., Sanders, R. J., Lucas, M. I., Moore, C. M., Mills, R. A., Statham, P. J., Allen, J. T., Baker, A. R., Bakker, D. C. E., Charette, M. A., Fielding, S., Fones, G. R., French, M., Hickman, A. E., Holland, R. J., Hughes, J. A., Jickells, T. D., Lampitt, R. S., Morris, P. J., Nédélec, F. H., Nielsdóttir, M., Planquette, H., Popova, E. E., Poulton, A. J., Read, J. F., Seeyave, S., Smith, T., Stinchcombe, M., Taylor, S., Thomalla, S., Venables, H. J., Williamson, R., Zubkov, M. V.: Southern Ocean deep-water carbon export enhanced by natural iron fertilization. *Nature* 457, 577–580. doi:10.1038/nature07716, 2009.

Pondaven, P., Ragueneau, O., Tréguer, P., Hauvespre, A., Dezileau, L., Reyss, J. L.: Resolving the 'opal paradox' in the Southern Ocean, *Nature*, 405, 168-172, 2000.

Quéguiner, B.: Iron fertilization and the structure of planktonic communities in high nutrient regions of the Southern Ocean, *Deep-Sea Research II*, 90, 43-54, 2013.

Quéroué, F., Sarthou, G., Planquette, H. F., Bucciarelli, E., Chever, F., van der Merwe, P., Lannuzel, D., Townsend, A. T., Cheize, M., Blain, S., d'Ovidio, F., Bowie, A. R.: High variability in dissolved iron concentrations in the vicinity of the Kerguelen Islands (Southern Ocean), *Biogeosciences*, 12, 3869-3883, doi:10.5194/bg-12-3869-2015, 2015.

Ragueneau, O., Tréguer, P., Leynaert, A., Anderson, R. F., Brzezinski, M. A., DeMaster, D. J., Dugdale, R. C., Dymond, J., Fischer, G., François, R., Heinze, C., Maier-Reimer, E., Martin-Jézéquel, V., Nelson, D. M., Quéguiner, B.: A review of the Si cycle in the modern ocean: recent progress and missing gaps in the application of biogenic opal as a paleoproductivity proxy, *Global and Planetary Change*, 26, 317-365, 2000.

Ragueneau, O., Savoye, N., Del Amo, Y., Cotton, J., Tardiveau, B., Leynaert, A.: A new method for the measurement of biogenic silica in suspended matter of coastal waters: using Si:Al ratios to correct for the mineral interference, *Continental Shelf Research*, 25, 697-710, 2005.

Rembauville, M., Blain, S., Salter, I.: Export fluxes in a naturally fertilized area of the Southern Ocean – Part 1: Seasonal dynamics of particulate organic carbon export from a moored sediment trap, *Biogeosciences*, 12, 3171-3195, doi:10.5194/bg-12-3171-2015, 2015a.

Rembauville, M., Blain, S., Armand, L., Quéguiner, B., Salter, I.: Export fluxes in a naturally iron-fertilized area of the Southern Ocean – Part 2 : Importance of diatom resting spores and faecal pellets for export, *Biogeosciences*, 12, 3171-3195, doi:10.5194/bg-12-3171-2015, 2015b.

Rembauville, M., Blain, S., Caparros, J., Salter, I.: Particulate matter stoichiometry driven by microplankton community structure in summer in the Indian sector of the Southern Ocean, *Limnology and Oceanography*, doi:10.1002/lno.10291, 2016.

Reynolds, B. C., Frank, M., Halliday, A. N.: Silicon isotope fractionation during nutrient utilization in the North Pacific, *Earth and Planetary Science Letters*, 244, 431-443, 2006.

Reynolds, B. C., Aggarwal, J., André, L., Baxter, D., Beucher, C., Brzezinski, M. A., Engström, E., Georg, R. B., Land, M., Leng, M. J., Opfergelt, S., Rodushkin, I., Sloane, H. J., van den Boorn, S. H. J. M., Vroon, P. Z., Cardinal, D.: An inter-laboratory comparison of Si isotope reference materials, *Journal of Analytical Atomic Spectrometry*, 22, 561-568, doi:10.1039/b616755a, 2007.

Roquet, F., Park, Y.-H., Guinet, C., Bailleul, F., Charrassin, J.-B.: Observations of the Fawn Trough Current over the Kerguelen Plateau from instrumented elephant seals, *Journal of Marine Systems*, 78, 377-393, 2009.

Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace, D. W. R., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., Rios, A. F.: The Ocean Sink for Anthropogenic CO₂, *Science*, 305, 367-371, 2004.

- Sanial, V., van Beek, P., Lansard, B., Souhaut, M., Kestenare, E., d'Ovidio, F., Zhou, M., Blain, S.: Use of Ra isotopes to deduce rapid transfer of sediment-derived inputs off Kerguelen, *Biogeosciences*, 12, 1415-1430, doi:10.5194/bg-12-1415-2015, 2015.
- Sarmiento, J. L., Gurber, N., Brzezinski, M. A., Dunne, J. P.: High-latitude controls of thermocline nutrients and low latitude biological productivity, *Nature*, 427, 56-60, 2004.
- Sutton, J. N., Varela, D. E., Brzezinski, M. A., Beucher, C. P.: Species-dependent silicon isotope fractionation by marine diatoms, *Geochimica et Cosmochimica Acta*, 104, 300-309, 2013.
- Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich, G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M., Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., Hoppema, M., Olafsson, J., Arnarson, T. S., Tilbrook, B., Johannessen, T., Olsen, A., Bellerby, R., Wong, C. S., Delille, B., Bates, N. R., de Baar, H. J. W.: Climatological mean and decadal change in surface ocean pCO₂, and net sea-air CO₂ flux over the global oceans, *Deep-Sea Research II*, 56, 554-577, doi: 10.1016/j.dsr2.2008.12.009, 2009.
- Takeda, S.: Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters, *Nature*, 393, 774-777, 1998.
- Tréguer, P. J. and De La Rocha, C. L.: The World Ocean Silica Cycle, *Annual Review of Marine Science*, 5, 477-501, 2013.
- Trull, T., Rintoul, S. R., Hadfield, M., Abraham, E. R.: Circulation and seasonal evolution of polar waters south of Australia: Implications for iron fertilization of the Southern Ocean, *Deep-Sea Research II*, 48, 2439-2466, 2001.
- Trull, T., Davies, D. M., Dehairs, F., Cavagna, A.-J., Lasbleiz, M., Laurenceau-Cornec, E. C., d'Ovidio, F., Planchon, F., Leblanc, K., Quéguiner, B., Blain, S.: Chemometric perspectives on plankton community responses to natural iron fertilization over and downstream of the Kerguelen Plateau in the Southern Ocean, *Biogeosciences*, 12, 1029-1056, doi:10.5194/bg-12-1029-2015, 2015.
- Uitz, J., Claustre, H., Griffiths, F. B., Ras, J., Garcia, N., Sandroni, V.: A phytoplankton class-specific primary production model applied to the Kerguelen Islands region (Southern Ocean), *Deep-Sea Research I*, 56, 541-560, 2009.
- Varela, D. E., Pride, C. J., Brzezinski, M. A.: Biological fractionation of silicon isotopes in Southern Ocean surface waters, *Global biogeochemical cycles*, 18(GB1047), doi:10.1029/2003GB002140, 2004.
- Van den Boorn, S. H. J. M., Vroon, P. Z., Van Bergen, M. J.: Sulfur-induced offsets in MC-ICP-MS silicon-isotope measurements, *Journal of Analytical Atomic Spectrometry*, 24, 1111-1114, 2009.
- Wetzel, F., de Souza, G. F., Reynolds, B. C.: What controls silicon isotope fractionation during dissolution of diatom opal?, *Geochimica et Cosmochimica Acta*, 131, 128-137, doi:10.1016/j.gca.2014.01.028, 2014.

Zhou, M., Zhu, Y., d'Ovidio, F., Park, Y.-H., Durand, I., Kestenare, E., Sanial, V., Van-Beek, P., Quéguiner, B., Carlotti, F., Blain, S.: Surface currents and upwelling in Kerguelen Plateau regions, *Biogeosciences Discussion*, 11, 6845-6876, 2014.

Table 1: Characteristics of the stations sampled during KEOPS-2 and discussed in this study. MLD represent the mixed layer depth (from Park et al., 2014).

Station	Zone	Position		Date	MLD (m)
		Latitude	Longitude		
A3-1	Plateau	50°37.77'S	72°04.82'E	20 Oct	160
TNS6	Meander	48°46.78'S	72°16.77'E	22 Oct	67
R2	HNLC	50°21.52'S	66°43.00'E	26 Oct	110
E1	Meander	48°29.88'S	72°10.66'E	30 Oct	66
E2	Meander	48°31.41'S	72°04.63'E	1 Nov	43
E3	Meander	48°42.07'S	71°58.02'E	4 Nov	37
F	Polar Front Zone	48°37.27'S	74°48.44'E	7 Nov	40
E4W	Plateau	48°47.93'S	71°25.50'E	12 Nov	55
E4E	Meander	48°42.93'S	72°33.76'E	14 Nov	79
A3-2	Plateau	50°37.46'S	72°03.34'E	16 Nov	165
E5	Meander	48°24.69'S	71°53.99'E	18 Nov	35

Figure captions

Figure 1: Map of the KEOPS-2 cruise area (Indian sector of the Southern Ocean) showing the location of stations. Colored dots represent the different regions discussed in this study: yellow for HNLC, green for Plateau, red for Meander, black for Polar Front and white for coastal stations. The general surface circulation (from Park et al., 2014) is shown with arrows; blue arrow is the Antarctic Surface Waters, red arrow is the Polar Front Surface Water. The Polar Front and the Sub-Antarctic Front are represented by black dotted lines. The color represents the detailed map of the satellite-derived surface chlorophyll *a* concentration (MODIS level 3 product) averaged over the cruise period and black lines are 500 m and 1000 m isobaths.

Figure 2: Vertical profiles of biogenic silica and silicic acid concentration ($\mu\text{mol L}^{-1}$) and isotopic composition (‰) of dissolved (panels a. to d.) and particulate silicon (panel e. to h.) for the main contrasted KEOPS-2 stations (A3, R2, E4W and F-L) and for the Meander stations (TNS6, E1, E2, E3, E4E and E5). Horizontal dashed lines represent the MLD for each station.

Figure 3: Seasonal evolution of the dissolved and particulate silicon isotope composition (a. and c.) and concentration (b. and d.) in the upper 500 m of the iron-fertilized and HNLC reference stations. Summer isotopic signatures (KEOPS-1, Fripiat et al., 2011a) are in grey and black colors; results from the spring period (this study) are in red and blue color.

Figure 4: Silicon isotopic composition vs. silicic acid concentration for the different ML Si-reservoir in the iron-fertilized areas of the Kerguelen Plateau. A unique source (averaged Plateau-WW, blue dot, $32.9 \pm 2.2 \mu\text{mol Si L}^{-1}$ and $1.71 \pm 0.03 \text{‰}$) was used for all Plateau and Meander stations. ML dissolved Si-pools above the Plateau (purple dots) and in the Meander (red dots), fit well with a steady state fractionation law (grey straight line) but not with a Rayleigh distillation law (grey broken line). Grey dotted line represents the 0.2 ‰ sd of the fractionation factor. Note that given the small uncertainty of the source DSi and $\delta^{30}\text{Si}$ characteristics, we have neglected this uncertainty, which is minor compared to the s.d. on $\delta^{30}\text{Si}$. ML particulate Si-pools above the Plateau (black dots) and in the Meander (green dots) lie between the instantaneous and the accumulated products predicted by a Rayleigh distillation model (dotted and broken lines respectively). Straight black line represents the products in the steady state model. Summer isotopic signatures (K1 for A3 KEOPS-1) are from Fripiat et al. (2011a).

Figure 5: Potential mixing curves between TNS6-WW and the different Meander ML water masses, WW are in blue and ML are in red. The two end-members are identified by black curves. The two grey dashed lines represent the mixing curves between TNS6-WW and TNS6-ML and TNS6-WW and E1-ML which may explain the Si properties of E1-WW and E2-WW, respectively. The ML averages and error bars were calculated based on the median values and interquartile range for each station while the « central value » was defined using the salinity threshold method for WW.

Figure 6: Surface chlorophyll *a* concentration measured by satellite (green line), BSi fluxes collected in the sediment trap at 289 m (blue bars), and silicon isotopic composition of settling diatoms (black diamonds) at the A3 station.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6