

**January 22, 2018**

**Reference:** Submission bg-2017-350

**Dear Dr. Stoy,**

We thank the two anonymous reviewers for providing recommendations and appreciate the time and effort they put into this study. We have made all changes requested by the referees.

Changes made to the text and responses to comments are in detail discussed below. Changes made to our manuscript include:

- Improvement of Figure 4
- Modification of the 3<sup>rd</sup> paragraph in the Introduction section (starting on page 3, line 12)
- Added details about the study area and methods
- Reorganization of several discussion subsections
- Addition in the Supplementary Material file of a new figure (Fig. S7 in Supplement S7) with annual burn rates
- Editorial revision for grammar and spelling performed by the federal editorial service of Natural Resources Canada

A version of the manuscript with tracked changes is enclosed with this submission.

We thank you for giving us the opportunity to improve our manuscript.

Sincerely yours,

Emeline Chaste and coauthors

*Referee #1 comment #1:* I interpret their results and discussion to say that the increase in fire activity was due to both an increase in lightning frequency and drier climate. I recommend emphasizing this aspect of their results more clearly and explicitly.

*Authors #1:* We changed one sentence in the abstract and sentences in the 3<sup>rd</sup> paragraph, section “4.2 History of fire in the eastern boreal forest of Canada described by LPJ-LMfire” to improve this aspect.

*Referee #1 comment #2:* It appears that there were no additional modifications made to the fire routine in LPJ-LMfire, such as changes to fuel limits on fire activity, beyond the new PFT parameterization. If true, I would encourage the authors to write a sentence confirming this.

*Authors #2:* We added the sentence “No other modifications were made to the Pfeiffer et al. (2013) version of LPJ-LMfire.” at the end of the section “2.5 PFT definitions and LPJ-LMfire model modifications”.

*Referee #1 comment #3:* Presenting the results of Figure 4 in a manner similar to Figure 2 would make it easier for the reader to see the spatial patterns of agreement and disagreement between model and observations.

*Authors #3:* We presented the results of Figure 4 in the format of Figure 2. Maps reporting differences between mean aboveground biomass simulated and observed were added on the Figure 4.

*Referee #1 comment #4:* Given the results shown in Figure 3, the interpretation that "heavy and intense rain events that occur later in the summer decrease the probability of starting fires, despite more lightning" does not seem well-founded.

*Authors #4:* We rewrote the 3<sup>rd</sup> paragraph, section “4.2 History of fire in the eastern boreal forest of Canada described by LPJ-LMfire” to improve the idea that both fuel conditions and ignition sources have an influence on annual burn rates simulated by LPJ-LMfire. Heavy and intense rain events that occur later in summer increase the moisture of fuel and consequently decrease the probability of starting fires despite high density of lightning strikes.

*Referee #1 comment #5:* There are multiple instances of incorrect grammar.

*Authors #5:* The final version was reedited for grammar and spelling by a technical editor.

*Referee #2 comment #1:* I would like to see a more thorough treatment of the differences between the “Climate + CO2” and “Climate only” simulations, specifically with respect to how fire activity differed between the two scenarios. It was not clear to me whether the fire activity results reported in section “3.2.1 Fire activity” and Fig. 5 were from the “Climate only” simulation or the “Climate+CO2” simulations. Because the authors emphasize that CO2 fertilization increases NPP but that this is offset by increased fire activity, I think the results for fire activity with and without CO2 need to be more clearly presented. The effects of CO2 on NPP are discussed in section 3.2.2. Fuels, but I do not see a similar discussion of the fire results.

*Authors #6:* All figures in the manuscript were from the ‘Climate + CO<sub>2</sub>’ experiment; this aspect is now more clearly stated in the manuscript. We compared the NPP simulated by LPJ-LMfire with and without the CO<sub>2</sub> effect in supplement S7 Fig. S6 and explained the results in the 1<sup>st</sup> paragraph of the section “3.2.2 fuels”. We also added in supplementary materials a new figure with a comparison of annual burn rates simulated by LPJ-LMfire for both ‘Climate-only’ and ‘Climate + CO<sub>2</sub>’ experiments (Fig. S7 in Supplement S7). Results from this comparison were explained in section “3.2.1”. In the discussion, we specified that the CO<sub>2</sub>-induced enhancement of NPP had an influence on the annual burn rates by increasing the availability of fuel.

*Referee #2 comment #2:* Section 2.2 LPJ-LMfire model: Does the LPJ-LMfire model allow for cell to cell spread of fire? It is my understanding that the base LPJ model has no cell to cell interaction, a major limitation for simulating fire. It was my understanding that the SPITFIRE model did incorporate cell-cell interactions, but this needs to be explicitly stated in the manuscript.

*Authors #7:* No fire model currently used to simulate fire over continental to global-scale domains includes a representation of cell-to-cell fire spread (Pfeiffer et al., 2013; Hantson et al., 2016; Rabin et al., 2017). The basic concept in global fire modeling is that most fires are not as large as a single gridcell and so it is possible to imagine that an individual fire event occurs only within the size of an individual gridcell. We added the sentence “As in the original version of SPITFIRE and nearly all other large-scale fire models, LMfire does not simulate the cell-to-cell spread of fire (Hantson et al., 2016; Pfeiffer et al., 2013; Rabin et al., 2017).” at the end of the section “2.2 LPJ-LMfire model”.

*Referee #2 comment #3:* Study area description: I think it would benefit the reader if the authors included some rough description of fire regimes (general fire frequencies and whether historical fires were predominantly low, mixed, or high-severity, for example) for each of the ecoregions. As written, only the Boreal Shield ecoregions is characterized as having “typically high” fire frequency, but no estimate is given as to what that means in terms of mean fire return interval.

*Authors #8:* We added some information about the fire regimes for each ecozones in section “2.1 Study area”: Within the study area, high-intensity crown fires are the most common type of fire events (Flannigan et al., 2016). Fire regimes are heterogeneous, but generally follow a declining trend along a southwest-northeast gradient (Boulanger et al., 2012). During the period 1961–1990, the highest burn rates occurred in the western part of the BS ecozone ( $> 1\% \text{ yr}^{-1}$ ), while they were the lowest in the TSE ecozone ( $< 0.2\% \text{ yr}^{-1}$ ) (Boulanger et al., 2014). Annual burn rates in the BP ecozone and in the eastern part of the BS ecozone varied from 0.2 to 0.5%  $\text{yr}^{-1}$ , whereas it varied from 0.5 to 1.0%  $\text{yr}^{-1}$  in the HP ecozone (Boulanger et al., 2014).

*Referee #2 comment #4:* On p14 L20-23, the authors suggest that strong correlations between simulated and observed annual burn rates indicate fire suppression efforts have had little affect on fire behavior. Although this makes sense overall, the authors also note on p13 lines 25-30 that overestimates of simulated burn rates in Ontario and Quebec were expected because the model doesn’t consider anthropogenic effects and that forest management has influenced forest

composition/fuels. Perhaps these paragraphs need to be combined and the nuances of how fire and forest management have varied spatially be expanded upon?

*Authors #9:* As proposed we combined these paragraphs at the end of the section "4.3 Uncertainties and future perspectives" and nuanced these sentences to clarify that the effect of humans on management and suppression is not taken into account in LPJ-LMfire. However, similar temporal patterns of simulated and observed annual burn rates showed there does not seem to be a detectable effect of human activities on fire history.

*Referee #2 comment #5:* Table 1. The datasets and references for the temperature, precipitation, cloud cover and convective available potential energy were listed in the text, but I found it odd for them not be listed in the table when that information is listed in the table for the remaining variables.

*Authors #10:* There was a problem during the conversion process to PDF and missing data were present in Table 1 and 2 of the submitted version. We will make sure that this problem does not happen again for the final version.

*Referee #2 comment #6:* P3 Lines 11-16. I find it odd that the authors claim a “lack of quantitative knowledge regarding the relative changes in fire regimes that are attributable to human impacts”. Perhaps this is true of boreal Canada, I am more familiar with fire history studies in the U.S. Southwest and US Pacific northwest. Especially in the U.S. Southwest, fire history and forest age/structure and species composition from dendroecological evidence have been used to quantify how human activities (logging, grazing, fire exclusion) have impacted fire regimes (see studies by Thomas Swetnam, Peter Fule, Peter Brown, among others). These statements need to be qualified.

*Authors #11:* We clarified the 3rd paragraph in the section "Introduction".

*Referee #2 comment #7:* P1 Line 33-34: Change the sentence to read: “Agreement between the spatiotemporal patterns of fire frequency and the observed data confirmed that fire in the study area is strongly ignition-limited.”

*Authors #12:* We changed the sentence in P1 line 33-34 as proposed.

## References

Boulanger, Y., Gauthier, S., Burton, P. J. and Vaillancourt, M.-A.: An alternative fire regime zonation for Canada, *Int. J. Wildland Fire*, 21(8), 1052–1064, 2012.

Boulanger, Y., Gauthier, S. and Burton, P. J.: A refinement of models projecting future Canadian fire regimes using homogeneous fire regime zones, *Can. J. For. Res.*, 44(4), 365–376, doi:10.1139/cjfr-2013-0372, 2014.

Flannigan, M. D., Wotton, B. M., Marshall, G. A., Groot, W. J. de, Johnston, J., Jurko, N. and Cantin, A. S.: Fuel moisture sensitivity to temperature and precipitation: climate change implications, *Clim. Change*, 134(1–2), 59–71, doi:10.1007/s10584-015-1521-0, 2016.

Hantson, S., Arneeth, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P. and others: The status and challenge of global fire modelling, *Biogeosciences*, 13(11), 3359–3375, doi:10.5194/bg-13-3359-2016, 2016.

Pfeiffer, M., Spessa, A. and Kaplan, J. O.: A model for global biomass burning in preindustrial time: LPJ-LMfire (v1. 0), *Geosci. Model Dev.*, 6(3), 643–685, doi:10.5194/gmd-6-643-2013, 2013.

Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzik, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S. and Arneeth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, *Geosci Model Dev*, 10(3), 1175–1197, doi:10.5194/gmd-10-1175-2017, 2017.

# The pyrogeography of eastern boreal Canada from 1901 to 2012 simulated with the LPJ-LMfire model

Emeline Chaste<sup>1,2</sup>, Martin P. Girardin<sup>1,3</sup>, Jed O. Kaplan<sup>4,5,6</sup>, Jeanne Portier<sup>1</sup>, Yves Bergeron<sup>1,7~~5~~</sup>, Christelle Hély<sup>2,7~~5~~</sup>

5 <sup>1</sup>Département des Sciences Biologiques, Université du Québec à Montréal and Centre for Forest Research, Case postale 8888, Succursale Centre-ville, Montréal, QC H3C 3P8, Canada

<sup>2</sup>EPHE, PSL Research University, ISEM, Univ. Montpellier, CNRS, IRD, CIRAD, INRAP, UMR 5554, F-34095 Montpellier, FRANCE

10 <sup>3</sup>Natural Resources Canada, Canadian Forest Service, Laurentian Forestry Centre, 1055 du PEPS, P.O. Box 10380, Stn. Sainte-Foy, Québec, QC G1V 4C7, Canada

<sup>4</sup>[ARVE Research SARM, 1009 Pully, Switzerland](#)

<sup>5</sup>[Max Planck Institute for the Science of Human History, 07743 Jena, Germany](#)

<sup>6</sup>[Environmental Change Institute, School of Geography and the Environment, University of Oxford, OX1 3QY, UK](#)

<sup>4</sup>[Institute of Earth Surface Dynamics, University of Lausanne, Geopolis, Quartier Mouline, 1015 Lausanne, Switzerland](#)

15 <sup>7~~5~~</sup>Forest Research Institute, Université du Québec en Abitibi-Témiscamingue, 445 boul. de l'Université, Rouyn-Noranda, QC J9X 5E4, Canada

Correspondence to: Emeline Chaste (emeline.chaste@canada.ca)

**Abstract.** Wildland fires are the main natural disturbance shaping forest structure and composition in eastern boreal Canada. On average, more than 700,000 ha of forest burns annually, and causes as much as C\$2.9 million worth of damage. Although we know that occurrence of fires depends upon the coincidence of favourable conditions for fire ignition, propagation and fuel availability, the interplay between these three drivers in shaping spatiotemporal patterns of fires in eastern Canada remains to be evaluated. The goal of this study was to reconstruct the spatiotemporal patterns of fire activity during the last century in eastern Canada's boreal forest as a function of changes in lightning ignition, climate and vegetation. We addressed this objective using the dynamic global vegetation model LPJ-LMfire, which we parametrized for four Plant Functional Types (PFTs) that correspond to the prevalent tree genera in eastern boreal Canada (*Picea*, *Abies*, *Pinus*, *Populus*). LPJ-LMfire was run with a monthly time-step from 1901 to 2012 on a 10-km<sup>2</sup> resolution grid covering the boreal forest from Manitoba to Newfoundland. Outputs of LPJ-LMfire were analyzed in terms of fire frequency, net primary productivity (NPP), and aboveground biomass. The predictive skills of LPJ-LMfire were examined by comparing our simulations of annual burn rates and biomass with independent data sets. The simulation adequately reproduced the latitudinal gradient in fire frequency in Manitoba and the longitudinal gradient from Manitoba towards southern Ontario, as well as the temporal patterns present in independent fire histories. ~~Nevertheless~~ However, the simulation led to ~~the~~ underestimation and overestimation of ~~the~~ fire frequency at both the northern and southern limits of the boreal forest in Quebec. The general pattern of simulated total tree biomass also agreed well with observations, with the notable exception of overestimated biomass at the northern treeline, mainly for ~~PFT~~ *Picea* ~~PFT~~. In these northern areas, the predictive ability of LPJ-LMfire is likely being affected by ~~the~~ low density of weather stations, which ~~has~~ led to underestimation of the strength of fire-weather interactions during extreme fire

20  
25  
30  
35

years and, therefore, vegetation consumption. Agreement ~~of-between~~ the spatiotemporal patterns of fire frequency ~~and with~~ the observed data ~~across a vast portion of the study area~~ confirmed that fire ~~in the study area therein~~ is strongly ignition-limited. ~~A Drier climate and coupled with an increase in lightning frequency during the second half of the 20th Century notably, had leaded to an increase in fire activity. Overall, climate and lightning ignition variability at multi-decadal and annual time-~~  
5 ~~scales was the primary driver of fire activity since the beginning of the 20th C. However~~ Finally, our simulations highlighted the importance of both climate and ~~vegetation-fire on firevegetation~~: despite an overarching CO<sub>2</sub>-induced enhancement of ~~NPP net primary productivity (NPP)~~ in LPJ-LMfire, forest biomass was relatively stable because of ~~the~~ compensatory effects of increasing fire activity.

## 1 Introduction

10 Wildland fires are the main natural disturbance shaping forest structure and composition in eastern boreal Canada (Bergeron et al., 1998, 2014). On average, more than 0.7 Mha burn annually across Manitoba, Ontario, Quebec and the Maritime Provinces, ~~and-which~~ causes as much as \$C2.9 million worth of damage and property losses (Canadian Council of Forest Ministers, 2017). About 97-% of these burned areas are generated by a small proportion (3-%) of large fires (fires > 200 ha in  
15 area; Stocks et al., 2003). For example, a fire burned 583,000 ha within a few days in 2013 near the aboriginal community of Eastmain (Province of Quebec), which is the equivalent of 31-% of the total area burned during that year in Quebec (Erni et al., 2017). Studies of the spatial distribution of wildland fires in the past have highlighted that the frequency and size of fires in Canada have continuously increased over the last 50 years or so in response to ~~the~~ on-going global warming (e.g., Kasischke and Turetsky, 2006; Hessl, 2011; Girardin and Terrier, 2015). Concerns are now being raised about the increasing frequency/severity of extreme climatic events with further warming, which could lead to an increasing concentration of  
20 numerous large fires in time and space (Wang et al., 2015). Given these observations and projections, there is growing concern about the capacity of the boreal forest to recover from disturbances (Bond et al., 2004; IPCC, 2013; Kurz et al., 2013; Rogers et al., 2013).

Wildland fire regimes are described by several attributes including the frequency, size, intensity, seasonality, type and severity of fires (Keeley, 2009). The spatiotemporal variability of a fire regime depends upon the coincidence of favourable  
25 conditions for fire ignition, fire propagation and fuel availability, which are controlled by ignition agents, weather/climate, and vegetation (Flannigan et al., 2009; Moritz et al., 2010). Almost half of the fires that occur in eastern boreal Canada are ignited by lightning and represent 81-% of the total area burned (Canadian Forest Service, 2016), while the remaining fires originate from human activities. The capacity of a fire to grow into a large fire is determined by many factors, which include weather and fuel. High temperature, low precipitation, high wind velocity and low atmospheric humidity can increase the growth of  
30 these fires (Flannigan et al., 2000). The intensity, severity and size of fires are further influenced by ~~the~~ species composition within ~~the~~ landscape, with needle-leaf species being more fire-prone than broad-leaf species owing to their high flammability (Hély et al., 2001). Physical variables such as slope, surficial deposits and soil moisture can also have significant effects on

the rate at which fires spread by influencing fuel moisture or creating natural fire breaks (Hély et al., 2001; Mansuy et al., 2011; Hantson et al., 2016). Climate change scenarios for Canada indicate an increase in both temperature and precipitation in the coming decades. Yet, the increase in precipitation should not compensate for the increase in temperature (IPCC, 2013), and a greater moisture deficit is expected compared to the current state. Warmer springs and winters that lead to an earlier start of the fire season are anticipated, together with an increase in the frequency of extreme drought years due to more frequent and persistent high-pressure blocking systems (Girardin and Mudelsee, 2008). These phenomena are expected to lead to an increase in the frequency and size of fires in eastern boreal Canada in response to the on-going global warming (Ali et al., 2012). Effects of these changes in seasonal onset and dryness are such that the average size of spring wildfires could be multiplied by a factor of three for each additional 1°C of warming (Ali et al., 2012; Girardin et al., 2013a; Price et al., 2013). An increase in the areas burned would affect both plans for forest management plans and fire suppression strategies. It could also have subsequent feedbacks on the global carbon cycle, given that the substantial quantities of carbon currently being stored in these landscapes could be re-emitted back into the atmosphere (Pan et al., 2011).

A number of uncertainties persist concerning future fire trajectories-projections, and biases still exist regarding our current understanding of the natural variability in fire regimes. ~~Forest management and fire suppression since the 1970s have contributed to modifying fire patterns and vegetation attributes in Canada (Gauthier et al., 2014). Climate also has been rapidly changing in recent decades with the expansion of human activities. All of these changes have altered the interactions between fire regimes and their various forms of controls (Bergeron et al., 2004b). Most of the fire history studies are based upon observations collected over relatively short time intervals (< 100 years), and reliable observations were often unnot-available for many boreal regions prior to the late 1960s (Podur et al., 2002). However~~ Moreover, ~~F~~forest management and active fire suppression since the 1970s have contributed to modifying fire patterns and vegetation attributes in Canada (Gauthier et al., 2014). Therefore, it is difficult to determine the contribution of climate alone on fire activity in studies using observations collected since the second half of the 20th century.C. ~~Climate also has been rapidly changing in recent decades with the expansion of human activities. All of these changes have altered the interactions between fire regimes and their various controls (Bergeron et al., 2004b). For instance, a lack of quantitative knowledge regarding the relative changes in fire regimes that are attributable to human impacts prevents modellers from adequately taking into account their influence in future fire trajectories. Although it is also possible to investigate recent changes in fire regimes that are based upon observations collected over relatively short time intervals (< 100 years), reliable observations were not available in many boreal regions prior to the late 1960s (Podur et al., 2002).~~ Furthermore, fire history studies rarely consider the feedbacks of fires on vegetation, mostly because historical data about vegetation composition are lacking (Danneyrolles et al., 2016). This is particularly true in the case ~~for~~of studies dealing with reconstructions of fire activity using dendrochronological evidence (e.g., Girardin et al., 2006) or by adjusting empirical data sets (Van Wagner et al., 1987). This problem may be circumvented by investigating past fire regimes over long periods of time through the analysis of charcoal and pollen in soil layers or lacustrine deposits (Payette et al., 2008; Ali et al., 2009). However, these paleoecological methods are costly, time-consuming, and incapable of do not make it possible

to capturing the overall spatial variability of fire regimes at annual- to decadal-scale resolutions. Faced with these gaps, increasing our knowledge of the spatiotemporal patterns of past fires is necessary to perform better predictions in the future.

Simulations using dynamic global vegetation models (DGVMs) allow make it possible to estimate the spatiotemporal distribution of fires to be estimated relative to climate and vegetation (Yue et al., 2015; Hantson et al., 2016). Indeed, these models simulate shifts in potential vegetation composition and related fire activity in response to changes in climate or environmental constraints (Smith et al., 2001). Experiments can be conducted at fine- to broad-spatial scales and validated at relatively short- to medium- timescales. Validation can be realized done in regions where human activities are sufficiently low to allow comparisons with natural potential vegetation, by comparing simulation results with high-resolution satellite products, such as MODIS, at global scales, such as MODIS (Tang et al., 2010). DGVM simulations also may be validated at decadal- to millennial- timescales by comparing them with historical records of vegetation or fire activity that have been reconstructed using indicators derived from pollen and charcoal, amongst others, which are deposited in lacustrine sediments (Molinari et al., 2013). One of these models, the Lund-Postdam-Jena (LPJ) model, has been the subject of numerous refinements over time, especially concerning simulations of fire patterns (Thonicke et al., 2010; Pfeiffer and Kaplan, 2012), and it has been validated in many regions worldwide, excluding eastern boreal Canada for example (e.g., Prentice et al., 2011; Pfeiffer et al., 2013; Yue et al., 2015; Knorr et al., 2016).

Here, we used the LPJ-LMfire model that was developed by Pfeiffer et al. (2013) to perform a simulation experiment that targeted the boreal forest of eastern Canada and which covered the last century, with customized parameterization to capture for prevalent tree genera present in eastern boreal Canada. The DGVM explicitly simulates fire ignition from lightning and; hence, it is particularly adapted to the largely ignition-limited fire regimes in our study region. The objectives of this study were (1) to calibrate the LPJ-LMfire model for boreal forests in eastern Canada, (2) to verify assess the predictive skills of the model with independent data sets from eastern Canada's boreal forests, (3) to reconstruct fire activity, net primary productivity NPP and aboveground biomass during the last century, and (4) to determine how the spatiotemporal pattern of these three components has evolved conjointly in relation to changes in climate variables.

## 2 Model, Experimental Set-up and Methods

### 2.1 Study area

The study area encompasses eastern Canada's boreal forest (Brandt, 2009) from Manitoba to Newfoundland, which ranges from 102.86°W to 52.64°W and from 46.61°N to 64.71°N (Fig. 1). The most common needle-leaf tree species present in this region are black spruce (*Picea mariana* (Mill.) B.S.P.), white spruce (*Picea glauca* (Moench) Voss), balsam fir (*Abies balsamea* (L.) Mill.), jack pine (*Pinus banksiana* Lamb.), white pine (*Pinus strobus* L.), red pine (*Pinus resinosa* Ait.), eastern larch (*Larix laricina* [Du Roi] K. Koch), and eastern white cedar (*Thuja occidentalis* L.). The main broad-leaf tree species are trembling aspen (*Populus tremuloides* Michx.) and white or paper birch (*Betula papyrifera* Marsh.) (Ecological Stratification Working Group, 1996; Brandt, 2009; Shorohova et al., 2011). The study area is divided from south to north into

four ecozones (Fig. 1; Ecological Stratification Working Group, 1996). 1) The Boreal Shield (BS) ecozone is characterized by rocky and rugged landscapes influenced by a continental climate (long and cold winters; short and warm summers) and by the cold air masses flowing out from Hudson Bay. ~~Fire frequency is typically high.~~ Landscapes are dominated by needle-leaf tree species in the westernmost areas, and ~~codominated by nee-needleleaf and~~ deciduous tree species in temperate eastern areas.

5 2) The Boreal Plain (BP) ecozone corresponds to drier areas that are characterized by glacial deposits of variable thickness on flat or slightly rolling terrain. Forests are dominated by mixed boreal species, mainly represented by black spruce, trembling aspen and jack pine. 3) The Hudson Plain (HP) ecozone is characterized by a sparser vegetation, which is dominated by *Sphagnum* and shrubs. Poor drainage conditions constrain southern trees to establish at drier, higher elevations. 4) The Taiga Shield (TS) ecozone, which is split into Eastern (TSE) and Western (TSW) parts, is characterized by colder climate conditions.

10 The landscape becomes more open along a latitudinal gradient from south to north. ~~In all regions, the d~~ Dominant tree species ~~include-are~~ black spruce and jack pine. ~~Within the study area, high-intensity crown fires are the most common type of fire events (Flannigan et al., 2016). Fire regimes are heterogeneous, but generally follow a declining trend along a southwest-northeast gradient (Boulanger et al., 2012). During the period 1961–1990, the highest burn rates occurred in the western part of the BS ecozone (> 1% yr<sup>-1</sup>), while they were the lowest in the TSE ecozone (< 0.2% yr<sup>-1</sup>) (Boulanger et al., 2014). Annual~~

15 ~~burn rates in the BP ecozone and in the eastern part of the BS ecozone varied from 0.2 to 0.5% yr<sup>-1</sup>, whereas it varied from 0.5 to 1.0% yr<sup>-1</sup> in the HP ecozone (Boulanger et al., 2014).~~

## 2.2 LPJ-LMfire model

Simulations of the terrestrial ecosystem were carried out using the dynamic global vegetation model LPJ-LMfire, which ~~contains-includes~~ updates of both LPJ and the SPread and InTensity of the FIRE (SPITFIRE) wildfire module (Thonicke et al., 2010). The model has been ~~most-~~extensively evaluated for boreal forests (Pfeiffer et al., 2013). LPJ-LMfire is designed to simulate regional ecosystem dynamics, structure and composition, with vegetation and fire events as responses to changes in climate and carbon dioxide ~~concentration-~~(CO<sub>2</sub>) concentration (Sitch et al., 2003). LPJ-LMfire describes the state of an ecosystem in terms of annual carbon stocks (living biomass, litter and soil), ~~Net Primary Productivity-(NPP),~~ net biome productivity, evapotranspiration, heterotrophic respiration, soil moisture fraction, ~~and-as well as~~ forest structure and vertical profile (cover fraction, individual density, crown area, leaf area index). In the present study, changes in the vegetation state are described in terms of NPP and total carbon stocks in ~~the~~ living aboveground biomass. In LPJ-LMfire, vegetation is defined by up to nine Plant Functional Types (PFTs). Each PFT represents one or several species sharing the same physiology and dynamics, governed by a short list of vital attributes, and constrained by bioclimatic limits (Sitch et al., 2003). Vegetation dynamics are updated annually based on the simulation of daily and annual processes. Daily processes ~~were-are~~ defined in

20 25 30 terms of photosynthesis, stomatal regulation, soil hydrology, autotrophic respiration, leaf and root phenology and decomposition. Annual processes ~~were-are~~ defined in terms of several sources of mortality, seedling establishment, reproduction, allocation and tissue turnover (Smith et al., 2001; Sitch et al., 2003). The computational core of SPITFIRE is based upon Rothermel-type surface fire behaviour models (Rothermel, 1972; Andrews et al., 2008) and is designed to simulate

processes of natural fires and their ~~implications-impacts~~ on vegetation mortality and fire emissions (Thonicke et al., 2010). The LMfire module simulates lightning ignitions based upon a daily time-step, and uses fuel bulk density and fuel moisture to calculate the fire's rate of spread, intensity, and fire-related mortality. It allows fires to burn over multiple days and simulates fire extinction from changes in weather and fuel (Pfeiffer et al., 2013). [As in the original version of SPITFIRE and nearly all other large-scale fire models, LMfire does not simulate the cell-to-cell spread of fire \(Hantson et al., 2016; Pfeiffer et al., 2013; Rabin et al., 2017\).](#)

### 2.3 Simulation protocol

LPJ-LMfire was run monthly from 1901 to 2012 on ~~a~~  $10 \times 10$  km equal-area grids covering eastern boreal Canada from Manitoba to Newfoundland. Driver data sets were prepared in netCDF format and are described in Table 1. Climate data were compiled at a monthly time-step, while atmospheric CO<sub>2</sub> concentrations (~~{CO<sub>2</sub>}~~) were compiled at an annual time-step (see Section 2.4). A 1120-year spin-up period was prescribed to equilibrate vegetation and carbon pools with climate at the beginning of the study period (Smith et al., 2001), and to ensure that forest biomass ~~and fire disturbances were was~~ in stable condition ~~with fire disturbances~~ (Tang et al., 2010). This spin-up run was made using linearly detrended 1901-2012 climate data, ~~and~~ repeated 10 times.

### 2.4 Environmental input data sets

#### 2.4.1 Climate

Monthly means of temperature, diurnal temperature range, precipitation, number of days with precipitation, and wind speed were extracted ~~between-for the~~ 1901-2012 ~~period~~ from Environment Canada's historical climate database (Environment Canada, 2013) using BioSIM software (v.10.3.2; Régnière et al., 2014). Gridded climate data were prepared in BioSIM by interpolating weather data from the four weather stations that were closest to each  $10 \times 10$  km grid, adjusted for elevation and location differentials with regional gradients, and averaged using inverse distance weighting ( $1/d^2$ , where  $d$  is distance). Missing wind speed values between 1901 and 1968, and those for 2010-2012 were set to the monthly 1969-2010 averages.

Monthly means of total cloud cover percentage for the entire atmosphere and convective available potential energy (CAPE) were interpolated on our grid from the NOAA-CIRES 20th Century Reanalysis v2 data set at ~~a~~  $\sim 2.0^\circ$  latitude and  $1.75^\circ$  longitude resolution (Compo et al., 2011). For a given grid cell, the annual monthly CAPE anomaly was calculated as the difference between the annual value and the monthly normal for CAPE, which was computed between 1961 and 1990.

#### 2.4.2 Lightning

The Canadian lightning detection network (CLDN) data set, covering the ~~period~~ 1999-2010 ~~period~~ (Orville et al., 2011), was used to reconstruct the monthly cloud-to-ground lightning strike density (number/day/km<sup>2</sup>) between 1901 to 2012. Given the strong correlation between lightning strikes and the product of CAPE and precipitation (e.g., Peterson et al., 2010; Romps et

al., 2014), we computed daily strike density using CAPE data and distributed the lightning strikes over the daily fraction of monthly rainy days (Pfeiffer et al., 2013). Across Canada and within our study area, July was the month with the maximum number of strikes between 1999 and 2010 ([supplement S1-Fig. S1-A in Supplement S1](#)); and, in turn, inter-annual strike variability (hereafter, referred to as min-to-mean and max-to-mean ratios) ranged from 0.1 to 7.5 times the July mean ([supplement S1-Fig. S1-B in Supplement S1](#)). This inter-annual variability in lightning strikes was preserved in our reconstruction by applying these two ratios to normalize values between -1 and +1 of CAPE anomalies (following Pfeiffer et al., 2013; see [supplement-Supplement S1](#) for further details), and were directly added to the 1999–2010 flash climatology.

### 2.4.3 Soils

The volume fraction of coarse fragments, together with the 0–100 cm [depth-deep](#) soil texture fractions of sand and clay, were interpolated on the 10 × 10 km grids from the 1-km resolution ISRIC – World Soil Information data set (Hengl et al., 2014).

For topography, we interpolated the 30 arc-second gridded digital elevation model (DEM) of Canada (Natural Resources Canada, 2007). We calculated slopes in degrees at 30 arc-seconds with the DEM map using ArcGIS 10.4.1 and interpolated the data to our 10 km<sup>2</sup> grids. To calculate the percentage of land (i.e., removing lakes and water course areas) in each grid cell, we rasterized the water fraction of the National Hydro Network (NHN) data set at 100-m resolution (Natural Resources Canada, 2010). We calculated [the](#) water fraction at [a](#) 10-km resolution from [100-m resolution](#) grid cells [at 100-m resolution](#) that had a percentage of water fraction > 50%. The land fraction was defined as the inverse of the water fraction. Roads, power lines, dams, mines, and other human-made structures, and areas of bare rock, were not considered in this study.

### 2.4.4 Atmospheric CO<sub>2</sub> concentration

Monthly mean atmospheric [carbon dioxide \(CO<sub>2</sub>\)](#) concentrations covering the periods [from](#) 1901 to 1980 and [from](#) 1981 to 2012 were obtained from Pfeiffer et al. (2013) and the Mauna Loa data set (Keeling et al., 2009), respectively. Annual mean atmospheric CO<sub>2</sub> [concentration](#) varied from 296.23 ppm in 1901 to 392.48 ppm in 2012, which corresponds to an increase of 32.5%.

### 2.5 PFT definitions and LPJ-LMfire model modifications

LPJ-LMfire was calibrated for four [Plant Functional Types \(PFTs\)](#) that corresponded to the predominant tree genera currently present in the boreal forest of Canada: *Picea*, *Abies*, *Pinus* and *Populus*. PFT-related parameters, e.g., fraction of roots in [the](#) upper soil layer or minimum and maximum temperatures of the coldest month for establishment, were assigned values from the published literature or global databases (see [Table 1 in supplement-Supplement S2](#) for further details).

### 2.5.1 Edaphic limits to establishment

Establishment and growth of boreal tree species are influenced by a wide range of soil properties that ~~were~~ are related to soil nutrient availability, ~~and~~ which included ~~pH~~, parent material, soil particle size, and water content, among others (Girardin et al., 2001; Beauguard and de Blois, 2014; Gewehr et al., 2014). Not all ecosystem processes linking these properties to tree establishment are simulated in the current version of LPJ-LMfire. Notably, the model does not simulate the development of peatlands or the process of paludification, and it does not include a complete module of biogeochemical cycling in soils that would emulate processes leading to acidification, for instance. As ~~has been~~ proposed by Beauguard and de Blois (2014), however, some edaphic variables may be indicative of certain soil processes at the stand level. In this study, correlations between the abundance of specific tree genera and soil clay content led to the implementation of a simple scheme to limit tree establishment in LPJ-LMfire (~~supplement S3~~ Fig. S2-A in Supplement S3). Edaphic limits to establishment were defined here in the same way that bioclimatic limits are used in LPJ. The correlations between the genus-specific tree cover fraction from Beaudoin et al. (2014) and clay volume fraction from Hengl et al. (2014) were analyzed at a 10-km resolution. For each PFT, the percentage of clay corresponding to the upper limit of the 90-% confidence interval (CI) of its distribution, for grid cells with at least 10-% of PFT cover, was used in the model as a threshold, above which the given PFT could not establish. The upper limit of the 90-% CI of the clay percentage distribution was 20-%, 13-%, 18-% and 23-% for *Picea*, *Abies*, *Pinus* and *Populus*, respectively (~~supplement S3~~ Fig. S2-~~AB~~ in Supplement S3). The 20-% threshold essentially results in the exclusion of the *Picea* and *Populus* PFTs in the Hudson PlainHP ecozone (~~supplement S3~~ Fig. S2-~~B and C~~ in Supplement S3), while the threshold of 13-% leads to the additional exclusion of other PFTs, especially *Pinus*, in the Midwestern Boreal Shield (MBS) and Boreal PlainBP ecozones (~~supplement S3~~ Fig. S2-~~B and C~~ in Supplement S3).

### 20 2.5.2 Post-fire recruitment

Recruitment of *Pinus banksiana* requires the heat of fires to release seeds from serotinous cones (Gauthier et al., 1996). This condition was implemented in the current LPJ-LMfire version specifically for the *Pinus* PFT by inhibiting seedling establishment during years without fire. Such fire effects on seed dispersal are also observed for *Picea mariana*, which has semi-serotinous cones. Given that black spruce cones can open gradually over time in the absence of fire (Messaoud et al., 2007), *Picea* PFT establishment was not constrained by fire occurrence, ~~neither- was that of~~ Establishment of the *Abies* and *Populus* PFTs ~~are also not constrained by fire occurrence~~. No other modifications were made to the Pfeiffer et al. (2013) version of LPJ-LMfire.

### 2.6 Model evaluation

30 We assessed the performance of our customized LPJ-LMfire by comparing simulation results with previously published data sets on fire and maps of genus-specific aboveground biomass for Canada's forests.

## 2.6.1 Fire activity

The simulated burned area fraction was evaluated against three fire data products. First, annual burn rates ~~during for~~ 1980–2012 were compiled from the Natural Resources Canada fire database (M.A. Parisien, personal communication) using ~~the~~ Canada’s national fire polygons with the hexagonal cells approach from Héon et al. (2014), but extended to our study area. We used 365  
5 hexagonal cells to cover our study area and to compute the 1980 to 2012 simulated mean annual burn rates with 95-% ~~confidence intervals (95-% CI)~~ for each hexagonal cell. The second fire data product originated from stand-replacing fire history studies. Here, historical annual proportions of burned areas were obtained for 26 locations (~~supplement S4~~-Fig. S3 ~~in Supplement S4~~) using post-fire stand initiation reconstructions based upon field and archival data that were digitized and ~~which were~~ included in GIS databases (Girardin et al., 2013b; Héon et al., 2014; Portier et al., 2016). Using a 100-km radius around  
10 each location centroid, we calculated the simulated mean annual burn rates between 1911 and 2012, together with the 95-% CI. Differences between our simulated ~~estimate~~ 95-% CIs ~~estimates~~ and these two fire data products were considered qualitatively as ‘not different’ if the observed annual burn rate fell within the 95-% CI of the simulated mean burn rate. Note that as the period covered by the historical fire data often extended further back in time into the 19th or 18th centuries for southern locations (~~supplement S4~~-Table S2 ~~in Supplement S4~~), some important differences could be expected in the  
15 comparison process. Finally, a third validation of fire simulations was made by comparing time-series of total simulated annual burned areas in boreal forests of Manitoba, Ontario and Quebec with provincial fire statistics (point data) from the Canadian National Fire Database (CNFDB; Canadian Forest Service, 2016) covering the 1959-2012 period. Human-caused fires were excluded from these analyses. Spearman’s rank correlation ( $r_s$ ) was used to quantify the agreement between observed and simulated data. The agreement between simulation and observation was further evaluated in terms of fire seasonality by  
20 comparing their respective distributions of mean monthly areas that ~~were~~ burned from 1959 to 2012.

## 2.6.2 Aboveground biomass

Published maps of total aboveground biomass at the genus level (Beaudoin et al., 2014) were used to evaluate model simulations. Maps that were created by Beaudoin et al. (2014) were constructed at ~~a~~ 250-m spatial resolution using remote sensing MODIS data sets, combined with photo-plot observations of Canada’s National Forest Inventory (NFI), mainly in the  
25 southern areas (see non-hatched area on Fig. 4). We aggregated the 250-m data to a 10-km resolution and applied a correction for the vegetated treed fraction of the landscape, as defined by Beaudoin et al. (2014). ~~The v~~Vegetated treed fraction corresponds to the fraction of the grid cells that are covered by tree species of any size on at least 10-% of the grid cell.

Total aboveground biomass, ~~that was~~ estimated by two other methods reported by Margolis et al. (2015), was used for a second evaluation of model simulations for the ~~5~~ five ecozones under study. The ~~Boreal Shield~~BS ecozone was divided into  
30 three ecoregions for comparison ~~purposes~~ (Fig. 1); ecoregions corresponding to the classification of ecological regions ~~on~~ at a finer scale than ecozones. The first method of biomass estimation is based upon the ‘Geosciences Lidar Altimetry System’ (GLAS) method, which estimates total aboveground biomass from the waveforms recorded over vegetated land using Lidar

instruments. The second method is based upon NFI photo-plot estimates of total aboveground biomass using allometric equations.

## 2.7 History of the eastern boreal forest of Canada described by LPJ-LMfire

5 The outputs of LPJ-LMfire for the eastern boreal forest of Canada were analyzed in terms of annual burn rates, NPP and total aboveground biomass. Significant changes in each temporal series were highlighted by a regime shift calculation developed by Rodionov (2004, 2006). A sequential application of Student's  $t$ -test on 1,000 randomly chosen grid cells was used (Rodionov, 2004, 2006). To be statistically significant at  $P = 0.10$ , the difference (*diff*) between mean values of two subsequent periods that was determined according to Student's  $t$ -test should satisfy the condition:

$$[1] \text{diff} = t\sqrt{2\sigma_i^2/l},$$

10 where  $t$  is the value from the  $t$ -distribution with  $2l - 2$  degrees of freedom at the given probability level  $P$ ,  $l$  is the cut-off length of the growth phase to be determined (hereafter, set to periods of 20 years), and  $\sigma_i^2$  is the average variance for running  $l$ -year intervals. The sample proportion, representing the fraction of  $k$  cells ( $\text{an-integer} \geq 0$ ) of a given population  $N$  ( $\text{an-integer} > 0$ ), which was identified positively as recording a growth decline (or release), a biomass reduction (or biomass increase) and an increase of fire activity (or decrease), was computed for each sampled year from 1920 to 2007.

## 15 2.8 Sensitivity analysis to CO<sub>2</sub> fertilization

In terrestrial ecosystem models, changes in atmospheric [CO<sub>2</sub> concentration \[CO<sub>2</sub>\]](#) in the recent past and future often have a more important influence on vegetation than does climate change (Girardin et al., 2011). Therefore, their inclusion has a very important effect on simulated changes in productivity. Here, the effect of [CO<sub>2</sub> \[CO<sub>2</sub>\]](#) fertilization was explored using two [simulation experiments](#). In the first [simulation experiment](#), 'Climate + CO<sub>2</sub>', ~~we ran the model with increases in the CO<sub>2</sub> concentration [CO<sub>2</sub>] as presented in section 2.4.4. This experiment was used throughout our [analyze evaluation of LPJ-LMfire model evaluations simulations. and history of the eastern boreal forest of Canada described by LPJ-LMfire.](#) In the second experiment, 'Climate-only', we ran the model with a constant [CO<sub>2</sub> concentration \[CO<sub>2</sub>\]](#) from 1901 to 2012, which was fixed at 296.23 ppm (year 1901 value). In this case, there was no response of vegetation gross primary production (GPP) or fire to changes in [CO<sub>2</sub> concentration \[CO<sub>2</sub>\]](#), i.e., 'Climate-only' experiment. ~~In the second simulation, we ran the model with increases in the [CO<sub>2</sub>]~~ ('Climate + CO<sub>2</sub>' experiment). For further details, see section 2.4.4. [The effect of CO<sub>2</sub> fertilization on vegetation was determined by the difference between simulations 'Climate + CO<sub>2</sub>' and 'Climate-only'. Due to the post-fire recruitment rules established in LPJ-LMfire \(see section 2.5.2\), the effect of CO<sub>2</sub> fertilization on fire was only determined by comparing the spatial pattern of annual burn rates simulated with the 'Climate + CO<sub>2</sub>' and 'Climate-only' experiments.](#)~~

20

25

### 3 Results

We report on the evaluation of ~~the~~ process-based model performance in adequately simulating spatial patterns of fire frequency and fuel conditions (as indicated by the aboveground biomass of the four PFTs and total ~~net primary productivity,~~ NPP) in eastern boreal Canada. We also report on changes in ~~the~~ fire activity during the last century as simulated by LPJ-LMfire, with associated changes in vegetation features.

#### 3.1 Predictive skills of the LPJ-LMfire model

##### 3.1.1 Fire activity

For the recent ~~period~~ 1980-2012 period, mean and maximum simulated annual burn rates were ~~respectively~~ 0.36-% yr<sup>-1</sup> and 1.49-% yr<sup>-1</sup>, respectively (Fig. 2-B), while ~~the~~ mean and maximum observed annual burn rates were 0.28-% yr<sup>-1</sup> and 2.03-% yr<sup>-1</sup> (Fig. 2-A). Observed and simulated burn rates were not significantly different in more than 80-% of the studied hexagonal cells (295 out of 365; Fig. 2-C). Therefore, LPJ-LMfire was able ,therefore, to capture the amplitude of interregional variation. Decreases in fire activity observed along both the latitudinal gradient in Manitoba, ~~and~~ the longitudinal gradient from Manitoba to southern Ontario were well reproduced by the simulation (Fig. 2-A and B). Furthermore, more than half of the observed historical annual burn rates fell within the 95-% CI of their corresponding simulated annual burn rates (for further details, see supplementary S4-Table S2 and Fig. S3 in Supplement S4). LPJ-LMfire overestimated annual burn rates from south of the Hudson Bay in Ontario to southwestern Quebec (Fig. 2-C), while it underestimated ~~the~~ annual burn rates in the western areas of the central boreal forest in Quebec (Fig. 2-C). Spearman's correlation coefficients ( $r_s$ ) of time-series of observed versus simulated areas burned are 0.41 for Quebec and 0.50 for Ontario and Manitoba (Fig. 3-A). As revealed by these coefficients, LPJ-LMfire was also able to emulate year-to-year variability in annual areas that were burned in Manitoba and Ontario, but less so in Quebec. High fire activity years over the temporal series were also captured in the simulations, including 1961, 1968, 2003 and 2005, mostly in Manitoba and Ontario (Fig. 3-A). However, three extreme fire years were not reproduced: 1983, 1989 and 2002 (Fig. 3-A). Based upon the comparison of monthly percentage of total areas that were burned between 1959 and 2012 in eastern boreal Canada, the simulated fire season generally started one month earlier than what was observed (Fig. 3-B).

##### 3.1.2 Fuels

Overall, the general latitudinal pattern of simulated total tree biomass (Fig. 4 A2) agreed with the pattern of observed total tree biomass (Fig. 4-A1). Median simulated total tree biomass (with 90-% CI) in the southern areas (non-hatched) was 77 T ha<sup>-1</sup> (33-108 T ha<sup>-1</sup>), while median observed total tree biomass in the same areas was 73 T ha<sup>-1</sup> (36-100 T ha<sup>-1</sup>). In the Boreal ShieldS ecozone, percentage differences between mean total tree biomass that was simulated and that which was estimated by using NFI-based and GLAS-based methods were ~~respectively~~ 31-% and -7.8-%, respectively, and decreased along a westward gradient from Quebec to Manitoba (Table 2). We greatly overestimated mean total tree biomass in the Boreal Plain ecozone

because these differences were -60% and -50%, respectively. For the ~~Taiga-Shield~~ ecozone in Quebec and Manitoba, ~~which that~~ corresponds to less intensively sampled northern regions (hatched areas), total tree biomass was largely overestimated, mostly in Quebec, due to ~~the~~ high genus-specific biomass of the *Picea* PFT (Fig. 4-B). In this ecozone, relative differences with GLAS-based estimates ranged from 1.3% in ~~the~~ west to 63.6% in ~~the~~ east, whereas it was only 1.6% in comparison with NFI-based estimates (Table 2). Greater relative differences were observed in the ~~Hudson-Plain~~ ecozone (Table 2), where we overestimated total tree biomass for grid cells in which edaphic limits were not too restrictive and where vegetation could establish (Fig. 4-A2). This overestimation was mainly due to ~~the~~ high biomass of ~~the~~ *Picea* and *Populus* PFTs (Fig. 4-B). Despite ~~local-scale-local~~ overestimates, the range of genus-specific biomass variability of ~~the~~ *Abies* and *Populus* PFTs was well captured.

## 10 3.2 Fire history simulated by LPJ-LMfire

### 3.2.1 Fire activity

Simulated burn rates displayed multi-decadal variation over the ~~20th 0<sup>th</sup>-Century~~, ~~notably~~ mostly in Manitoba and Ontario (Fig. 5-A). ~~The h~~High fire activity that was reported ~~for in~~ the ~~period-1910-1930~~ period was followed by a decrease in fire activity until the 1970s, and then increased to levels similar to those of the early 20th ~~century~~ (Fig. 5-A). Since the 1970s, annual burn rates ~~have~~ increased in central Manitoba and western Ontario, and ~~in the~~ south-central areas of Quebec (Fig. 5-A). Episodes of successive years of intense fire activity ~~have~~ occurred in 1908-1910, 1919-1923, 1995-1998 and 2002-2007 (~~supplement S6-Fig. S5-A in Supplement S6~~). ~~SA similar temporal pattern of annual burn rates between 1901–2012 was reported within the ‘Climate-only’ experiment, but with lower annual burn rates (Fig. S7 in Supplement S7).~~

\_\_\_\_\_The simulated fire season was not stationary: a fire seasonality index (FSI) was computed as the percentage ~~of~~ difference between spring and summer total burned areas (~~supplement S6-Fig. S5-B in Supplement S6~~), and varied between 0.17% and 83%. The period ~~extending~~ from the end of the 1960s to end of the 1990s ~~corresponded~~ corresponds to a period during which several years of high FSI were observed compared ~~to-with~~ the entire time series. ~~A FSI that was~~ greater than 50% was calculated for 1968, 1977, 1980 and 1993 (~~supplement S6-Fig. S5-B in Supplement S6~~). May and June were consistently the spring months with the largest burned areas, while summer months recorded fewer and fewer burned areas over the course of the 20th ~~century~~.

### 3.2.2 Fuels

For the ‘Climate + CO<sub>2</sub>’ experiment, ~~the~~ simulated annual NPP ~~that was~~ averaged over the entire study region and ~~the~~ whole period was 5.4 T ha<sup>-1</sup>, with a minimum of 4.2 T ha<sup>-1</sup> in 1907 and a maximum of 7.1 T ha<sup>-1</sup> in 2003 (Fig. 5-B). Both sequential *t*-test analysis and temporal time-series showed that NPP ~~has~~ increased since the 1970s (Fig. 6-A and B), ~~largely~~ mostly in southern areas of Quebec and in eastern Ontario (Fig. 5-B). This constant increase in NPP since the 1970s was not observed in Manitoba and western Ontario, where a significant increase in annual burn rates was observed (Fig. 5-A). Some regions in

south-central Ontario showed a decline in NPP during the early 20th ~~century~~, and the same trend ~~was~~ observed in south-central Quebec since the 1980s. The proportion of cells recording a decline ~~of~~ in NPP was particularly noteworthy in 2004 and 2006 (Fig. 6-A and B).

~~Differences in NPP between the simulations ‘Climate + CO<sub>2</sub>’ and ‘Climate-only’ experiments have highlighted that~~

5 ~~a~~ Annual simulated NPP, averaged over the whole area, was positively correlated with annual atmospheric ~~CO<sub>2</sub> concentration~~ ~~[CO<sub>2</sub>]~~ ( $r^2 = 0.767$ ,  $P < 0.001$ ). Mean percentage ~~of~~ increase in NPP that incurred by rising ~~CO<sub>2</sub> concentration~~ ~~[CO<sub>2</sub>]~~ for our ~~five time~~ periods was 2.7%, 5.5%, 8.9%, 16.7% and 27.6% (~~supplement S7~~ Fig. S6 ~~in Supplement S7~~), while it was 18% for the entire period. An even larger effect of CO<sub>2</sub> fertilization was simulated in the extreme southern and northern parts of the study region (~~supplement S7~~ Fig. S6-C ~~in Supplement S7~~).

10 Mean total aboveground biomass averaged 66.4 T ha<sup>-1</sup> in eastern boreal Canada over the ~~period~~ 1901-2012 ~~period~~. Mean total aboveground biomass decreased slightly from the beginning of the 20th ~~century~~, until the 1930s, and then increased until 1995, after which it reached a stable level (Fig. 5-C). Periods of total aboveground biomass loss ~~that~~ were recorded at the beginning of the 20th ~~century~~, corresponded ~~with~~ high fire activity, as previously mentioned (Fig. 5-A). Sequential *t*-test analysis of total aboveground biomass time-series showed that biomass increase and reduction, ~~respectively~~, followed the same

15 trends that were observed for growth releases and declines, ~~respectively~~, until ~~the~~ year 2000 (Fig. 6-C and D). Genus-specific aboveground biomass of ~~the~~ *Picea*, *Pinus* and *Populus* PFTs showed the same increasing trends over the past century, whereas *Abies* PFT aboveground biomass decreased until ~~the~~ year 1960, before regaining the value it had at the beginning of the 20th ~~century~~ (~~supplement S8~~ Fig. S78-A ~~in Supplement S8~~). The strongest variation in total aboveground biomass occurred for the *Picea* PFT; ~~which~~ varied from a minimum of 27.8 T ha<sup>-1</sup> in 1910 to a maximum of 36.7 T ha<sup>-1</sup> in 2003 (~~supplement S8~~

20 Fig. S78-A ~~in Supplement S8~~). Conversely, genus-specific aboveground biomass of *Abies*, *Pinus* and *Populus* PFTs varied by less than 1 T ha<sup>-1</sup>, 2 T ha<sup>-1</sup> and 3 T ha<sup>-1</sup>, respectively, over the same period (~~supplement S8~~ Fig. S78-A ~~in Supplement S8~~). The ratio of mean genus-specific aboveground biomass in the recent ~~period of~~ 1991-2012 ~~period~~, when compared ~~to~~ with the past period ~~of~~ 1911-1930, was 1.23, 1.04, 1.13 and 1.31 for ~~the~~ *Picea*, *Abies*, *Pinus* and *Populus* PFTs, respectively. The highest ratios for each PFT were found in the northern areas (~~supplement S8~~ Fig. S78-B ~~in Supplement S8~~).

## 25 4 Discussion

### 4.1 Agreements and disagreements in fire activity and forest growth

We used LPJ-LMfire, which was driven by gridded climatology, atmospheric ~~CO<sub>2</sub> concentration~~ ~~[CO<sub>2</sub>]~~, and an estimate of lightning strike density, to study the pyrogeography of eastern Canada's boreal forest. Compared with ~~the~~ previous modelling efforts that ~~were~~ had been conducted by Pfeiffer et al. (2013) using the original LPJ-LMfire model, the results that are reported

30 here showed substantial improvements in the capacity of the DGVM to simulate fire ignition in the Canadian boreal forest. The use of a high-quality lightning strike data ~~set~~ instead of the low-resolution LIS/OTD global data ~~set~~ that was used by Pfeiffer et al. (2013) allowed us to capture the spatial gradient of fire activity in a substantially better manner (Baker et al.,

2016). The results confirmed that fire in the study area is strongly ignition-limited, while most fire models have simply assumed that fire ~~will~~would always occur under appropriate weather and fuel conditions, e.g., SIMFIRE (Hantson et al. 2016). LPJ-LMfire simulations confirmed the necessity of simulating fire in a model as the product of the probabilities that are associated with fuel, moisture and ignition.

5 ~~The~~ Inter-annual variation in lightning strike density was more faithfully reproduced when weighting the mean flash climatology with the CAPE variable to predict lightning-induced fire ignitions and their variability (Peterson et al., 2010). However, this variation is still constrained by the short temporal depth of the 11 years of record in the CLDN lightning strike data\_set (Orville et al., 2002; Kochtubajda and Burrows, 2010). Synchronicity in major fire activity years across provinces (e.g., 1961, 2005, 2007) was consistent with several studies on fire history, suggesting that changes in forest fire activity ~~were~~  
10 ~~have been~~ observed ~~con~~jointly over vast areas since the 1900s (e.g., Bergeron et al., 2004b; Macias Fauria and Johnson, 2008).

Annual burn rates (recent and historical) were underestimated in many areas of northern Quebec. It appears that the simulation could not capture the expression of a climate type that is encountered in the Clay Belt of northwestern Quebec, where periodic drought is known to occur. This likely may reflect some limitation that is imposed by the low density of weather stations north of 49°N. ~~Lack~~~~The lack~~ of station replication can create excessively smoothed climate records, thereby reducing  
15 the possibility of correctly emulating the relationship between climate and forest fire activity during extreme drought and fire years (Girardin et al., 2006b, 2009; Xiao and Zhuang, 2007). For example, 1989 is known as a drought year, which was induced by changes in atmospheric circulation ~~and~~ that ~~was~~ere at the origin of numerous large fires (> 50,000 ha) in Manitoba and Quebec (~~supplement S5~~ Fig. S4 ~~in Supplement S5~~; Goetz et al., 2006; Xiao and Zhuang, 2007). Other large fires exceeding 50,000 ha were observed in northern Quebec in 1983 and 2002 (~~supplement S5~~ Fig. S4 ~~in Supplement S5~~). However, these  
20 extreme weather conditions were not reproduced in our input data\_set and, consequently, the model could not simulate these very large fires. These underestimates may also result, in part, from the lack of lightning strike records in these northernmost regions, which prevents fire ignition from being simulated there. Polarity or energy of lightning was not taken into account in our simulations. Positive lightning strikes (transfers of positive charges to the ground) mainly occur in the north and correspond to 10-% of all strikes (Morissette and Gauthier, 2008), with the remaining strikes being negative. Positive strikes correspond  
25 to an exchange of energy between the highest part of ~~the~~ clouds and the soil, while negative strikes are triggered in a lower part of the clouds. For this reason, positive strikes are more likely to start fires because they carry higher energy owing to the greater travelling distance between ~~the~~ clouds and the soil (Flannigan and Wotton, 1991). As previously mentioned, the number of lightning sensors in northern regions (hatched areas on Fig. 4) is also limited (Orville et al., 2011), leading to ~~a decrease~~in  
in detection efficiency ~~over~~~~at~~ these latitudes (Morissette and Gauthier, 2008). Thus, 10-% of positive strikes are not  
30 appropriately captured and, consequently, the probability of fire ignition ~~also~~ is ~~also~~ likely to be underestimated in these areas. Underestimation of fire activity in northern areas had consequences for the simulation results. Amongst other things, simulated tree mortality was underestimated and, hence, biomass proliferated (as can be noted in Fig. 4 with ~~the~~ *Picea* PFT).

~~—Overestimates of simulated burn rates were reported for southern Ontario and Quebec. Indeed, some significant differences were expected with observed data in southern areas because they take into account anthropogenic effects. Forest management~~

~~has contributed to modifying the composition and distribution of forest fuels (Girardin and Terrier, 2015; Danneyrolles et al., 2016). However, alterations in fuel composition and structure that were induced by human activities are not implemented in the present LPJ-LMfire simulations. The strong correlations between our simulated annual burn rates and observed data further suggest that active fire suppression efforts and forest management since about the 1950s (Le Goff et al., 2008; Lefort et al., 2003) have not contributed much to shifting fire behaviour trajectories in our study region, which admittedly has very low densities of both population and infrastructure.~~

## 4.2 History of fire in the eastern boreal forest of Canada described by LPJ-LMfire

Based upon the above preliminary agreement and despite some disagreements, the temporal patterns of annual burn rates that were simulated by LPJ-LMfire were strongly consistent with [the](#) forest fire histories that have been reconstructed in many studies (e.g., Stocks et al., 2003; Bergeron et al., 2004a; Girardin et al., 2006a). Multi-decadal temporal changes in annual burn rates reflect the underlying influence of climate variability and extreme fire weather (Macias Fauria and Johnson, 2008; Girardin et al., 2009); these multi-decadal temporal changes were well represented in the input climate data\_sets. An increase in temperatures and stability ~~of in~~ precipitation between 1916 and 1924 ([supplement S9 Fig. S8S9 in Supplement S9](#)) could be [at](#) the origin of a high frequency of fire occurrence during those years, marking a pause in the decline of fire activity that had been observed since the 1850s (Bergeron et al., 2004a). Advection of humid air masses over eastern Canada between 1940 and 1970 contributed to the creation of moister conditions ([supplement S9 Fig. S8](#)), which can lessen the capacity of a fire to spread after a lightning-induced fire ignition (Macias Fauria and Johnson, 2008). Both inter-annual variation and unsynchronized trends in climatic variables may have brought about changes in fire activity and could have affected the fire\_ season, as ~~it is has been~~ proposed to have occurred over millennial timescales during the Holocene (Ali et al., 2012). For example, during [the](#) years 1977 and 1980, an increase in spring temperatures was observed, whereas spring precipitation decreased, which resulted in the total areas that were burned in spring being 50-% greater than in summer ([supplement S9 Fig. S89 in Supplement S9](#)).

Correlations between simulated and observed provincial annual burn rates were slightly higher than ~~have what has been~~ typically [been](#) encountered in past studies of fire-climate relationships over the region (e.g., Girardin et al., 2004, 2006a, 2009). For example, Girardin et al. (2009) reported that about 35-% of the variance in the annual areas that were burned in the provinces of Ontario and Quebec was explained by summer moisture availability. In our modelling experiment, we obtained values between 41-% and 50-% for these same provinces, without empirical adjustments (e.g., through regression analysis). The improvements that were made here reinforce the idea that aside from “top-down” climate control on fire activity, other factors such as lightning, fuel availability and composition can influence fire statistics (Podur et al., 2002). This highlights the necessity of reconstructing fire history in a complex system that is related to climate and vegetation by taking into account several feedbacks (Hantson et al., 2016).~~The strong correlations between our simulated annual burn rates and observed data further suggest that active fire suppression efforts and forest management since about the 1950s (Le Goff et al., 2008; Lefort~~

et al., 2003) have not contributed much to shifting fire behaviour trajectories in our study region, which admittedly has very low densities of both population and infrastructure.

LPJ-LMfire simulations provide evidence for ~~correctly simulated the~~ the combined influence of fuel conditions and ignition sources ~~to simulate~~ on fire within our study areas. Indeed, an increase in precipitation around the 1930s constrained fire activity, despite a very high lightning strike density (supplement S9 Fig. S8 in Supplement S9). Conversely, at the end of the century, an increase in lightning strike density and a drier climate (Fig. S8 in Supplement S9) resulted in an increase of in annual burn rates, in the presence of a relatively constant amount of precipitation (supplement S9 Fig. S8). The season during which precipitation events and lightning ignitions occurred ~~are~~ were also found to be ~~also~~ important. Notably, ~~Moreover, LPJ-LMfire doesid~~ not simulate the core of the fire season between June and August when the highest density of lightning strikes ~~takesooktakes~~ place (Morissette and Gauthier, 2008). This phenomenon finds an explanation in that ~~H~~heavy and intense rain events that ~~occur~~ occurring later ~~induring~~ during the summer decrease the probability of starting fires; ~~Indeed, low burn rate during the summer despite high lightning density could also result from the weather being~~ becomes less conducive to fire due to higher amounts of precipitation between July and September ~~than between~~ in comparison with April ~~to~~ and June (supplement S9 Fig. S8). ~~That being said,~~

~~Moreover, LPJ-LMfire does not simulate the core of the fire season between June and August when the highest density of lightning strikes takes place (Morissette and Gauthier, 2008). This result showed that heavy and intense rain events that occur later in the summer decrease the probability of starting fires, despite more lightning.~~

~~Our~~ simulation was biased with regard to the onset of fire seasonality. LPJ-LMfire simulated the core of the fire season earlier than ~~in~~ what is actually observed. LMfire excludes fire ignition when snow cover is present (Pfeiffer et al., 2013). However, detailed investigations at the grid-cell level ~~on~~ in our study area revealed that the Fire Danger Index, which was calculated by the LMfire module, was high as soon as all snow had melted in May and June. This index estimates the probability that an ignition event will start a fire, depending upon both fuel moisture and fire weather conditions (Thonicke et al., 2010). As suggested by Pfeiffer et al. (2013), LPJ-LMfire simulates a very quick drying-out of soils in spring when the snow cover has disappeared or snowmelt has occurred prematurely. This phenomenon may be the reason why ~~we were able to~~ simulated the occurrence of fire season onset earlier than what is observed in Canada's eastern boreal forest.

CO<sub>2</sub>-induced enhancement of NPP (Norby et al., 2005; Huang et al., 2007) was clearly emulated in LPJ-LMfire. Our simulated 18% growth enhancement, with a 50% increase in ~~{CO<sub>2}}</sub>~~ CO<sub>2</sub> concentration between 1901 and 2012, was higher than the 15% and 14% growth increases that have been proposed by ~~Hickler et al., 2008~~ Hickler et al. (2008) and Girardin et al. (2011), respectively. LPJ-LMfire is highly sensitive to atmospheric ~~{CO<sub>2}}</sub>~~ CO<sub>2</sub> concentration and interpreting its impacts must be made with caution (~~Girardin et al. 2011~~)(Girardin et al., 2011). That being said, our results suggest that CO<sub>2</sub>-induced enhancement of forest productivity can be offset by fires and climate, which is consistent with the results of Hayes et al. (2011) and Kelly et al. (2016). Despite strong CO<sub>2</sub>-induced enhancement of forest productivity in LPJ-LMfire, the total amount of aboveground biomass and forest composition did not ~~indeed~~ change significantly during the course of the simulation period.

The CO<sub>2</sub>-induced enhancement of NPP had a ~~positive~~ influence on ~~the annual burn rates by increasing the availability of fuel.~~

Under very dry conditions, such as ~~in~~ 1971-1990 and 1991-2012, an increase in fire activity led to a decrease in growth and biomass. Drier conditions during the past few decades provided indications for an increase in growth decline events and in biomass reduction ~~that was~~ related to an increase in fire activity. A similar trend in such conditions was observed around the  
5 1920s, but the range of these negative events during ~~the~~ past decades exceeded ~~sed~~ the historical range of variability ~~that was~~ recorded by the simulated forest. Fires had a non-negligible influence on the state of the boreal forest in eastern Canada, especially during the last few decades, but our results also confirmed ~~ed~~ the relative influence of climate alone on the forest in northern regions. Indeed, in northern areas in Quebec and Manitoba, biomass has not significantly increased, despite a very strong effect of CO<sub>2</sub>-induced enhancement (~~supplement S7~~ Fig. S6 ~~in Supplement S7~~). We hypothesize that with ongoing  
10 global warming, growth decline events could increase substantially, given that the positive effect of CO<sub>2</sub> ~~concentration~~ on the growth of forests ~~s~~ may not be strong enough to compensate for ~~the losses~~ of biomass to fires and climate change (Kurz et al., 2008), which could lead to the opening up of landscapes.

### 4.3 Uncertainties and future perspectives

The present study ~~has~~ demonstrated that LPJ-LMfire is generally able to capture fire history and forest growth trends in the  
15 eastern boreal forest of Canada. However, several uncertainties persist. First, forest establishment and the start of growth during the “spin~~up~~” phase was simulated using a detrended version of modern climate, as is usually performed in DGVM runs (Prentice et al., 2011; Pfeiffer et al., 2013; Yue et al., 2015; Knorr et al., 2016). This initial condition assumes that past relationships between climate, fire, and vegetation have been stationary through time and that variability of modern climate is representative of all variability that has been recorded over the past 1200 years (time of spin~~up~~ phase + 112 years of  
20 simulation). Yet, it has been increasingly recognized that such an assumption is invalid and that modern observations are not a good analogue for prehistoric variability (Kelly et al., 2016; Hudiburg et al., 2017). For example, fire activity over much of the Holocene was higher in terms of frequency and fire sizes ~~than are the~~ current levels across broad areas of eastern Canada (Girardin et al., 2013a; Remy et al., 2017). It is likely that not accounting for such variability may introduce biases in forest productivity dynamics and levels, more specifically on soil carbon dynamics (Hudiburg et al., 2017). This may be less  
25 problematic when studying fire and forest dynamics over the last century because the mean age of the major part of eastern boreal forest is less than 100 years (Bergeron et al., 2002).

The non-negligible influence of forest composition on fire regimes (Hély et al., 2001) is limited in the model to the representation of three needle-leaf PFTs and one broad-leaf PFT. Improving LPJ-LMfire's representation of biodiversity with further broad-leaf PFT genera could counterbalance or offset overestimates of fire activity in southern areas since these species  
30 are less flammable than needle-leaf species. Similarly, improving LPJ-LMfire parametrization to account for mosses could reduce overestimation of ~~the quantities~~ of fuel ~~that was~~ available in northern areas. In the Clay Belt, ~~the~~ poor drainage conditions ~~that were~~ induced by the presence of an impermeable clay substrate, flat topography and a cold climate facilitate the accumulation of thick layers of organic soil, an edaphic process that is ~~described~~ ~~referred to~~ as paludification (Fenton et al.,

2006). Once *Sphagnum* species increase on the forest floor, the depth of burn varies only slightly in response to changes in weather conditions, owing to very low fluctuations in the degree of water saturation of the organic layer (Fenton et al., 2006).

In the present study, simulations are limited by the relatively low accuracy of soil attributes in databases for Canada's boreal forest (Hengl et al., 2014). The input data set of soil attributes that was used in our simulations tended to underestimate clay and sand percentages ~~on~~in our study area when compared to point observations (~~supplement S10~~Fig. S910 in Supplement S10). These effects add up to other weaknesses in ~~the~~ physiological constraints, such as cold climate not being sufficiently restrictive and allowing *Picea* to become overly abundant in the simulation runs. While a previous study showed that the abundance of *Picea* decreases with latitude in the Tundra region and is coupled with the occurrence of dwarf shrubs in the *Ericaceae* and herbs (Gajewski et al., 1993), such species were not parametrized in the current version of LPJ-LMfire, due to a lack of information on their physiological and biogeographical preferences. Future research could incorporate recently developed parameterizations for boreal shrubs and non-vascular plants into LPJ-LMfire (Druel, 2017; Druel et al., 2017).

Forest stand structure and successional dynamics (age classes), together with processes leading to the formation of peatlands, are not included in the present version of LPJ-LMfire. Yet, all of these aspects are important determinants of fire ignition and propagation under a given climate (Hély et al., 2001) and ~~also~~ also influence the distinction between crown and surface fires, which affect tree mortality differently (Hély et al., 2003; Yue et al., 2015). Moreover, LPJ-LMfire, like most DGVMs, does not consider constraints on species migrations, phenotypic plasticity, and local adaptation of species (Morin and Thuiller, 2009). The simulation results may be overly optimistic in terms of the capacity of southern species to colonize newly available areas in northern regions as the climate warms. As previously mentioned by Morin and Thuiller (2009), species colonization in northern regions could be limited by forest attributes, such as fragmented landscapes or high competition levels from existing species, or through migrational lag (Epstein et al., 2007).

Wildland fires are the most important natural disturbances in Canada's eastern boreal forest, but non-fire and human disturbances also have considerable effects (Price et al., 2013) and may influence fire activity trajectories indirectly. Integrating a range of forest disturbances into a DGVM could improve the accuracy of forecasting and modelling ~~of~~ climate change effects on Canada's eastern boreal forest. For instance, insect damage (MacLean, 2016), and outbreaks of eastern spruce budworm (*Choristoneura fumiferana*) in particular (Zhang et al., 2014, James et al., 2017), represent ~~a~~ significant forest disturbances by the way they temporarily altering ing forest structure by affecting specific tree growth, tree survival, regeneration and succession. These disturbances can also have an important impact on fire activity, ~~in turn,~~ by modifying fuel distribution and connectivity (James et al., 2017). Additionally, successive fires that take place over a short period before the trees have attained maturity have ~~can~~ lead to complete regeneration failure (Girard et al., 2008). Such events in young, unproductive stands can also lead to modified forest composition (Girard et al., 2008) and could exert a strong feedback on ecosystem structure by generating changes in temporal fire patterns ~~on~~ over long timescales. Finally, ~~T~~the effects of human activities, such as forest management and active fire suppression efforts, on the composition and distribution of forest fuels ~~were~~ not implemented in the present LPJ-LMfire simulations. Nonetheless, ~~t~~the strong correlations between our simulated annual burn rates and observed data, ~~further~~ nonetheless, suggests that active fire suppression efforts and forest management since about the 1950s (Le Goff et al.,

2008; Lefort et al., 2003) have not contributed much to shifting fire behaviour trajectories in our study region, which admittedly has very low densities of both population and infrastructure in comparison with other populated areas such as in the United States (e.g., Syphard et al., 2017; Syphard et al., 2017).

## **5: Conclusions**

5 In this study, we used LPJ-LMfire to simulate fire activity from 1901 to 2012 in Canada's eastern boreal forest, at a 10-km resolution. LPJ-LMfire was parametrized for the predominant forest tree genera that ~~were~~are present in our study region, ~~viz.i.e.~~viz. i.e. *Picea*, *Abies*, *Pinus* and *Populus*. The predictive skill of the model to simulate fire activity was determined by comparing our model simulations with published data. LPJ-LMfire was able to simulate inter-annual- to decadal-scale fire variability from the beginning of the 20th ~~century~~century. However, the low density of weather stations in northern areas likely  
10 limited the model's ability to capture some extreme fire years. Our study highlights the importance of changes in climate variables at multi-decadal and annual timescales in strongly controlling spatiotemporal patterns of fire that were simulated by LPJ-LMfire. Spatiotemporal patterns were well captured, based upon our climate data inputs. Despite an overarching CO<sub>2</sub>-induced enhancement of NPP in LPJ-LMfire, aboveground biomass was relatively stable because of the compensatory effects of increasing fire activity. This study helps reduce uncertainties in our knowledge regarding fire patterns in the recent past and  
15 confirms that fires ~~were~~have been a dominant driver of boreal forest in eastern Canada during the last century. We further provide a new tool to refine predictions of future fire risks and effects of ongoing climate change in these forests to better inform management and improve risk mitigation strategies.

### **Code availability.**

20 The source code of LPJ-LMfire is available at: <https://github.com/ARVE-Research/LPJ-LMfire/tree/canada/src>.

### **Authors' ~~contributors~~ contribution.**

E. Chaste, M. Girardin, Y. Bergeron and C. Hély conceived and designed the study. E. Chaste performed the simulations and ~~preparation~~prepared of input data sets with the help of J. Kaplan. J. Portier performed statistical calculations of annual burn  
25 rates for our simulation period according to the protocol ~~that is~~ described by Portier et al. (2016). E. Chaste, M. Girardin, J. Kaplan, Y. Bergeron and C. Hély interpreted the results. E. C. Chaste prepared the manuscript with contributions from all co-authors.

## *Competing financial interests.*

The authors declare that they have no conflict of interest.

## *Acknowledgements.*

- 5 This study was made possible thanks to the financial support provided by the European IRSES NEWFOREST project, the Forest Complexity Modelling (FCM) program and the NSERC Strategic and Discovery programs. Jed Kaplan was supported by the European Research Council (COEVOLVE 313797). This research was conducted as part of the International Associated Laboratory MONTABOR (LIA France-Canada) and the International Research Group on Cold Forests. We thank Melanie Desrochers and Xiao Jing Guo for their help with mapping and computation for this project. We also thank Daniel Stubbs from
- 10 Calcul Quebec and Compute Canada for help with the Fortran language help and server space facilities for running LPJ-LMfire. We also thank W.F.J. Parsons and Isabelle Lamarre for English-language editing of the draft of this manuscript and the two anonymous reviewers for comments on its earlier version.

## **References**

- 15 Ali, A. A., Carcaillet, C. and Bergeron, Y.: Long-term fire frequency variability in the eastern Canadian boreal forest: the influences of climate vs. local factors, *Glob. Change Biol.*, 15(5), 1230–1241, doi:10.1111/j.1365-2486.2009.01842.x, 2009.
- Ali, A. A., Blarquez, O., Girardin, M. P., Hely, C., Tinquaut, F., El Guellab, A., Valsecchi, V., Terrier, A., Bremond, L., Genries, A., Gauthier, S. and Bergeron, Y.: Control of the multimillennial wildfire size in boreal North America by spring climatic conditions, *Proc. Natl. Acad. Sci. USA*, 109(51), 20966–20970, doi:10.1073/pnas.1203467109, 2012.
- 20 Andrews, P. L., Bevins, C. D. and Seli, R. C.: BehavePlus fire modeling system, version 4.0: User's Guide, [online] Available from: [https://www.fs.fed.us/rm/pubs/rmrs\\_gtr106.pdf](https://www.fs.fed.us/rm/pubs/rmrs_gtr106.pdf) (Accessed 24 July 2017), 2008.
- Baker, D. J., Hartley, A. J., Butchart, S. H. M. and Willis, S. G.: Choice of baseline climate data impacts projected species' responses to climate change, *Glob. Change Biol.*, 22(7), 2392–2404, doi:10.1111/gcb.13273, 2016.
- 25 Beaudoin, A., Bernier, P. Y., Guindon, L., Villemaire, P., Guo, X. J., Stinson, G., Bergeron, T., Magnussen, S. and Hall, R. J.: Mapping attributes of Canada's forests at moderate resolution through kNN and MODIS imagery, *Can. J. For. Res.*, 44(5), 521–532, doi:10.1139/cjfr-2013-0401, 2014.
- Beauregard, F. and de Blois, S.: Beyond a climate-centric view of plant distribution: edaphic variables add value to distribution models, *PLoS One*, 9(3), e92642, doi:10.1371/journal.pone.0092642, 2014.
- Bergeron, Y., Richard, P. J., Carcaillet, C., Gauthier, S., Flannigan, M. and Prairie, Y.: Variability in fire frequency and forest composition in Canada's southeastern boreal forest: a challenge for sustainable forest management, *Conserv. Ecol.*, 2(2), 1998.
- 30 Bergeron, Y., Leduc, A., Harvey, B. D. and Gauthier, S.: Natural fire regime: A guide for sustainable management of the Canadian boreal forest., *Silva Fenn.*, 36(1), 81–95, 2002.
- Bergeron, Y., Gauthier, S., Flannigan, M. and Kafka, V.: Fire regimes at the transition between mixedwood and coniferous boreal forest in northwestern Quebec, *Ecology*, 85(7), 1916–1932, doi:10.1890/02-0716, 2004a.

- Bergeron, Y., Flannigan, M., Gauthier, S., Leduc, A. and Lefort, P.: Past, current and future fire frequency in the Canadian boreal forest: implications for sustainable forest management, *Ambio*, 33(6), 356–360, doi:10.1579/0044-7447-33.6.356, 2004b.
- 5 Bergeron, Y., Chen, H. Y., Kenkel, N. C., Leduc, A. L. and Macdonald, S. E.: Boreal mixedwood stand dynamics: ecological processes underlying multiple pathways, *For. Chron.*, 90(2), 202–213, doi:10.5558/tfc2014-039, 2014.
- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H. and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res. Atmospheres*, 109, D14203, doi:10.1029/2003JD003697, 2004.
- 10 Boulanger, Y., Gauthier, S., Burton, P. J. and Vaillancourt, M.-A.: An alternative fire regime zonation for Canada, *Int. J. Wildland Fire*, 21(8), 1052–1064, 2012.
- Boulanger, Y., Gauthier, S. and Burton, P. J.: A refinement of models projecting future Canadian fire regimes using homogeneous fire regime zones, *Can. J. For. Res.*, 44(4), 365–376, doi:10.1139/cjfr-2013-0372, 2014.
- Brandt, J. P.: The extent of the North American boreal zone, *Environ. Rev.*, 17, 101–161, doi:10.1139/A09-004, 2009.
- 15 Canadian Council of Forest Ministers: National Forestry Database. Forest Fire Statistics by Province/Territory/Agency, 1990-2016, [online] Available from: [http://nfdp.ccfm.org/data/compendium/html/comp\\_31e.html](http://nfdp.ccfm.org/data/compendium/html/comp_31e.html), 2017.
- Canadian Forest Service: National Fire Database – Agency Fire Data. Natural Resources Canada, Canadian Forest Service, Northern Forestry Centre, Edmonton, Alberta., [online] Available from: <http://cwfis.cfs.nrcan.gc.ca/ah/nfdb> (Accessed 6 March 2017), 2016.
- 20 Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Matsui, N., Allan, R. J., Yin, X., Gleason, B. E., Vose, R. S., Rutledge, G., Bessemoulin, P., Brönnimann, S., Brunet, M., Crouthamel, R. I., Grant, A. N., Groisman, P. Y., Jones, P. D., Kruk, M. C., Kruger, A. C., Marshall, G. J., Maugeri, M., Mok, H. Y., Nordli, Ø., Ross, T. F., Trigo, R. M., Wang, X. L., Woodruff, S. D. and Worley, S. J.: The Twentieth Century Reanalysis Project, *Q. J. R. Meteorol. Soc.*, 137(654), 1–28, doi:10.1002/qj.776, 2011.
- 25 Danneyrolles, V., Arseneault, D. and Bergeron, Y.: Pre-industrial landscape composition patterns and post-industrial changes at the temperate–boreal forest interface in western Quebec, Canada, *J. Veg. Sci.*, 27(3), 470–481, doi:10.1111/jvs.12373, 2016.
- Druel, A.: Modélisation de la végétation boréale et de sa dynamique dans le modèle de surface continentale ORCHIDEE, Université Grenoble Alpes, Grenoble., 2017.
- 30 Druel, A., Peylin, P., Krinner, G., Ciais, P., Viovy, N., Peregon, A., Bastrikov, V., Kosykh, N. and Mironycheva-Tokareva, N.: Towards a more detailed representation of high-latitude vegetation in the global land surface model ORCHIDEE (ORCHID-VEGv1.0), *Geosci Model Dev Discuss*, 2017, 1–51, doi:10.5194/gmd-2017-65, 2017.
- Ecological Stratification Working Group: A national ecological framework for Canada. [online] Available from: [http://sis.agr.gc.ca/cansis/references/1996ew\\_a.html](http://sis.agr.gc.ca/cansis/references/1996ew_a.html) (Accessed 7 February 2017), 1996.
- Environment Canada: National Climate Data and Information Archive. Available at [climate.weatheroffice.gc.ca/](http://climate.weatheroffice.gc.ca/), [online] Available from: <http://climate.weatheroffice.gc.ca/> (Accessed 27 February 2017), 2013.
- 35 Epstein, H. E., Yu, Q., Kaplan, J. O. and Lischke, H.: Simulating future changes in arctic and subarctic vegetation, *Comput. Sci. Eng.*, 9(4), 12–23, doi:10.1109/MCSE.2007.84, 2007.

- Erni, S., Arseneault, D., Parisien, M.-A. and Bégin, Y.: Spatial and temporal dimensions of fire activity in the fire-prone eastern Canadian taiga, *Glob. Change Biol.*, 23(3), 1152–1166, doi:10.1111/gcb.13461, 2017.
- Fenton, N., Légaré, S., Bergeron, Y., Paré, D. and others: Soil oxygen within boreal forests across an age gradient, *Can. J. Soil Sci.*, 86(1), 1–10, doi:10.4141/S05-004, 2006.
- 5 Flannigan, M., Stocks, B., Turetsky, M. and Wotton, M.: Impacts of climate change on fire activity and fire management in the circumboreal forest, *Glob. Change Biol.*, 15(3), 549–560, doi:10.1111/j.1365-2486.2008.01660.x, 2009.
- Flannigan, M. D. and Wotton, B. M.: Lightning-ignited forest fires in northwestern Ontario, *Can. J. For. Res.*, 21(3), 277–287, 1991.
- 10 Flannigan, M. D., Stocks, B. J. and Wotton, B. M.: Climate change and forest fires, *Sci. Total Environ.*, 262(3), 221–229, 2000.
- Flannigan, M. D., Wotton, B. M., Marshall, G. A., Groot, W. J. de, Johnston, J., Jurko, N. and Cantin, A. S.: Fuel moisture sensitivity to temperature and precipitation: climate change implications, *Clim. Change*, 134(1–2), 59–71, doi:10.1007/s10584-015-1521-0, 2016.
- 15 Gajewski, K., Payette, S. and Ritchie, J. C.: Holocene vegetation history at the boreal-forest–shrub-tundra transition in North-Western Quebec, *J. Ecol.*, 81(3), 433–443, 1993.
- Gauthier, S., Bergeron, Y. and Simon, J.-P.: Effects of fire regime on the serotiny level of jack pine, *J. Ecol.*, 84(4), 539, doi:10.2307/2261476, 1996.
- 20 Gauthier, S., Bernier, P., Burton, P. J., Edwards, J., Isaac, K., Isabel, N., Jayen, K., Le Goff, H. and Nelson, E. A.: Climate change vulnerability and adaptation in the managed Canadian boreal forest, *Environ. Rev.*, 22(3), 256–285, doi:10.1139/er-2013-0064, 2014.
- Gewehr, S., Drobyshev, I., Berninger, F. and Bergeron, Y.: Soil characteristics mediate the distribution and response of boreal trees to climatic variability, *Can. J. For. Res.*, 44(5), 487–498, doi:10.1139/cjfr-2013-0481, 2014.
- 25 Girard, F., Payette, S. and Gagnon, R.: Rapid expansion of lichen woodlands within the closed-crown boreal forest zone over the last 50 years caused by stand disturbances in eastern Canada, *J. Biogeogr.*, 35(3), 529–537, doi:10.1111/j.1365-2699.2007.01816.x, 2008.
- Girardin, M. P. and Mudelsee, M.: Past and future changes in Canadian boreal wildfire activity, *Ecol. Appl.*, 18(2), 391–406, doi:10.1890/07-0747.1, 2008.
- Girardin, M. P. and Terrier, A.: Mitigating risks of future wildfires by management of the forest composition: an analysis of the offsetting potential through boreal Canada, *Clim. Change*, 130(4), 587–601, doi:10.1007/s10584-015-1373-7, 2015.
- 30 Girardin, M. P., Bergeron, Y., Tardif, J. C., Gauthier, S., Flannigan, M. D. and Mudelsee, M.: A 229-year dendroclimatic-inferred record of forest fire activity for the Boreal Shield of Canada, *Int. J. Wildland Fire*, 15(3), 375–388, doi:10.1071/WF05065, 2006a.
- 35 Girardin, M. P., Tardif, J. C., Flannigan, M. D. and Bergeron, Y.: Forest fire-conducive drought variability in the southern Canadian boreal forest and associated climatology inferred from tree rings, *Can. Water Resour. J.*, 31(4), 275–296, doi:10.4296/cwrj3104275, 2006b.

- Girardin, M. P., Ali, A. A., Carcaillet, C., Mudelsee, M., Drobyshev, I., Hély, C. and Bergeron, Y.: Heterogeneous response of circumboreal wildfire risk to climate change since the early 1900s, *Glob. Change Biol.*, 15(11), 2751–2769, doi:10.1111/j.1365-2486.2009.01869.x, 2009.
- 5 Girardin, M. P., Bernier, P. Y., Raulier, F., Tardif, J. C., Conciatori, F. and Guo, X. J.: Testing for a CO<sub>2</sub> fertilization effect on growth of Canadian boreal forests, *J. Geophys. Res. Biogeosciences*, 116, G01012, doi:10.1029/2010JG001287, 2011.
- Girardin, M. P., Ali, A. A., Carcaillet, C., Gauthier, S., Hély, C., Le Goff, H., Terrier, A. and Bergeron, Y.: Fire in managed forests of eastern Canada: Risks and options, *For. Ecol. Manag.*, 294, 238–249, doi:10.1016/j.foreco.2012.07.005, 2013a.
- Girardin, M. P., Ali, A. A., Carcaillet, C., Blarquez, O., Hély, C., Terrier, A., Genries, A. and Bergeron, Y.: Vegetation limits the impact of a warm climate on boreal wildfires, *New Phytol.*, 199(4), 1001–1011, doi:10.1111/nph.12322, 2013b.
- 10 Girardin, M.-P., Tardif, J. and Bergeron, Y.: Gradient analysis of *Larix laricina* dominated wetlands in Canada's southeastern boreal forest, *Can. J. Bot.*, 79(4), 444–456, doi:10.1139/b01-023, 2001.
- Girardin, M.-P., Tardif, J., Flannigan, M. D., Wotton, B. M. and Bergeron, Y.: Trends and periodicities in the Canadian Drought Code and their relationships with atmospheric circulation for the southern Canadian boreal forest, *Can. J. For. Res.*, 34(1), 103–119, doi:10.1139/x03-195, 2004.
- 15 Goetz, S. J., Fiske, G. J. and Bunn, A. G.: Using satellite time-series data sets to analyze fire disturbance and forest recovery across Canada, *Remote Sens. Environ.*, 101(3), 352–365, doi:10.1016/j.rse.2006.01.011, 2006.
- Hantson, S., Arneith, A., Harrison, S. P., Kelley, D. I., Prentice, I. C., Rabin, S. S., Archibald, S., Mouillot, F., Arnold, S. R., Artaxo, P. and others: The status and challenge of global fire modelling, *Biogeosciences*, 13(11), 3359–3375, doi:10.5194/bg-13-3359-2016, 2016.
- 20 Hayes, D. J., McGuire, A. D., Kicklighter, D. W., Gurney, K. R., Burnside, T. J. and Melillo, J. M.: Is the northern high-latitude land-based CO<sub>2</sub> sink weakening?, *Glob. Biogeochem. Cycles*, 25(3), GB3018, doi:10.1029/2010GB003813, 2011.
- Hély, C., Flannigan, M., Bergeron, Y. and McRae, D.: Role of vegetation and weather on fire behavior in the Canadian mixedwood boreal forest using two fire behavior prediction systems, *Can. J. For. Res.*, 31(3), 430–441, doi:10.1139/cjfr-31-3-430, 2001.
- 25 Hély, C., Flannigan, M. and Bergeron, Y.: Modeling tree mortality following wildfire in the southeastern Canadian mixed-wood boreal forest, *For. Sci.*, 49(4), 566–576, 2003.
- Hengl, T., de Jesus, J. M., MacMillan, R. A., Batjes, N. H., Heuvelink, G. B. M., Ribeiro, E., Samuel-Rosa, A., Kempen, B., Leenaars, J. G. B., Walsh, M. G. and Gonzalez, M. R.: SoilGrids1km — Global soil information based on automated mapping, *PLoS One*, 9(8), e105992, doi:10.1371/journal.pone.0105992, 2014.
- 30 Héon, J., Arseneault, D. and Parisien, M.-A.: Resistance of the boreal forest to high burn rates, *Proc. Natl. Acad. Sci.*, 111(38), 13888–13893, doi:10.1073/pnas.1409316111, 2014.
- Hessl, A. E.: Pathways for climate change effects on fire: Models, data, and uncertainties, *Prog. Phys. Geogr.*, 35(3), 393–407, doi:10.1177/0309133311407654, 2011.
- 35 Hickler, T., Smith, B., Prentice, I. C., Mjöfors, K., Miller, P., Arneith, A. and Sykes, M. T.: CO<sub>2</sub> fertilization in temperate FACE experiments not representative of boreal and tropical forests, *Glob. Change Biol.*, 14(7), 1531–1542, doi:10.1111/j.1365-2486.2008.01598.x, 2008.


- Huang, J.-G., Bergeron, Y., Denneler, B., Berninger, F. and Tardif, J.: Response of forest trees to increased atmospheric CO<sub>2</sub>, *Crit. Rev. Plant Sci.*, 26(5–6), 265–283, doi:10.1080/07352680701626978, 2007.
- Hudiburg, T. W., Higuera, P. E. and Hicke, J. A.: Fire-regime variability impacts forest carbon dynamics for centuries to millennia, *Biogeosciences Discuss.*, 1–24, doi:10.5194/bg-2017-42, 2017.
- 5 IPCC, Ed.: *Climate Change 2013 - The Physical Science Basis: Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge., 2013.
- James, P. M. A., Robert, L.-E., Wotton, B. M., Martell, D. L. and Fleming, R. A.: Lagged cumulative spruce budworm defoliation affects the risk of fire ignition in Ontario, Canada, *Ecol. Appl.*, 27(2), 532–544, doi:10.1002/eap.1463, 2017.
- 10 Kasischke, E. S. and Turetsky, M. R.: Recent changes in the fire regime across the North American boreal region—Spatial and temporal patterns of burning across Canada and Alaska, *Geophys. Res. Lett.*, 33(9), L09703, doi:10.1029/2006GL025677, 2006.
- Keeley, J. E.: Fire intensity, fire severity and burn severity: a brief review and suggested usage, *Int. J. Wildland Fire*, 18(1), 116, doi:10.1071/WF07049, 2009.
- 15 Keeling, R. F., Piper, S. C., Bollenbacher, A. F. and Walker, J. S.: Atmospheric CO<sub>2</sub> records from sites in the SIO air sampling network. In *Trends: A Compendium of Data on Global Change. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN, U.S.A.*, , doi:10.3334/CDIAC/atg.035, 2009.
- Kelly, R., Genet, H., McGuire, A. D. and Hu, F. S.: Palaeodata-informed modelling of large carbon losses from recent burning of boreal forests, *Nat. Clim. Change*, 6(1), 79–82, doi:10.1038/nclimate2832, 2016.
- 20 Knorr, W., Arneeth, A. and Jiang, L.: Demographic controls of future global fire risk, *Nat. Clim. Change*, 6(8), 781–785, doi:10.1038/nclimate2999, 2016.
- Kochtubajda, B. and Burrows, W.: *Cloud-to-Ground Lightning in Canada: 1999 - 2008*, vol. 12, p. 1280. [online] Available from: <http://adsabs.harvard.edu/abs/2010EGUGA..12.1280K> (Accessed 31 July 2017), 2010.
- Kurz, W. A., Stinson, G. and Rampley, G.: Could increased boreal forest ecosystem productivity offset carbon losses from increased disturbances?, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 363(1501), 2259–2268, doi:10.1098/rstb.2007.2198, 2008.
- 25 Kurz, W. A., Shaw, C. H., Boisvenue, C., Stinson, G., Metsaranta, J., Leckie, D., Dyk, A., Smyth, C. and Neilson, E. T.: Carbon in Canada’s boreal forest — A synthesis, *Environ. Rev.*, 21(4), 260–292, doi:10.1139/er-2013-0041, 2013.
- Le Goff, H., Girardin, M. P., Flannigan, M. D. and Bergeron, Y.: Dendroclimatic inference of wildfire activity in Quebec over the 20th century and implications for natural disturbance-based forest management at the northern limit of the commercial forest, *Int. J. Wildland Fire*, 17(3), 348–362, doi:10.1071/WF07080, 2008.
- 30 Lefort, P., Gauthier, S. and Bergeron, Y.: The influence of fire weather and land use on the fire activity of the Lake Abitibi area, eastern Canada, *For. Sci.*, 49(4), 509–521, 2003.
- Macias Fauria, M. and Johnson, E. A.: Climate and wildfires in the North American boreal forest, *Philos. Trans. R. Soc. B Biol. Sci.*, 363(1501), 2315–2327, doi:10.1098/rstb.2007.2202, 2008.
- 35 MacLean, D. A.: Impacts of insect outbreaks on tree mortality, productivity, and stand development, *Can. Entomol.*, 148, S138 – S159, doi:10.4039/tce.2015.24, 2016.

- Mansuy, N., Gauthier, S., Robitaille, A. and Bergeron, Y.: The effects of surficial deposit–drainage combinations on spatial variations of fire cycles in the boreal forest of eastern Canada, *Int. J. Wildland Fire*, 19(8), 1083–1098, doi:10.1071/WF09144, 2011.
- 5 Margolis, H. A., Nelson, R. F., Montesano, P. M., Beaudoin, A., Sun, G., Andersen, H.-E. and Wulder, M. A.: Combining satellite lidar, airborne lidar, and ground plots to estimate the amount and distribution of aboveground biomass in the boreal forest of North America I, *Can. J. For. Res.*, 45(7), 838–855, doi:10.1139/cjfr-2015-0006, 2015.
- Messaoud, Y., Bergeron, Y. and Asselin, H.: Reproductive potential of balsam fir (*Abies balsamea*), white spruce (*Picea glauca*), and black spruce (*P. mariana*) at the ecotone between mixedwood and coniferous forests in the boreal zone of western Quebec, *Am. J. Bot.*, 94(5), 746–754, doi:10.3732/ajb.94.5.746, 2007.
- 10 Molinari, C., Lehsten, V., Bradshaw, R. H. W., Power, M. J., Harmand, P., Arneith, A., Kaplan, J. O., Vannière, B. and Sykes, M. T.: Exploring potential drivers of European biomass burning over the Holocene: a data-model analysis, *Glob. Ecol. Biogeogr.*, 22(12), 1248–1260, doi:10.1111/geb.12090, 2013.
- Morin, X. and Thuiller, W.: Comparing niche- and process-based models to reduce prediction uncertainty in species range shifts under climate change, *Ecology*, 90(5), 1301–1313, doi:10.1890/08-0134.1, 2009.
- 15 Morissette, J. and Gauthier, S.: Study of cloud-to-ground lightning in Quebec: 1996–2005, *Atmosphere-Ocean*, 46(4), 443–454, doi:10.3137/AO919.2008, 2008.
- Moritz, M. A., Krawchuk, M. A., Parisien, M.-A. and others: Pyrogeography: Understanding the ecological niche of fire, *PAGES Newsl.*, 18(2), 83–85, 2010.
- Natural Resources Canada: Geobase - Canadian digital elevation data, Geogratis [online] Available from: 20 [http://ftp.geogratis.gc.ca/pub/nrcan\\_rncan/elevation/canada3d/](http://ftp.geogratis.gc.ca/pub/nrcan_rncan/elevation/canada3d/), 2007.
- Natural Resources Canada: Geobase - National hydro network, Geogratis [online] Available from: [http://ftp.geogratis.gc.ca/pub/nrcan\\_rncan/vector/geobase\\_nhn\\_rhn/](http://ftp.geogratis.gc.ca/pub/nrcan_rncan/vector/geobase_nhn_rhn/), 2010.
- Norby, R. J., DeLucia, E. H., Gielen, B., Calfapietra, C., Giardina, C. P., King, J. S., Ledford, J., McCarthy, H. R., Moore, D. J. P., Ceulemans, R., De Angelis, P., Finzi, A. C., Karnosky, D. F., Kubiske, M. E., Lukac, M., Pregitzer, K. S., Scarascia- 25 Mugnozza, G. E., Schlesinger, W. H. and Oren, R.: Forest response to elevated CO<sub>2</sub> is conserved across a broad range of productivity, *Proc. Natl. Acad. Sci. USA*, 102(50), 18052–18056, doi:10.1073/pnas.0509478102, 2005.
- Orville, R. E., Huffines, G. R., Burrows, W. R., Holle, R. L. and Cummins, K. L.: The North American lightning detection network (NALDN)—First results: 1998–2000, *Mon. Weather Rev.*, 130(8), 2098–2109, doi:10.1175/1520-0493(2002)130<2098:TNALDN>2.0.CO;2, 2002.
- 30 Orville, R. E., Huffines, G. R., Burrows, W. R. and Cummins, K. L.: The North American Lightning Detection Network (NALDN)—Analysis of flash data: 2001–09, *Mon. Weather Rev.*, 139(5), 1305–1322, doi:10.1175/2010MWR3452.1, 2011.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S. W., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S. and Hayes, D.: A large and persistent carbon sink in the world’s forests, *Science*, 333(6045), 988–993, doi:10.1126/science.1201609, 2011.
- 35 Payette, S., Filion, L. and Delwaide, A.: Spatially explicit fire-climate history of the boreal forest-tundra (Eastern Canada) over the last 2000 years, *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 363(1501), 2299–2314, doi:10.1098/rstb.2007.2201, 2008.


- Peterson, D., Wang, J., Ichoku, C. and Remer, L. A.: Effects of lightning and other meteorological factors on fire activity in the North American boreal forest: implications for fire weather forecasting, *Atmospheric Chem. Phys.*, 10(14), 6873–6888, doi:10.5194/acp-10-6873-2010, 2010.
- Pfeiffer, M. and Kaplan, J. O.: SPITFIRE-2: an improved fire module for Dynamic Global Vegetation Models, *Geosci. Model Dev. Discuss.*, 5, 2347–2443, 2012.
- Pfeiffer, M., Spessa, A. and Kaplan, J. O.: A model for global biomass burning in preindustrial time: LPJ-LMfire (v1. 0), *Geosci. Model Dev.*, 6(3), 643–685, doi:10.5194/gmd-6-643-2013, 2013.
- Podur, J., Martell, D. L. and Knight, K.: Statistical quality control analysis of forest fire activity in Canada, *Can. J. For. Res.*, 32(2), 195–205, doi:10.1139/X01-183, 2002.
- 10 Portier, J., Gauthier, S., Leduc, A., Arseneault, D. and Bergeron, Y.: Fire regime along latitudinal gradients of continuous to discontinuous coniferous boreal forests in eastern Canada, *Forests*, 7(10), 211, doi:10.3390/f7100211, 2016.
- Prentice, I. C., Kelley, D. I., Foster, P. N., Friedlingstein, P., Harrison, S. P. and Bartlein, P. J.: Modeling fire and the terrestrial carbon balance, *Glob. Biogeochem. Cycles*, 25(3), GB3005, doi:10.1029/2010GB003906, 2011.
- Price, D. T., Alfaro, R. I., Brown, K. J., Flannigan, M. D., Fleming, R. A., Hogg, E. H., Girardin, M.-P., Lakusta, T., Johnston, M. H., McKenney, D. W., Pedlar, J. H., Stratton, T., Sturrock, R. N., Thompson, I. D., Trofymow, J. A. and Venier, L. A.: Anticipating the consequences of climate change for Canada’s boreal forest ecosystems., *Environ. Rev.*, 21(4), 322–365, doi:10.1139/er-2013-0042, 2013.
- 15 Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D. S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzick, L., Spessa, A., Folberth, G. A., Sheehan, T., Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S. and Arneth, A.: The Fire Modeling Intercomparison Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, *Geosci Model Dev*, 10(3), 1175–1197, doi:10.5194/gmd-10-1175-2017, 2017.
- 20 Régnière, J., Saint-Amant, R., Béchard, A. and others: BioSIM 10 – Guide de l'utilisateur., *Ressour. Nat. Can., Serv. can. for., Cent. for. Laurentides, Québec (Québec)*. [online] Available from: <http://www.cfs.nrcan.gc.ca/publications/?id=34817> (Accessed 31 July 2017), 2014.
- 25 Remy, C. C., Hély, C., Blarquez, O., Maignan, G., Bergeron, Y., Lavoie, M. and Ali, A. A.: Different regional climatic drivers of Holocene large wildfires in boreal forests of northeastern America, *Environ. Res. Lett.*, 12(3), 035005, doi:10.1088/1748-9326/aa5aff, 2017.
- Rodionov, S. N.: A sequential algorithm for testing climate regime shifts, *Geophys. Res. Lett.*, 31(9), L09204, doi:10.1029/2004GL019448, 2004.
- 30 Rodionov, S. N.: Use of prewhitening in climate regime shift detection, *Geophys. Res. Lett.*, 33(12), doi:10.1029/2006GL025904, 2006.
- Rogers, B. M., Randerson, J. T. and Bonan, G. B.: High-latitude cooling associated with landscape changes from North American boreal forest fires, *Biogeosciences*, 10(2), 699–718, doi:10.5194/bg-10-699-2013, 2013.
- 35 Romps, D. M., Seeley, J. T., Vollaro, D. and Molinari, J.: Projected increase in lightning strikes in the United States due to global warming, *Science*, 346(6211), 851–854, doi:10.1126/science.1259100, 2014.

- Rothermel, R. C.: A Mathematical Model for Predicting Fire Spread in Wildland Fuels, Intermountain Forest and Range Experiment Station, Forest Service, U.S. Department of Agriculture, Ogden, Utah., 1972.
- Shorohova, E., Kneeshaw, D., Kuuluvainen, T., Gauthier, S. and others: Variability and dynamics of old-growth forests in the circumboreal zone: implications for conservation, restoration and management, *Silva Fenn.*, 45(5), 785–806, 2011.
- 5 Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K. and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9(2), 161–185, doi:10.1046/j.1365-2486.2003.00569.x, 2003.
- Smith, B., Prentice, I. C. and Sykes, M. T.: Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space, *Glob. Ecol. Biogeogr.*, 10(6), 621–637, doi:10.1046/j.1466-822X.2001.t01-1-00256.x, 2001.
- 10 Stocks, B. J., Mason, J. A., Todd, J. B., Bosch, E. M., Wotton, B. M., Amiro, B. D., Flannigan, M. D., Hirsch, K. G., Logan, K. A., Martell, D. L. and Skinner, W. R.: Large forest fires in Canada, 1959–1997, *J. Geophys. Res. Atmospheres*, 107(D1), 8149, doi:10.1029/2001JD000484, 2003.
- Syphard, A. D., Keeley, J. E., Pfaff, A. H. and Ferschweiler, K.: Human presence diminishes the importance of climate in driving fire activity across the United States, *Proc. Natl. Acad. Sci.*, 114(52), 13750–13755, doi:10.1073/pnas.1713885114, 2017.
- 15 Tang, G., Beckage, B., Smith, B. and Miller, P. A.: Estimating potential forest NPP, biomass and their climatic sensitivity in New England using a dynamic ecosystem model, *Ecosphere*, 1(6), 1–20, doi:10.1890/ES10-00087.1, 2010.
- Thonicke, K., Spessa, A., Prentice, I. C., Harrison, S. P., Dong, L. and Carmona-Moreno, C.: The influence of vegetation, fire spread and fire behaviour on biomass burning and trace gas emissions: results from a process-based model, *Biogeosciences*, 7(6), 1991–2011, doi:10.5194/bg-7-1991-2010, 2010.
- 20 Van Wagner, C. E., Forest, P. and others: Development and Structure of the Canadian Forest FireWeather Index System, in *Can. For. Serv., Forestry Tech. Rep., Citeseer*. [online] Available from: <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.460.3231> (Accessed 26 April 2017), 1987.
- 25 Wang, X., Thompson, D. K., Marshall, G. A., Tymstra, C., Carr, R. and Flannigan, M. D.: Increasing frequency of extreme fire weather in Canada with climate change, *Clim. Change*, 130(4), 573–586, doi:10.1007/s10584-015-1375-5, 2015.
- Xiao, J. and Zhuang, Q.: Drought effects on large fire activity in Canadian and Alaskan forests, *Environ. Res. Lett.*, 2(4), 044003, doi:10.1088/1748-9326/2/4/044003, 2007.
- Yue, C., Ciais, P., Zhu, D., Wang, T., Peng, S. S. and Piao, S. L.: How past fire disturbances have contributed to the current carbon balance of boreal ecosystems?, *Biogeosciences Discuss.*, 12, 14833–14867, doi:10.5194/bgd-12-14833-2015, 2015.
- 30 Zhang, X., Lei, Y., Ma, Z., Kneeshaw, D. and Peng, C.: Insect-induced tree mortality of boreal forests in eastern Canada under a changing climate, *Ecol. Evol.*, 4(12), 2384–2394, doi:10.1002/ece3.988, 2014.


**Captions**




**Figure 1** Map of eastern Canada's boreal forest from Manitoba to Newfoundland showing ecoregions in colour. The ecozone "Boreal Shield" is divided into three ecoregions: eastern Canadian forests, central Canadian Shield forests and midwestern Canadian Shield forests.




5 **Figure 2**— Observed versus LPJ-LMfire-simulated annual burn rates across eastern boreal Canada. (A) Observed annual burn rates computed for 365 hexagonal cells between 1980 and 2012 (data from Natural Resources Canada, 2017). (B) LPJ-LMfire simulated annual burn rates computed over the same period and hexagonal cells. (C) Percent age of difference between observed and simulated annual burn rates. (D) Percent age of difference between historical annual burn rates reconstructed from stand-replacing fire history studies (data from Girardin et al., 2013b; Héon et al., 2014; Portier et al., 2016) and LPJ-LMfire-simulated annual burn rates between 1911 and to 2012 (see supplement-Supplement S4 for further details). White points indicate where the observed (and historical) annual burn rate lies outside the 95-% confidence interval (95-% CI) of the averaged annual burn rates in hexagonal cells simulated by LPJ-LMfire.




**Figure 3** (A) Observed versus simulated annual total annual areas burned in three provinces of eastern Canada. Observed data (1959 to 2012) are from the Canadian National Fire Database (CNFDB). The Spearman's rank correlation between data is shown (correlations are significant at  $P < 0.05$  for Quebec and  $P < 0.001$  for the other provinces). (B) Monthly percentage of total areas burned between 1959 and 2012 in eastern boreal Canada.



**Figure 4.** (A) Observed, LPJ-LMfire simulated, and differences (%) in mean total aboveground biomass (T/ha) between 2000 and 2006 across eastern boreal Canada. (B) Genus-specific aboveground biomasses. The observed aboveground biomass maps across Canada were predicted and validated with photo-plot information in the southern areas (non-hatched areas) and data published by Beaudoin et al. (2014). Median (m) tree aboveground biomass values were also mentioned indicated for each maps; these and were calculated for the non-hatched areas at a 10-km resolution.



**Figure 5.** LPJ-LMfire simulated (A) annual burn rates (%), (B) net primary productivity (T/ha/yr), and (C) total aboveground biomass (T/ha) across eastern boreal Canada for five periods between 1911 and 2012.



**Figure 6.** (A) Annual (A) and (B) decadal (smoothed over 10-year sums; B) proportions of cells showing a significant decline and/or release in NPP with 90-% confidence intervals (error bars) computed using Bayes' method. (C) Annual (C) and (D) decadal (smoothed over 10-year sums; D) proportions of cells showing a significant reduction and/or increase in biomass total aboveground with 90-% confidence intervals (error bars) computed using the same method.

## Tables

**Table 1.** Climate and other data sets used to drive LPJ-LMfire.

	<b>Variables (units)</b>	<b>Period</b>	<b>Data sets</b>	<b>References</b>
<b>CLIMATE DATA</b>	Monthly mean temperature (°C) and monthly mean diurnal temperature range (°C)	1901-2012	Model “climatic monthly”, Software BIOSIM	(Environment Canada, 2013)
	Monthly mean precipitation (mm) and number of days per month with precipitation			
	Monthly mean of wind speed (m.s <sup>-1</sup> )	1969-2010		
	Monthly mean of total cloud cover percentage	1901-2012	20th <a href="#">Century-century</a> reanalysis	(Compo et al., 2011)
	Monthly mean convectible available potential energy (J.kg <sup>-1</sup> )			
Lightning flashes (number/day/km <sup>2</sup> )	1999-2010	Canadian lightning detection network	(Orville et al., 2011)	
<b>ENVIRONMENTAL CONSTRAINTS</b>	Soil particle size distribution and volume fraction of coarse fragments (%)	-	ISRIC - World Soil Information	(Hengl et al., 2014)
	Elevation (meters) and slope (degrees)	-	Canada 3D	(Natural Resources Canada, 2007)
	Water fraction	-	National Hydro Network (NHN)	(Natural Resources Canada, 2010)
	Atmospheric CO <sub>2</sub> concentrations (ppm)	-	Composite CO <sub>2</sub> time-series	(Keeling et al., 2009; Pfeiffer et al., 2013)

**Table 2-** LPJ-LMfire vs Margolis et al. (2015) mean total aboveground biomass estimates (with standard deviations) between 2000 and 2006 across 5-five ecozones in eastern boreal Canada. The Boreal Shield ecozone was divided into three ecoregions (ecozone subdivisions) for comparison.

Zone	Ecozones	Ecoregions	Mean TAB (T.ha <sup>-1</sup> )			
			LPJ-LMfire	GLAS	NFI	kNN
<b>North</b>	Taiga shield east	-	72.8 (30.0)	44.5	54.8	39.8
	Taiga shield west	-	38.6 (33.2)	38.1		25
	Hudson plain	-	59.0 (43.1)	26.1	24.4	37.2
<b>South</b>		Eastern Canadian forests	88.7 (17.7)	67.9		64.7
	Boreal shield	Central Canadian <u>Boreal</u> Shield forests	78.8 (17.3)	68.4	81.4	67.8
		Midwestern Canadian Shield forests	57.6 (15.1)	56.4		52.8
	Boreal plain	-	31.9 (23.5)	64.0	79.9	55.5

5