

## **Improving global paleogeography since the late Paleozoic using paleobiology**

Wenchao Cao<sup>\*,1</sup>, Sabin Zahirovic<sup>1</sup>, Nicolas Flament<sup>†,1</sup>, Simon Williams<sup>1</sup>, Jan Golonka<sup>2</sup> and R. Dietmar Müller<sup>1</sup>

<sup>1</sup> EarthByte Group, School of Geosciences, The University of Sydney, NSW 2006, Australia

<sup>2</sup> Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Mickiewicza 30, 30-059 Kraków, Poland

*\*Correspondence to:* Wenchao Cao ([wenchao.cao@sydney.edu.au](mailto:wenchao.cao@sydney.edu.au))

*†*Current address: School of Earth and Environmental Sciences, University of Wollongong, Northfields Avenue, Wollongong, New South Wales 2522, Australia

This file contains the following content.

- (1) Responses to comments from Referee #3 - Shanan Peters (Pages 2-5)
- (2) Responses to comments from Anonymous Referee #4 (Pages 6-8)
- (3) A marked-up version of the revised manuscript (Pages 9-32)

### Referee #3: Shanan Peters, [peters@geology.wisc.edu](mailto:peters@geology.wisc.edu)

**Reviewer:** Accurate paleogeographic reconstructions are required to test a wide range of hypotheses in the geosciences and they are a critical foundation upon which to build next-generation 4D Earth systems models. In order to produce the best possible reconstructions, no data can be left behind. This paper represents an important step towards a general method by which paleogeographically sensitive proxy data (e.g., fossils, sediments) can be used to test and improve existing paleogeographic reconstructions. This type of reconstruction model-data comparison is critical and the scale of the problem is such that algorithmic solutions and automation are of great added value. The paper is well written, the examples given are clear and powerful, and the potential of the approach and scientific outcomes are substantial. I have only a few suggestions, detailed below.

#### 1) Discrete timescales of reconstructions and the mapping of proxy data therein.

Table 1 outlines the timescale used for the Golonka reconstructions and this represents the first major cull of Paleobiology Database data. Only those collections that are temporally resolved in the PBDB to fall entirely within the bins specified by the reconstruction timescale are included. An example bin used is the “late Cenomanian-early Campanian.” According to the methods description, this means that any PBDB fossil collection assigned to a time interval of “Cenomanian” or “Campanian” would be omitted from the analysis and only fossil collections resolved to a finer biostratigraphic zone within these two international ages would be included. This is an understandable convention given the discrete bins of the original Golonka reconstructions, but this protocol results in a large cull of PBDB data. International ages (e.g., Cenomanian) are generally defined by our ability to consistently correlate globally. It is certainly the case that biostratigraphic zonation can be more precise, particularly in the marine record, but a stage-level age assignment for a given collection is something that PBDB data enterers would be very satisfied with in the majority of cases. Indeed, in the example of the Cenomanian and Campanian above, there are more than 2,200 collections resolved to these two intervals and these would not be included. Similar numbers of collections are probably omitted from all such divided international ages (this is somewhat conveyed in the differences between curves in Fig. 11). Does this cull matter? Given that the Cenomanian-Turonian sea level high stand is likely captured by at least some collections that are resolved only to “Cenomanian,” its possible that it does. There are a few statistical approaches one could take to overcoming this problem (in addition to improving the PBDB ages constraints, see below).

Finally, it is noted that PBDB data were “downloaded” from the database on a given date. This is a useful description, but the PBDB API allows for very specific definition of download protocols in the form of a URL. I strongly recommend that these details either be specified or, better still, that the URL for the API request be provided (it might also be good to include citations that describe these PBDB resources). Requesting an official PBDB publication number, should the manuscript be accepted, and including that in the Acknowledgements would be appropriate as well. John Alroy and company’s original vision and the many PBDB data contributors should be recognized in this capacity.

**Authors:** Yes, the abundance of fossil collections selected from the PBDB is sensitive to temporal resolution. Please refer to 5.3 “Marine fossil collection abundances in two different time scales” in the manuscript. In this study, we chose a conservative approach, only using fossil collections with temporal ranges lying entirely within the corresponding time intervals, to ensure that the resulting PBDB data do not unnecessarily “smear” the paleogeographic boundaries, although many fossil collections were ignored due to this selection criterion. One solution as the reviewer suggested here is improving the PBDB age constraints.

We specified the details of the downloaded information in the Supplementary Materials as the old PBDB API did not provide a URL for the paleobiology data when we requested them on 7 September 2016. We have acknowledged John Alroy et al.’s original vision and all the PBDB data contributors, and added an official PBDB publication number in the Acknowledgements.

**Reviewer:** 2) The “500 km test”

Fossil collections deviating by more than 500 km from previously interpreted paleoshorelines are excluded herein. Why 500km? Why not 397 km? Or 193 km? With few exceptions, whenever there is some value arbitrarily defined as a threshold there is a more interesting and principled approach. I would find the distribution of deviations between “previously interpreted” shorelines and PBDB collections fascinating, both in aggregate and on an interval-by-interval and/or a plate-by-plate basis. These distributions would have statistical utility of several different types, including defining a principled threshold criterion for data rejection. To me, this would be single most interesting analytical addition to the paper.

One potential utility of adopting an approach that leveraged the statistical nature of the distribution of deviations is that this could be used to add quantitative estimates of error (mostly of a random nature) that is introduced at every step, from the original coordinates of PBDB collections (some of which are most certainly not accurately located) to the assignment of an age.

**Authors:** We estimate the distances of marine fossil collections from the paleo-coastlines derived from the original paleogeographic maps of Golonka et al. (2006) since the Cretaceous period (see Figure 1 below). The result indicates that most marine fossil collections are within 500 km from the paleo-coastlines. We also have clarified this in the manuscript. In addition, we are trying to avoid the use of fossil data in cases where there may have been local/regional lakes or inland seas, that may not have been captured by the starting paleogeography. Please see the red points on Figure 4c in the manuscript and a set of PBDB test maps in the Supplementary Materials.




Figure 1. Cumulative frequency of the distance of the marine fossil collections from PBDB to paleo-coastlines derived from the paleogeographic maps of Golonka et al. (2006) since the Cretaceous period. Note that fossil collections located more than ~500 km away from paleo-coastlines represent outliers of their distribution.

**Reviewer:** 3) Minor points/questions: Use of fossils in pre-Triassic reconstructions, plates that appear during Phanerozoic, and next steps.

For geologically obvious reasons, not all plates can be tracked backwards in time to the Paleozoic. Are such plates excluded here? Does this matter to any of the results presented herein?

**Authors:** Yes, such plates and continental fragments are excluded. Excluding the plates that do not exist for the whole time period leads to discarding the fossil collections that are located on these plates.

**Reviewer:** Prior to the constraints on rotations provided by sea floor data, there is considerable uncertainty in the paleopositions of the continents, particularly with respect to longitude. At such times, the fossil record becomes increasingly important, not just for shoreline reconstructions but also for continent positions. I have previously detected a quantitative signature of PBDB biogeographic patterns in Paleozoic reconstructions and have concluded that at least some of the signal reflects the fact the the fossil record was relied on more heavily in the Paleozoic than in the post Paleozoic. Does this matter? Is a potentially changing role for fossils in deriving paleopositions detectable in any way here?

**Authors:** This point is interesting but in our view out of the scope of this study. This study mainly provides a workflow to revise the paleo-coastline locations and paleogeographic geometries using paleo-environmental information indicated by the marine fossil collections from the PBDB.

**Reviewer:** I'm excited by the possibility of completely closing the loop, from aggregation of paleogeographically-useful data in PBDB and Macrostrat-type resources to testing and improving paleogeographic reconstructions, in a largely automated, algorithmic fashion. Certainly there are things we are working on now to improve the temporal resolution of PBDB fossil collections, notably by integrating them with stratigraphic models such as those in Macrostrat. I'd be very interested in discussing how to improve this process and to fully capitalize on paleogeographic reconstructions in a way that avoids circularity problems and that makes the data as useful as possible to the GPlates team, and vice versa. This paper is an important and welcome first step.

**Authors:** We think it is a good idea to integrate stratigraphic data from Macrostrat Database to further constrain the paleogeographic reconstructions in a more automated and algorithmic fashion. We recognise the reviewer's ongoing work of improving the temporal resolution of PBDB fossil collections, which will greatly improve the availability of PBDB data. We welcome the reviewer's appreciation of our work.

## Anonymous Referee #4

**Reviewer:** There is a growing appreciation of the wide and constructive uses of paleogeographic maps for many areas of the geosciences. The community is transitioning from “static” (as the authors call them) or non digital to digital format paleogeographic maps. As with many aspects of our science, the effort is somewhat scattered and in this case the University of Sydney group is a pioneer in this area.

The authors stated main contribution is to “develop a workflow to restore the ancient paleogeographic geometries back to their modern coordinates”. Their longer term goal is to provide the “first step towards the construction of paleogeographic maps with flexible spatial and temporal resolutions that are more easily testable and expandable with the incorporation of new paleo-environmental datasets...”. These are worthy goals and I think this manuscript can make a contribution. But I found myself asking almost philosophical questions on the approach here rather than specific comments on the results. I think this manuscript is concise in its explanation of the methods and main results, but it needs expansion of the Discussion. My points below can mainly be addressed in an expansion of the 5.4 Limitations of the Workflow section into a more general Discussion section.

One major point I have is this. As the earlier reviewer #1 implies and I ask – should we use older paleogeographic reconstructions as is being done here, or essentially start over with more quantitative local to regional data and build up to global scale maps? Related to this – are the revised coastlines presented here an improvement over the Golonka 2006 coastlines, and how do we really know? A more general question is how to transition from the largely pre-digital era and publications with little (or at least much less) meta data and documentation of sources of data to the fully digital era? The lesser documentation of earlier paleogeographic research is mainly inevitable in that most (all?) older primary publications with proposed paleoenvironmental features (for example, coastlines) did not give quantitative spatial data in the modern sense. Note for example that the authors in their response to review #1 admit that: “Since the original data that were used to estimate the coastlines are not available for us, it is difficult to give the weight to the paleobiology data. The coastlines drawn on the original maps represent maximum transgression surfaces and we do not know much about their errors.” This latter point is made by the authors despite the well-respected Professor Golonka as the lead author of the synthesis of the coastlines (his 2006 paper) used in the methods in the present paper. In summary, the current community that produces quantitative tectonic reconstructions and paleogeographic maps may have to admit that it might have to essentially “start over” with regional-scale studies in which paleobiological and paleoenvironmental data are entered on GIS-based or other spatially quantitative base maps rather than try to “improve” older maps.

**Authors:** Rebuilding a global scale paleogeographic maps requires access to the original data that were used to estimate paleogeographic maps. However, these original data are not available in the scientific community as they are confidential industrial data, even though Professor Golonka is a co-author of this study. The consistency ratio between the original paleogeographic maps of Golonka et al. (2006) and the paleo-environmental data from the PBDB used in this study is ~75% on average since the Devonian period. This indicates a good

quality and reliability of paleo-coastlines from Golonka et al. (2006). Ideally, we would build the paleogeographic maps from scratch, using a variety of data types with high spatial and temporal resolution. This is currently not possible, and is beyond the scope of our work.

As we note in the manuscript, the consistency ratio between the revised paleogeography and the paleo-environments indicated by the marine fossil collections is increased to nearly full consistency (100%). This indicates the revised coastlines are improved over the original coastlines of Golonka et al. (2006).

**Reviewer:** Another point of discussion – there are serious problems with the variability of the temporal resolution of the various paleoenvironmental data used here and this is only partly addressed. For example, maps attempting to show maximum transgressions imply that we know the coastline location at very specific times in the past (especially during Icehouse periods) with perhaps thousands to a few tens thousands years resolution. I agree with reviewer #1 that it might be best, and more appropriate to the data available, to locate only marginal marine zones or belts that suggest much wider temporal and spatial spans involved (10s to a few 100 thousands years). For example, individual fossil locations in PBDB have a temporal resolution dependent on the type of index fossil and where it lies in the paleontological record. Many index fossils have a resolution of many 100 thousands to 1-2 million years. My point is not that this variability of data resolution invalidates using these different data sets, but that there needs to be more discussion here of what exactly is being depicted on the paleogeographic maps and what a particular paleogeographic feature means in terms of temporal and spatial resolution.

**Authors:** Again, due to the inaccessibility of original data that were used to build the paleogeographic maps, we cannot estimate the temporal resolution of the coastline locations on the paleogeographic maps. A fossil location from the PBDB does have a temporal resolution. We have added some clarification in the 5.4 “Limitation of the workflow” section of the manuscript.

**Reviewer:** Finally, I think these points should be addressed with more discussion in the 5.4 Limitations of the Workflow section. How can the community best move to developing quantitative paleogeography maps? And what have the authors learned by trying to update older paleogeographic maps? The current discussion in section 5.4 of the PBDB is good and provides valid suggestions. Here is an example of the need for more discussion: you state in section 5.4 that “A remaining question is how to provide a continuous representation of paleogeographic change that combines continuous plate motion models with paleogeographic maps that do not explicitly capture changes at the same temporal resolution.” I think the authors who just completed this workflow project should make suggestions on how to answer this question. I would also like the authors to directly address the earlier point I made on the value of revising older paleogeographic maps (Golonka or Blakey as examples) versus building new maps from more fragmentary, but spatially and temporally more quantitative, data from the PBDB, StratDB and Macrostrat.

**Authors:** From this study, we have learnt that (1) the paleogeographic maps of Golonka et al. (2006) are good estimates indicated by a high consistency with paleo-environmental data from the PBDB; (2) paleogeographically sensitive proxy data, such as paleo-environmental

data from the PBDB used in this study, can be used to test and improve existing paleogeographic reconstructions, which is the first step towards the construction of paleogeographic maps with flexible spatial and temporal resolutions.

In order to produce more quantitative paleogeographic maps, especially for local or regional paleogeographic reconstructions, it is important to integrate various data such as paleo-environment and paleo-lithofacies data, stratigraphic data from Macrostrat Database and StratDB Database, and tectonic settings.

# Improving global paleogeography since the late Paleozoic using paleobiology

Wenchao Cao<sup>\*1</sup>, Sabin Zahirovic<sup>1</sup>, Nicolas Flament<sup>†,1</sup>, Simon Williams<sup>1</sup>, Jan Golonka<sup>2</sup> and R. Dietmar Müller<sup>1</sup>

5 <sup>1</sup> EarthByte Group, School of Geosciences, The University of Sydney, NSW 2006, Australia

<sup>2</sup> Faculty of Geology, Geophysics and Environmental Protection, AGH University of Science and Technology, Mickiewicza 30, 30-059 Kraków, Poland

*\*Correspondence to:* Wenchao Cao (wenchao.cao@sydney.edu.au)

10 <sup>†</sup>Current address: School of Earth and Environmental Sciences, University of Wollongong, Northfields Avenue, Wollongong, New South Wales 2522, Australia

**Abstract.** Paleogeographic reconstructions are important to understand Earth's tectonic evolution, past eustatic and regional sea level change, paleoclimate and ocean circulation, deep Earth resources, and to constrain and interpret the dynamic topography predicted by mantle convection models. Global paleogeographic maps have been compiled and published, but they are generally presented as static maps with varying map projections, different time intervals represented by the maps, and different plate motion models that underlie the paleogeographic reconstructions. This makes it difficult to convert the maps into a digital form and link them to alternative digital plate tectonic reconstructions. To address this limitation, we develop a workflow to restore global paleogeographic maps to their present-day coordinates and enable them to be linked to a different tectonic reconstruction. We use marine fossil collections from the Paleobiology Database to identify inconsistencies between their indicative paleo-environments and published paleogeographic maps, and revise the locations of inferred paleo-coastlines that represent the estimated maximum transgression surfaces by resolving these inconsistencies. As a result, the consistency ratio between the paleogeography and the paleo-environments indicated by the marine fossil collections is increased from an average 75% to nearly full consistency (100%). The paleogeography in the main regions of North America, South America, Europe and Africa is significantly revised, especially in Late Carboniferous, Middle Permian, Triassic, Jurassic, Late Cretaceous and most of Cenozoic times. The global flooded continental areas since Early Devonian times calculated from the revised paleogeography in this study are generally consistent with results derived from other paleo-environment and paleo-lithofacies data and with the strontium isotope record in marine carbonates. We also estimate the terrestrial areal change over time associated with transferring reconstruction, filling gaps and modifying the paleogeographic geometries based on the paleobiology test. This indicates that the variation of the underlying plate reconstruction is the main factor that contributes to the terrestrial areal change, and the effect of revising paleogeographic geometries based on paleobiology is secondary.

## 1 Introduction

40

Paleogeography, describing the ancient distribution of highlands, lowlands, shallow seas, and deep ocean basins, is widely used in a range of fields including paleoclimatology, plate tectonic reconstructions, paleobiogeography, resource exploration and geodynamics. Global deep-time paleogeographic compilations have been published (e.g. Blakey, 2008; Golonka et al., 2006; Ronov, et al., 1984, 1989; Scotese, 2001, 2004; Smith et al., 1994). However, they are generally presented as static paleogeographic snapshots with varying map projections and different time intervals represented by the maps, and are tied to different plate motion models. This makes it difficult to convert the maps into a digital format, link them to alternative digital plate tectonic reconstructions, and update them when plate motion models are improved. It is therefore challenging to use paleogeographic maps to help constrain or interpret numerical models of mantle convection that predict long-wavelength topography (Gurnis et al., 1998; Spasojevic and Gurnis, 2012) based on different tectonic reconstructions, or as an input to models of past ocean and atmosphere circulation/climate (Godd ris et al., 2014; Golonka et al., 1994) and models of past erosion/sedimentation (Salles et al., 2017).

45

50

55

In order to address these issues, we develop a workflow to restore the ancient paleogeographic geometries back to their modern coordinates so that the geometries could be attached to a different plate motion model. This is the first step towards the construction of paleogeographic maps with flexible spatial and temporal resolutions that are more easily testable and expandable with the incorporation of new paleo-environmental datasets (e.g. Wright et al., 2013). In this study, we use a set of global paleogeographic maps (Golonka et al., 2006) covering the entire Phanerozoic time period as the base paleogeographic model. Coastlines on these paleogeographic maps represent estimated maximum marine transgression surfaces (KieSSLing et al., 2003). We first restore the global paleogeographic geometries of Golonka et al. (2006) to their present-day coordinates by reversing the sign of the rotation angle, and then reconstruct them to geological times using a different plate motion model of Matthews et al. (2016). We then use paleo-environmental information from marine fossil collections from the Paleobiology Database to modify the inferred paleo-coastline locations and paleogeographic geometries. Next, we use the revised paleogeography to estimate the surface areas of global paleogeographic features including deep oceans, shallow marine environments, landmasses, mountains and ice sheets. In addition, we compare the global flooded continental areas since the Devonian Period calculated from the revised paleogeography with other results derived from other paleo-environment and paleo-lithofacies maps (Ronov, 1994; Smith et al., 1994; Walker et al., 2002; Blakey, 2003, 2008; Golonka, 2007b, 2009, 2012) or from the Strontium isotope record (van der Meer et al., 2017). We estimate the terrestrial areal change over time associated with transferring reconstruction, filling gaps and modifying the paleogeographic geometries based on consistency test. Finally, we test the marine fossil collection dataset used in this study for fossil abundances over time using different time scales of 2016 time scale of the International Commission on Stratigraphy (ICS2016) and Golonka (2000) and discuss the limitations of the workflow we develop in this study.

60

65

70

75

## **2 Data and Paleogeographic Model**

80

The data used in this study are global paleogeographic maps and paleo-environmental data for the last 402 million years (Myr), which originate from the set of paleogeographic maps produced by Golonka et al. (2006) and the Paleobiology Database (PBDB, paleobiodb.org), respectively. The global paleogeographic compilation extending back to Early Devonian times of Golonka et al. (2006) is  
85 divided into 24 time-interval maps using the time scale of Golonka (2000) which is based on the original time scale of Sloss (1988) (Table 1). Each map is a compilation of paleo-lithofacies and paleo-environments for each geological time interval. These paleogeographic reconstructions illustrate the changing configuration of ice sheets, mountains, landmasses, shallow marine environments (inclusive of shallow seas and continental slopes) and deep oceans over the last ~400 Myr.

90

[Insert Table 1]

The paleogeographic maps of Golonka et al. (2006) are constructed using a plate tectonic model available in the Supplement of Golonka (2007a), where relative plate motions are described. In this  
95 rotation model, paleomagnetic data are used to constrain the paleolatitudinal positions of continents and rotation of plates, and hot spots, where applicable, are used as reference points to calculate paleolongitudes (Golonka, 2007a). This rotation model is necessary to restore these paleogeographic geometries (Golonka et al., 2006) to their present-day coordinates so that they can be attached to a different plate motion model. The relative plate motions of Golonka (2006, 2007a) are based on the  
100 reconstruction of Scotese (1997, 2004).

Here, we use a global plate kinematic model to reconstruct paleogeographies back in time from present-day locations. The global tectonic reconstruction of Matthews et al. (2016), with continuously closing plate boundaries from 410-0 Ma, is primarily constructed from a Mesozoic and Cenozoic plate  
105 model (230-0 Ma) (Müller et al., 2016) and a Paleozoic model (410-250 Ma) (Domeier and Torsvik, 2014). This model is a relative plate motion model that is ultimately tied to Earth's spin axis through a paleomagnetic reference frame for times before 70 Ma, and a moving hotspot reference frame for younger times (Matthews et al., 2016).

110

[Insert Figure 1]

The PBDB is a compilation of global fossil data covering deep geological time. All fossil collections in the database contain detailed metadata, including the time range (typically biostratigraphic age), present-day geographic coordinates, host lithology, and paleo-environment. Figure 1 represents  
115 distributions of the global fossil collections at present-day coordinates and shows their numbers since the Devonian Period. The recorded fossil collections are unevenly distributed both spatially and temporally, largely due to the differences in fossil preservation, the spatial sampling biases of fossil localities and the uneven entry of fossil data to the PBDB (Alroy, 2010). For this study, a total of

120 57,854 fossil collections with temporal and paleo-environmental assignments from 402 Ma to 2 Ma  
were downloaded from the database on 7 September 2016.

### 3 Methods

[Insert Figure 2]

125

The methodology can be divided into three main steps: (1) the original paleogeographic geometries are restored to present-day coordinates by applying the inverse of the rotations used to make the reconstruction, (2) these restored geometries are then rotated to new locations using the plate tectonic model of Matthews et al. (2016), (3) the paleo-coastline locations and paleogeographic geometries are adjusted using paleo-environmental data from the PBDB. Figure 2 illustrates the generalized workflow that can be applied to a different paleogeography model. In order to represent the paleogeographic maps as digital geographic geometries, they are first georeferenced using the original projection and coordinate system (global Mollweide in Golonka et al., 2006), and then reprojected into the WGS84 geographic coordinate system. The resulting maps are then attached to the original rotation model using the open-source and cross-platform plate reconstruction software *GPlates* (gplates.org). Every plate is then assigned a unique plate ID that defines the rotation of the tectonic elements so that the paleogeographic geometries can be rotated back to their present-day coordinates (see example in Figs 3a, b). We use present-day coastlines and terrane boundaries with the plate IDs of Golonka (2007a) as a reference to refine the rotations and ensure that the paleogeographic geometries are restored accurately to their present-day locations.

140

When the paleogeographic geometries in present-day coordinates are attached to a new reconstruction model, as Matthews et al. (2016) used in this study, the resulting paleogeographies result in gaps (Fig. 3c, pink) and overlaps between neighbouring polygons, when compared to the original reconstruction (Fig. 3a). These gaps and overlaps essentially arise from the differences in the reconstructions described in Matthews et al. (2016) and Golonka et al. (2006). The reconstruction of Golonka et al. (2006) has a tighter fit of the major continents within Pangea prior to the supercontinent breakup. In addition, this reconstruction contains a different plate motion history and block boundary definitions in regions of complex continental deformation, for example along active continental margins (e.g. Himalayas, western North America, Fig. 3c).

145

150

The gaps and overlaps cause changes in the total terrestrial or oceanic paleogeographic areas at different time intervals, becoming larger or smaller, when compared with the original paleogeographic maps (Golonka et al., 2006). The gaps can be fixed by interactively extending the outlines of the polygons in a GIS platform to make the plates connect as in the original paleogeographic maps (Fig. 3a, c, d). Changes in the extent of total terrestrial or oceanic area of the paleogeographies with filled gaps are compared with the original paleogeographies in Fig. 3d (Golonka et al., 2006).

155

[Insert Figure 3]

160

Once the gaps are filled, the reconstructed paleogeographic features are compared with the paleo-environments indicated by the marine fossil collections from the PBDB. These comparisons aim to identify the differences between the mapped paleogeography and the marine fossil collection environments in order to revise the paleo-coastline locations and paleogeographic geometries. Fossil collections belonging to each time interval (Table 1, Golonka, 2000) are first extracted from the dataset downloaded from the PBDB. Only the fossil collections with temporal ranges lying entirely within the corresponding time intervals are selected, as opposed to including the fossil collections that have larger temporal ranges. Fossil collections with temporal ranges crossing any time-interval boundary are not taken into consideration. As a result, a minimum number of fossil collections are selected for each time interval. The selected fossil collections are classified into either terrestrial or marine setting category, according to a lookup table (Table 2).

165

170

[Insert Table 2]

175

Marine fossil collections are then attached to the plate motion model of Matthews et al. (2016) so they can be reconstructed at each time interval. Subsequently, a point-in-polygon test is used to determine whether the indicated marine fossil collection is within the appropriate marine paleogeographic polygon. The results of these tests is discussed in the following section.

180

In the next step, we modify the paleo-coastline locations and paleogeographic geometries based on the test (Fig. 4, 5 and Supplement). Modifications are made according to the following rules: (1) Marine fossil collections from the PBDB are presumed to be well-dated, constrained geographically, not reworked and representative of their broader paleo-environments. Their indicative environments are assumed to be correct. (2) Only marine fossil collections within 500 km of the nearest paleo-coastlines are taken into account as most marine fossil collections used in this study are located within 500 km from the paleo-coastlines (see Figure S1 in the Supplementary Materials). (3) The paleo-coastlines and paleogeographic geometries are modified until they are consistent with the marine fossil collection environments and at the same time remain about 30 km distance from the fossil points used (Fig. 5c, f, l). (4) The adjacent paleo-coastlines are accordingly adjusted and smoothed (Fig. 4, 5). (5) The modified area (Fig. 5b, e, k, blue) resulting from shifting the coastline is filled using the shallow marine environment. These rules are designed to maximize the use of the paleo-environmental information obtained from the marine fossil collections to improve the coastline locations and paleogeography while attempting to minimize spurious modifications.

185

190

195

[Insert Figure 4]

[Insert Figure 5]

However, in some rare cases, outlier marine fossil data may be a deceptive recorder of paleogeography. For instance, Wichura et al. (2015) discussed the discovery of a ~17 Myr old beaked whale fossil 740 km inland from the present-day coastline of the Indian Ocean in the East Africa. The authors found evidence to suggest that this whale could have travelled inland from the Indian Ocean along an eastward-directed fluvial (terrestrial) drainage system and was stranded there, rather than representing a marine setting that would be implied under our assumptions. Therefore, theoretically, when using the fossil collections to improve paleogeography, additional concerns about living habits of fossils and associated geological settings should be taken into account. In this study, we have removed this misleading fossil whale from the dataset. Such instances of deceptive fossil data are a potential limitation within our workflow, which we seek to minimise by excluding inconsistent fossils more than 500 km away from previously interpreted paleoshorelines described above.

## 210 **4 Results**

### **4.1 Paleo-environmental tests**

Global reconstructed paleogeographic maps from 402 Ma to 2 Ma are tested against paleo-environments indicated by the marine fossil collections that are reconstructed in the same rotation model (Matthews et al., 2016). The consistency ratio is defined by the marine fossil collections within shallow marine or deep ocean paleogeographic polygons as a percentage of all marine fossil collections at the time interval, and in contrast, the inconsistency ratio, by the marine fossil collections not within shallow marine or deep ocean paleogeography as a percentage of all marine fossil collections. Heine et al. (2015) used a similar metric to evaluate global paleoshoreline models since the Cretaceous.

The inconsistent marine fossil collections are used to modify coastlines and paleogeographic geometries according to the rules outlined in the Methods section. The consistency ratios of marine fossil collections during 402-2 Ma are all over 55%, with an average of 75% (Fig. 6a, shaded area) although with large fluctuations over time (Fig. 6). This indicates that the paleogeography of Golonka et al. (2006) has relatively high consistency with the fossil records. However, 52 fossil collections over all time intervals cannot be resolved as they are over 500 km distant from the nearest coastline (For example, red points on Fig. 5c, l). Therefore, in some cases, the paleogeography cannot be fully reconciled with the paleobiology (see Supplement). The results since the Cretaceous are similar to that of Heine et al. (2015).

[Insert Figure 6]

The sums of marine fossil collections change significantly over time (Fig. 6b), for example, more than 4000 in total within 269-248 Ma but only 20 during 37-29 Ma. These variations are due to the spatiotemporal sampling bias and incompleteness of the fossil record (Benton et al., 2000; Benson and Upchurch, 2013; Smith et al., 2012; Valentine et al., 2006, Wright et al., 2013), biota extinction and recovery (Hallam and Wignall, 1997; Hart, 1996), the uneven entry of fossil data to the PBDB (Alroy,

2010) and our temporal selection criterion. In addition, the differences in the duration of geological time subdivisions lead to some time-intervals having shorter time spans that contain fewer fossil records, which we discuss in a later section. As for the time intervals during which fossil data are scarce, the fossil collections are of limited use in improving paleogeography. However, additional records in the future will increase the usefulness of the PBDB in such instances.

#### 4.2 Revised global reconstructed paleogeography

Based on the PBDB test results at all the time intervals, we can revise the inferred paleo-coastlines and paleogeographic geometries using the approach described in the Methods section. As a result, the revised paleo-coastlines and paleogeographies are significantly improved, mainly in the regions of North America, South America, Europe and Africa during Late Carboniferous, Middle Permian, Triassic, Jurassic, Late Cretaceous and most of Cenozoic times (Figs 4, 5, 6 and Supplement). The resulting improved global paleogeographic maps since Devonian times are presented in Figure 7. They provide improved paleo-coastlines that are important to constrain past changes in sea level and long-wavelength dynamic topography.

[Insert Figure 7]

We subsequently calculate the area covered by each paleogeographic feature as a percentage of Earth's total surface area at each time interval from 402 Ma to 2 Ma (Fig. 8), using the HEALPix pixelization method that results in equal sampling of data on a sphere (Górski et al., 2005) and therefore equal sampling of surface areas. This method effectively excludes the effect of overlaps between paleogeographic geometries.

[Insert Figure 8]

As a result, the areas of landmass, mountain and ice sheet generally indicate increasing trends, while shallow marine and deep ocean areas show decreasing trends through geological time (Fig. 8). Overall, the computed areas increase in the order of ice sheet (average 1.0% of Earth surface), mountain belts (3.4%), shallow marine (14.3%), landmass (21.3%) and deep ocean (60.1%). Only during the time interval of 323-296 Ma, landmass and shallow marine areas are nearly equal at about 14.0%, and only during 359-285 Ma, ice sheet areas exceed mountain areas but ice sheets only exist during 380-285 Ma, 81-58 Ma, and 37-2 Ma. With Pangea formation during the latest Carboniferous or the Early Permian and breakup initiation in the Early Jurassic (Blakey, 2003; Domeier et al., 2012; Lenardic, 2016; Stampfli et al., 2013; Vai, 2003; Veevers, 2004; Yeh and Shellnutt, 2016), these paleogeographic feature areas significantly change over time (Fig. 8). During 323-296 Ma (Late Carboniferous-the earliest Permian), the landmass extent reaches their smallest area (13.6%) and subsequently undergoes a rapid increase until they peak at 26.6% between 224-203 Ma (Late Triassic). In contrast, ice sheets reach their largest area (7.2%) between 323-296 Ma. In the Early Jurassic of Pangea breakup, landmass

areas rapidly decrease from 26.6% between 224-203 Ma to 23.5% between 203-179 Ma but shallow marine areas increase by 3.7%.

280

## 5 Discussions

### 5.1 Global flooded continental areas

[Insert Figure 9]

285

We estimate the global flooded continental areas since Early Devonian times from the revised paleogeography in this study (Fig. 9, pink solid line) and from the original paleogeographic maps of Golonka et al. (2006) (Fig. 9, grey solid line). Both sets of results are similar, with a decrease during Pangea amalgamation from the late Devonian Period until the Late Carboniferous Period, increase from Early Jurassic times with the breakup of Pangea until Late Cretaceous times, and then decrease again until Pleistocene times. We compare the two curves (Fig. 9, pink solid line, grey solid line) to the results of other studies (Fig. 9, Ronov, 1994; Smith et al., 1994; Walker et al., 2002; Blakey, 2003, 2008; Golonka, 2007b, 2009, 2012) derived from independent paleo-environment and paleo-lithofacies data. The results are generally consistent, except for the periods 338-269 Ma and 248-203 Ma during which the flooded continental areas for this study and Golonka et al. (2006) are smaller, which reflects smaller extent of transgression in these times. van der Meer et al. (2017, green line on Fig. 9) derived sea level and continental flooding from the strontium isotope record of marine carbonates. These results are generally consistent with the estimates from paleo-environment and paleo-lithofacies data, except during the Permian and the Late Jurassic-early Cretaceous times, during which van der Meer et al. (2017) predict larger extent of flooding than others (Fig. 9). This could indicate that the evolution of  $^{87}\text{Sr}/^{86}\text{Sr}$  reflects variations in the composition of emergent continental crust (Bataille et al., 2017; Flament et al., 2013) as well as global weathering rates (e.g. Flament et al., 2013, V erard et al., 2015, van der Meer et al., 2017).

290

295

300

305

### 5.2 Terrestrial areal change associated with transferring reconstruction, filling gaps and revising paleogeography

[Insert Figure 10]

310

We estimate the terrestrial areas, including ice sheets, mountains and landmasses, as percentages of Earth's surface area, from the original paleogeography of Golonka et al. (2006) (Fig. 10, green), from the paleogeography reconstructed using a different plate motion model of Matthews et al. (2016) and gaps filled (Fig. 10, red), and from the paleogeography with gaps fixed and revised using the paleo-environmental information indicated by marine fossil collections from the PBDB (Fig. 10, blue). These three curves are similar and generally indicate a reverse changing trend to the flooded continental areal curves over time (Fig. 9), as expected. We also calculate the areas of the terrestrial paleogeographic geometries after transferring the reconstruction but before filling gaps and the results are nearly

315

identical to the original terrestrial paleogeographic areas of Golonka et al. (2006). This is because the reconstruction of Golonka et al. (2006) has a tighter fit of the major continents within Pangea prior to the supercontinent breakup than the reconstruction of Matthews et al. (2016), so that transferring the paleogeographic geometries mainly produces gaps rather than overlaps. Comparing between the three curves (Fig. 10), filling gaps results in a larger terrestrial areal change than revising paleogeographic geometries based on PBDB test. Therefore, variation of the underlying plate reconstruction is the main factor that contributes to the terrestrial areal change (Fig. 10, red and green), and the effect of revising paleogeographic geometries based on paleobiology is secondary (Fig. 10, blue).

### 5.3 Marine fossil collection abundances in two different time scales

[Insert Figure 11]

We test the marine fossil collection dataset used in this study for fossil abundances over time with two different time scales: ICS2016 and Golonka (2000) (Table 1). The results indicate the abundances of the dataset in the two time scales are significantly different in most time intervals (Fig. 11). Generally, shorter time spans generally contain fewer data, for instance, there are about 400 marine fossil collections between 224-203 Ma using the Golonka (2000) time scale (Fig. 11, red) while there are over 1,300 collections during 232-200 Ma using the ICS2016 time scale (Fig. 11, blue). In addition, the difference of the start age and end age of the time interval could remarkably affect the fossil abundance, so that there are over 2000 marine fossil collections between 387.7-365.6 Ma in ICS2016 but less than 300 collections between 380-359 Ma using the Golonka (2000) time scale. As a result, the time scale applied to the paleobiology could significantly affect the fossil collection abundance being assigned to paleogeographic time intervals.

### 5.4 Limitations of the workflow

The workflow we develop in this study illustrates transferring paleogeographic geometries from one plate motion model to another and then using paleo-environmental information indicated by marine fossil collections from the PBDB to improve the paleo-coastline locations and paleogeographic geometries. However, the methodology still has some limitations. Transferring paleogeographic geometries to a different reconstruction inevitably results in gaps and/or overlaps, which can only be addressed using presently laborious methods. In addition, revising the coastlines and paleogeographic geometries based on the PBDB test is also currently achieved manually, and could be automated in the future.

Paleogeographic maps such as those considered here typically represent discrete time periods of many millions of years, whereas global plate motion models, even though also based on tectonic stages, provide a somewhat more continuous description of evolving plate configurations. A remaining question is how to provide a continuous representation of paleogeographic change that combines

continuous plate motion models with paleogeographic maps that do not explicitly capture changes at the same temporal resolution. In addition, it is currently difficult to apply a time scale to the raw paleobiology data from the PBDB that is currently not tied to any time scale. The paleo-environmental data used here have variable temporal resolutions, but the paleo-coastlines representing maximum transgressions are presented in a location at specific times. However, due to the inaccessibility of the original data that were used to build the paleogeographic maps, we are not in a position to estimate the temporal resolution of the original coastlines and paleogeographic maps.

The PBDB is a widely used resource (e.g., Wright et al., 2013; Finnegan et al., 2015; Heim et al., 2015; Mannion et al., 2015; Nicolson et al., 2015; Fischer et al., 2016; Tennant et al., 2016; Close et al., 2017; Zaffos et al., 2017), yet, the spatial coverage of data is still highly heterogeneous, with relatively few data points across large areas of the globe for some time periods. Hence, it is important to combine with other geological data, such as stratigraphic data from StratDB Database (<http://sil.usask.ca>) and Macrostrat Database (<https://macrostrat.org/>) and other sources of paleo-environment and paleo-lithofacies data, to further constrain the paleogeographic reconstructions.

## 6 Conclusions

Our study highlights the flexibility of digital paleogeographic models linked to a plate tectonic reconstructions in order to better understand the interplay of continental growth and eustasy, with wider implications for understanding Earth's paleotopography, ocean circulation, and the role of mantle convection in shaping long-wavelength topography. We present a workflow that enables the construction of paleogeographic maps with variable spatial and temporal resolutions, while also becoming more testable and expandable with the incorporation of new paleo-environmental datasets.

We develop an approach to revise the paleo-coastline locations and paleogeographic geometries using paleo-environmental information indicated by the marine fossil collections from the PBDB. Using this approach, the consistency ratio between the paleogeography and the paleobiology records since the Devonian is increased from an average 75% to nearly full consistency. The paleogeography in the main regions of North America, South America, Europe and Africa is significantly improved, especially in the Late Carboniferous, Middle Permian, Triassic, Jurassic, Late Cretaceous and most portions of the Cenozoic. The flooded continental areas since the late Devonian inferred from the revised global paleogeography in this study are generally consistent with the results derived from other paleo-environment and paleo-lithofacies data or from the strontium isotope record in marine carbonates.

Comparing the terrestrial areal change over time associated with transferring the reconstruction and filling gaps, and revising paleogeographic geometries using the paleo-environmental data from the PBDB, indicates that reconstruction difference is a main factor to result in the paleogeographic areal change comparing with the original maps, and revising paleogeographic geometries based on PBDB test is secondary.

## Supplementary data

400 We provide two sets of digital global paleogeographic maps during 402-2 Ma: the paleogeography  
reconstructed using the plate motion model of Matthews et al., (2016) and revised using paleo-  
environmental information indicated by the marine fossil collections from the PBDB and the original  
paleogeography of Golonka et al. (2006), an original rotation file of Golonka et al. (2006), a set of  
paleogeographic maps illustrating the PBDB test and revision of paleo-coastlines and paleogeographic  
405 geometries, a set of GeoTiff files of all revised paleogeographic maps, paleobiology data in shapefile  
used in this study separated into two sets of consistent marine fossil collections and inconsistent marine  
fossil collections, an animation for the revised global paleogeographic maps, and a README file  
outlined the workflow of this study. All supplementary material can be downloaded from the link  
410 ([https://www.dropbox.com/s/pwlygntz5mgria5/bg-2017-94-  
supplement.zip?dl=0](https://www.dropbox.com/s/pwlygntz5mgria5/bg-2017-94-supplement.zip?dl=0)~~[https://www.dropbox.com/s/91qhwdvmlbevmhp/bg-2017-94-  
supplement.zip?dl=0](https://www.dropbox.com/s/91qhwdvmlbevmhp/bg-2017-94-<br/>supplement.zip?dl=0)~~).

## Acknowledgements

This work was supported by Australian Research Council grants ARC grants IH130200012 (RDM, SZ),  
DE160101020 (NF) and SIEF RP 04-174 (SW). We thank Julia Sheehan and Logan Yeo for digitizing  
415 these paleogeographic maps, and John Cannon and Michael Chin for help with GPlates and pyGPlates.  
We sincerely ~~appreciate~~thank Christopher Scotese, Shanan Peters and ~~an~~two anonymous reviewers  
for their constructive reviews and suggestions. We thank Natascha Töpfer for editing the manuscript.  
We also thank John Alroy et al.'s original vision and all PBDB data contributors. This is Paleobiology  
Database Publication 296.

## 420 References

- Amante, C., Eakins, B., and Boulder, C.: ETOPO1 1 arc-minute global relief model: Procedures, data  
sources and analysis, NOAA Technical Memorandum, 2009.
- 425 Alroy, J.: Geographical, environmental and intrinsic biotic controls on Phanerozoic marine  
diversification. *Palaeontology* 53: 1211–1235, 2010.
- Bataille, C. P., Willis, Amy., Yang, X., and Liu, X. M.: Continental igneous rock composition: A major  
control of past global chemical weathering, *Science Advances*, 3, 1–16, 2017.
- Benson, R. B. J. and Upchurch, P.: Diversity trends in the establishment of terrestrial vertebrate eco-  
systems: interactions between spatial and temporal sampling biases, *Geology*, 41, 43–46, 2013.
- 430 Benton, M. J., Wills, M. A., and Hitchin, R.: Quality of the fossil record through time, *Nature*, 403,  
534–537, 2000.
- Blakey, R.: Carboniferous Permian global paleogeography of the assembly of Pangaea. In: Symposium  
on Global Correlations and Their Implications for the Assembly of Pangea, Utrecht (August 10-

- 16, 2003), International Congress on Carboniferous and Permian Stratigraphy, p. 57. International  
 435 Commission on Stratigraphy, 2003.
- Blakey, R. C.: Gondwana paleogeography from assembly to breakup—a 500 m.y. odyssey. *Geol. Soc. Am. Spec. Pap.* 441:1–28. [http://dx.doi.org/10.1130/2008.2441\(01\)](http://dx.doi.org/10.1130/2008.2441(01)), 2008.
- Close, R. A., Benson, R. B. J., Upchurch, P., and Butler, R. J.: Controlling for the species-area effect supports constrained long-term Mesozoic terrestrial vertebrate diversification. *Nature*  
 440 *Communications*, 8:15381. DOI: 10.1038/ncomms15381, 2017.
- Domeier, M. and Torsvik, T. H.: Plate tectonics in the late Paleozoic. *Geosci. Front.* 5, 303–350, 2014.
- Domeier, M., Van der Voo, R., and Torsvik, T. H.: Paleomagnetism and Pangea: the road to reconciliation, *Tectonophysics*, 514–517, pp. 14–43, 2012.
- Finnegan, S., Anderson, S. C., Harnik, P. G., Simpson, C., Byrnes, J. E., Tittensor, D. P., Finkel, Z. V.,  
 445 Lindberg, D. R., Liow, L. H., Lockwood, R., Lotze, H. K., McClain, C. R., McGuire, J. L., O’Dea, A., and Pandolfi, J. M.: Paleontological baselines for evaluating extinction risk in the modern oceans, *Science* 348:567–570. DOI: 10.1126/science.aaa6635, 2015.
- Fischer, V., Bardet, N., Benson, R. B. J., Arkhangelsky, M. S., and Friedman, M.: Extinction of fish-shaped marine reptiles associated with reduced evolutionary rates and global environmental  
 450 volatility. *Nature Communications*, 7:10825. DOI: 10.1038/ncomms10825, 2016.
- Flament, N., Coltice, N., and Rey, P. F.: The evolution of the  $^{87}\text{Sr}/^{86}\text{Sr}$  of marine carbonates does not constrain continental growth, *Precambrian Research*, 229, 177–188, 2013.
- Goddéris, Y., Donnadiou, Y., Le, Hir. G., and Lefebvre, V.: The role of palaeogeography in the Phanerozoic history of atmospheric CO<sub>2</sub> and climate, *Earth-Science Reviews*, 128, 122–138, 2014.
- 455 Golonka, J.: Late Triassic and Early Jurassic palaeogeography of the world, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 244, 297–307, 2007a.
- Golonka, J.: Phanerozoic paleoenvironment and paleolithofacies maps: Mesozoic. *Geol. Akad. Gór.-Hut. Im Stanisława Staszica W Krakowie T.* 33, z. 2, 211–264, 2007b.
- Golonka, J.: Phanerozoic paleoenvironment and paleolithofacies maps: Cenozoic. *Geol. Akad. Gór.-Hut. Im Stanisława Staszica W Krakowie T.* 35, z. 4, 507–587, 2009.
- 460 Golonka, J.: *Paleozoic Paleoenvironment and Paleolithofacies Maps of Gondwana*. AGH University of Science and Technology Press, Kraków, 2012.
- Golonka, J., Krobicki, M., Pajak, J., Giang, N. V., and Zuchiewicz, W.: *Global Plate Tectonics and Paleogeography of Southeast Asia*, Faculty of Geology, Geophysics and Environmental  
 465 Protection, AGH University of Science and Technology, Arkadia, Krakow, Poland, 2006.
- Golonka, J., Ross, M.I., and Scotese, C.R.: Phanerozoic paleogeographic and paleoclimatic modeling maps, in: Embry, A.F., Beauchamp, B., Glass, D.J. (Eds.), *Pangea: Global Environment and Resources*, *Memoir-Canadian Society of Petroleum Geologists*, vol. 17, pp. 1–47, 1994.
- Górski, K. M., Hivon, E., Banday, A. J., Wandelt, B. D., Hansen, F. K., Reinecke, M., and Bartelmann, M.: HEALPix: A Framework for High-Resolution Discretization and Fast Analysis of Data  
 470 Distributed on the Sphere, *The Astrophysical Journal*, 622: 759–771, 2005.
- Gurnis, M., Müller R. D., and Moresi, L.: Dynamics of Cretaceous to the present vertical motion of Australia and the Origin of the Australian- Antarctic Discordance, *Science*, 279, 1499–1504, 1998.

- 475 Hallam, A. and Wignall, P.B.: Mass extinctions and their aftermath. Oxford: Oxford University Press, 320p, 1997.
- Hart, M. B.: Biotic recovery from mass extinction events. Geological Society of London Special Publication 102, 1996.
- Heim, N. A., Knope, M. L., Schaal, E. K., Wang, S. C., and Payne, J. L.: Cope's Rule in the evolution of marine animals, *Science*, 347, 867–870. DOI: 10.1126/science.1260065, 2015.
- 480 Heine, C., Yeo, L. G., and Müller, R. D.: Evaluating global paleoshoreline models for the Cretaceous and Cenozoic. *Aust. J. Earth Sci.*, 62, 275–287, 2015.
- Kiessling, W., Flügel, E., and Golonka, J.: Patterns of Phanerozoic carbonate platform sedimentation, *Lethaia*, 36: 195–226, 2003.
- Lenardic, A.: Plate tectonics: A supercontinental boost, *Nature Geoscience*, doi:10.1038/ngeo2862, 485 2016.
- Tennant, J. P., Mannion, P. D., Upchurch, P.: Sea level regulated tetrapod diversity dynamics through the Jurassic/Cretaceous interval, *Nature Communications*, 7:12737. DOI: 10.1038/ncomms12737, 2016.
- 490 Matthews, K. J., Maloney, K. T., Zahirovic, S., Williams, S. E., Seton, M. and Müller, R. D.: Global plate boundary evolution and kinematics since the late Paleozoic. *Global and Planetary Change*, 146, 226–250, 2016.
- Müller, R. D., Seton, M., Zahirovic, S., Williams, S. E., Matthews, K. J., Wright, N. M., Shephard, G. E., Maloney, K. T., Barnett-Moore, N., Hosseinpour, M., Dan, J. B., and John, C.: Ocean basin evolution and global-scale reorganization events since Pangea breakup, *Annual Review of Earth and Planetary Science Letters*, 44, 107-138, 2016.
- 495 Nicholson, D. B., Holroyd, P. A., Benson, R. B. J., and Barrett, P. M.: Climate mediated diversification of turtles in the Cretaceous, *Nature Communications*, 6:7848. DOI: 10.1038/ncomms8848, 2015.
- Ronov, A. B.: Phanerozoic transgressions and regressions on the continents; a quantitative approach based on areas flooded by the sea and areas of marine and continental deposition. *Am. J. Sci.* 500 294:777–801, 1994.
- Ronov, A., Khain, V., and Balukhovskiy, A.: Atlas of Lithological-Paleogeographical Maps of the World, Mesozoic and Cenozoic of Continents and Oceans, U.S.S.R. Academy of Sciences, Leningrad, 79 pp, 1989.
- Ronov, A., Khain, V., and Sestlavinsky, K.: Atlas of Lithological-Paleogeographical Maps of the World, 505 Late Precambrian and Paleozoic of Continents, U.S.S.R. Academy of Sciences, Leningrad, 70 pp, 1984.
- Salles, T., Flament, N., and Müller, D.: Influence of mantle flow on the drainage of eastern Australia since the Jurassic Period, *Geochem. Geophys. Geosyst.*, 18, doi:10.1002/2016GC006617, 2017.
- Scotese, C. R.: Paleogeographic Atlas. PALEOMAP project, Arlington, TX, 1997.
- 510 Scotese, C. R.: Atlas of Earth History, Volume 1, Paleogeography, PALEOMAP Project, Arlington, Texas, 52 pp. [https://www.researchgate.net/publication/264741875\\_Atlas\\_of\\_Earth\\_History](https://www.researchgate.net/publication/264741875_Atlas_of_Earth_History), 2001.

- Scotese, C.: A continental drift flipbook, *The Journal of Geology*, 112, 729–741, doi: 10.1086/424867, 2004.
- 515 Sloss, L.: Tectonic evolution of the craton in Phanerozoic time, *The Geology of North America*, 2, 25–51, 1988.
- Smith, A. B., Lloyd, G. T., and McGowan, A. J.: Phanerozoic marine diversity: rock record modelling provides an independent test of large-scale trends, *Proceedings of the Royal Society B: Biological Sciences* 279, 4489–4495, 2012.
- 520 Smith, A. G., Smith, D. G., and Funnell, B. M.: *Atlas of Mesozoic and Cenozoic Coastlines*, Cambridge University Press, Cambridge, 99 pp, 1994.
- Spasojevic, S. and Gurnis, M.: Sea level and vertical motion of continents from dynamic Earth models since the Late Cretaceous, *American Association of Petroleum Geologists Bulletin*, 96, 2037–2064, doi:10.1306/03261211121, 2012.
- 525 Stampfli, G. M., Hochard, C., Vérard, C., Wilhem, C. and vonRaumer, J.: The formation of Pangea. *Tectonophysics*, 593, 1–19, 2013.
- Tennant, J. P., Mannion, P. D., Upchurch, P.: Sea level regulated tetrapod diversity dynamics through the Jurassic/Cretaceous interval. *Nature Communications*, 7:12737. DOI: 10.1038/ncomms12737, 2016.
- 530 Vai, G. B.: Development of the palaeogeography of Pangaea from Late Carboniferous to Early Permian, *Paleogeography, Palaeoclimatology, Palaeoecology*, 196, 125–155, 2003.
- Valentine, J. W., Jablonski, D., Kidwell, S., and Roy, K.: Assessing the fidelity of the fossil record by using marine bivalves, *Proceedings of the National Academy of Sciences*, 103, 6599–6604, 2006.
- van der Meer, D. G., van den Berg van Saparoea, A. P. H., van Hinsbergen, D. J. J., van de Weg, R. M. B., Godderis, Y., Le Hir, G., and Donnadieu, Y.: Reconstructing first-order changes in sea level during the Phanerozoic and Neoproterozoic using strontium isotopes, *Gondwana Research*, 44, 22–34, 2017.
- 535 Veevers, J. J.: Gondwanaland from 650–500 Ma assembly through 320 Ma merger in Pangea to 185–100 Ma breakup: supercontinental tectonics via stratigraphy and radiometric dating, *Earth-Science Reviews*, 68, 1–132, 2004.
- 540 Vérard, C., Hochard, C., Baumgartner, P. O., and Stampfli, G. M.: 3D palaeogeographic reconstructions of the Phanerozoic versus sea-level and Sr-ratio variations, *Journal of Palaeogeography*. 4(2), 167–188, 2015.
- Walker, L. J., Wilkinson, B. H., and Ivany, L. C.: Continental drift and Phanerozoic carbonate accumulation in shallow-shelf and deep-marine settings. *J. Geol.* 110:75–87. <http://dx.doi.org/10.1086/324318>, 2002.
- 545 Wichura, H., Jacobs, L. L., Lin, A., Polcyn, M. J., Manthi, F. K., Winkler, D. A., Strecker, M. R., and Clemens, M.: A 17-My-old whale constrains onset of uplift and climate change in east Africa, *Proc. Natl. Acad. Sci. U.S.A.*, 112(13), 3910–3915, 2015.
- 550 Wright, N., Zahirovic, S., Müller, R. D., and Seton, M.: Towards community-driven paleogeographic reconstructions: integrating open-access paleogeographic and paleobiology data with plate tectonics, *Biogeosciences*, 10, 1529–1541, doi:10.5194/bg-10-1529-2013, 2013.

Yeh, M. W. and Shellnutt, J. G.: The initial break-up of Pangæa elicited by Late Palæozoic deglaciation, *Scientific Reports*, 6: 31442, doi: 10.1038/srep31442, 2016.


555 Zaffos, A., Finnegan, S., and Peters, S. E.: Plate tectonic regulation of global marine animal diversity, *Proceedings of the National Academy of Sciences USA*. DOI: 10.1073/pnas.1702297114, 2017.

560

**Table 1. Time scale since Early Devonian times (Golonka, 2000) used in Golonka et al. (2006)'s paleogeographic maps, the original time scale of Sloss (1988), and 2016 time scale of the International Commission on Stratigraphy (ICS2016). Ages in italics are obtained by linear interpolation between subdivisions.**

565


Era	Sloss (1988)			Golonka (2000)				ICS2016		
	Subsequence	Start (Ma)	End (Ma)	Time Slice	Epoch/Age	Start (Ma)	End (Ma)	Reconstruction Time (Ma)	Start (Ma)	End (Ma)
Cenozoic	Tejas III	29	0	Late Tejas III	Tortonian – Gelasian	11	2	6	11.63	1.80
				Late Tejas II	Burdigigalian – Serravallian	20	11	14	20.44	11.63
				Late Tejas I	Chattian – Aquitanian	29	20	22	28.1	20.44
	Tejas II	39	29	Early Tejas III	Priabonian – Rupelian	37	29	33	37.8	28.1
	Tejas I	60	39	Early Tejas II	Lutetian – Bartonian	49	37	45	47.8	37.8
				Early Tejas I	Thanetian – Ypresian	58	49	53	59.2	47.8
	Mesozoic	Zuni III	96	60	Late Zuni IV	middle Campanian – Selandian (Late Cretaceous – earliest Paleogene)	81	58	76	79.8
Late Zuni III					late Cenomanian – early Campanian (Late Cretaceous)	94	81	90	96.1	79.8
Zuni II		134	96	Late Zuni II	late Aptian – middle Cenomanian (Early Cretaceous – earliest Late Cretaceous)	117	94	105	119.0	96.1
				Late Zuni I	late Valanginian – early Aptian (Early Cretaceous)	135	117	126	136.4	119.0
Zuni I		186	134	Early Zuni III	late Tithonian – early Valanginian (latest Late Jurassic – earliest Early Cretaceous)	146	135	140	147.4	136.4
				Early Zuni II	late Bathonian – middle Tithonian (earliest Middle Jurassic – Late Jurassic)	166	146	152	166.8	147.4
				Early Zuni I	middle Aalenian – middle Bathonian (Middle Jurassic)	179	166	169	172.8	166.8
Absaroka III		245	186	Late Absaroka III	late Hettangian – early Aalenian (Early Jurassic – earliest Middle Jurassic)	203	179	195	200.0	172.8
				Late Absaroka II	late Carnian – middle Hettangian (Late Triassic – earliest Jurassic)	224	203	218	232	200.0
				Late Absaroka I	Induan – early Carnian (Early – earliest Late Triassic)	248	224	232	252.17	232
Paleozoic	Absaroka II	268	245	Early Absaroka IV	Roadian – Changhsingian (Late Permian)	269	248	255	272.3	252.17
				Early Absaroka III	Sakmarian – Kungurian (Early Permian)	285	269	277	295.0	272.3
				Early Absaroka II	Gzhelian – Asselian (latest Carboniferous – earliest Permian)	296	285	287	303.7	295.0
	Absaroka I	330	268	Early Absaroka I	Bashkirian – Kasimovian (Late Carboniferous)	323	296	302	323.2	303.7
				Kaskaskia IV	middle Visean – Serpukhovian (Lower Carboniferous)	338	323	328	341.4	323.2
	Kaskaskia II	362	330	Kaskaskia III	late Fammenian – early Visean (latest Devonian – Early Carboniferous)	359	338	348	365.6	341.4
				Kaskaskia I	Givetian – early Fammenian (Middle – Late Devonian)	380	359	368	387.7	365.6
	Kaskaskia I	401	362	Kaskaskia I	late Pragjan – Eifelian (Early – Middle Devonian)	402	380	396	408.7	387.7



570


**Fig. 1. Global distributions and number of fossil collections since the Devonian Period. The greyscale background shows global present-day topography ETOPO1 (Amante and Eakins, 2009) with lighter shades corresponding to increasing elevation. Fossil collections from the PBDB are colored following the standard used by the International Commission on Stratigraphy.**

575



**Fig. 2. Workflow used to transfer a set of paleogeographic geometries from one reconstruction to another, followed by revision using paleo-environmental information indicated by marine fossil collections from the Paleobiology Database (PBDB).**

580



585 **Fig. 3. (a) Original global paleogeographic map from Golonka et al. (2006) at 126 Ma. (b) Global paleogeographic geometries at 126 Ma in present-day coordinates. (c) Global paleogeography at 126 Ma reconstructed using the plate motion model of Matthews et al. (2016). Gaps are highlighted in pink. (d) Global paleogeography at 126 Ma reconstructed using the reconstruction of Matthews et al. (2016) with gaps fixed by filling with adjacent paleo-environment attributes. Grey lines indicate reconstructed present-day coastlines and terrane boundaries. Mollweide projection with 0°E central meridian.**


590

**Table 2. Lookup table to classify fossil data indicating different paleo-environments into marine or terrestrial settings and their corresponding paleogeographic types presented in Golonka et al. (2006).**

595 **Terrestrial fossil paleo-environments correspond to paleogeographic features of landmasses, mountains or ice sheets, and marine fossil paleo-environments to shallow marine environments or deep oceans.**

Marine		Terrestrial/Transitional Zone	
Paleogeography	Fossil Paleo-environments	Paleogeography	Fossil Paleo-environments
Shallow marine environments/Deep oceans	marine indet.	slope	terrestrial indet.
	carbonate indet.	basinal (carbonate)	fluvial indet.
	peritidal	basinal (siliceous)	alluvial fan
	shallow subtidal indet.	marginal marine indet.	channel lag
	open shallow subtidal	coastal indet.	coarse channel fill
	lagoonal/restricted shallow subtidal	estuary/bay	fine channel fill
	sand shoal	lagoonal	channel
	reef, buildup or bioherm	paralic indet.	wet floodplain
	perireef or subreef	interdistributary bay	dry floodplain
	intrashelf/intraplatform reef	delta front	floodplain
	platform/shelf-margin reef	prodelta	crevasse splay
	slope/ramp reef	deltaic indet.	levee
	basin reef	foreshore	mire/swamp
	deep subtidal ramp	shoreface	fluvial-lacustrine indet.
	deep subtidal shelf	transition zone/lower shoreface	delta plain
	deep subtidal indet.	offshore	fluvial-deltaic indet.
offshore ramp	submarine fan	lacustrine - large	
offshore shelf	basinal (siliciclastic)	lacustrine - small	
offshore indet.	deep-water indet.	glacial	
		Ice sheets	
		Landmasses/Mountains	pond
			crater lake
			lacustrine delta plain
			lacustrine interdistributary bay
			lacustrine delta front
			lacustrine prodelta
			lacustrine deltaic indet.
			lacustrine indet.
			dune
			interdune
			loess
			eolian indet.
			cave
			fissure fill
			sinkhole
			karst indet.
			tar
			spring

600



605

**Fig. 4. (a) Test between the global paleogeography at 76 Ma reconstructed using the plate motion model of Matthews et al. (2016) with gaps fixed and the paleo-environments indicated by the marine fossil collections from the PBDB. (b) Area modified (blue) to resolve the test inconsistencies. (c) Test between the revised**

paleogeography at 76 Ma and the same marine fossil collections. Mollweide projection with 0°E central meridian.

610




Fig. 5. Test between unrevised and revised paleogeography at 76 Ma respectively and paleo-environments indicated by the marine fossil collections from the PBDB, and revision of the paleo-coastlines and paleogeographic geometries based on the test results, for southern North America (a, b, c), southern South America (d, e, f), northern Africa (g, h, i) and India (j, k, l). Regional Mollweide projection.

615






Fig. 6. (a) Consistency ratios between global paleogeography with gap filled, but before PBDB test for the period 402-2 Ma, reconstructed using the plate motion model of Matthews et al. (2016) and the paleoenvironments indicated by the marine fossil collections from the PBDB. (b) Numbers of consistent (light grey) and inconsistent (dark grey) marine fossil collections used in the tests for each time interval from 402 Ma and 2 Ma.

620

625






**Fig. 7. Global paleogeography from 402 Ma to 2 Ma reconstructed using the plate motion model of Matthews et al. (2016) and revised using paleo-environmental data from the PBDB. Black toothed lines indicate subduction zones, and other black lines denote mid-ocean ridges and transforms. Grey outlines delineate reconstructed present-day coastlines and terranes. Mollweide projection with 0°E central meridian.**


630

635




**Fig. 8. Global paleogeographic feature areas as percentages of Earth's total surface area estimated from the revised paleogeographic maps from 402 Ma to 2 Ma.**

640



**Fig. 9. Global flooded continental area since the Early Devonian Period from the original paleogeographic maps of Golonka et al. (2006) (grey solid line) and from the revised paleogeography in this study (pink line). Results for Blakey (2003, 2008), Golonka (2007b, 2009, 2012), Ronov (1994), Smith et al. (2004), Walker et al. (2002) are as in van der Meer et al. (2017). The van der Meer et al. (2017) curve (green line) is derived from the strontium isotope record of marine carbonates.**


645



650

**Fig. 10. Terrestrial areal change due to filling gaps and modifying the paleo-coastlines and paleogeographic geometries over time. Green: based on the original paleogeographic maps of Golonka et al. (2006); Red: based on paleogeography reconstructed using a different plate motion model of Matthews et al. (2016) and gaps filled; Blue: based on paleogeography with gaps fixed and revised using the paleo-environments indicated by marine fossil collections from the PBDB.**

655



660

**Fig. 11. Fossil abundance test on the marine fossil collection dataset used in this study with two different time scales: Golonka (2000) and ICS2016 (Table 1).**