

# Point-by-point Reply to all Referees

November 1, 2018

Biogeosciences Discussion Manuscript bg-2018-379 (submitted 14 Aug 2018):  
**Iron fertilization efficiency and the number of past and future regenerations of iron in the ocean**

Authors:

**Benoît Pasquier** and **Mark Holzer**

Interactive discussion available at:

<https://www.biogeosciences-discuss.net/bg-2018-379/>

Dear Dr. Middelburg,

Please find the comments from the three referees and our point-by-point replies from the interactive discussion reproduced below. Along with our replies, we have added to this letter in green the corresponding revisions that were made to the manuscript.

The manuscript version with tracked changes, produced with the `latexdiff` package, is appended and shows the deleted parts struck through in red and the revised parts in blue.

In addition, we have re-arranged the title of our manuscript to read

**The number of past and future regenerations of iron in the ocean and its intrinsic fertilization efficiency**

so that it better reflects the paper's content and emphasis.

Sincerely,

Benoît Pasquier and Mark Holzer

## Reply to Anonymous Referee #1

**Referee #1 General Comments:** This is a review of “Iron fertilization efficiency and the number of past and future regenerations of iron in the ocean” by B. Pasquier and M. Holzer. The authors present a novel technique to track the life-cycle of dissolved iron (DFe) in the ocean. The authors apply the technique to multiple instances of data-constrained representations of the iron cycle and investigate how many cycles DFe parcels experience before and after participating in the biological pump. The authors use their technique to quantify the efficiency of iron fertilization on biological export, one of the motivations for their study. The manuscript is fairly well written, laying out clearly their novel technique, and they present well-designed experiments to utilize their technique. I think this manuscript will be a useful addition to the literature. I am making suggestions for some minor changes and/or additions.

**Authors’ response:** We thank Referee #1 for these general comments.

*No changes to the manuscript in response to these general comments.*

No changes.

**Referee #1 Minor Point 1:** The abstract doesn’t mention that the model used by the authors includes multiple types of external sources of DFe. I think it would be useful to mention in the abstract the types of external sources of DFe under consideration.

**Authors’ response:** We agree that it would be good to mention the modelled external iron sources in the abstract.

*In response, we will revise the abstract to explicitly state that aeolian, sedimentary, and hydrothermal iron sources are modelled.*

The abstract now starts with:

Iron fertilization is explored by tracking dissolved iron (DFe) through its life cycle from injection by aeolian, sedimentary, and hydrothermal sources (birth) to burial in the sediments (death).

**Referee #1 Minor Point 2:** The notation for the nonlinear model in Section 2.1 deviates from the notation in the author’s previous work (Pasquier & Holzer, Biogeosciences, 2017) (e.g., removal of the Redfield ratio for the uptake terms, and changed notation for the particle transport terms). I suggest that the authors either use the same notation as their previous work, or describe how and why the notation in the current work differs from the previous work.

**Authors’ response:** Referee #1 is correct that we changed notation from what we used to describe our fully coupled Fe–P–Si model [Pasquier and Holzer, 2017]. Here, we use simplified notation for extra clarity and readability, exploiting the fact that not all the complexity of the fully coupled model is needed to develop our new iron-cycle diagnostics.

*In response, we will add a brief statement to Section 2.1 which points this out and makes the connection with the corresponding symbols in Pasquier and Holzer, [2017].*

We have added the corresponding symbols of Pasquier and Holzer [2017] in the new Glossary (Appendix A, Table A1), which is referred to in an added parenthetical

sentence at the end of the paragraph describing Equation (1):

(Note that we have simplified the notation of the fully coupled nonlinear model of Pasquier and Holzer (2017) for clarity and readability — see Appendix Table A1 for the symbol correspondence.)

**Referee #1 Minor Point 3:** There is no motivation given for the definition of the equivalent linear model in Section 2.2. I think the paper would benefit from having a paragraph describing what the goals/requirements of the equivalent linear model are, and how goals/requirements lead to the model that the authors are using.

**Authors' response:** The opening paragraph of Section 2.3 (“In order to track iron from its birth at the source to its eventual death. . .”) was meant to motivate the need for the equivalent linear model, but we agree with Referee #1 that we do not explicitly point out why the nonlinear model itself cannot be used for tracing *partitions* of DFe or, equivalently, iron *labels*. The iron labels, unlike iron itself, are passive tracers and obey a linear equation of motion. Mathematically, only these linear tracers allow for correct partitioning of the DFe distribution because the superposition principle only applies to linear systems. This is often overlooked in the bigeoscience literature where studies typically evaluate the contribution of a given process by computing the anomaly that results from the removal of the process (e.g., removing a source). Such an anomaly approach results in errors that scale with the degree of nonlinearity of the system. For example, *Holzer et al.* [GBC, 2016] showed that omitting an iron source to evaluate its contribution to the DFe distribution underestimates the true contribution (evaluated using the equivalent linear model) by a factor of  $\sim 2$ . For the iron cycle, the nonlinearities of the uptake and scavenging processes are the reason why an equivalent linear model is required.

*In response, we will add a paragraph to the beginning of Section 2.3 that includes the arguments above to explain the necessity of the equivalent nonlinear model.*

We have revised the paragraph starting Section 2.3 to now read:

In order to track iron from its birth at the source to its eventual death (via the irreversible part of scavenging), we consider a labelling tracer that we can think of as being attached to the nonlinearly evolving DFe. This labelling tracer has the same concentration as DFe but satisfies a linear evolution equation in which the nonlinear uptake and scavenging are replaced by linear processes. These linear processes are diagnosed from the nonlinear steady-state solution of Eq. (1) to provide identical uptake and scavenging rates and hence identical tracer solutions. It is necessary to employ such iron labelling tracers because their linear equations satisfy the superposition principle, which allows us to rigorously partition the iron concentration according to source type, number of regenerations, and so on. (The underlying parent model cannot directly be used for this purpose because of its nonlinearities.)

**Referee #1 Minor Point 4:** While some symbols chosen for the various terms in the nonlinear and equivalent linear models do seem related to the processes being represented by the terms (e.g.,  $U$  for uptake,  $\mathcal{R}$  for regeneration,  $\mathcal{D}$  for death), not all of the connections are clear (e.g.,  $J$  for scavenging,  $\mathcal{L}$  for uptake

## Reply to Anonymous Referee #1

that gets exported). This makes it hard for the reader to keep track of which terms mean what. I suggest adding a table that describes, in terms of processes, what each symbol denotes.

**Authors' response:** We strongly agree with Referee #1 that good notation is important, and when possible we do use symbols whose meaning is self-evident. However, the desire for simple notation needs to be balanced with precision and clarity. For example, we need to use different symbols for linear and nonlinear operators, and we do not want to use an "s"-based symbol for both particle transport ("sinking") and scavenging. We think the notation in our manuscript is a reasonable compromise. We agree that a table describing our symbols could be helpful to the reader.

*In response, we will add a short glossary of symbols as suggested by Referee #1. We have added a glossary of symbols in new Appendix A.*

**Referee #1 Minor Point 5:** Have you considered how a particular instance of the nonlinear model would respond to a substantial change to aeolian input, such as would happen in the LGM or a future climate change scenario. Does the technique presented shed light on how the nonlinear model would respond to this change in forcing? This could be mentioned in the Discussion section.

**Authors' response:** Thank you for the suggestion. We think that exploring the response of the iron cycle to changes in iron input is a rich subject deserving a separate study (and beyond the scope of the current manuscript). The diagnostics developed here probe the iron cycle without perturbing it and are therefore by themselves insufficient to infer the response to source changes. To examine the steady-state response to source changes, e.g., aeolian input for an LGM or future climate scenario, one would first have to use the fully coupled Fe—P—Si model of *Pasquier and Holzer* [2017] to solve for the perturbed steady-state nutrient cycles. Once these have been calculated, our diagnostics can readily be applied to the perturbed iron cycle to elucidate, for example, what the mean number of iron passages through the biological pump was during the LGM.

*In response, we will add some discussion to the Discussion and Caveats section to explain that while we diagnosed the unperturbed iron cycle, one can also apply the diagnostics to perturbed states resulting from added iron, e.g., to shed light on paleo or future climate scenarios.*

We have added the following sentence to the last paragraph of the Conclusions section:

We also note that the approach of using a linear equivalent linear model to partition iron and diagnose its life cycle can also be applied to perturbed states (e.g., due to iron addition or changes in circulation) to shed light on how the iron cycle operates for various paleo or future climate scenarios. Finally, the concepts and methods employed here can be applied to other nutrients for a more complete picture of how the interaction between the biological pump and the physical transport shapes their distributions and cycling rates; we plan to do so in future work.

## Reply to Dr. Lauderdale (Referee #2)

**Dr. Lauderdale’s (Referee #2) General Comments:** In this manuscript, Pasquier and Holzer present a series of diagnostics to document the “life cycle” of dissolved iron in the ocean. Depending on the total source of iron (from an ensemble of nearly 300 solutions that “equally well” resemble oceanic nutrient distributions) they find that the majority of iron molecules are scavenged permanently from the ocean before they have had a chance to be biologically utilized. Of those that are taken up by phytoplankton, the majority will only have one circuit of the “ferrous wheel” before they too are permanently buried in the sediments. I thought this was a really interesting paper, that certainly fits the criteria for publication in Biogeosciences. I would like to suggest a few points that the authors might consider.

**Authors’ response:** We thank Dr. Lauderdale (Referee #2) for these positive remarks.

*No changes to the manuscript in response to the General Comments.*

No changes.

**Referee #2 Minor Point 1:** Although fairly well written overall, in places I found the manuscript overly technical. For example, on page 9: lines 14–19 where there are 4 equivalences in as many lines, and only the last one (or two) are relevant. Perhaps there is a way to simplify? Furthermore, I appreciated where the authors had split their prose to identify the “physical” cause or effect and then the “mathematical” proof (page 7: line 20–22). Can this clarity be afforded elsewhere in the manuscript?

**Authors’ response:** We think the equations in this instance are useful and should not be avoided because they define and clarify convenient notation (the  $\chi$ -weighted global averages). This notation is used later on multiple occasions in both the manuscript and figures and enhances subsequent readability. However, we do agree with Referee #2 that our manuscript need not be overly technical.

*In response, we will rephrase the second last sentence of this passage using more succinct expressions to ensure its point is clearly made with a minimum of symbols: “Note that this fraction can be considered to be the  $\chi_k$ -weighted global average of the local unused fraction  $f_k^{0\downarrow} \equiv \chi_k^{0\downarrow}/\chi_k$ . This weighted average is defined as  $\langle f_k^{0\downarrow} \rangle_{\chi_k} = \langle f_k^{0\downarrow} \chi_k \rangle / \langle \chi_k \rangle$ , where we introduced the  $\langle \cdot \rangle_{\chi_k}$  notation, which will be used throughout.”*

*We will check the revised manuscript everywhere for clarity and employ the “split-prose” approach where appropriate.*

We have revised the passage, which now reads:

To quantify the amount of iron that was not regenerated in the past, we now ask how the unused fraction of the global DFe inventory varies with total iron source strength,  $\sigma_{\text{tot}}$ . Note that this fraction can be considered to be the  $\chi_k$ -weighted global average of the local unused fraction  $f_k^{0\downarrow} \equiv \chi_k^{0\downarrow}/\chi_k$ . This weighted average is defined as  $\langle f_k^{0\downarrow} \rangle_{\chi_k} = \langle f_k^{0\downarrow} \chi_k \rangle / \langle \chi_k \rangle$ , where we introduced the  $\langle \cdot \rangle_{\chi_k}$  notation, which will be used throughout. The unused fractional DFe inventory regardless of source type is given by  $\langle f^{0\downarrow} \rangle_{\chi}$ , where  $f^{0\downarrow} \equiv$

$$\sum_k \chi_k^{0\downarrow} / \chi.$$

**Referee #2 Minor Point 2:** Adding to the slightly overwhelming number of symbols used in this manuscript, I did come across something that looks like a mistake, or maybe requires clarification: Figure 1 suggests that  $\mathcal{D}$  is the reversible scavenging process — after the first regeneration, iron is transported to the near-surface where it is scavenged onto a sinking particle and released at depth to be then transported into the euphotic layer and biologically utilized. Similarly,  $\tilde{\mathcal{D}}$  is used for future reversible scavenging. However, section 2.3 defines  $\mathcal{D}$  as “iron scavenging minus redissolution of scavenged iron” and the “permanent loss of iron due to burial in the sediments” (page 4: lines 6–7), which appears to correspond to  $d$  in the schematic.

**Authors’ response:** We did try to minimize the number of symbols and always introduce new notation only to make the manuscript clearer or to provide precision where we think it is important. However, we agree with Referee #2 that we should have been clearer when introducing the operator  $\mathcal{D}$ . (The details of  $\mathcal{D}$  are provided in Appendix A.)  $\mathcal{D}$  is the linear integral operator that, applied to the DFe concentration field  $\chi$ , gives the local rate of scavenging minus the local rate with which scavenged iron is redissolved. The net local rate  $(-\mathcal{D}\chi)(\mathbf{r})$  in Equation (2) can thus be locally positive (net DFe re-dissolution) but its vertical integral is always negative (a finite fraction is scavenged out of the system). Hence,  $\mathcal{D}$  provides both the scavenging-pump transport for DFe (conservative “reversible scavenging”) as well as the permanent DFe sink (non-conservative “death”). The field  $d$  is the local iron death rate, which is calculated from  $\mathcal{D}\chi$  as detailed in Appendix A and introduced later in Section 4.1.  $d(\mathbf{r})$  is the rate at which DFe is removed at  $\mathbf{r}$  by scavenging *and* instant sediment burial. Thus,  $d$  (unlike  $\mathcal{D}$ ) does not capture the recycling of scavenged DFe.

*In response, we will revise the text to clarify the action of  $\mathcal{D}$  where it is introduced. For clarity and simplicity, we will also revise Figure 1 slightly by removing the adjoint operator symbols (they come too early for this introductory schematic) and to use the same type of  $\mathcal{D}$ -labelled arrow for both mid-stream reversible scavenging and for permanent burial, as  $\mathcal{D}$  accomplishes both. We will also label the uptake process in the euphotic zone with  $\mathcal{L}$ , so that all physical processes are labelled on the figure.* We have reworded the description of  $\mathcal{D}$  in the manuscript after Equation (2) to read:

..., and  $\mathcal{D} \equiv \sum_j (1 - \mathcal{S}_j) \gamma_j$  is the reversible scavenging operator. More precisely,  $\mathcal{D}$  is the linear integral operator that, applied to the DFe concentration field  $\chi$ , gives the local rate of scavenging minus the local rate with which scavenged iron is redissolved. Thus,  $\mathcal{D}$  provides both the transport of the “scavenging pump” (conservative “reversible” scavenging) as well as the permanent iron sink due to burial in the sediments (non-conservative “death”).

We have accordingly revised Figure 1 and its caption.

**Referee #2 Minor Point 3:** The phrase “Southern Ocean nutrient trapping”

is frequently used, and I wondered if the authors could check that all uses are appropriate. For example, page 7: lines 17–19, I think the authors have the correct explanation that hydrothermal iron is added to density classes that upwell in the Southern Ocean, but is this really “nutrient trapping” and not just transport?

**Authors’ response:** We think that we use “Southern Ocean nutrient trapping” correctly here because we are considering hydrothermal DFe that has already been regenerated once ( $n = 1$ ). We agree with Referee #2 that hydrothermal DFe is first “just transported” to the Southern Ocean surface where the density layers of the hydrothermal vents outcrop, but subsequently part of this hydrothermal DFe is utilized and trapped in the Southern Ocean (the trapping mechanism is described in the cited references). The plots of Figure 2 for hydrothermal DFe (column 3) show that hydrothermal DFe with  $n \geq 1$  is found in the Southern Ocean with the characteristic pattern of Southern Ocean nutrient trapping.

*In response, we will double check the revised manuscript to ensure that all other occurrences of “Southern Ocean nutrient trapping” are appropriate.*  
No changes.

**Referee #2 Minor Point 4:** Another paper that considered the iron fertilization efficiency was *Dutkiewicz et al.* (2006; GRL; doi:10.1029/2005GL024987). Using an adjoint of the MITgcm biogeochemistry model, they found a similar pattern of tropical-Pacific-dominated primary production and carbon uptake when iron is added to the ocean.

**Authors’ response:** We thank Referee #2 for reminding us about this paper. We agree that we should have referenced it and that it will provide additional interesting context for our study. It is important, however, to appreciate that the experiments described by *Dutkiewicz et al.* (2006) are finite-amplitude perturbations, while our study quantifies the “natural” fertilization efficiency of the *unperturbed* iron cycle.

*In response, we will add references to Dutkiewicz et al. (2006) where relevant.*  
We have added the following sentences at the beginning of the introduction where we cover previous work:

Dutkiewicz et al. (2006) used an adjoint technique and a model of the coupled carbon, phosphorus, and iron cycles to quantify the sensitivity of global biological production and air-sea carbon fluxes to local perturbations in the aeolian iron source. They found that both quantities were most sensitive to iron addition in the central and eastern tropical Pacific.

In the second paragraph of Section 5.1, we have added the following sentences in Section 5.1 (Export supported per unit DFe injection at  $\mathbf{r}$ ):

Furthermore, the local efficiency  $\bar{m}^P(\mathbf{r})$  is defined regardless of whether an iron source is actually present at  $\mathbf{r}$ . The efficiency  $\bar{m}^P(\mathbf{r})$  quantifies the global phosphorus export rate *per unit DFe source rate* at  $\mathbf{r}$ . At all points  $\mathbf{r}$ , even where there is no actual source in the system,  $\bar{m}^P(\mathbf{r})$  can be considered to be the “sensitivity” of the linear equivalent system to the insertion of the arbitrary test source  $s_k(\mathbf{r})$ : Equation (14) shows that  $\bar{m}^P(\mathbf{r})$  is the proportionality between the export response  $\Phi_k^P(\mathbf{r})$  and the test source  $s_k(\mathbf{r})$ .

In this sense, the fertilization efficiency  $\overline{m}^P$  is a close, but distinct, cousin of the sensitivity to small-amplitude perturbations considered by Dutkiewicz et al. (2006)

We have also added the following sentences in Section 5.1 where we discuss the geographic pattern of the natural fertilization efficiency:

Interestingly, Dutkiewicz et al. (2006) found that the sensitivity of their coupled model to aeolian iron addition is also largest in the central tropical and eastern Pacific. This suggests that the intrinsic fertilization efficiency of the unperturbed state is shaped by similar processes as the sensitivity to small-amplitude perturbations.

Finally, we have added a comparison of the magnitude of our intrinsic fertilization efficiency to the fertilization efficiency estimates of Dutkiewicz et al. (2006) at the of Section 5.1:

The sensitivity of global production to perturbations in the local aeolian source estimated by Dutkiewicz et al. (2006) has a spatial distribution with a range of about  $20\text{--}180\text{ g C m}^{-2}\text{ yr}^{-1}$  for a  $0.02\text{ mmol Fe m}^{-2}\text{ yr}^{-1}$  perturbation. The model of Dutkiewicz et al. (2006) exports 1/3 of its production as POP so that these sensitivities translate to a POP export per added DFe molecule of about  $28\text{--}250\text{ mol C (mmol Fe)}^{-1}$ , a lower bound on the full carbon export and about a third of the intrinsic fertilization efficiency estimated here. We emphasize that differences between these various estimates are not only due to uncertainties in the iron cycle (as expressed by our range of values), but also due to differences in the definition of fertilization efficiency, not to mention due to differences among models and methodologies.

**Referee #2 Minor Point 5:** Finally, I wonder if the authors could comment on the caveat that their biogeochemical model may not capture the full array of interactions that might lead to enhanced iron regenerations through grazing by zooplankton, or bacteria/virus interactions, for instance. This is in regards to the “ferrous wheel” idea where recycling of iron is considered important (e.g. *Kirchman, 1996, Nature, doi:10.1038/383303a0*; *Maldonado et al., 2005, GBC, doi:10.1029/2005GB002481*; *Strzepek et al., 2005, GBC, doi:10.1029/2005GB002490*; *Boyd et al., 2017, Nature Geoscience, doi:10.1038/ngeo2876*). Maybe these views can be reconciled, with reference to figure 3?

**Authors’ response:** We thank Referee #2 for bringing these studies to our attention. While the details of the ferrous wheel are beyond the scope of our study (as is their bearing on Figure 3), we agree that it would be appropriate to briefly reference these papers where we comment on the associated issues in relation to our model. Specifically, our simple formulation of the Fe:C uptake ratio may well be unrealistic (e.g., *Kirchman, 1996*; *Strzepek et al., 2005*), as we acknowledged in the Discussion and Caveats section where we discuss the work of *Rafter et al., (2017)*. Similarly, we acknowledged (with reference to *Twining et al., 2014*) that different remineralization lengthscales for iron and macronutrients (*Boyd et al., 2017*) are not modelled. The effect of ligands on iron bioavailability (*Maldonado et al., 2005*)

is also not represented by our model, which does not have dynamic ligands but instead prescribes a ligand distribution.

Importantly, we would like to note that not every process thought to operate on DFe needs to be explicitly modelled for a useful representation of the iron cycle. The inverse model of *Pasquier and Holzer (2017)* is of intermediate complexity, and any effect of the above issues is captured implicitly when parameters are optimized to fit the observed nutrient and phytoplankton fields. Explicit modelling of these complexities may be important for models that try to predict how the system will change in the future, but we do not think this is necessary for constraining and diagnosing the large-scale cycling of DFe in the current state of the ocean as we do here.

*In response, we will mention the effect of ligands on iron bioavailability (which was missing in the submitted version), and we will add references to the suggested papers where we discuss the associated issues in the Discussion and Caveats section.*

*We have added the following sentences on ligands in Section 6 (Discussion and caveats):*

Regarding ligands, the inverse model of Pasquier and Holzer (2017) prescribes an optimized distribution of a single type of ligand that is enhanced in hydrothermal plumes and old waters, with optimized parameters. Ligands are not dynamically transported and the effect of different ligand types on iron bioavailability (Maldonado et al., 2005) is neglected.

*We have also revised the passage on the Fe:P stoichiometry in Section 6 to include the references suggested:*

A key control on our model results is the Fe:P stoichiometry. The model approximates the iron dependence of the Fe:P uptake ratio by a Monod function with a half saturation constant that is the same for all phytoplankton classes. We acknowledge that this may not be realistic. For example, Kirchman (1996) suggested that including the microbial “ferrous wheel”, which operates with different stoichiometric ratios, could affect iron budgets in the euphotic zone. Similarly, Strzepek et al. (2005) explored variations in the iron regeneration rates of different organisms and cautioned modelers to pay careful attention to the details of the Fe:C stoichiometry. Relatedly, our model simply remineralizes iron in the same Fe:P ratio with which it was utilized, so that the vertical profiles of iron and phosphate remineralization have identical shapes. However, measurements by Twining et al. (2014) show that, at least for some phytoplankton species, iron is remineralized more slowly than phosphate, suggesting that our remineralization profile for iron could be too shallow. Different remineralization lengthscales for iron and phosphate were also emphasized by Boyd et al. (2017).

*Finally, we have added a paragraph concluding the Discussion and caveats section to briefly discuss the role of complexity in models.*

Importantly, we would like to note that not every process thought to influence DFe needs to be explicitly modelled for a useful representation of the iron cycle. The model of Pasquier and Holzer (2017) is of intermediate

Reply to Dr. Lauderdale (Referee #2)

complexity, and effects due to processes not explicitly modelled are captured implicitly when parameters are optimized to fit the observed nutrient and phytoplankton fields. Explicit modelling of all known processes in their full complexity may be important for models that try to predict how the system will change in the future, but this is neither necessary nor desirable for constraining and diagnosing the large-scale cycling of DFe in the current state of the ocean as we do here.

## Reply to Dr. Völker (Referee #3)

**Dr. Völker (Referee #3) Introductory Comment 1:** Quite seldom, when reviewing manuscripts, I encounter a study where I wished I had had the idea myself. This manuscript, of which I have already reviewed an earlier version for another journal, is one of them. It introduces several new diagnostics for the iron cycle that help to understand the fate of iron stemming from different external sources as it is advected through the ocean and every now and then gets taken up by a phytoplankton cell. Over the last decade or so it has been realized that other sources of iron besides dust (sediments, hydrothermal sources, volcanos,...) contribute substantially to the inventory of iron in the ocean, and estimates of iron's residence time had to be corrected downwards. But there remain major uncertainties on the relative magnitude of these sources, and models with very different source strengths and residence times have been equally, albeit only moderately, successful in modelling the measured iron distribution. At this state of things it is an important step forward to have tools at hand that allow to quantify the biological effect of different iron sources, even if it is only for one specific iron model and assumes steady state.

**Authors' response:** We are delighted by Dr. Völker's (Referee #3) comments.

*No changes to the manuscript in response to these Introductory Comments.*

No changes.

**Referee #3 General Comment 1:** The new diagnostics are based on a linearization of optimized steady-state solutions from a global ocean biogeochemical model to separate the effect of the different iron sources. The idea is to take the resulting iron distribution, and to linearize the nonlinear parts in the evolution equation for iron around that solution. The linearized equations are not only the basis for separating the effect of the different iron sources, but also allow, by splitting the linear operators, to separate in a second step, how often a specific concentration of iron has passed through biological uptake and remineralization, and how often it will do so in the future.

**Authors' response:** Just to clarify, we do not linearize our nonlinear nutrient model in the usual sense, but instead construct an equivalent linear model for tracer labels as explained in our response to Referee #1, Minor Point 3. There is a fundamental difference: Linearization usually refers to the first-order Taylor expansion of the nonlinear model around a suitable base state, which captures the system behaviour for small perturbations about the base state. Here, we instead construct the equations for passive tracers that follow the DFe and that have the same solution as the nonlinear equations. The resulting model is linear in the sense that the labels participate in the physical processes in proportion to the local DFe abundance; hence all nonlinear processes are replaced by the DFe-label concentration multiplied by a rate coefficient diagnosed so that the linear and nonlinear models have identical solutions. The passive labelling tracers of the linear model then allow a rigorous partitioning according to source type, number of regenerations, and so on, because the superposition principle applies to the linear equivalent model.

*In response, we will add a sentence or two to the section on the linear equivalent model that cautions the reader not to mistake the linear model as a linearization of*

*the nonlinear model.*

We have added the following short paragraph at the end of Section 2.3 (Equivalent Linear Model: Iron Labelling Tracers):

We caution the reader that the equivalent linear model (2) is not a linearization of the nonlinear parent model (1) in the usual sense. The equivalent linear model is constructed to partition DFe in the *unperturbed* system. By contrast, linearization usually refers to the first-order Taylor expansion of the nonlinear model around a base state, which captures the system behaviour for small perturbations about that base state.

**Referee #3 General Comment 2:** It is important that this exercise is not done for one specific model run, but for a whole family of model runs, differing mostly in the relative strength of iron sources, and correspondingly, in the timescale for scavenging, all reproducing iron observations about equally well (or badly). Of course this means that the linearized equations are different for every member of the ensemble, and the results obtained that way also differ to some extent. Nevertheless, the authors show quite convincingly that some results are quite robust and consistent between the different ensemble members.

**Authors' response:** Agreed.

*No change to the manuscript in response to this comment.*

No changes.

**Referee #3 Minor Point 1:** The main results obtained in that way are that for reasonable strength of external iron sources, the average number of past and future passages through biological uptake of a given iron concentration is less than one, meaning that most iron has not or will not pass through biology before getting scavenged. This number is significantly higher for dust-deposited iron, since it enters the ocean closer to the place of biological uptake. Regardless of source the pattern of iron concentration that has passed through several uptake-recycling cycles converges towards a Southern Ocean-trapped pattern. Not all results are equally important or surprising: for example the result that “total DFe is more likely to have been regenerated in the past than it is to be regenerated in the future” (page 17, line 30) is fairly obvious from the fact that remineralisation happens deeper in the water column than e.g. dust deposition.

**Authors' response:** We agree that not all our results are equally important and some may seem, after the fact, “fairly obvious”. The likelihood of passing through the biological pump depends subtly on how transport to surface uptake from either external source or from internal regeneration samples the scavenging field. Our diagnostics allow this to be quantified rigorously for our model. The results can be rationalized after the fact, but we think that both qualitatively and certainly quantitatively, the finding of more likely past than future regeneration is far from obvious before one does the calculation. For example, it is not obvious that sedimentary iron, which has deep sources, is more likely to have passed through the pump in the past than in the future.

*No change to the manuscript in response to this minor comment.*

No changes.

**Referee #3 Minor Point 2:** Some results may also have to do with the specific iron model: that iron fertilization is most efficient in the equatorial Pacific may also have to do with the particular parameterisation of a variable Fe:P quota in the model by Pasquier and Holzer, which basically follows a Michaelis-Menten-type curve, meaning that for Fe tending towards zero, the Fe:P ratio will also tend towards zero; the linearisation of the iron uptake would then ascribe a very high ratio of P to Fe export in regions with very low Fe. Most iron models produce an extremely low surface Fe concentration in the equatorial Pacific, which is far away from dust sources and where the upwelling waters are quite old, meaning they are low in Fe.

The dependency on Fe:P is mentioned on page 22, line 11, and the sensitivity of the results on this is discussed briefly in the subsequent paragraph. In the discussion section this is however, discussed maybe a bit too briefly (page 25, line 30).

**Authors' response:** We agree that the DFe dependence of the Fe:P uptake ratio is a major control on our model results. We would not characterize our discussion of the issue on page 22 as particularly brief: An entire paragraph (lines 13–21) is devoted to it. We agree, however, that we should reiterate the importance of the Fe:P ratio in the Discussion and Caveats section.

*In response, we will add a sentence or two to the Discussion and Caveats section reiterating the importance of the Fe:P ratio for shaping the pattern of the diagnosed natural iron fertilization efficiency.*

We have revised the passage in Section 6 (Discussion and caveats) to emphasize the importance of the Fe:P ratios:

A key control on our model results is the Fe:P stoichiometry. The model approximates the iron dependence of the Fe:P uptake ratio by a Monod function with a half saturation constant that is the same for all phytoplankton classes. We acknowledge that this may not be realistic. For example, Kirchman (1996) suggested that including the microbial “ferrous wheel”, which operates with different stoichiometric ratios, could affect iron budgets in the euphotic zone. Similarly, Strzepek et al. (2005) explored variations in the iron regeneration rates of different organisms and cautioned modelers to pay careful attention to the details of the Fe:C stoichiometry. Relatedly, our model simply remineralizes iron in the same Fe:P ratio with which it was utilized, so that the vertical profiles of iron and phosphate remineralization have identical shapes. However, measurements by Twining et al. (2014) show that, at least for some phytoplankton species, iron is remineralized more slowly than phosphate, suggesting that our remineralization profile for iron could be too shallow. Different remineralization lengthscales for iron and phosphate were also emphasized by Boyd et al. (2017). Because our model is optimized to fit the DFe observations, with an emphasis on deep profiles relative to surface measurements Pasquier and Holzer (2017), a potentially too shallow remineralization of iron would be compensated by an increased strength of the biological pump. Furthermore, the relative amount of scavenging by opal and POP particles is optimizable in our model, so that deeper iron remineralization can be achieved by increasing

the scavenging by opal. We acknowledge, however, that when optimizing the match to observed DFe, the model may produce biases in the relative contributions of the biological and scavenging pumps, which would affect our estimates of the number of passages through the biological pump.

**Referee #3 Minor Point 3:** In summary I think that this is quite a significant paper for understanding the iron cycle in the ocean, and it should be published in Biogeosciences after minor revision.

The paper is quite well written (although it probably appeals more to a reader with some background in linear algebra) and I have checked the mathematical argumentation in depth and it is clear and correct. The authors attempt, and usually succeed in connecting the mathematically rigorous description of their results with what these results mean in terms of biogeochemistry. Nevertheless, here and there, the authors could do a bit more to make the explanations more palatable to the readership of Biogeosciences.

One example that I have is in the beginning of the section on future passages through the biological pump, where the authors explain that “the natural way to formulate the necessary equations is to consider the time-reversed adjoint flow... The adjoints are defined for the volume-weighted inner product.” While this is probably clear to a mathematically trained physicist, it may less be so for the average reader of Biogeosciences. Maybe the authors could add a few lines here on what the adjoints are, what the inner product.

My small remarks to the earlier version of this manuscript have been taken into account already, so I stop here.

**Authors’ response:** We are delighted that Dr. Völker appreciates our work, and we certainly agree that our paper should be as accessible as possible to the Biogeosciences audience.

*In response, we will rephrase the introduction at the start of Section 4 (Future Contributions to Export) to explain and motivate why we use the adjoint, which is to track DFe backwards in time “from” death with computational efficiency.*

We have revised the passage on adjoints and the inner product at the start of Section 4 to read:

These adjoints are defined in terms of the volume weighted inner product  $\langle x, y \rangle \equiv \int x(\mathbf{r})y(\mathbf{r})d^3\mathbf{r}$ , where the integral ranges over the entire ocean volume. Thus,  $\tilde{\mathcal{H}}$ , the adjoint of  $\mathcal{H}$ , is defined as usual so that  $\langle \tilde{\mathcal{H}}x, y \rangle = \langle x, \mathcal{H}y \rangle$ . In the time-reversed adjoint flow, the death operator becomes a source of labels that we then track through sequential regenerations backward in time to the present, analogously to what we did in the previous section for the usual time-forward flow. The use of adjoint operators allows for numerically efficient evaluation of our diagnostics because a single tracer injected at death into the time-reversed adjoint flow suffices to produce the full three-dimensional concentration fields of interest at the present.

# Iron fertilization efficiency and the number of past and future regenerations of iron in the ocean and its intrinsic fertilization efficiency

Benoît Pasquier<sup>1,2</sup> and Mark Holzer<sup>1</sup>

<sup>1</sup>Department of Applied Mathematics, School of Mathematics and Statistics, University of New South Wales, Sydney, NSW, Australia.

<sup>2</sup>Now at Department of Earth System Science, University of California, Irvine, CA, USA.

**Correspondence:** Benoît Pasquier (pasquieb@uci.edu)

**Abstract.** Iron fertilization is explored by tracking dissolved iron (DFe) through its life cycle from injection by ~~external-aeo-~~ ~~lian, sedimentary, and hydrothermal~~ sources (birth) to burial in the sediments (death). We develop new diagnostic equations that count iron and phosphate regenerations with each passage through the biological pump and partition the ocean's DFe concentration according to the number of its past and future regenerations. We apply these diagnostics to a family of data-constrained estimates of the iron cycle with sources  $\sigma_{\text{tot}}$  in the range 1.9–41 Gmol yr<sup>-1</sup>. We find that for states with  $\sigma_{\text{tot}} > 7$  Gmol yr<sup>-1</sup>, 50% of the DFe inventory has not been regenerated in the past and 85% will not be regenerated in the future. The globally averaged mean number of past and future regenerations scale with the bulk iron lifetime  $\tau \sim \sigma_{\text{tot}}^{-1}$  and have ranges of 0.05–2.2 and 0.01–1.4, respectively. Memory of birth location fades rapidly with each regeneration, and DFe regenerated more than  $\sim 5$  times is found in a pattern shaped by Southern Ocean nutrient trapping. We quantify the ~~natural fertilization efficiency-intrinsic~~ ~~fertilization efficiency of the unperturbed system~~ at any point  $r$  in the ocean as the global export production resulting from the DFe at  $r$ , per iron molecule. We show that this efficiency is closely related to the mean number of future regenerations that the iron will experience. At the surface, the ~~natural-intrinsic~~ fertilization efficiency has a global mean in the range 0.7–7 mol P (mmol Fe)<sup>-1</sup> across our family of state estimates and is largest in the central tropical Pacific, with the Southern Ocean having comparable importance only for high iron-source scenarios.

## 15 1 Introduction

Iron is an essential micronutrient for phytoplankton photosynthesis (e.g., Raven, 1988). In the ocean, dissolved iron (DFe) is a trace element that has been shown to limit biological production over vast high-nutrient-low-chlorophyll (HNLC) regions (e.g., Boyd et al., 2007; Aumont and Bopp, 2006). In HNLC regions, ample phosphate and nitrate are available but phytoplankton are not able to completely utilize these macronutrients because of a lack of sufficient DFe. Iron thus exerts a major influence over the marine carbon cycle and hence over the global climate system.

Many studies have been devoted to quantifying the extent to which biological productivity and ocean carbon uptake can be influenced by altering the iron supply through geoengineering (e.g., Aumont and Bopp, 2006; Boyd et al., 2007). Relatedly, it

is natural to ask how efficient iron is in supporting the export of organic matter in the current state of the system and what the relative efficiencies of the different iron sources are (e.g., aeolian versus sedimentary). Zeebe and Archer (2005) suggested that artificial iron fertilization would not sufficiently impact atmospheric CO<sub>2</sub> to mitigate anthropogenic global warming. Zeebe and Archer (2005)'s argument relied, among other things, on quantifying the efficiency of [artificial](#) iron fertilization, which was  
5 inferred from the Southern Ocean Iron Experiment (SOFEX). Similarly, de Baar et al. (2008) used data from localized artificial iron fertilization experiments to explore the per-iron-molecule efficiency of supporting carbon export, defined by the change in local dissolved inorganic carbon per unit added DFe. From this, de Baar et al. (2008) estimated the “natural” fertilization efficiency (i.e., the efficiency of [the unperturbed system perturbing DFe only](#)) by correcting for the lack of ligand protection in artificial iron fertilization. [Dutkiewicz et al. \(2006\) used an adjoint technique and a model of the coupled carbon, phosphorus,  
10 and iron cycles to quantify the sensitivity of global biological production and air-sea carbon fluxes to local perturbations in the aeolian iron source. They found that both quantities were most sensitive to iron addition in the central and eastern tropical Pacific.](#)

Here we develop new diagnostics for following an iron molecule's passages through the biological pump from its “birth” by source injection to its “death” by scavenging and burial in the sediments. We show that the number of times that an  
15 iron molecule is biologically utilized and regenerated at depth during its birth-to-death journey is a fundamental metric for understanding the marine biosphere's iron fertilization efficiency. Using Green-function techniques, we track DFe through past regenerations back to its birth and through future regenerations forward to its death. We apply our new diagnostics to a family of data-constrained state estimates of the iron cycle and quantify the [natural-intrinsic](#) iron fertilization efficiency — i.e., [the fertilization efficiency of DFe molecules in the unperturbed system](#) — in three dimensions throughout the global ocean. To the  
20 best of our knowledge, this has never done before. Previous studies, like those of Zeebe and Archer (2005) and de Baar et al. (2008), estimated the fertilization efficiency only for the specific surface regions where iron was artificially injected into the euphotic zone.

Our analysis uses the 287 optimized state estimates of Pasquier and Holzer (2017), which were obtained from a steady-state inverse model of the ocean's coupled iron, phosphorus, and silicon cycles, embedded in the data-assimilated steady global  
25 ocean circulation of Primeau et al. (2013). The members of our family of state estimates correspond to different iron source strengths, given that the actual values for the real ocean are still highly uncertain (e.g., Tagliabue et al., 2016). All these optimized states fit the available observed DFe and nutrient concentrations about equally well, with a total iron source that ranges from 1.9 to 41 Gmol Fe yr<sup>-1</sup> across our family. (Atmospheric models estimate an aeolian source of soluble iron to the ocean of ~6 Gmol Fe yr<sup>-1</sup> (Luo et al., 2008).) Having states for a wide range of iron source scenarios enables us to quantify  
30 systematic variations with source strength.

While recent studies of the iron cycle have begun to quantify the distribution and pathways of regenerated iron (Tagliabue et al., 2014; Qin et al., 2016; Achterberg et al., 2018), to the best of our knowledge the biological cycling of iron during its full birth-to-death lifetime has previously not been quantified. In particular, the future passages of a given DFe molecule through the biological pump have not been systematically considered, thus missing a potentially important part of its contribution to  
35 supporting carbon export. In fact, de Baar et al. (2008) dismissed the importance of iron regenerated at depth for subsequently

fertilizing production altogether, based on the assumption that all DFe that is regenerated at depth would be lost to scavenging. However, scavenging is weaker at depth so that deep regenerated DFe molecules are not necessarily scavenged and instead may be transported back to the surface where they may support production multiple times before being scavenged out of the system.

5 Here, for the first time, the entire birth-to-death journey of DFe is considered. Specifically, we aim to address the following questions:

1. What fraction of the DFe distribution in the current state of the ocean has passed  $n$  times through the biological pump in the past, and what fraction will pass  $m$  times through the biological pump in the future, for any given  $n$  and  $m$ ?
2. How much does the DFe present at any given location in the current state of the ocean contribute to the global export  
10 production per DFe molecule?
3. How do the mean number of past and future passages through the biological pump, and the closely related iron fertilization efficiency, depend on the uncertain iron source strengths?

We find that for states with global iron sources above  $7 \text{ Gmol Fe yr}^{-1}$ , most of the DFe gets scavenged out of the system without participating in the biological pump. Future passages through the biological pump are less likely than past passages,  
15 but for sources around  $7 \text{ Gmol Fe yr}^{-1}$ , roughly 10% of the iron inventory will still participate in future biological production before death. DFe that has been regenerated more than about 5 times in the past can be found in a characteristic pattern (mathematically an eigenmode) that bears a strong signature of Southern Ocean nutrient trapping. DFe that will be regenerated many times in the future is similarly found in a Southern-Ocean-intensified eigenmode.

We link the mean number of future regenerations to the ~~natural~~-intrinsic iron fertilization efficiency, which we are able to  
20 quantify at any point in the ocean. At the surface, which is most relevant to geoengineering, we find that this fertilization efficiency is largest in the central tropical Pacific, with the Southern Ocean having comparable efficiency only for states with a high total iron source.

In Section 2, we briefly detail the salient features of the iron model and introduce our diagnostic framework. The DFe distribution is partitioned according to the number of past and future passages through the biological pump in Sections 3 and  
25 4, respectively. In Section 5, we develop the connection with the iron fertilization efficiency. Caveats of our approach are discussed in Section 6, and we present conclusions in Section 7.

## 2 Iron model

### 2.1 Nonlinear model

Following Pasquier and Holzer (2017), we write the nonlinear tracer equation for DFe concentration  $\chi$  as

$$(\partial_t + \mathcal{T})\chi = \sum_c (\mathcal{S}_c - 1)U_c + \sum_j (\mathcal{S}_j - 1)J_j + \sum_k s_k \quad , \quad (1)$$

**Table 1.** Iron source strengths in units of  $\text{Gmol Fe yr}^{-1}$  for the typical state and four other states sampling our family of solutions. Source strengths are listed for aeolian ( $\sigma_A$ ), sedimentary ( $\sigma_S$ ), hydrothermal ( $\sigma_H$ ), and total ( $\sigma_{\text{tot}}$ ) iron. The corresponding bulk iron lifetime,  $\tau$ , in units of years, is also tabulated. ( $\tau$  is the ratio of the global DFe inventory to  $\sigma_{\text{tot}}$ .)

Source scenario	Acronym	$\sigma_A$	$\sigma_S$	$\sigma_H$	$\sigma_{\text{tot}}$	$\tau$
Low-hydrothermal	LoH	5.3	1.7	0.15	7.2	113
High-sedimentary-low-aeolian	HiS-LoA	1.8	6.2	0.87	8.9	96
Typical	TYP	5.4	1.7	0.88	8.0	104
High-aeolian-low-sedimentary	HiA-LoS	15.	0.45	0.88	17.	45
High-hydrothermal	HiH	6.3	2.0	2.3	11.	80

where  $\mathcal{T}$  is the advective-diffusive transport operator of the data-assimilated steady ocean circulation of Primeau et al. (2013).  $U_c$  is the iron uptake rate of phytoplankton functional class  $c$ ,  $J_j$  is the iron scavenging rate for particle type  $j$ , and  $s_k$  is the source of dissolved iron of type  $k$ , with  $k = A, S, \text{ or } H$ , for aeolian, sedimentary, and hydrothermal iron, respectively. The subscript  $c$  ranges over small, large, and diatom phytoplankton functional classes, and the subscript  $j$  ranges over particulate  
5 organic phosphorus (POP), biogenic silica (**B**Si**b**Si), and dust particle types. (Note that we have simplified the notation for the fully coupled nonlinear model of Pasquier and Holzer (2017) for clarity and readability — see Appendix Table A1 for the symbol correspondence.)

The uptake  $U_c$  depends nonlinearly on the concentration of DFe, phosphate, and silicic acid and hence couples the iron, phosphorus, and silicon cycles. The remineralization rate of DFe taken up by phytoplankton class  $c$  is modelled by  $\mathcal{S}_c U_c$ ,  
10 where the linear “source” operator  $\mathcal{S}_c$  accomplishes both the in situ remineralization in the euphotic zone and the biogenic transport followed by remineralization at depth. (Remineralization at depth is modelled as instantaneous with the divergence of a Martin power-law POP flux profile (Martin et al., 1987).) Similarly, the redissolution rate of scavenged DFe is modelled by  $\mathcal{S}_j J_j$  where the source operator  $\mathcal{S}_j$  represents the instantaneous transport to depth of iron scavenged by particles of type  $j$ . The detailed model formulation of the coupled iron–phosphorus–silicon cycles, including the construction of the operators  $\mathcal{S}_c$   
15 and  $\mathcal{S}_j$ , has been published by Pasquier and Holzer (2017).

## 2.2 Family of optimal state estimates

Pasquier and Holzer (2017) coupled the iron cycle to the global phosphorus and silicon cycles and determined the steady state of the system using an efficient Newton solver. The biogeochemical model parameters were systematically optimized by minimizing the quadratic mismatch between modelled and observed nutrient and phytoplankton concentrations. This approach  
20 led to a family of 287 possible state estimates, which correspond to widely different iron-source scenarios and a range in total iron source strength of 1.9 to 41  $\text{Gmol Fe yr}^{-1}$ . (The true magnitude of the ocean’s iron sources is still highly uncertain (e.g., Tagliabue et al., 2016).) All state estimates have very similar fidelity to the observational constraints.

By performing our analyses for a family of states spanning a wide range of possible iron sources, we are able to establish features that are insensitive to the iron-source scenario, while also being able to quantify systematic variations of key metrics with the uncertain iron source strengths. Below we will show spatial patterns for a typical state estimate and for four other states that are representative of the variations across our family of estimates. The source strengths and bulk iron lifetimes of these five representative states are collected in Table 1. When emphasizing systematic variations of a specific metric with iron source strength, we will plot the metric for all 287 states.

### 2.3 Equivalent linear model: iron labelling tracers


In order to track iron from its birth at the source to its eventual death (via the irreversible part of scavenging), we consider a tracer  $\chi$  that labelling tracer that we can think of as being attached to the nonlinearly evolving DFe. This labelling tracer has the same concentration as DFe, but for but satisfies a linear evolution equation in which the nonlinear uptake and scavenging are replaced by linear processes. These linear processes are diagnosed from the nonlinear steady-state solution of Eq. (1) to provide identical uptake and scavenging rates and hence identical tracer solutions. Specifically, it is necessary to employ such iron labelling tracers because their linear equations satisfy the superposition principle, which allows us to rigorously partition the iron concentration according to source type, number of regenerations, and so on. (The underlying parent model cannot directly be used for this purpose because of its nonlinearities.)

To construct the linear processes for the iron labels, we diagnose the local rate constants  $\gamma_c(\mathbf{r}) = U_c(\mathbf{r})/\chi(\mathbf{r})$  and  $\gamma_j(\mathbf{r}) = J_j(\mathbf{r})/\chi(\mathbf{r})$ , so that  $U_c$  and  $J_j$  can be replaced by  $\gamma_c\chi$  and  $\gamma_j\chi$ , which are linear in  $\chi$ . Following Pasquier and Holzer (2017), we write  $\mathcal{S}_c = (1 - f_c) + \mathcal{B}f_c$  to separate the remineralization rate into the fraction  $(1 - f_c)$  that is remineralized in situ in the euphotic zone and the detrital fraction  $f_c$  that is exported to depth by the biogenic transport and remineralization operator  $\mathcal{B}$ . Substituting the linear forms of  $U_c$  and  $J_j$  into Eq. (1) and reorganizing terms, we obtain

$$(\partial_t + \mathcal{T})\chi = \mathcal{R}\chi - \mathcal{L}\chi - \mathcal{D}\chi + \sum_k s_k, \quad (2)$$

where  $\mathcal{L} \equiv \sum_c f_c \gamma_c$  is the uptake operator for DFe that gets exported,  $\mathcal{R} \equiv \mathcal{B}\mathcal{L}$  is the regeneration operator, and  $\mathcal{D} \equiv \sum_j (1 - \mathcal{S}_j)\gamma_j$  is the reversible scavenging operator. The operator  $\mathcal{D}$  represents iron scavenging minus redissolution of scavenged iron. More precisely,  $\mathcal{D}$  is the linear integral operator that, applied to the DFe concentration field  $\chi$ , gives the local rate of scavenging minus the local rate with which scavenged iron is redissolved. Thus,  $\mathcal{D}$  provides the death process for DFe, which is the net permanent loss both the transport of the “scavenging pump” (conservative “reversible” scavenging) as well as the permanent iron sink due to burial in the sediments (non-conservative “death”). The operator  $\mathcal{B}$  represents conservative biogenic particle transport and subsequent regeneration. In Eq. (2), DFe enters the biological pump with the utilization rate  $\mathcal{L}\chi$  and exits the biological pump with the regeneration rate  $\mathcal{R}\chi$ . Throughout, “regeneration” refers to remineralization in the aphotic zone following a passage through the biological pump. (We define the regeneration operator  $\mathcal{R}$  such that in the euphotic zone  $\mathcal{R}\chi = 0$  so that the biological pump’s intake ( $\mathcal{L}\chi$ ) and output ( $\mathcal{R}\chi$ ) are cleanly separated.)

The source  $s_k$  in Eq. (2) may be thought of as injecting tracer labels attached to DFe. These labels are eventually completely removed by  $\mathcal{D}$ . The uptake  $\mathcal{L}$  also removes labels from iron in the euphotic zone, but the regeneration  $\mathcal{R} = \mathcal{B}\mathcal{L}$  re-injects these



**Figure 1.** Schematic of the birth-to-death ~~life cycle~~ life cycle of a labeled DFe molecule (red dot). As captured by Eq. (2), the molecule makes several passages through the biological pump (green, uptake  $\mathcal{L}$  and regeneration  $\mathcal{R}$ ), through the scavenging pump (purple dotted,  $\mathcal{D}$ ) while being transported by the ocean circulation (blue,  $\mathcal{T}$ ) from its birth (red,  $s_k$ , here the aeolian source) to its death by eventual burial in sediments (red,  $\mathcal{D}$ ) following scavenging. This particular DFe molecule passed  $n=2$  times through the biological pump since its birth, and will pass  $m=1$  times through the biological pump until its death. We consider the DFe concentration at location  $\mathbf{r}$  and at present time  $t$ . Our diagnostics partition DFe at  $(\mathbf{r}, t)$  into the fractions that have undergone  $n$  regenerations since birth and that will undergo  $m$  regenerations until death. Computationally, we track DFe from its birth to the present using the usual time-forward flow, while we track DFe backward in time from its death to the present using the time-reversed adjoint operators ( $\tilde{\mathcal{T}}$ ,  $\tilde{\mathcal{R}}$ , and so on) flow.

labels throughout the water column below without any losses ( $\mathcal{B}$  is conservative). A schematic of the transport and cycling of iron labels by these operators is provided in Figure 1. For convenience, we bring all the operators to the left-hand-side of the equation so that (2) can be written compactly as  $(\partial_t + \mathcal{H})\chi = \sum_k s_k$ , where the complete linear system operator is given by  $\mathcal{H} \equiv \mathcal{T} - \mathcal{R} + \mathcal{L} + \mathcal{D}$ . Below it will be useful to consider the iron cycle without regeneration, whose evolution is governed by

5  $\mathcal{F} \equiv \mathcal{T} + \mathcal{L} + \mathcal{D}$ . (Note that  $\mathcal{F} = \mathcal{H} + \mathcal{R}$ . The definitions and units of the linear operators are collected in Appendix Table A1.)

~~We emphasize that the linear equivalent model (2) has solutions that are identical to the solutions of nonlinear model (1). The linear model (2) is not a linearization of the full nonlinear model but, instead, tracks passive labelling tracers that faithfully follow the DFe through the nonlinear biological and scavenging processes.~~

~~The linearity of~~ Because the superposition principle applies to the labelling tracers ~~allows us to~~, we can consider the DFe concentrations to be the sum of the concentrations of different iron “source types”, that is  $\chi = \sum_k \chi_k$ , where  $\chi_k$  is the concentration due to source  $s_k$ . In steady state, we can therefore calculate the concentration  $\chi_k$  of each source type by solving (Holzer et al., 2016)

$$\mathcal{H}\chi_k = s_k \quad . \quad (3)$$

5 We caution the reader that the equivalent linear model (2) is not a linearization of the nonlinear parent model (1) in the usual sense. The equivalent linear model is constructed to partition DFe in the *unperturbed* system. By contrast, linearization usually refers to the first-order Taylor expansion of the nonlinear model around a base state, which captures the system behaviour for small perturbations about that base state.

### 3 Past contributions to export

10 To establish how much a given iron molecule has contributed to organic-matter export since its birth, we partition the DFe concentration according to the number of its past passages through the biological pump.

#### 3.1 Iron concentration regenerated $n$ times since birth

We first consider the concentration of DFe that has never been regenerated since it was injected (born) by the source. This concentration, denoted by  $\chi_k^{0\downarrow}$ , can simply be computed from our equivalent linear system by injecting iron labels with source  $s_k$ , but not permitting them to be regenerated. We therefore remove the  $\mathcal{R}$  term from Eq. (3), which is equivalent to replacing  $\mathcal{H}$  with  $\mathcal{F}$ , to obtain

$$\mathcal{F}\chi_k^{0\downarrow} = s_k \quad . \quad (4)$$


We use arrow superscripts to indicate past processes that occurred since injection *into* ( $\downarrow$ ) the ocean, or future processes that will occur until removal *out of* ( $\uparrow$ ) the ocean.

We can now calculate the concentration of DFe from source  $k$  that has been regenerated exactly one time,  $\chi_k^{1\downarrow}$ , as follows. The source of labels for  $\chi_k^{1\downarrow}$  is the rate of first regeneration of DFe, which is given by  $\mathcal{R}\chi_k^{0\downarrow}$ . We simply allow the system to cycle these labels but remove them on uptake using  $\mathcal{F}$  (no second regeneration). Thus, in steady state,  $\chi_k^{1\downarrow}$  obeys  $\mathcal{F}\chi_k^{1\downarrow} = \mathcal{R}\chi_k^{0\downarrow}$ . Similarly, the source of DFe that has been regenerated exactly  $n + 1$  times since birth is the rate of  $(n + 1)$ st regeneration. This gives the recursion relation

$$\mathcal{F}\chi_k^{(n+1)\downarrow} = \mathcal{R}\chi_k^{n\downarrow} \quad , \quad (5)$$

with (4) providing the starting point of the recursion. Note that the  $\chi_k^{n\downarrow}$  partition the DFe concentration exactly, with  $\sum_{n=0}^{\infty} \chi_k^{n\downarrow} = \chi_k$ , as shown in Appendix B.

Figure 2 shows global zonal averages of the iron concentration for our typical state, partitioned by source type (aeolian, sedimentary, hydrothermal) and according to the number  $n$  of past regenerations since birth. To emphasize the spatial patterns,



**Figure 2.** Global zonal averages of the concentration of DFe that was regenerated  $n$  times since birth, normalized by its global mean, for  $n=0,1,2,3$ , and 10. The normalized concentrations of the three source types (aeolian, sedimentary, hydrothermal) are shown as well as their sum, the total concentration regardless of source type. This figure is for our typical state.


each  $\chi_k^{n\downarrow}(\mathbf{r})$  field has been normalized by its global volume-weighted average,  $\langle \chi_k^{n\downarrow} \rangle$ . Iron that has never been regenerated ( $n=0$ ) carries a strong source signature, with peak concentrations where the ~~largest sources are sources~~ are largest. The patterns change dramatically with one passage through the biological pump ( $n=1$ ). Because iron is regenerated at depth with a Martin power-law profile, the ~~peak zonally-averaged aeolian DFe concentration~~ concentration of aeolian DFe that has been

regenerated  $n = 1$  times has a subsurface maximum at roughly 1500 m depth. Similarly, sedimentary iron that passed once through the biological pump has a mid-depth maximum and is no longer concentrated near the ~~sea floor~~seafloor. Sedimentary iron that has been regenerated  $n = 1$  times is concentrated ~~into~~in the Arctic, presumably because the model has already shallow sedimentary sources there. However, the model’s Arctic circulation is poorly constrained (Primeau et al., 2013) and this particular feature may not be robust. Hydrothermal iron that has been regenerated  $n = 1$  times shows the signature of Southern Ocean nutrient trapping (Primeau et al., 2013; Holzer et al., 2014) as expected from the fact that hydrothermal iron is injected at seawater densities that outcrop in the Southern Ocean.

As the number  $n$  of past regenerations increases,  $\chi_k^{n\downarrow}$  rapidly converges to a pattern that is independent of source type. Physically, this is because the memory of birth place quickly fades with each passage through the biological pump. Mathematically, this can be seen from recursion relation (5), which gives  $\chi_k^{n\downarrow} = \mathcal{A}^n \chi_k^{0\downarrow}$ , where  $\mathcal{A} \equiv \mathcal{F}^{-1}\mathcal{R}$ . Thus, for sufficiently large  $n$ , the concentration  $\chi_k^{n\downarrow}$  becomes proportional to the eigenmode of  $\mathcal{A}$  with the largest eigenvalue  $\lambda$ , and the amplitude of the pattern decays exponentially like  $\lambda^n = \exp(-n/n^*)$ , where  $n^* \equiv -1/\log(\lambda)$ . (Note that  $\lambda < 1$ , as must be the case for  $\sum_{n=0}^{\infty} \chi_k^{n\downarrow}$  to converge to  $\chi_k$ .) In other words, as  $\mathcal{A}$  is applied repeatedly, only the projection of  $\chi_k^{0\downarrow}$  ~~on~~onto the eigenmode of  $\mathcal{A}$  with the gravest eigenvalue survives. Because  $\mathcal{A}$  is independent of source type, all source types approach the same large- $n$  asymptotic eigen pattern. Convergence to this pattern is remarkably rapid: For aeolian and sedimentary iron, the pattern has emerged after about 5 regenerations (not shown), and for hydrothermal iron even sooner. By  $n = 10$  the asymptotic pattern is well established and indistinguishable for the different iron source types (bottom plots of Figure 2).

How robust is the large- $n$  eigen pattern of  $\chi_k^{n\downarrow}$  across our family of solutions? To quantify the approach to the exponentially decaying eigenmode, we plot in Figure 3a the global mean concentration that has been regenerated  $n$  times since birth as a function of  $n$  for the 5 representative state estimates of Table 1. Figure 3a shows that the  $e$ -folding scale  $n^*$  is independent of source type as expected (same large- $n$  slope for all source types on the semilog plots). For all 5 states the convergence to exponential decay is quickest for hydrothermal iron. The value of the  $e$ -folding scale  $n^*$  depends on the state of the iron cycle. For our 5 representative states, the smallest  $n^*$  values occur for the high-source states ( $n^* \sim 0.6$  for HiA-LoS and 0.8 for HiH), while the largest  $e$ -folding scales occur for the low-source states (1.4 for both HiS-LoA and LoH), with  $n^* \sim 1.2$  for our typical state. Across the entire family of state estimates,  $n^*$  ranges from about 0.4 to 2.8. The dependence of  $n^*$  on source scenario makes sense when we consider that all these states are optimized against the observed DFe concentrations: High aeolian input must be countered by vigorous removal from the surface ocean, and hence a large proportion of the iron in the ocean is accounted for by the first few regenerations since birth, leaving little DFe for multiple regenerations. Conversely, low aeolian input corresponds to less vigorous biological cycling so that a larger portion of the iron is available to pass through the biological pump repeatedly. Similar consideration hold for hydrothermal and sedimentary iron. For all states, asymptotic behavior is well established for  $n \gtrsim 5$ .


The sedimentary and hydrothermal iron concentrations get their largest contributions from iron that has never been regenerated ( $n = 0$ ), again pointing to significant permanent removal of these iron types before they can reach the euphotic zone following injection (birth) at depth. For aeolian iron in the typical, LoH, and HiS-LoA states, the largest contribution comes from iron that has been regenerated  $n = 1$  times as all freshly born aeolian iron is immediately available for uptake and regen-



**Figure 3.** (a) Global mean DFe concentration regenerated  $n$  times in the past,  $\langle \chi_k^{n\downarrow} \rangle$ , as a function of  $n$ , for five representative states of the iron cycle. (b) Mean concentration regenerated  $m$  times in the future,  $\langle \chi_k^{m\uparrow} \rangle$ , as a function of  $m$ .

eration. However, for the high-source states (HiA-LoS and HiH), the largest contribution still comes from iron that was never regenerated because the increased scavenging necessary to balance the high sources prevents most aeolian DFe from being utilized even though it is deposited directly into the euphotic zone.

Percentage of DFe inventory not regenerated



**Figure 4.** (a) Percentage of the total inventory of DFe that has not passed through the biological pump in the past (i.e., since birth), as a function of total source strength,  $\sigma_{\text{tot}}$ , for the family of state estimates of Pasquier and Holzer (2017). (b) Percentage of the total inventory of DFe that will not pass through the biological pump in the future (i.e., until death). Note the logarithmic abscissa. Shown are the fractional inventories of the individual source types of DFe (color coded), and well as the total fractional inventory regardless of source type (black).

Figure 3a shows that DFe that has not passed through the biological pump since birth ( $n = 0$ , unused iron) generally has the highest global mean concentration, except for aeolian DFe under some source scenarios. To quantify the amount of iron that was not regenerated in the past, we now ask how the unused fraction  $f_{\text{d}^3\text{r}\chi_k}^{0\downarrow}(\mathbf{r})/f_{\text{d}^3\text{r}\chi_k}(\mathbf{r})$  of the global DFe inventory varies with total iron source strength,  $\sigma_{\text{tot}}$ . ~~This fraction may be written as  $\langle f_k^{0\downarrow} \rangle_{\chi_k} = f_{\text{d}^3\text{r}f_k^{0\downarrow}(\mathbf{r})\chi_k(\mathbf{r})/f_{\text{d}^3\text{r}\chi_k(\mathbf{r})}$~~

5 ~~and therefore~~ Note that this fraction can be considered to be the  $\chi_k$ -weighted global average of the local unused fraction  $f_k^{0\downarrow}(\mathbf{r}) \equiv \chi_k^{0\downarrow}(\mathbf{r})/\chi_k(\mathbf{r})$ . This weighted average is defined as  $\langle f_k^{0\downarrow} \rangle_{\chi_k} = \langle f_k^{0\downarrow} \chi_k \rangle / \langle \chi_k \rangle$ , where we introduced the  $\langle \cdot \rangle_{\chi_k}$  notation, which will be used throughout. The unused fractional DFe inventory regardless of source type is given by  $\langle f^{0\downarrow} \rangle_{\chi}$ , where  $f^{0\downarrow}(\mathbf{r}) \equiv \sum_k \chi_k^{0\downarrow}(\mathbf{r})/\chi(\mathbf{r})$  and  $f^{0\downarrow} \equiv \sum_k \chi_k^{0\downarrow}/\chi$ .

Figure 4a shows the fractional unused DFe inventories  $\langle f_k^{0\downarrow} \rangle_{\chi_k}$  and  $\langle f^{0\downarrow} \rangle_{\chi}$  as a function of  $\sigma_{\text{tot}}$  for every member of our family of state estimates. We emphasize that Figure 4 does *not* show the response of the iron cycle to changes in  $\sigma_{\text{tot}}$ , but instead shows the fraction of the unused DFe inventory for distinct equilibrium states, each of which was optimized against observations under a different prescribed iron source scenario (Pasquier and Holzer, 2017). Because the total iron inventory is well constrained across the family, the bulk iron lifetime  $\tau$ , given by the ratio of inventory to source (see also Table 1), is inversely proportional to  $\sigma_{\text{tot}}$ , and hence the mean iron age also scales with  $(\sigma_{\text{tot}})^{-1}$  (Holzer et al., 2016). The systematic increase of the fractional inventory of unused iron, and its approach to saturation at 100 % for the largest sources seen in


Figure 4a, reflects the fact that the probability of past regeneration decreases with the mean iron age until the age becomes so short that very little iron has a chance to pass through the biological pump before having to be scavenged out of the system to match the observed DFe concentrations and inventory. For our smallest total source of  $\sigma_{\text{tot}} \sim 2 \text{ Gmol Fe yr}^{-1}$ , DFe molecules have lived long enough to be regenerated at least once with  $\langle f^{0\downarrow} \rangle_{\chi} \sim 35\%$ , while for our largest total source of  $\sigma_{\text{tot}} \sim 40 \text{ Gmol Fe yr}^{-1}$ , most DFe molecules have not had sufficient time to be regenerated and  $\langle f^{0\downarrow} \rangle_{\chi} \sim 95\%$ . For state estimates with  $\sigma_{\text{tot}} > 7 \text{ Gmol Fe yr}^{-1}$ , more than half of the DFe in the ocean has never passed through the biological pump, i.e.,  $\langle f^{0\downarrow} \rangle_{\chi} > 50\%$ .

Figure 4a also shows that the fraction  $\langle f_k^{0\downarrow} \rangle_{\chi_k}$  varies significantly with iron source type  $k$ . This is because the probability of a past passage through the biological pump is strongly dependent on birth location. The fraction that has not been regenerated since birth is lower for aeolian DFe than for benthic DFe. Benthic DFe must first be transported to the euphotic zone to participate in biological production and is therefore more likely to be scavenged en route compared to aeolian DFe, which is directly injected into the surface. Hydrothermal DFe is the least likely to have passed through the biological pump, with  $\langle f_{\text{H}}^{0\downarrow} \rangle_{\chi_{\text{H}}} \gtrsim 80\%$  for all states and  $\langle f_{\text{H}}^{0\downarrow} \rangle_{\chi_{\text{H}}} \gtrsim 95\%$  for states with  $\sigma_{\text{tot}} > 7 \text{ Gmol Fe yr}^{-1}$ . For sedimentary DFe,  $\langle f_{\text{S}}^{0\downarrow} \rangle_{\chi_{\text{S}}} \gtrsim 45\%$  for all states, and  $\langle f_{\text{S}}^{0\downarrow} \rangle_{\chi_{\text{S}}} \gtrsim 60\%$  for states with  $\sigma_{\text{tot}} > 7 \text{ Gmol Fe yr}^{-1}$ . As the mean age becomes ever smaller with increasing  $\sigma_{\text{tot}}$ , the unused fractional benthic DFe inventory saturates faster to 100% than the unused fractional aeolian DFe inventory, which reaches only  $\sim 90\%$  even for  $\sigma_{\text{tot}} \sim 40 \text{ Gmol Fe yr}^{-1}$ , again reflecting greater probability of biological utilization for surface-injected iron. We will return to Figure 4b below where we explore future passages through the biological pump.

To explore the variations of the asymptotic eigenmode pattern across our different state estimates, Figure 5 shows the zonally averaged patterns of  $\chi^{n\downarrow} = \sum_k \chi_k^{n\downarrow}$  for  $n = 10$ , which is well within the asymptotic regime, for our 5 representative states. (Recall that individual source types all converge to the same pattern, which is hence also the pattern of the total iron concentration.) While there is a signature of Southern Ocean nutrient trapping for all states, there is also significant variation across the state estimates. Broadly, the Southern Ocean trapping has a stronger influence on the eigenmodes of  $\mathcal{F}^{-1}\mathcal{R}$  for the high-source states than for the low-source states. This is likely due to the fact that high-source states are able to match the observed DFe concentrations by having both more active biological iron pumps and scavenging. The transport to depth by both the biological pump and by scavenging particles (the “scavenging pump”) enhances the Southern Ocean trapping. The scavenging is largest at tropical and high latitudes, so that stronger scavenging will tend to remove iron from low and high latitudes (see Appendix C for zonally averaged death rates). At high latitudes this is counteracted by stronger trapping, but at low latitudes the death rate dominates and the concentration of iron that passes many times through the biological pump is strongly diminished. Conversely, for the low-source states, reduced scavenging allows iron to pass many times through the biological pump without being scavenged out of the system at low latitudes. Consequently, the low-source states have iron that has been regenerated many times flowing out of the Southern Ocean with mode and intermediate waters well into the Northern Hemisphere.

Eigenmode for past regenerations:

$$\chi^{10\downarrow} / \langle \chi^{10\downarrow} \rangle$$



**Figure 5.** Normalized global zonal averages of the concentration of DFe that has passed  $n=10$  times through the biological pump since birth, regardless of source type,  $\chi^{10\downarrow}$ , to show the eigen patterns of our five representative state estimates (Table 1). The zonal averages are normalized by the global mean.

### 3.2 Mean number of regenerations since birth

A key metric of how much the DFe field has contributed in the past to organic-matter export is the mean number of times,  $\bar{n}_k$ , that a given iron molecule has been regenerated since its birth. From the local fraction of the total DFe that was regenerated  $n$  times since birth,  $f_k^{n\downarrow}(\mathbf{r})$ , the mean number of past regenerations is by definition given by  $\bar{n}_k(\mathbf{r}) \equiv \sum_{n=0}^{\infty} n f_k^{n\downarrow}(\mathbf{r})$ . **As**

5 **Mathematically**, as shown in Appendix B, it follows from Eq. (5) that  $\bar{n}_k$  obeys

$$\mathcal{H}(\bar{n}_k \chi_k) = \mathcal{R} \chi_k \quad . \quad (6)$$

**Equation-Physically**, Eq. (6) may be interpreted as the equation for a labelling tracer  $\bar{n}_k \chi_k$  that is cycled just like DFe by the  $\mathcal{H}$  operator, but whose numerical value accumulates  $\bar{n}_k$ -fold with  $\bar{n}_k$  past regenerations. **Solving-Computationally**, solving (6) provides an efficient means of finding  $\bar{n}_k$  that avoids first finding and explicitly summing  $f_k^{n\downarrow}(\mathbf{r})$ .

A given iron molecule at  $\mathbf{r}$  is responsible, on average, for the export of  $\bar{n}_k(\mathbf{r})$  iron molecules of source type  $k$  since its birth.


10 The corresponding organic-matter export is quantified by the mean number of phosphorus molecules that are exported along with the iron. The **recycling-regeneration** operator for phosphorus is given by  $\mathcal{R}^P = \sum_c \mathcal{B} f_c \gamma_c^P$  where  $\gamma_c^P(\mathbf{r}) \equiv U_c^P(\mathbf{r}) / \chi(\mathbf{r})$  is diagnosed from the optimized phosphorus uptake  $U_c^P(\mathbf{r})$  of the underlying model (Pasquier and Holzer, 2017). The mean number of phosphorus molecules,  $\bar{n}_k^P(\mathbf{r})$ , globally exported and remineralized in the past, per DFe molecule that is currently at  $\mathbf{r}$ , obeys

$$\mathcal{H}(\bar{n}_k^P \chi_k) = \mathcal{R}^P \chi_k \quad . \quad (7)$$

15 In other words, on average, one of the type- $k$  DFe molecule currently at  $\mathbf{r}$  has supported the export of  $\bar{n}_k^P(\mathbf{r})$  phosphorus molecules since its birth. In analogy with **equation-Eq.** (6), we interpret (7) as defining a labelling tracer  $\bar{n}_k^P \chi_k$  that is cycled like DFe by the  $\mathcal{H}$  operator, but whose numerical value counts the number of phosphate molecules that were regenerated in the past via the phosphate regeneration operator  $\mathcal{R}^P$ .

Figure 6 shows the global zonal averages of  $\bar{n}_k(\mathbf{r})$  for our typical state estimate. Note that the maximum values of  $\bar{n}_k$  are 20 order unity and, on average, most of the iron in the ocean has been regenerated less than once since its birth. Iron with the largest  $\bar{n}_k$  is found near the Southern Ocean surface and in mode and intermediate waters flowing out of the Southern Ocean, which is presumably a signature of Southern Ocean nutrient trapping. In terms of source types, aeolian iron has been most active in the biological pump with  $\bar{n}_A$  exceeding 0.6 throughout most of the ocean interior and exceeding 1.0 throughout much of the Southern Ocean water column. In the surface waters of the low-production subtropical gyres  $\bar{n}_A$  approaches zero. The 25 mean number of regenerations since birth is much smaller for sedimentary and hydrothermal iron, again reflecting the greater scavenging hazard for benthic iron. For sedimentary and hydrothermal iron, values of  $\bar{n}_k$  greater than 0.5 extend from the subantarctic Southern Ocean surface into mode and intermediate waters where scavenging is relatively weak (Appendix C).


The zonally averaged *patterns* of  $\bar{n}_k$  are remarkably similar across our family of estimates. As shown in Appendix D, the spatial patterns are also insensitive to the value of the recyclable fraction of scavenged iron,  $f_{\text{rec}}$ , although their amplitude can 30 vary by a factor of  $\sim 2$  between the  $f_{\text{rec}} = 0$  and  $f_{\text{rec}} = 1$  extremes. The patterns of the global zonal averages of  $\bar{n}_k^P$  (not shown) are nearly identical to those of  $\bar{n}_k$ .



**Figure 6.** Zonal averages of the mean number of past passages through the biological pump (i.e., since birth),  $\bar{n}_k$ . This figure is for our typical state.

We expect significant variations in the amplitudes of  $\bar{n}_k$  and  $\bar{n}$  (the mean number regardless of source type) across the family of state estimates. Because the number of regenerations that are possible during the lifetime of an iron molecule should be proportional to the bulk iron lifetime  $\tau \propto \sigma_{\text{tot}}^{-1}$ , we expect  $\bar{n} \propto \sigma_{\text{tot}}^{-1}$ . To test this and to quantify the range of possible variations, Figure 7a shows the global average  $\langle \bar{n} \rangle$  as a function of  $\sigma_{\text{tot}}$  on a log-log plot and Figure 7b shows the corresponding

### Mean number of regenerations across family



**Figure 7.** (a) The globally averaged mean number of past and future DFe passages through the biological pump per injected DFe molecule (regardless of source type),  $\langle \bar{n} \rangle$  and  $\langle \bar{m} \rangle$ , as a function of the total iron source strength,  $\sigma_{\text{tot}}$ , for the family of state estimates of Pasquier and Holzer (2017). (b) The corresponding globally averaged mean number of phosphorus molecules exported per DFe molecule,  $\langle \bar{n}^{\text{P}} \rangle$  and  $\langle \bar{m}^{\text{P}} \rangle$ . Note the log-log axes to highlight the approximate inverse relationship with  $\sigma_{\text{tot}}$ . To guide the eye, dashed grey lines indicate an exact  $(\sigma_{\text{tot}})^{-1}$  power law.

behavior for  $\langle \bar{n}^{\text{P}} \rangle$ . Both  $\langle \bar{n} \rangle$  and  $\langle \bar{n}^{\text{P}} \rangle$  exhibit the expected approximate inverse relation with  $\sigma_{\text{tot}}$ . Note that the  $\langle \bar{n} \rangle \propto (\sigma_{\text{tot}})^{-1}$  scaling is not exact; instead Figure 7a suggests that there are  $(\sigma_{\text{tot}})^{-1}$  scaling regimes for a low-source cluster and for a high-source cluster, with a transition for intermediate source strengths. This approximate nature of the scaling reflects the fact that the timescale for a passage through the biological pump is not merely set by the prescribed circulation and (in our model) instant particle transport and remineralization, but also depends on the spatial distribution of the scavenging, which varies with the optimized source scenarios. The detailed timescales that link circulation, scavenging, and pumping frequency, will be explored in a future publication. The numerical values of  $\langle \bar{n} \rangle$  range from 0.05 for  $\sigma_{\text{tot}} = 41 \text{ Gmol yr}^{-1}$  to 2.2 for  $\sigma_{\text{tot}} = 1.9 \text{ Gmol yr}^{-1}$ , while  $\langle \bar{n}^{\text{P}} \rangle$  correspondingly ranges from approximately 0.4 to  $4.3 \text{ mol P (mmol Fe)}^{-1}$  (i.e., from 40 to  $460 \text{ mol C (mmol Fe)}^{-1}$  using a simple uniform Redfield C:P ratio of 106:1 for unit conversion).

## 4 Future contributions to export

We now ask how many times a given DFe molecule in the ocean will get regenerated in the future before eventually being permanently scavenged out of the system. The natural way to formulate the necessary equations is to consider the time-reversed adjoint flow (Holzer and Hall, 2000), for which the system is governed by the adjoint operators  $\tilde{\mathcal{H}}$ ,  $\tilde{\mathcal{F}}$ ,  $\tilde{\mathcal{R}}$ , and  $\tilde{\mathcal{D}}$ .

- 5 These adjoints are defined ~~for the volume-weighted inner product~~ in terms of the volume-weighted inner product  $\langle x, y \rangle \equiv \int x(\mathbf{r})y(\mathbf{r})d^3\mathbf{r}$ , where the integral ranges over the entire ocean volume. Thus,  $\tilde{\mathcal{H}}$ , the adjoint of  $\mathcal{H}$ , is defined as usual so that  $\langle \tilde{\mathcal{H}}x, y \rangle = \langle x, \mathcal{H}y \rangle$ . In the time-reversed adjoint flow, the death operator becomes a source of labels that we then track through sequential regenerations (see Figure 1) backward in time to the present, analogously to what we did in the previous section for the usual time-forward flow. The use of adjoint operators allows for numerically efficient evaluation of our diagnostics because
- 10 a single tracer injected at death into the time-reversed adjoint flow suffices to produce the full three-dimensional concentration fields of interest at the present. Here we provide the key equations — their derivation in terms of Green functions is detailed in Appendix E.

### 4.1 Number of regenerations until death

- To compute the number of future regenerations, it is useful to calculate the ~~specific~~ death rate  $d(\mathbf{r})$  ~~and its linear equivalent~~  $\gamma^{\mathcal{D}}(\mathbf{r})\chi(\mathbf{r})$ , with which iron is permanently (non-reversibly) removed at point  $\mathbf{r}$ . This death rate can be diagnosed from the scavenging operator  $\mathcal{D}$  and the DFe concentration  $\chi(\mathbf{r})$  as detailed in Appendix C, where we also show basin zonal averages of  $d(\mathbf{r})$ . The corresponding rate constant  $\gamma^{\mathcal{D}}$  for the linear equivalent model is defined such that  $d(\mathbf{r}) = \gamma^{\mathcal{D}}(\mathbf{r})\chi(\mathbf{r})$ .

One can show that the local fraction  $f^\uparrow(\mathbf{r})$  that eventually dies obeys

$$\tilde{\mathcal{H}}f^\uparrow = \gamma^{\mathcal{D}} \quad , \quad (8)$$

where  $f^\uparrow(\mathbf{r}) = 1$  uniformly everywhere because all ~~the DFe at any point~~ DFe must eventually be scavenged out of the system.

- 20 In analogy with (4), the fraction  $f^{0\uparrow}$  that is regenerated zero times until death obeys

$$\tilde{\mathcal{F}}f^{0\uparrow} = \gamma^{\mathcal{D}}. \quad (9)$$

By using the regeneration rate of DFe that will be regenerated  $m$  times until death and going back in time by one regeneration, we obtain the fraction that will be regenerated  $(m+1)$  times until death, giving us the recursion relation

$$\tilde{\mathcal{F}}f^{(m+1)\uparrow} = \tilde{\mathcal{R}}f^{m\uparrow} \quad . \quad (10)$$

Note that  $\sum_{m=0}^{\infty} f^{m\uparrow}(\mathbf{r}) = f^\uparrow = 1$  as can be verified from (9) and (10). The mean number of regenerations until death,  $\bar{m} = \sum_{m=1}^{\infty} m f^{m\uparrow}$ , can be shown to obey

$$\tilde{\mathcal{H}}\bar{m} = \tilde{\mathcal{R}}f^\uparrow \quad . \quad (11)$$

The mean number of phosphorus molecules,  $\overline{m}^P(\mathbf{r})$ , globally exported and remineralized in the future, per DFe molecule that is currently at  $\mathbf{r}$ , obeys

$$\tilde{\mathcal{H}}(\underline{\overline{m}^P}) = \tilde{\mathcal{R}}^P f^\uparrow . \quad (12)$$

In other words, on average, ~~one of the a~~ DFe molecule currently at  $\mathbf{r}$  will support the export of  $\overline{m}^P$  phosphorus molecules ~~until before~~ it is buried in the sediments.


5 Figure 8 shows the zonally averaged fraction  $f^{m\uparrow}$  of DFe that will undergo  $m$  regenerations in the future, normalized by the global average,  $\langle f^{m\uparrow} \rangle$ , to emphasize changes in the pattern with  $m$ . Note that this fraction is the same for all source types because the future of a given DFe molecule is independent of its past. Below the thermocline, the pattern of  $f^{0\uparrow}$  is nearly uniform at a value just above its global mean but drops to below 50 % of the global mean near the surface where the probability of further biological ~~utilizations-utilization~~ and regeneration is largest (careful inspection is required to see this in Figure 8).  
 10 The pattern of  $f^{1\uparrow}$  has its largest amplitude at the surface and in the Southern Hemisphere, where it spreads deepest from the surface, reflecting the relatively low Southern-Hemisphere death rates (Appendix C). As  $m$  increases,  $f^{m\uparrow}$  becomes rapidly proportional to the gravest eigenmode of  $\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$  [cf., [Equation-Eq.\(10\)](#)]. As this eigenmode is approached, the pattern of  $f^{m\uparrow}$  contracts into the Southern Ocean, presumably because iron that will be regenerated many times before death can only be found where there is both a low death rate and efficient nutrient trapping.

15 To quantify the approach of  $f^{m\uparrow}$  to the gravest eigenmode of  $\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$ , we return to Figure 3b, which shows the global averages of the corresponding DFe concentrations  $\chi_k^{m\uparrow} \equiv \chi_k f^{m\uparrow}$  as a function of  $m$ . Note that  $\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$  and  $\mathcal{F}^{-1}\mathcal{R}$  have the same eigenvalues so that  $\langle \chi_k^{m\uparrow} \rangle$  and  $\langle \chi_k^{n\downarrow} \rangle$  approach the same exponential decay (Figures 3a and 3b).

We return to Figure 4b to consider the fractional DFe inventories that will not pass through the biological pump in the future for all our state estimates. These inventories for source type  $k$  and regardless of source type are given by  $\langle f^{0\uparrow} \rangle_{\chi_k}$  and  
 20  $\langle f^{0\uparrow} \rangle_{\chi}$ . We expect the probability of future regenerations to increase with the remaining lifetime of DFe, which should also scale like  $\sigma_{\text{tot}}^{-1}$  given a well-constrained global DFe inventory. This is confirmed in Figure 4b by the systematic increase and approach to 100 % saturation of  $\langle f^{0\uparrow} \rangle_{\chi_k}$  and  $\langle f^{0\uparrow} \rangle_{\chi}$  with increasing  $\sigma_{\text{tot}}$ . (Note that in the theoretical  $\sigma_{\text{tot}} \rightarrow 0$  limit, we expect  $\langle f^{0\uparrow} \rangle_{\chi_k} \rightarrow 0$  and  $\langle f^{0\uparrow} \rangle_{\chi} \rightarrow 0$ , although our lowest sources are not nearly small enough to exhibit this limiting behavior.)

Figure 4 shows striking asymmetries between unused past and future inventories. The fractional inventory of iron that will  
 25 not be utilized in the future (Figure 4b) is nearly independent of iron type  $k$ , in sharp contrast with the fractional inventory that was not utilized in the past (Figure 4a). The insensitivity of  $\langle f^{0\uparrow} \rangle_{\chi_k}$  to source type is due to the independence of  $f^{0\uparrow}$  on source type so that the  $\chi_k$ -weighted global average is only sensitive to changes in the *pattern* of  $\chi_k$ , which varies little across the family of states (Pasquier and Holzer, 2017). Note that for a given  $\sigma_{\text{tot}}$ , the inventory of total DFe unused in the future,  $\langle f^{0\uparrow} \rangle_{\chi}$ , is significantly larger than the inventory of total DFe unused in the past,  $\langle f^{0\downarrow} \rangle_{\chi}$ . In other words, total DFe is more  
 30 likely to have been regenerated in the past than it is to be regenerated in the future. This asymmetry stems from the fact that  $f^{0\downarrow} = \sum_k \frac{\chi_k}{\chi} f_k^{0\downarrow}$  is dominated by the relatively small unused aeolian fraction  $f_A^{0\downarrow}$  (cf. Figure 4a). In terms of individual source types, aeolian and sedimentary DFe are also more likely to have been regenerated in the past than in the future, but the reverse is true for hydrothermal DFe for which the scavenging hazard following birth is greatest. Thus, for a hypothetical state in which

Future regenerations:  $f^{m\uparrow} / \langle f^{m\uparrow} \rangle$



**Figure 8.** Global zonal averages of the fraction of DFe that will be regenerated  $m$  times,  $f^{m\uparrow}$ , normalized by its global mean value,  $\langle f^{m\uparrow} \rangle$ , for  $m = 0, 1, 2, 3$ , and 10. This figure is for our typical state.

hydrothermal iron dominates the total DFe inventory, it is possible that one could get  $\langle f^{0\downarrow} \rangle_{\chi} > \langle f^{0\uparrow} \rangle_{\chi}$  — however, none of our states fits this scenario.

We now return to the asymptotic eigen patterns of  $f^{m\uparrow}$ . How robust are these eigen patterns across our family of state estimates? Figure 9 shows  $f^{10\uparrow}$  normalized by its global mean and zonally averaged over the global ocean for our 5 representative states (for  $m = 10$ ,  $f^{m\uparrow}$  is an excellent approximation to the gravest eigenmode of  $\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$ ). While there is some variation across the family, the patterns are qualitatively similar. Because the natural quantity that keeps track of future regenerations is the source-type independent *fraction* of DFe that will undergo  $m$  regenerations, the spread in the large- $m$  pattern of  $f^{m\uparrow}$  across the family of states is much smaller than the spread in the large- $n$  pattern of  $\chi_k^{n\downarrow}$  ( $\chi_k^{n\downarrow}$  being the natural quantity for keeping track of past regenerations).

Figure 10a shows the global zonal average of the mean number of future regenerations  $\bar{m}(\mathbf{r})$ , which like  $f^{m\uparrow}$  is independent of source type. The pattern of  $\bar{m}$  is similar to the pattern of  $f^{1\uparrow}$ , which dominates the rapidly converging sum  $\bar{m} = \sum_{m=1}^{\infty} m f^{m\uparrow}$ . The  $\bar{m}$  pattern is surface intensified and largest in the Southern Ocean.  $\bar{m}$  decays rapidly with depth reflecting the fact that the deeper the DFe, the harder it will be to escape death on the way to the surface to participate in the biological pump. These qualitative features are robust across the family of state estimates (not shown). Figure 10b shows that the pattern of the global zonal average of  $\bar{m}^{\text{P}}(\mathbf{r})$  is almost indistinguishable from that of  $\bar{m}(\mathbf{r})$ . This is because the patterns of  $\bar{m}^{\text{P}}$  and  $\bar{m}$  are dominated by the phosphorus and iron export productions, respectively, which are similar despite the substantial spatial variations of the Fe:P uptake ratio (Pasquier and Holzer, 2017).

Returning to Figure 7b, we see that both  $\langle \bar{m} \rangle$  and  $\langle \bar{m}^{\text{P}} \rangle$  are again approximately proportional to  $(\sigma_{\text{tot}})^{-1}$ , as expected. The magnitude of  $\langle \bar{m} \rangle$  remains at or below order unity and ranges from 0.01 to 1.4 as  $\sigma_{\text{tot}}$  varies from 41 to 1.9 Gmol yr<sup>-1</sup>. Correspondingly,  $\langle \bar{m}^{\text{P}} \rangle$  ranges from 0.07 to 2.8 mol P (mmol Fe)<sup>-1</sup>, or from 7 to 290 mol C (mmol Fe)<sup>-1</sup> when converted using a C:P ratio of = 106:1.

## 5 Intrinsic iron fertilization efficiencies


### 5.1 Export supported per unit DFe injection at $\mathbf{r}$

We begin by asking how much an arbitrary (“test”) source  $s_k(\mathbf{r})$  contributes to the globally integrated export and remineralization of iron and phosphate in organic matter. (In our model all exported phosphorus is instantly respired with the divergence of a Martin POP flux profile — DOP is not explicitly modelled.) We first consider the global export that will be supported per unit injection of DFe at point  $\mathbf{r}$ . The concentration response to a unit injection is the Green function associated with operator  $\mathcal{H}$ , and the global export due to this response is simply obtained as the global integral of the regeneration operator  $\mathcal{R}$  acting on this response. As shown in Appendix F, one obtains that DFe injection into volume element  $d^3\mathbf{r}$  at rate  $s_k(\mathbf{r})d^3\mathbf{r}$  supports a globally integrated iron export rate,  $\Phi_k(\mathbf{r})$ , given by

$$\Phi_k(\mathbf{r}) = \bar{m}(\mathbf{r}) s_k(\mathbf{r}) \quad , \quad (13)$$


Eigenmode for future regenerations:

$$f^{10\uparrow} / \langle f^{10\uparrow} \rangle$$



**Figure 9.** Global zonal averages of  $f^{10\uparrow}$ , normalized by their global means,  $\langle f^{10\uparrow} \rangle$ , to show the approximate asymptotic eigen patterns for five representative state estimates.

## Intrinsic fertilization efficiency at any point: zonal averages



**Figure 10.** Three-dimensional **intrinsic** fertilization efficiency metrics for our typical state, zonally averaged across the global oceans. (a) The mean number of iron molecules,  $\bar{m}(\mathbf{r})$ , that will be regenerated per iron molecule at  $\mathbf{r}$  during that molecule’s lifetime. (b) The corresponding mean number of phosphorus molecules,  $\bar{m}^P(\mathbf{r})$ , that will be regenerated.

while the corresponding phosphorus export,  $\Phi_k^P(\mathbf{r})$ , is given by

$$\Phi_k^P(\mathbf{r}) = \bar{m}^P(\mathbf{r}) s_k(\mathbf{r}) . \quad (14)$$

These equations have a straightforward **physical** interpretation: The mean number of future regenerations,  $\bar{m}(\mathbf{r})$ , is simply the number of DFe molecules that will be exported per DFe molecule injected at point  $\mathbf{r}$ , and  $\bar{m}^P(\mathbf{r})$  is the corresponding number of phosphorus molecules that will be exported. The quantities  $\bar{m}(\mathbf{r})$  and  $\bar{m}^P(\mathbf{r})$  are hence measures of the efficiency of **natural** iron fertilization at point  $\mathbf{r}$ . Note that this local efficiency is independent of source type; the efficiency is determined by the biological pump and the transport of water, neither of which depend on the iron source type. **Indeed** **Furthermore**, the local efficiency  $\bar{m}^P(\mathbf{r})$  **does not depend on whether there is even a non-zero source** is defined regardless of whether an iron source is actually present at  $\mathbf{r}$  **in the current state of the iron cycle** — it quantifies the export that would result if DFe were injected at. The efficiency  $\bar{m}^P(\mathbf{r})$  quantifies the global phosphorus export rate *per unit DFe source rate* at  $\mathbf{r}$ . At all points  $\mathbf{r}$ , even where there is no actual source in the system,  $\bar{m}^P(\mathbf{r})$  can be considered to be the “sensitivity” of the linear equivalent system to the insertion of the arbitrary test source  $s_k(\mathbf{r})$ : Equation (14) shows that  $\bar{m}^P(\mathbf{r})$  is the proportionality between the export response  $\Phi_k^P(\mathbf{r})$  and the test source  $s_k(\mathbf{r})$ . In this sense, the fertilization efficiency  $\bar{m}^P$  is a close, but distinct, cousin of the sensitivity to small-amplitude perturbations considered by Dutkiewicz et al. (2006).

Put-Viewed another way, by construction  $\overline{m}(\mathbf{r})$  and  $\overline{m}^P(\mathbf{r})$  quantify the per-molecule fertilization efficiency of the are the mean number of DFe and phosphorus molecules exported per molecule of iron that is present at  $\mathbf{r}$  in the current state of the ocean. These diagnostics, before that molecule is buried. Thus,  $\overline{m}(\mathbf{r})$  and  $\overline{m}^P(\mathbf{r})$  again have the natural interpretation of being fertilization efficiencies. We emphasize that  $\overline{m}(\mathbf{r})$  and  $\overline{m}^P(\mathbf{r})$  are defined without perturbing the system and are hence non-invasive measures of the metrics of what we call the *natural/intrinsic* iron fertilization efficiency. Thus, the The plots of the zonally averaged  $\overline{m}(\mathbf{r})$  and  $\overline{m}^P(\mathbf{r})$  in Figure 10 may thus be interpreted as zonal averages of this intrinsic, three-dimensional iron fertilization efficiency, which can be seen to be highest at the surface and rapidly diminishes with depth. The zonal averages of  $\overline{m}$  and  $\overline{m}^P$  are dominated by the Southern Ocean. However, we will see that the Southern Ocean only plays a secondary role for the fertilization efficiency at the surface.

Figure 11a shows that the surface patterns of  $\overline{m}$  are qualitatively similar across the 5 representative states, with a broad maximum in the central tropical Pacific and with secondary maxima in the subpolar oceans. The states with the largest global mean surface fertilization efficiency,  $\langle \overline{m} \rangle_{\text{surf}}$ , (values provided on Figure 11a) are the low-source states as one would expect: The less iron is in the system, the more the biological pump is expected to benefit from the iron present. For the HiS-LoA state, a typical molecule of DFe at the surface leads to a globally integrated export of  $\langle \overline{m} \rangle_{\text{surf}} \sim 1.4$  molecules of iron. For the high-source states (HiA-LoS and HiH), both the tropical Pacific and the Southern Ocean are locally most prominent, but the surface mean efficiencies are only  $\langle \overline{m} \rangle_{\text{surf}} \sim 0.2$  (HiA-LoS) and 0.5 (HiH) molecules of iron exported per typical surface molecule.

Figure 11b shows the surface patterns of the efficiency for fertilizing organic matter export,  $\overline{m}^P(\mathbf{r})$ . The global surface average,  $\langle \overline{m}^P \rangle_{\text{surf}}$ , is again highest for the low-source states. However,  $\langle \overline{m}^P \rangle_{\text{surf}}$  varies less across our representative states than  $\langle \overline{m} \rangle_{\text{surf}}$  (a range of  $\sim 3$  compared to  $\sim 7$ ). This is consistent with the fact that all states have very similar phosphorus exports (well constrained by the data used in the optimizations), but widely differing iron exports and Fe:P uptake ratios (Pasquier and Holzer, 2017).

The surface patterns of  $\overline{m}^P$  and  $\overline{m}$  are similar, with very similar systematic variations across the different states. However, because of the iron dependence of the Fe:P uptake ratio,  $\overline{m}^P$  has sharper gradients than  $\overline{m}$ , with a more pronounced contrast between the low-efficiency subtropical gyres and the high-efficiency tropical Pacific. The Fe:P uptake ratio in our model is proportional to  $\chi/(\chi+k)$ , which has its lowest values of around 0.1 in the central and eastern tropical Pacific for all our states. The corresponding P:Fe uptake ratio (not shown) has a global pattern that is broadly similar to the pattern of the intrinsic fertilization efficiency  $\overline{m}^P$  because large P:Fe means that a relatively large number of P molecules are taken up (and hence exported) per utilized DFe molecule. The correspondence is not exact, but the P:Fe uptake ratio plays a central role in shaping  $\overline{m}^P$  at the surface. At depth, the effect of the P:Fe ratio on  $\overline{m}^P$  is less important because, depending on where the DFe re-emerges into the euphotic zone, deep DFe will not necessarily be utilized in regions of high P:Fe.

The key result of Figure 11 is that regardless of state, the tropical Pacific is where iron has its highest *natural-intrinsic* fertilization efficiency. For high-source states, fertilizing the fertilization efficiency of the Southern Ocean can be equally as effective large, although the global surface mean efficiency is much lower than for low-source states. The prominence of the tropical Pacific is likely due to the fact that this region tends to be iron-stressed and the equatorial upwelling tends to bring


~~remineralized DFe back to the surface where it is needed to support vigorous biological production.~~ The efficiency is not highest in the eastern tropical Pacific where production is highest because of the associated large scavenging rate due to organic particles. The sweet spot between upwelling-fertilized production and relatively low scavenging lies in the central tropical Pacific to the west of the highest productivity. The state-dependent relative importance of the Southern Ocean compared to the tropical Pacific is a complicated function of how our optimized states match the nutrient observations and reflects a delicate balance between fertilizing biological production and the resulting enhanced scavenging. Interestingly, Dutkiewicz et al. (2006) found that the sensitivity of their coupled model to aeolian iron addition is also largest in the central tropical and eastern Pacific. This suggests that the intrinsic fertilization efficiency of the unperturbed state is shaped by similar processes as the sensitivity to small-amplitude perturbations.

10 How do our estimates of ~~natural~~-intrinsic fertilization efficiency compare to previous estimates ~~in the literature~~ of differently defined fertilization efficiencies? Across our entire family of estimates,  $\langle \bar{m}^P \rangle_{\text{surf}}$  ~~ranges from~~ has a range of 0.7 to  $-7 \text{ mol P (mmol Fe)}^{-1}$ , which converts to ~~73 to~~  $-750 \text{ mol C (mmol Fe)}^{-1}$  using a simple uniform C:P ratio of 106:1. Thus, our estimate of the ~~natural~~-intrinsic fertilization efficiency is roughly one to two orders of magnitude larger than the estimate of about  $3.3 \text{ mol C (mmol Fe)}^{-1}$  by Zeebe and Archer (2005), who reported  $900 \text{ t C}$  exported ~~per~~ for  $1.26 \text{ added-t}$  of iron added during the SOFeX artificial fertilization experiment. There are ~~at least two~~-multiple reasons why our estimate of fertilization efficiency differs from that of Zeebe and Archer (2005). First, the latter is a regional estimate based on a localized artificial fertilization experiment, while the former is a global mean estimate of the unperturbed system. Second, Zeebe and Archer (2005) used the data from the iron fertilization experiments without any corrections for ligand protection in the natural unperturbed system, which leads to an underestimate of the natural fertilization efficiency. de Baar et al. (2008) correct for this in their analysis, giving an estimate of ~~2.6 to~~  $-100 \text{ mol C (mmol Fe)}^{-1}$  for the natural fertilization efficiency. ~~The~~ Despite being inferred from a finite-amplitude perturbation experiment, the estimate of de Baar et al. (2008) ~~thus~~ overlaps with the low end of our estimates ~~,~~ (which corresponds to the possibly more realistic high-source states). The sensitivity of global production to perturbations in the local aeolian source estimated by Dutkiewicz et al. (2006) has a spatial distribution with a range of about  $20\text{--}180 \text{ g C m}^{-2} \text{ yr}^{-1}$  for a  $0.02 \text{ mmol Fe m}^{-2} \text{ yr}^{-1}$  perturbation. The model of Dutkiewicz et al. (2006) exports  $1/3$  of its production as POP so that these sensitivities translate to a POP export per added DFe molecule of about  $28\text{--}250 \text{ mol C (mmol Fe)}^{-1}$ , a lower bound on the full carbon export and about a third of the intrinsic fertilization efficiency estimated here. We emphasize that differences between these various estimates are not only due to uncertainties in the iron cycle (as expressed by our range of values), but also due to differences in the definition of fertilization efficiency, not to mention due to differences among models and methodologies.

## 30 5.2 Relation to relative export efficiency

To compare the importance of the different iron source types in the current state of the ocean, Pasquier and Holzer (2017) defined the export-support efficiency of iron type  $k$  as  $\epsilon(s_k) \equiv \widehat{\Phi}_k / \widehat{\sigma}_k$ , where  $\widehat{\Phi}_k$  is the fraction of the global phosphorus export supported by source type  $k$  and  $\widehat{\sigma}_k$  is the fractional global source of type  $k$ . We found that a key metric that is robust across our family of state estimates is the relative export-support efficiency,  $e_k^P = \epsilon(s_k) / \epsilon(\widetilde{s}_k)$ , which is the ratio of the export-

Intrinsic fertilization efficiency at the surface: normalized patterns



**Figure 11.** Normalized patterns of our **intrinsic** fertilization efficiency metrics at the surface. (a) The mean number of iron molecules,  $\bar{m}(r)$ , that will be regenerated per iron molecule at surface point  $r$  during that molecule's lifetime. (b) The corresponding mean number of phosphorus molecules,  $\bar{m}^{\text{P}}(r)$ , that will be regenerated. To allow for a meaningful comparison of the patterns of different states,  $\bar{m}(r)$  and  $\bar{m}^{\text{P}}(r)$  have been normalized by their global surface averages,  $\langle\bar{m}\rangle_{\text{surf}}$  and  $\langle\bar{m}^{\text{P}}\rangle_{\text{surf}}$  (values are given in each plot).

support efficiency of source type  $k$  to the export-support efficiency of the other source types, whose combined source is the complement  $\tilde{s}_k \equiv s_{\text{tot}} - s_k$ .

It follows algebraically from (14) that the relative export efficiency can be expressed as

$$e_k^P = \frac{\langle \bar{m}^P \rangle_{s_k}}{\langle \bar{m}^P \rangle_{\tilde{s}_k}}, \quad (15)$$

where  $\langle \cdot \rangle_{s_k}$  is the  $s_k$ -weighted global mean, defined so that for any field  $x(\mathbf{r})$ ,  $\langle x \rangle_{s_k} \equiv \int d^3\mathbf{r} x(\mathbf{r}) s_k(\mathbf{r}) / \int d^3\mathbf{r} s_k(\mathbf{r})$ ,  $\langle x \rangle_{s_k} \equiv \langle x s_k \rangle / \langle s_k \rangle$  and  $\langle \cdot \rangle_{\tilde{s}_k}$  is the corresponding  $\tilde{s}_k$ -weighted global mean. Thus, the relative export-support efficiency is the ratio of the mean **intrinsic** fertilization efficiency of the source to the mean **intrinsic** fertilization efficiency of its complement.

- 5 Pasquier and Holzer (2017) found that all members of the family of state estimates share approximately the same relative export efficiencies of  $e_A^P = 3.1 \pm 0.8$ ,  $e_S^P = 0.4 \pm 0.2$ , and  $e_H^P = 0.3 \pm 0.1$ , where the uncertainties represent the scatter across the family. What is new here is relationship (15), which relates the global metric  $e_k^P$  to source-weighted averages of the local **intrinsic** fertilization efficiency,  $\bar{m}^P(\mathbf{r})$ .

## 6 Discussion and caveats

- 10 Our analysis comes with a number of caveats that should be kept in mind. ~~A number~~ Some of these were already identified in the work of Pasquier and Holzer (2017), ~~who designed the~~ where the design of our coupled Fe–P–Si inverse model ~~used~~ ~~hereis~~ presented in detail. In particular, the model used DFe data from the GEOTRACES Intermediate Data Product (IDP) 2014 (in addition to an older compilation by Tagliabue et al. (2012)), which did not contain the newer data from the Pacific and Southern Oceans made available only recently in the GEOTRACES IDP 2017. Although it is inevitable that assimilating
- 15 the additional constraints of the IDP 2017 would lead to some quantitative changes especially in the Pacific, we think that including the IDP 2017 data would not lead to qualitative changes. The states used here do feature strong Pacific DFe plumes of about the observed spatial extent to the west of the East Pacific Rise (EPR), but unlike the observations these plumes also extend to the east of the EPR. The unrealistic eastward plume would not be corrected by assimilating the IDP 2017 data
- 20 out. The deep Pacific circulation can be corrected by assimilating  $\delta^3\text{He}$  into a circulation inverse model (~~DeVries, personal communication~~)(DeVries and Holzer, 2019), but this is beyond the scope of the current study. Moreover, any quantitative differences due to additional constraints would very likely be much smaller than the variations across our family of state estimates, given its two-order-of-magnitude range in iron source strengths.

- While the state estimates used were optimized against the observations available at the time, we note that the sedimentary
- 25 sources of the underlying inverse model of Pasquier and Holzer (2017) are keyed to the flux of particulate organic matter into the bottom box where the organic matter is completely oxidized in our model of the **P-phosphorus** cycle. This parameterization of the sedimentary source is based on observations along the California coast of a correlation between the DFe flux from sediments and the flux of oxidized organic matter (Elrod et al., 2004). All the model’s sediment sources are thus reductive as is the case in many current iron models (e.g., the PISCES and BEC models, Aumont and Bopp, 2006; Moore and Braucher,
- 30 2008). However, the analysis of  $\delta^{56}\text{Fe}$  by Conway and John (2014) has highlighted the importance of non-reductive sources, which our inverse model does not include. Our state estimates feature sedimentary sources that are realistically dominated by

the continental shelves (Figure H1, Pasquier and Holzer, 2017), but they also include one-to-two orders of magnitude smaller sources below highly productive regions in the eastern tropical Pacific, where a sluggish abyssal circulation allows some DFe to accumulate at depth, as can be seen in Figure Fig. 2. Similarly, our hydrothermal sources inject DFe into the ocean where it is then protected from rapid scavenging by our enhanced ligand concentrations near hydrothermal vents. However, the recent work of Fitzsimmons et al. (2017) suggests that this model of hydrothermal iron needs to be revised in the future to include reversible exchange between DFe and particulate iron, which is currently omitted in our scavenging parametrization. Regarding ligands, the inverse model of Pasquier and Holzer (2017) prescribes an optimized distribution of a single type of ligand that is enhanced in hydrothermal plumes and in old waters, with optimized parameters. Ligands are not dynamically transported and the effect of different ligand types on iron bioavailability (Maldonado et al., 2005) is neglected.

Other caveats relate to biogeochemical parameters that could not be optimized. The recyclable fraction,  $f_{\text{rec}}$ , of DFe scavenged by opal and POP ( ~~$f^{\text{Si}}$  and  $f^{\text{POP}}$  in the notation of Pasquier and Holzer (2017)~~), was prescribed to be 90% for all scavenging particles.  $f_{\text{rec}}$  is highly uncertain and in recent models its value has spanned the entire 0–100% range (e.g., Moore and Braucher, 2008; Galbraith et al., 2010; Frants et al., 2016). We established the sensitivity of our results to the value of  $f_{\text{rec}}$  by generating new state estimates from our typical state by prescribing different values of  $f_{\text{rec}}$  ranging from 0 to 100% and re-optimizing the scavenging and source parameters following the strategy of Pasquier and Holzer (2017) (see Appendix D for details). We find that the patterns of  $\bar{n}$  and  $\bar{m}$  are robust to changes in  $f_{\text{rec}}$ , although their global mean values, which are order unity for the reoptimized typical state with  $f_{\text{rec}} = 0$ , systematically decrease by roughly a factor of 2 when  $f_{\text{rec}}$  is increased to 100%. The mean number of past and future phosphorus molecules exported per DFe molecule,  $\bar{n}^{\text{P}}$  and  $\bar{m}^{\text{P}}$ , are robust to changes in  $f_{\text{rec}}$  in both pattern and magnitude. This robustness reflects the fact that the optimization can compensate changes in the scavenging pump with changes in the biological iron pump.

A key control on our model results is the Fe:P stoichiometry. The model approximates the iron dependence of the Fe:P uptake ratio by a Monod function with a half saturation constant that is the same for all phytoplankton classes. We acknowledge that this may not be realistic. ~~Furthermore, the model remineralizes iron with~~ For example, Kirchman (1996) suggested that including the microbial “ferrous wheel”, which operates with different stoichiometric ratios, could affect iron budgets in the euphotic zone. Similarly, Strzpek et al. (2005) explored variations in the iron regeneration rates of different organisms and cautioned modelers to pay careful attention to the details of the Fe:C stoichiometry. Relatedly, our model simply remineralizes iron in the same Fe:P stoichiometric ratio with which it was utilized, i.e., so that the vertical profiles of iron and phosphate remineralization have identical shapes. However, measurements by Twining et al. (2014) show that, at least for some phytoplankton species, iron is remineralized more slowly than phosphate, suggesting that our remineralization profile for iron could be too shallow. ~~Because the~~ Different remineralization lengthscales for iron and phosphate were also emphasized by Boyd et al. (2017). ~~Because our~~ model is optimized to fit the DFe observations, with an emphasis on deep profiles relative to surface measurements (Pasquier and Holzer, 2017), a potentially too shallow remineralization of iron would be compensated by an increased strength of the biological pump. Furthermore, the relative amount of scavenging by opal and POP particles is optimizable in our model, so that deeper iron remineralization can be achieved by increasing the scavenging by opal. We acknowledge, however, that when optimizing the match to observed DFe, the model may produce biases in the relative con-

tributions of the biological and scavenging pumps, which would affect our estimates of the number of passages through the biological pump.

The recent work of Rafter et al. (2017) suggests that in surface waters DFe must be preferentially recycled compared to nitrate in order to sustain the observed nitrate consumption in the iron-limited equatorial Pacific. While we do not model the nitrogen cycle, it is reasonable to assume that in the euphotic zone DFe may also be preferentially recycled compared to phosphate. At face value, this appears to contradict our assumption that the detrital fractions,  $f_c$ , are identical for Fe, P, and Si export. However, the uptake and export of DFe in our model are proportional to the product  $f_c R^{\text{Fe:P}}$ , and  $R^{\text{Fe:P}}$  is optimized, so that any difference between the iron and phosphate detrital fractions is simply absorbed into  $R^{\text{Fe:P}}$ . However, this does point to the need for caution when interpreting the optimized values of  $R^{\text{Fe:P}}$ .

Importantly, we would like to note that not every process thought to influence DFe needs to be explicitly modelled for a useful representation of the iron cycle. The model of Pasquier and Holzer (2017) is of intermediate complexity, and effects due to processes not explicitly modelled are captured implicitly when parameters are optimized to fit the observed nutrient and phytoplankton fields. Explicit modelling of all known processes in their full complexity may be important for models that try to predict how the system will change in the future, but this is neither necessary nor desirable for constraining and diagnosing the large-scale cycling of DFe in the current state of the ocean as we do here.

## 7 Conclusions

We have presented a new conceptual and mathematical framework for quantifying the contribution of DFe to the biological pump during its journey from birth by an external source to death by irreversible scavenging and burial. New diagnostics were developed to partition the DFe concentration into the fraction that was regenerated  $n$  times in the past or that will be regenerated  $m$  times in the future. These diagnostics include new tracer equations for the mean number of regenerations  $\bar{n}$  and  $\bar{m}$ , which afford numerically efficient computation. The mean number  $\bar{m}$  of future regenerations that iron at any point  $r$  will undergo is a measure of the **intrinsic** fertilization efficiency of iron at  $r$ , giving the number of iron and phosphorus molecules globally exported per DFe molecule at  $r$ .

We applied our new diagnostics to a family of optimized state estimates of the global coupled iron, phosphorus, and silicon cycles, assuming steady state. All states of the family match the observed nutrient concentrations about equally well despite spanning a large range of external iron source strengths. Performing our analyses across the family of states allowed us to identify aspects of the birth-to-death journey of DFe that are robust and to quantify systematic variations with iron source strength. Our key findings are as follows:

1. A large portion of the global DFe inventory never participates in the biological pump. For states with iron sources  $\sigma_{\text{tot}}$  larger than  $\sim 7 \text{ Gmol Fe yr}^{-1}$ , more than 50 % of the inventory has not passed through the pump since birth and more than 85 % will not pass through the pump before death. Because of its direct injection into the euphotic zone, a larger portion of aeolian iron passes through the biological pump than other source types. Both the mean number of past and future passages through the biological pump,  $\bar{n}$  and  $\bar{m}$ , are approximately proportional to the bulk iron lifetime  $\tau \propto \sigma_{\text{tot}}^{-1}$ .

For an increase of  $\sigma_{\text{tot}}$  from 1.9 to 40 Gmol Fe yr<sup>-1</sup>, the global average,  $\langle \bar{n} \rangle$ , decreases from 2.2 to 0.05, while  $\langle \bar{m} \rangle$  decreases from 1.4 to 0.01.

2. The three-dimensional distribution of  $\bar{n}_k(\mathbf{r})$  has its largest values in the Southern Ocean and a pattern that suggests nutrient trapping for all iron source types  $k$ , but with different vertical structure for different source types. The precise patterns of  $\bar{n}_k(\mathbf{r})$  are shaped by the delicate balance between regeneration and scavenging, which are processes with strong spatial overlap near the surface. For aeolian iron, the largest values of  $\bar{n}_k$  are found in surface, mode, intermediate, and bottom waters, while for sedimentary and hydrothermal iron  $\bar{n}_k$  is small in ~~bottom~~ deep waters because of the greater scavenging hazard during transit from benthic sources to first uptake in the euphotic zone.
3. The spatial distribution of DFe that was regenerated  $n$  times since birth varies with  $n$  because DFe is reorganized in the water column with each passage through the biological pump. Unused DFe ( $n = 0$ ) is generally concentrated near the sources. The concentration of unused aeolian DFe has a secondary maximum at  $\sim 2$  km depth and extends to the ~~sea floor~~ seafloor because of the action of the scavenging pump (scavenging and particle transport to depth followed by re-dissolution). Aeolian DFe regenerated exactly once since birth ( $n = 1$ ) has its maximum concentration well below the surface, and sedimentary DFe regenerated exactly once has its maximum concentration well above the bottom, with both maxima at  $\sim 1.5$  km depth. DFe ~~trapped in the Southern Ocean that~~ has undergone the largest number of regenerations since birth is trapped in the Southern Ocean.
4. The pattern of the large- $n$  DFe concentration is independent of iron source type because the memory of birth place quickly dissipates with successive passages through the biological pump, although the rate of convergence to the asymptotic large- $n$  pattern does depend on source type. The large- $n$  pattern corresponds to the gravest eigenmode of  $\mathcal{F}^{-1}\mathcal{R}$ , which represents the combined transport by the circulation and by sinking particles for one passage through the biological pump. For  $n$  larger than  $\sim 5$ , the concentration of DFe that was regenerated  $n$  times since birth decays exponentially with  $n$ . The  $e$ -folding scale  $n^*$  ranges from 0.4–2.8 across our family, with higher sources corresponding to smaller  $n^*$  (faster decay consistent with shorter lifetimes).
5. The fraction  $f^{m\uparrow}(\mathbf{r})$  of DFe currently at  $\mathbf{r}$  that will pass  $m$  times through the biological pump in the future is independent of iron source type. The local fraction that will not participate in biological production in the future,  $f^{0\uparrow}(\mathbf{r})$ , has a nearly spatially uniform distribution and makes the largest contribution to the DFe inventory. The local fraction that will be utilized exactly once before death,  $f^{1\uparrow}(\mathbf{r})$ , is typically concentrated near the surface, where the likelihood of passing through the biological pump before being scavenged is largest. Correspondingly, the three-dimensional distribution of  $\bar{m}(\mathbf{r})$  has its maximum near the surface and is Southern Ocean intensified.
6. The fraction of DFe that will be regenerated  $m$  times in the future again approaches an eigenmode for asymptotically large  $m$ . The amplitude of this eigenmode decays with the same  $e$ -folding scale  $n^*$  as the eigenmode associated with past regenerations. However, the eigenmodes associated with future and past regenerations have different patterns, with the ~~mode~~ eigenmode for future regenerations being much more surface intensified. Both past and future ~~modes~~ eigenmodes

are concentrated in the Southern Ocean because multiple regenerations during an iron molecule's lifetime are more likely where nutrients are effectively trapped (Primeau et al., 2013; Holzer et al., 2014).

7. We defined and quantified the ~~natural~~ **intrinsic** iron fertilization efficiency at ~~an arbitrary~~ point  $r$  in terms of the number of globally exported phosphorus molecules per DFe molecule at  $r$ . This is the first time that ~~the a metric of~~ iron fertilization efficiency has been estimated in three dimensions, which may ultimately prove useful for quantifying the importance of different iron reservoirs for supporting the ocean's global export production. At the surface, the ~~natural~~ **intrinsic** fertilization efficiency is largest in the central tropical Pacific with secondary maxima in the subpolar oceans. The relative importance of the Southern Ocean is greatest for high-source states. Globally averaged, the surface fertilization efficiency ranges from 0.7 to 7 mol P (mmol Fe)<sup>-1</sup> across our family of state estimates, with the low-source states having the largest efficiencies. ~~(In carbon units, the efficiency ranges from 73 to 750 using a C:P ratio of 106:1.)~~

Intimately connected to the mean number of past and future regenerations are the age and expected remaining lifetimes of DFe in the ocean. A full exploration of these timescales and the associated transit-time distributions is beyond the scope of this study. However, in future work, we plan to explore the timescales of the iron cycle and their connection to setting the efficiency with which iron fertilization achieves carbon sequestration (e.g., DeVries et al., 2012; Pasquier and Holzer, 2016). **We also note that the approach of using a linear equivalent linear model to partition iron and diagnose its life cycle can also be applied to perturbed states (e.g., due to iron addition or changes in circulation) to shed light on how the iron cycle operates for various paleo or future climate scenarios.** Finally, the concepts and methods employed here can be applied to other nutrients for a more complete picture of how the interaction between the biological pump and the physical transport shapes their distributions and cycling rates; we plan to do so in future work.

## 20 **Appendix A: Derivation of equation for $\bar{n}(r)$** **Glossary**

### **Appendix B: Derivation of $\bar{n}(r)$**

First we verify that  $\sum_{n=0}^{\infty} \chi_k^{n\downarrow} = \chi_k$ . From (4) and (5) we have

$$\chi_k^{n\downarrow} = \mathcal{A}^n \mathcal{F}^{-1} s_k \quad , \quad (\text{B1})$$

where  $\mathcal{A} \equiv \mathcal{F}^{-1} \mathcal{R}$ . Because  $\mathcal{A}$  must have a maximum eigenvalue less than unity for convergence, we can use the geometric operator sum  $\sum_{n=0}^{\infty} \mathcal{A}^n = (1 - \mathcal{A})^{-1}$ . Thus,

$$\sum_{n=0}^{\infty} \chi_k^{n\downarrow} = (1 - \mathcal{A})^{-1} \mathcal{F}^{-1} s_k \quad . \quad (\text{B2})$$

- 25 Applying  $\mathcal{F}(1 - \mathcal{A})$  from the left, and recognizing that  $\mathcal{F}(1 - \mathcal{A}) = \mathcal{F} - \mathcal{R} = \mathcal{H}$ , we have

$$\mathcal{H} \sum_{n=0}^{\infty} \chi_k^{n\downarrow} = s_k \quad . \quad (\text{B3})$$

**Table A1.** Glossary of mathematical symbols, their definitions, and SI units. PH17 refers to Pasquier and Holzer (2017).

Symbol	Definition	SI unit
$\mathcal{T}$	Advective eddy-diffusive transport operator: $(\mathcal{T}\chi)(\mathbf{r})$ is the flux divergence of $\chi$ at point $\mathbf{r}$	$\text{s}^{-1}$
$\mathcal{L}$	Uptake operator: $(\mathcal{L}\chi)(\mathbf{r})$ is the rate of uptake of $\chi$ at point $\mathbf{r}$	$\text{s}^{-1}$
$\mathcal{S}_c, \mathcal{S}_j$	Biogenic transport and remineralization operators for the full water column (including the euphotic zone) for phytoplankton class $c$ or particle type $j$ ( $\mathcal{S}_c^{\text{Fe}}, \mathcal{S}^{\text{s},j}$ in PH17)	dimensionless
$\mathcal{B}$	Biogenic transport and remineralization operator for export below the euphotic zone	dimensionless
$\mathcal{R}, \mathcal{R}^{\text{P}}$	Iron and phosphorus regeneration operators	$\text{s}^{-1}, \frac{\text{mol P}}{\text{mol Fe}} \text{s}^{-1}$
$\mathcal{D}$	Reversible scavenging operator: $(\mathcal{D}\chi)(\mathbf{r})$ is the rate of removal of $\chi$ by scavenging minus the rate with which scavenged iron is recycled at point $\mathbf{r}$	$\text{s}^{-1}$
$\mathcal{H}, \mathcal{F}$	Full system operators $\mathcal{H} \equiv \mathcal{T} - \mathcal{R} + \mathcal{L} + \mathcal{D}$ and $\mathcal{F} \equiv \mathcal{T} + \mathcal{L} + \mathcal{D}$	$\text{s}^{-1}$
$\mathcal{A}$	Recursion operator $\mathcal{A} \equiv \mathcal{F}^{-1}\mathcal{R}$ such that $\chi_k^{(n+1)\downarrow} = \mathcal{A}\chi_k^{n\downarrow}$	dimensionless
$\lambda$	Largest eigenvalue of $\mathcal{A}$	dimensionless
$n^*$	$e$ -folding scale for the decay of $\chi_k^{n\downarrow}$ and $\chi_k^{m\uparrow}$ with number of regenerations (independent of source type $k$ )	dimensionless
$\tilde{\mathcal{O}}$	Adjoint of $\mathcal{O}$ for any operator $\mathcal{O}$	-
$\mathbf{r}$	Position vector	m
$U_c(\mathbf{r}), U_c^{\text{P}}(\mathbf{r})$	Iron and phosphorus uptake rates for phytoplankton class $c$ ( $R^{\text{Fe:P}}U_c$ and $U_c$ in PH17)	$\text{mol m}^{-3} \text{s}^{-1}$
$J_j(\mathbf{r})$	DFe scavenging rate for particle type $j$	$\text{mol m}^{-3} \text{s}^{-1}$
$d(\mathbf{r})$	Death rate (rate of DFe scavenging that is not recycled and results in burial)	$\text{mol m}^{-3} \text{s}^{-1}$
$\gamma_c(\mathbf{r}), \gamma_j(\mathbf{r}), \gamma^{\text{D}}(\mathbf{r})$	Rate constants for iron uptake, scavenging, and death, respectively	$\text{s}^{-1}$
$f_c$	Detrital fraction for phytoplankton class $c$	dimensionless
$f_{\text{rec}}$	Fraction of scavenging that is recyclable ( $f^{\text{bSi}}$ and $f^{\text{POP}}$ in PH17)	dimensionless
$\chi(\mathbf{r})$	Concentration of total DFe ( $\chi_{\text{Fe}}$ in PH17)	$\text{mol m}^{-3}$
$\chi_k(\mathbf{r})$	Concentration of DFe of source type $k$ (Aeolian (A), Sedimentary (S), or Hydrothermal (H))	$\text{mol m}^{-3}$
$s_k(\mathbf{r})$	DFe source of type $k$ (global integral $\sigma_k$ )	$\text{mol m}^{-3} \text{s}^{-1}$
$\tau$	Bulk DFe lifetime	s
$\chi^{n\downarrow}(\mathbf{r}), \chi^{m\uparrow}(\mathbf{r})$	Concentration of DFe that passed $n$ times, or that will pass $m$ times, through the biological pump	$\text{mol m}^{-3}$
$f^{n\downarrow}(\mathbf{r}), f^{m\uparrow}(\mathbf{r})$	Fraction of DFe that passed $n$ times, or that will pass $m$ times, through the biological pump	dimensionless
$\bar{n}_k(\mathbf{r}), \bar{m}(\mathbf{r})$	Mean number of past ( $n$ ) or future ( $m$ ) passages of DFe through the biological pump	dimensionless
$\bar{n}^{\text{P}}(\mathbf{r}), \bar{m}^{\text{P}}(\mathbf{r})$	Mean number of past ( $n$ ) or future ( $m$ ) moles of P regenerated per mole of Fe	$\text{mol P} (\text{mol Fe})^{-1}$
$\Phi_k(\mathbf{r}), \Phi_k^{\text{P}}(\mathbf{r})$	Globally integrated iron or phosphorus export rate due to iron source $s_k(\mathbf{r})$	$\text{mol m}^{-3} \text{s}^{-1}$
$\langle \cdot \rangle, \langle \cdot \rangle_x, \langle \cdot \rangle_{\text{surf}}$	Global averages: volume-weighted, $x$ -weighted, and area-weighted surface	-

This shows that  $\sum_{n=0}^{\infty} \chi_k^{n\downarrow}$  and  $\chi_k$  obey the same equation and hence that they are equal.

The defining equation for  $\bar{n}_k(\mathbf{r})$  is

$$\chi_k(\mathbf{r})\bar{n}_k(\mathbf{r}) = \sum_{n=0}^{\infty} n\chi_k^{n\downarrow}(\mathbf{r}) . \quad (\text{B4})$$

From recursion relation (5), we have  $\chi_k^{n\downarrow} = \mathcal{A}^n \chi_k^{0\downarrow}$ . Substituting this, using the geometric operator sum  $\sum_{n=0}^{\infty} n\mathcal{A}^n = (1 - \mathcal{A})^{-1} \mathcal{A} (1 - \mathcal{A})^{-1}$ , and the fact that from (B1)  $\chi_k^{0\downarrow} = \mathcal{F}^{-1} s_k = \mathcal{F}^{-1} \mathcal{H} \mathcal{H}^{-1} s_k = (1 - \mathcal{A}) \chi_k$ , we obtain

$$\chi_k \bar{n}_k = (1 - \mathcal{A})^{-1} \mathcal{A} \chi_k . \quad (\text{B5})$$

5 Applying  $\mathcal{R} \mathcal{A}^{-1} (1 - \mathcal{A}) = \mathcal{F} - \mathcal{R} = \mathcal{H}$  from the left yields (6).


### Appendix C: Local iron death rate

To diagnose the local death rate with which iron is permanently removed from the ocean, we first rewrite the reversible scavenging operator  $\mathcal{D}$  in terms of a simplified local **specific**-death rate. The operator  $\mathcal{D}$  is an integral operator so that  $(\mathcal{D}\chi)(\mathbf{r}) = \int d^3\mathbf{r}' K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}') \chi(\mathbf{r}')$  with adjoint  $(\tilde{\mathcal{D}}\phi)(\mathbf{r}') = \int d^3\mathbf{r} \phi(\mathbf{r}') K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}')$ . The term  $K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}') \chi(\mathbf{r}')$  represents the rate  
 10 with which iron is scavenged at  $\mathbf{r}'$  minus the rate with which this scavenged iron is redistributed to points  $\mathbf{r}$  through particle transport and re-dissolution. By integrating over all destination points  $\mathbf{r}$ , we obtain the deficit between the rate of iron scavenging at  $\mathbf{r}'$  and the total water-column integrated rate of redissolution of that iron. Thus,  $d(\mathbf{r}') \equiv \int d^3\mathbf{r} K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}') \chi(\mathbf{r}')$  is the rate of permanent removal or death at  $\mathbf{r}'$ . **In terms of the equivalent linear specific death rate, we have  $d(\mathbf{r}') = \gamma^{\mathcal{D}}(\mathbf{r}') \chi(\mathbf{r}')$ , with  $\gamma^{\mathcal{D}}(\mathbf{r}') \equiv \int d^3\mathbf{r} K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}')$ .** For the local rate constant we define  $\gamma^{\mathcal{D}} \equiv \int d^3\mathbf{r} K_{\mathcal{D}}(\mathbf{r}|\mathbf{r}')$ , which gives  $d(\mathbf{r}') = \gamma^{\mathcal{D}}(\mathbf{r}') \chi(\mathbf{r}')$ .

15 For the interpretation of our diagnostics it is helpful to know how the local iron death rate,  $d(\mathbf{r}') = \gamma^{\mathcal{D}}(\mathbf{r}') \chi(\mathbf{r}')$ ,  $d(\mathbf{r}')$  is distributed through the ocean. Figure C1 shows basin and global zonal averages of the death rate for our five representative states. Key features common to all states are relatively low death rates in the subtropical gyres of both hemispheres where production and hence scavenging particle fluxes are low, and relatively high death rates at high latitudes and in the tropics where production is large. The highest death rates occur in the surface ocean where particle fluxes are highest. The near-  
 20 surface death rates are larger in the Northern Hemisphere than in the Southern Hemisphere, presumably because the DFe concentrations are higher in the Northern Hemisphere.

### Appendix D: Variation of key diagnostics with recyclable fraction parameter

To explore variations of our results with the non-optimized recyclable fraction  $f_{\text{rec}}$ , we changed the value of  $f_{\text{rec}}$  of our typical state (for which  $f_{\text{rec}} = 90\%$ ), and then re-optimized the biogeochemical sink and source parameters following the  
 25 optimization strategy described by Pasquier and Holzer (2017). This generated a set of 8 optimized state estimates with  $f_{\text{rec}} = 0, 10, 30, 50, 70, 80, 90$ , and 100%. Except for the  $f_{\text{rec}} = 100\%$  case, these states have similar iron source strengths ( $\sigma_{\text{A}}$  ranges from 5.3 to 5.7  $\text{Gmol yr}^{-1}$ ,  $\sigma_{\text{S}}$  from 1.7 to 1.9  $\text{Gmol yr}^{-1}$ , and  $\sigma_{\text{H}}$  is unchanged to 2 significant figures at




**Figure C1.** Basin and global zonal averages of the death rate  $d(r')$  for our five representative states. Note the logarithmic color scale.

0.88 Gmol yr<sup>-1</sup>). For the extreme case of  $f_{\text{rec}} = 100\%$ , when the only permanent iron sink in the system is due to the flux of scavenged iron that reaches the ocean bottom where we assume it is buried (in addition, iron scavenged by mineral dust is always assumed not to be recyclable, but dust scavenging is very small for our estimates), the optimized sedimentary iron source more than doubles and the other sources also undergo significant adjustments during the re-optimization ( $\sigma_A = 5.1$ ,  $\sigma_S = 3.7$ , and  $\sigma_H = 0.85$  Gmol yr<sup>-1</sup> for  $f_{\text{rec}} = 100\%$ ).

Figure D1 shows that the spatial pattern of the zonally averaged  $\bar{n}$  is very insensitive to the value of  $f_{\text{rec}}$  when the scavenging and source parameters are optimized. The patterns of the zonally averaged  $\bar{m}$  have the same qualitative features for all values of  $f_{\text{rec}}$ , but become more surface-intensified as  $f_{\text{rec}}$  increases to unity, presumably because a greater recycling fraction increases the chance that DFe is available near the euphotic zone to pass through the biological pump. The patterns of  $\bar{n}^P$  and  $\bar{m}^P$  (not shown) are similarly insensitive to  $f_{\text{rec}}$ .

Figure D2 shows how the volume-weighted global means of  $\bar{n}$ ,  $\bar{m}$ ,  $\bar{n}^P$ , and  $\bar{m}^P$  vary with the recyclable fraction,  $f_{\text{rec}}$ . We find that  $\langle \bar{n} \rangle$  and  $\langle \bar{m} \rangle$  are order unity for  $f_{\text{rec}} = 0\%$  and tend to decrease by roughly a factor of 2 to 3 when  $f_{\text{rec}}$  is increased from 0 to 100%. We emphasize that this is not simple sensitivity to  $f_{\text{rec}}$ , because the scavenging and source parameters have been reoptimized for each choice of  $f_{\text{rec}}$ . Broadly, a larger value of the number of regenerations means that more iron is transported




**Figure D1.** Top: The zonal means of  $\bar{n}(\mathbf{r})$ , normalized by the corresponding global averages  $\langle \bar{n} \rangle$ , for our typical state estimate re-optimized with three very different recyclable fractions:  $f_{\text{rec}} = 0, 50$ , and  $100\%$ . Bottom: The corresponding normalized zonal means of  $\bar{m}(\mathbf{r})$ . The values of the global averages,  $\langle \bar{n} \rangle$  and  $\langle \bar{m} \rangle$ , are given in the plot titles.

to depth with the biological pump. This is consistent with the fact that for small values of  $f_{\text{rec}}$ , less iron is transported to depth through scavenging and re-dissolution. Given the similar iron sources of these states, and hence similar global mean scavenging rates, reduced iron transport through scavenging must be compensated by increased transport through biological uptake and regeneration, as measured here by the mean number or iron regenerations. The mean number of phosphorus molecules that are regenerated in the past or the future per iron molecule,  $\bar{n}^{\text{P}}$  and  $\bar{m}^{\text{P}}$ , are much less sensitive to the value of  $f_{\text{rec}}$  in these optimized states. The reason for this insensitivity probably lies in the fact that the phosphorus export is well constrained by the nutrient and phytoplankton data against which we optimize (Pasquier and Holzer, 2017), and any changes in iron export are compensated by different optimized values of the Fe:P uptake ratio, which we also use as the Fe:P ratio for remineralization.

## Appendix E: Recursion relation for $f^{m\uparrow}$

### 10 E1 Green functions

To derive the recursion relation for the fraction of iron at a given location that is regenerated exactly  $m$  times before death, we use a Green function approach. The Green function associated with the equation  $(\partial_t + \mathcal{H})\chi = s$  is obtained by replacing the



**Figure D2.** (a) The globally averaged mean number of past and future DFe regenerations per injected DFe molecule,  $\langle \bar{n} \rangle$  and  $\langle \bar{m} \rangle$ , as a function of the recyclable fraction  $f_{\text{rec}}$ . The corresponding states were obtained by changing the value of  $f_{\text{rec}}$  of the typical state and re-optimizing all other source and scavenging parameters. The typical state has  $f_{\text{rec}} = 90\%$  and is indicated by the vertical dashed line. (b) As (a) for the past and future number of phosphorus molecules exported per injected DFe molecule,  $\langle \bar{n}^{\text{P}} \rangle$  and  $\langle \bar{m}^{\text{P}} \rangle$ .

source  $s$  with a Dirac delta function in space and time:

$$(\partial_t + \mathcal{H}_r)G_{\mathcal{H}}(\mathbf{r}, t | \mathbf{r}', t') = \delta(t - t')\delta^3(\mathbf{r} - \mathbf{r}') \quad , \quad (\text{E1})$$

where the subscript on  $\mathcal{H}_r$  reminds us that  $\mathcal{H}$  acts on the field-point coordinates  $\mathbf{r}$ . The Green function for the time-reversed adjoint flow (e.g., Holzer and Hall, 2000) obeys

$$(-\partial_{t'} + \tilde{\mathcal{H}}_{r'})G_{\tilde{\mathcal{H}}}(\mathbf{r}', t' | \mathbf{r}, t) = \delta(t - t')\delta^3(\mathbf{r} - \mathbf{r}') \quad , \quad (\text{E2})$$

where the adjoint Green function  $G_{\tilde{\mathcal{H}}}$  obeys the reciprocity relation  $G_{\tilde{\mathcal{H}}}(\mathbf{r}', t' | \mathbf{r}, t) = G_{\mathcal{H}}(\mathbf{r}, t | \mathbf{r}', t')$ . The Green functions associated with  $(\partial_t + \mathcal{F})\chi = s$  are defined in exactly the same manner.

All adjoints here are defined in terms of the volume-weighted inner product so that for linear operator  $\mathcal{H}$  and any two fields  $\phi$  and  $\psi$  we have  $\int d^3\mathbf{r} \phi(\mathbf{r})(\mathcal{H}\psi)(\mathbf{r}) = \int d^3\mathbf{r} (\tilde{\mathcal{H}}\phi)(\mathbf{r})\psi(\mathbf{r})$  and  $y$  we have  $\int d^3\mathbf{r} x(\mathbf{r})(\mathcal{H}y)(\mathbf{r}) = \int d^3\mathbf{r} (\tilde{\mathcal{H}}x)(\mathbf{r})y(\mathbf{r})$ . For computation, all linear operators are discretized on a numerical grid and organized into sparse matrices, e.g.,  $\mathcal{H}$  becomes matrix  $\mathbf{H}$  with adjoint  $\tilde{\mathbf{H}} = \mathbf{V}^{-1}\mathbf{H}^T\mathbf{V}$ , where  $\mathbf{V}$  is a diagonal matrix of the grid box volumes.

## E2 All DFe must die

Now consider the concentration of iron  $\chi(\mathbf{r}', t')$  in some volume  $d^3\mathbf{r}'$ . As the system evolves the mass  $\chi(\mathbf{r}', t') d^3\mathbf{r}'$  results in concentration  $\mathcal{X}(\mathbf{r}, t|\mathbf{r}', t') d^3\mathbf{r}'$  at  $(\mathbf{r}, t)$  which is obtained by propagating with  $G_{\mathcal{H}}$  so that

$$\mathcal{X}(\mathbf{r}, t|\mathbf{r}', t') d^3\mathbf{r}' = G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t') \chi(\mathbf{r}', t') d^3\mathbf{r}' \quad . \quad (\text{E3})$$

(Note that there is no integral as this is pointwise propagation.) The death rate per unit volume incurred by  $\mathcal{X}(\mathbf{r}, t|\mathbf{r}', t') d^3\mathbf{r}'$  at  $(\mathbf{r}, t)$  is given by  $\gamma^{\mathcal{D}}(\mathbf{r}) \mathcal{X}(\mathbf{r}, t|\mathbf{r}', t') d^3\mathbf{r}'$ . Integrating this death rate over all times  $t$  and over the entire ocean for  $\mathbf{r}$ , we must recover the initial mass  $\chi(\mathbf{r}', t') d^3\mathbf{r}'$ . Thus, we have

$$\chi(\mathbf{r}', t') d^3\mathbf{r}' \equiv \int dt \int d^3\mathbf{r} \gamma^{\mathcal{D}}(\mathbf{r}) G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t') \chi(\mathbf{r}', t') d^3\mathbf{r}' \quad , \quad (\text{E4})$$

where the initial volume  $d^3\mathbf{r}'$  is *not* integrated over. Because  $G_{\mathcal{H}}$  is just a function and not a differential operator, and because we are integrating with respect to  $(\mathbf{r}, t)$ , we can divide both sides by  $\chi(\mathbf{r}', t') d^3\mathbf{r}'$ , which gives

$$f^{\uparrow} \equiv \int dt \int d^3\mathbf{r} \gamma^{\mathcal{D}}(\mathbf{r}) G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t') \quad , \quad (\text{E5})$$

where  $f^{\uparrow}$  is just a spatially uniform field of unit value, i.e.,  $f^{\uparrow} = 1$ . Applying the time-reversed adjoint operator  $(-\partial_{t'} + \tilde{\mathcal{H}}_{\mathbf{r}'})$  from the left to (E5) and using the reciprocity relation of  $G_{\mathcal{H}}$ , we obtain (8). Note that  $\tilde{\mathcal{H}} = \tilde{\mathcal{T}} + (\tilde{\mathcal{L}} - \tilde{\mathcal{R}}) + \tilde{\mathcal{D}}$ . Because  $\mathcal{T}$  and  $\mathcal{L} - \mathcal{R}$  are mass-conserving operators,  $\tilde{\mathcal{T}} f^{\uparrow} = 0$  and  $(\tilde{\mathcal{L}} - \tilde{\mathcal{R}}) f^{\uparrow} = 0$  so that (8) is equivalent to  $\tilde{\mathcal{D}} f^{\uparrow} = \gamma^{\mathcal{D}}$ , which reproduces the definition of  $\gamma^{\mathcal{D}}$ .

## E3 Fraction not regenerated in the future

Similarly we can construct the concentration of DFe that has been regenerated exactly zero times since being at  $(\mathbf{r}', t')$ . By propagating the mass  $\chi(\mathbf{r}', t') d^3\mathbf{r}'$  with  $\mathcal{F}$  instead of  $\mathcal{H}$ , we obtain the resulting concentration at  $t$  that has not passed through the biological pump since  $t'$ :

$$\chi^{0\uparrow}(\mathbf{r}, t|\mathbf{r}', t') d^3\mathbf{r}' = G_{\mathcal{F}}(\mathbf{r}, t|\mathbf{r}', t') \chi(\mathbf{r}', t') d^3\mathbf{r}' \quad . \quad (\text{E6})$$

Calculating the death rate per unit volume by multiplying with  $\gamma^{\mathcal{D}}$  and integrating over all  $\mathbf{r}$  and  $t$  must give the mass in  $d^3\mathbf{r}'$  that will not be regenerated in the future, i.e.,  $\chi^{0\uparrow}(\mathbf{r}', t') d^3\mathbf{r}'$ . Dividing both sides by the starting mass  $\chi(\mathbf{r}', t') d^3\mathbf{r}'$  and defining the fraction  $f^{0\uparrow}(\mathbf{r}', t') \equiv \chi^{0\uparrow}(\mathbf{r}', t')/\chi(\mathbf{r}', t')$  gives

$$f^{0\uparrow}(\mathbf{r}', t') = \int dt \int d^3\mathbf{r} \gamma^{\mathcal{D}}(\mathbf{r}) G_{\mathcal{F}}(\mathbf{r}, t|\mathbf{r}', t') \quad . \quad (\text{E7})$$

Applying the time-reversed adjoint operator  $(\partial_{t'} + \tilde{\mathcal{F}}_{\mathbf{r}'})$  from the left gives (9).

#### E4 Recursion for the fraction regenerated $m$ times in the future

To construct the recursion equation for  $f^{m\uparrow}$ , it is useful to explicitly write the regeneration operator in terms of its integration kernel

$$(\mathcal{R}\chi)(\mathbf{r}) = \int d^3\mathbf{r}'' K_{\mathcal{R}}(\mathbf{r}|\mathbf{r}'')\chi(\mathbf{r}'', t') . \quad (\text{E8})$$

We again consider the concentration at  $(\mathbf{r}, t)$  resulting from the mass of DFe in  $d^3\mathbf{r}'$  at  $t'$  that was not regenerated since  $t'$ ,  $\mathcal{X}^{0\uparrow}(\mathbf{r}, t|\mathbf{r}', t')$   $d^3\mathbf{r}'$ , as defined by (E6). The rate with which  $\mathcal{X}^{0\uparrow}(\mathbf{r}, t|\mathbf{r}', t')$   $d^3\mathbf{r}'$  is regenerated (i.e., remineralized) in volume  $d^3\mathbf{r}''$  is given by  $\int d^3\mathbf{r} K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r})\mathcal{X}^{0\uparrow}(\mathbf{r}, t|\mathbf{r}', t')$   $d^3\mathbf{r}'$  (where  $d^3\mathbf{r}'$  is not integrated over) and the fraction of this rate that will get regenerated a further  $m$  times in the future is given by

$$f^{m\uparrow}(\mathbf{r}'') \int d^3\mathbf{r} K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r})G_{\mathcal{F}}(\mathbf{r}, t|\mathbf{r}', t')\chi(\mathbf{r}', t')d^3\mathbf{r}' . \quad (\text{E9})$$

Integrating this rate over all  $\mathbf{r}''$  and  $t$ , we must recover the fraction of the initial mass at  $t'$  that will be regenerated exactly  $m + 1$  times in the future. Hence we have

$$f^{(m+1)\uparrow}(\mathbf{r}')\chi(\mathbf{r}', t')d^3\mathbf{r}' = \int dt \int d^3\mathbf{r} \int d^3\mathbf{r}'' f^{m\uparrow}(\mathbf{r}'') \times K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r})G_{\mathcal{F}}(\mathbf{r}, t|\mathbf{r}', t')\chi(\mathbf{r}', t')d^3\mathbf{r}' . \quad (\text{E10})$$

10 Dividing both sides by the initial mass  $\chi(\mathbf{r}', t')d^3\mathbf{r}'$  and with the adjoint regeneration operator  $\tilde{\mathcal{R}}$  defined through  $(\tilde{\mathcal{R}}f)(\mathbf{r}) = \int d^3\mathbf{r}'' f(\mathbf{r}'')K_{\tilde{\mathcal{R}}}(\mathbf{r}|\mathbf{r}'')$ , where  $K_{\tilde{\mathcal{R}}}(\mathbf{r}|\mathbf{r}'') \equiv K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r})$ , we have

$$f^{(m+1)\uparrow}(\mathbf{r}') = \int dt \int d^3\mathbf{r}'' \int d^3\mathbf{r} G_{\tilde{\mathcal{F}}}(\mathbf{r}', t'|\mathbf{r}, t) \times K_{\tilde{\mathcal{R}}}(\mathbf{r}|\mathbf{r}'')f^{m\uparrow}(\mathbf{r}'') . \quad (\text{E11})$$

Note that (E11) propagates the fraction to be regenerated  $m$  times through one regeneration (modelled as instantaneous) and through the adjoint flow backward in time to the fraction that will be regenerated  $(m + 1)$  times. Applying  $(-\partial_{t'} + \tilde{\mathcal{F}})$  from the left gives the recursion equation (10). Note that this recursion equation can be written as  $f^{m\uparrow} = (\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}})^m f^{0\uparrow}$ . Using (9)

15 and regrouping the factors of  $\tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$ , this recursion relation can be re-written as

$$f^{m\uparrow} = \tilde{\mathcal{F}}^{-1}\tilde{\mathcal{A}}^m\gamma^{\mathcal{D}} , \quad (\text{E12})$$

where  $\tilde{\mathcal{A}} \equiv \tilde{\mathcal{F}}^{-1}\tilde{\mathcal{R}}$  as in Appendix B, with adjoint  $\tilde{\mathcal{A}} = \tilde{\mathcal{R}}\tilde{\mathcal{F}}^{-1}$ . Equation (E12) is the analog of [Equation–Eq. \(B1\)](#) for the time-reversed adjoint problem.

Using similar techniques as in Appendix B, it follows readily that  $\sum_{m=0}^{\infty} f^{m\uparrow} = f^{\uparrow} = 1$ , as must be the case. Equation (11) for  $\bar{m}$  is also derived analogously to  $\bar{n}$  in Appendix B.

## Appendix F: Relation between export and $\bar{m}$

Here we calculate the steady-state globally integrated export that results from a steady injection with source  $s(\mathbf{r}')$  (in mol Fe  $\text{m}^{-3} \text{s}^{-1}$ ) in some volume  $\text{d}^3\mathbf{r}'$  during  $\text{d}t'$ . Propagating the initial DFe mass  $s(\mathbf{r}')\text{d}^3\mathbf{r}'\text{d}t'$  with  $\mathcal{H}$  results in concentration  $G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t')s(\mathbf{r}')\text{d}^3\mathbf{r}'\text{d}t'$  at  $(\mathbf{r}, t)$ . This DFe is instantly regenerated at  $\mathbf{r}''$  with rate  $\int \text{d}^3\mathbf{r} K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r})G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t')s(\mathbf{r}')\text{d}^3\mathbf{r}'\text{d}t'$ . Integrating over all  $\mathbf{r}''$  and  $t$  must give the globally integrated export  $\Phi(\mathbf{r}')\text{d}^3\mathbf{r}'\text{d}t'$  that is due to the initial injection of the mass  $s(\mathbf{r}')\text{d}^3\mathbf{r}'\text{d}t'$ , and further dividing by  $\text{d}^3\mathbf{r}'\text{d}t'$  gives

$$\Phi(\mathbf{r}') = \int \text{d}^3\mathbf{r}'' \int \text{d}^3\mathbf{r} K_{\mathcal{R}}(\mathbf{r}''|\mathbf{r}) \int \text{d}t G_{\mathcal{H}}(\mathbf{r}, t|\mathbf{r}', t')s(\mathbf{r}') . \quad (\text{F1})$$

The globally integrated export per unit source injection at  $\mathbf{r}'$  is thus  $g(\mathbf{r}') \equiv \Phi(\mathbf{r}')/s(\mathbf{r}')$ . Dividing (F1) by  $s(\mathbf{r}')$  and applying  $(-\partial_{t'} + \tilde{\mathcal{H}})$  gives

$$\tilde{\mathcal{H}}g = \tilde{\mathcal{R}}f^\dagger , \quad (\text{F2})$$

which is the Green function that propagates a unit source to globally integrated export. Comparison with (11) shows that  $g = \bar{m}$  from which (13) of the main text follow by replacing our generic source  $s(\mathbf{r}')$  with one of the sources  $s_k(\mathbf{r}')$ . Equation (14) for the global phosphorus export due to DFe injection at  $\mathbf{r}'$  can be derived in exactly the same way by using regeneration operator  $\mathcal{R}^{\text{P}}$  instead of  $\mathcal{R}$ .

*Acknowledgements.* Benoît Pasquier acknowledges support from the Government of Monaco, the Scientific Centre of Monaco, the Frères Louis et Max Principale Foundation, the Cuomo Foundation, J. K. Moore (DOE Grant DE-SC0016539 and NSF Grant 1658380) and F. Primeau (NSF Grant 1658380). Mark Holzer acknowledges a UNSW Goldstar award. The state estimates of the iron cycle used here are available from the authors on request.

## References

- Achterberg, E. P., Steigenberger, S., Marsay, C. M., LeMoigne, F. A. C., Painter, S. C., Baker, A. R., Connelly, D. P., Moore, C. M., Tagliabue, A., and Tanhua, T.: Iron Biogeochemistry in the High Latitude North Atlantic Ocean, *Scientific Reports*, 8, 1283, <https://doi.org/10.1038/s41598-018-19472-1>, <https://doi.org/10.1038/s41598-018-19472-1>, 2018.
- 5 Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochemical Cycles*, 20, GB2017, <https://doi.org/10.1029/2005GB002591>, <http://dx.doi.org/10.1029/2005GB002591>, 2006.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale Iron Enrichment Experiments 1993–2005: Synthesis and Future Directions, *Science*, 315, 612–617, <https://doi.org/10.1126/science.1131669>, 2007.
- 10 Boyd, P. W., Ellwood, M. J., Tagliabue, A., and Twining, B. S.: Biotic and abiotic retention, recycling and remineralization of metals in the ocean, *Nature Geoscience*, 10, 167 EP –, <https://doi.org/10.1038/ngeo2876>, 2017.
- Conway, T. M. and John, S. G.: Quantification of dissolved iron sources to the North Atlantic Ocean, *Nature*, 511, 212 EP –, <http://dx.doi.org/10.1038/nature13482>, 2014.
- 15 de Baar, H., Gerringa, L., Laan, P., and Timmermans, K.: Efficiency of carbon removal per added iron in ocean iron fertilization, *Marine Ecology Progress Series*, 364, 269–282, <https://doi.org/10.3354/meps07548>, <https://doi.org/10.3354/meps07548>, 2008.
- DeVries, T. and Holzer, M.: Radiocarbon and helium isotope constraints on deep-ocean ventilation and mantle-<sup>3</sup>He sources, *Journal of Geophysical Research – Oceans* (submitted Oct. 2018), 2019.
- DeVries, T., Primeau, F., and Deutsch, C.: The sequestration efficiency of the biological pump, *Geophys. Res. Lett.*, 39, L13601, <https://doi.org/10.1029/2012GL051963>, 2012.
- 20 Dutkiewicz, S., Follows, M. J., Heimbach, P., and Marshall, J.: Controls on ocean productivity and air-sea carbon flux: An adjoint model sensitivity study, *Geophys. Res. Lett.*, 33, L02603, <https://doi.org/10.1029/2005GL024987>, 2006.
- Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: A missing source for global budgets, *Geophys. Res. Lett.*, 31, <https://doi.org/10.1029/2004GL020216>, L12307, 2004.
- 25 Fitzsimmons, J. N., John, S. G., Marsay, C. M., Hoffman, C. L., Nicholas, S. L., Toner, B. M., German, C. R., and Sherrell, R. M.: Iron persistence in a distal hydrothermal plume supported by dissolved–particulate exchange, *Nature Geoscience*, 10, 195 EP –, <http://dx.doi.org/10.1038/ngeo2900>, 2017.
- Frants, M., Holzer, M., DeVries, T., and Matear, R.: Constraints on the Global Marine Iron Cycle from a Simple Inverse Model, *Journal of Geophysical Research: Biogeosciences*, 121, 28–51, <https://doi.org/10.1002/2015jg003111>, 2016.
- 30 Galbraith, E. D., Gnanadesikan, A., Dunne, J. P., and Hiscock, M. R.: Regional Impacts of Iron-Light Colimitation in a Global Biogeochemical Model, *Biogeosciences*, 7, 1043–1064, <https://doi.org/10.5194/bg-7-1043-2010>, 2010.
- Holzer, M. and Hall, T. M.: Transit-time and tracer-age distributions in geophysical flows, *J. Atmos. Sci.*, 57, 3539–3558, [https://doi.org/10.1175/1520-0469\(2000\)057<3539:TTATAD>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<3539:TTATAD>2.0.CO;2), 2000.
- Holzer, M., Primeau, F. W., DeVries, T., and Matear, R.: The Southern Ocean silicon trap: Data-constrained estimates of regenerated silicic acid, trapping efficiencies, and global transport paths, *Journal of Geophysical Research: Oceans*, 119, 313–331, <https://doi.org/10.1002/2013JC009356>, 2014.
- 35

- Holzer, M., Frants, M., and Pasquier, B.: The age of iron and iron source attribution in the ocean, *Global Biogeochem. Cycles*, <https://doi.org/10.1002/2016GB005418>, 2016.
- Kirchman, D. L.: Microbial ferrous wheel, *Nature*, 383, 303 EP –, <https://doi.org/10.1038/383303a0>, 1996.
- Luo, C., Mahowald, N., Bond, T., Chuang, P. Y., Artaxo, P., Siefert, R., Chen, Y., and Schauer, J.: Combustion iron distribution and deposition, *Global Biogeochemical Cycles*, 22, <https://doi.org/10.1029/2007GB002964>, GB1012, 2008.
- 5 Maldonado, M. T., Strzepek, R. F., Sander, S., and Boyd, P. W.: Acquisition of iron bound to strong organic complexes, with different Fe binding groups and photochemical reactivities, by plankton communities in Fe-limited subantarctic waters, *Global Biogeochemical Cycles*, 19, GB4S23, <https://doi.org/10.1029/2005GB002481>, 2005.
- Martin, J. W., Knauer, G. A., Karl, D. M., and Broenkow, W. W.: VERTEX: Carbon cycling in the NE Pacific, *Deep-Sea Research*, 34, 267–285, 1987.
- 10 Moore, J. K. and Braucher, O.: Sedimentary and mineral dust sources of dissolved iron to the world ocean, *Biogeosciences*, 5, 631–656, <https://doi.org/10.5194/bg-5-631-2008>, 2008.
- Pasquier, B. and Holzer, M.: The plumbing of the global biological pump: Efficiency control through leaks, pathways, and time scales, *Journal of Geophysical Research: Oceans*, 121, 6367–6388, <https://doi.org/10.1002/2016JC011821>, 2016.
- 15 Pasquier, B. and Holzer, M.: Inverse-model estimates of the ocean’s coupled phosphorus, silicon, and iron cycles, *Biogeosciences*, 14, 4125–4159, <https://doi.org/10.5194/bg-14-4125-2017>, <https://www.biogeosciences.net/14/4125/2017/>, 2017.
- Primeau, F. W., Holzer, M., and DeVries, T.: Southern Ocean nutrient trapping and the efficiency of the biological pump, *J. Geophys. Res.*, 118, 2547–2564, <https://doi.org/10.1002/jgrc.20181>, 2013.
- Qin, X., Menviel, L., Gupta, A. S., and Sebillie, E.: Iron sources and pathways into the Pacific Equatorial Undercurrent, *Geophysical Research Letters*, 43, 9843–9851, <https://doi.org/10.1002/2016GL070501>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL070501>, 2016.
- 20 Rafter, P. A., Sigman, D. M., and Mackey, K. R. M.: Recycled iron fuels new production in the eastern equatorial Pacific Ocean, *Nature Communications*, 8, 1100, <https://doi.org/10.1038/s41467-017-01219-7>, <https://doi.org/10.1038/s41467-017-01219-7>, 2017.
- Raven, J. A.: The iron and molybdenum use efficiencies of plant growth with different energy, carbon and nitrogen sources, *New Phytologist*, 109, 279–287, <https://doi.org/10.1111/j.1469-8137.1988.tb04196.x>, <http://dx.doi.org/10.1111/j.1469-8137.1988.tb04196.x>, 1988.
- Strzepek, R. F., Maldonado, M. T., Higgins, J. L., Hall, J., Safi, K., Wilhelm, S. W., and Boyd, P. W.: Spinning the “Ferrous Wheel”: The importance of the microbial community in an iron budget during the FeCycle experiment, *Global Biogeochemical Cycles*, 19, <https://doi.org/10.1029/2005GB002490>, 2005.
- Tagliabue, A., Mtshali, T., Aumont, O., Bowie, A. R., Klunder, M. B., Roychoudhury, A. N., and Swart, S.: A global compilation of dissolved iron measurements: focus on distributions and processes in the Southern Ocean, *Biogeosciences*, 9, 2333–2349, <https://doi.org/10.5194/bg-9-2333-2012>, 2012.
- Tagliabue, A., Williams, R. G., Rogan, N., Achterberg, E. P., and Boyd, P. W.: A ventilation-based framework to explain the regeneration-scavenging balance of iron in the ocean, *Geophysical Research Letters*, 41, 7227–7236, <https://doi.org/10.1002/2014GL061066>, 2014GL061066, 2014.
- 35 Tagliabue, A., Aumont, O., DeAth, R., Dunne, J. P., Dutkiewicz, S., Galbraith, E., Misumi, K., Moore, J. K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., and Yool, A.: How Well do Global Ocean Biogeochemistry Models Simulate Dissolved Iron Distributions?, *Global Biogeochem. Cycles*, 30, 149–174, <https://doi.org/10.1002/2015gb005289>, 2016.

Twining, B. S., Nodder, S. D., King, A. L., Hutchins, D. A., LeClerc, G. R., DeBruyn, J. M., Maas, E. W., Vogt, S., Wilhelm, S. W., and Boyd, P. W.: Differential remineralization of major and trace elements in sinking diatoms, *Limnol. Oceanogr.*, 59, 689–704, <https://doi.org/10.4319/lo.2014.59.3.0689>, 2014.

5 Zeebe, R. E. and Archer, D.: Feasibility of ocean fertilization and its impact on future atmospheric CO<sub>2</sub> levels, *Geophysical Research Letters*, 32, L09703, <https://doi.org/10.1029/2005GL022449>, <http://dx.doi.org/10.1029/2005GL022449>, 2005.