



Stand age and species composition effects on surface albedo in a mixedwood boreal forest

Mohammad Abdul Halim^{1,2}, Han Y. H. Chen³, Sean C. Thomas¹

5 ¹University of Toronto, Faculty of Forestry, 33 Willcocks Street, M5S 3B3, ON, Canada

²Shahjalal University of Science and Technology, Dept. of Forestry and Environmental Science, Sylhet-3114, Bangladesh

³Lakehead University, Natural Resources Management, 955 Oliver Road, Thunder Bay, P7B 5E1, ON, Canada

Correspondence to: Mohammad Abdul Halim (abdul.halim@mail.utoronto.ca)

10

Abstract. Surface albedo is one of the most important processes governing climate forcing in the boreal forest and is directly affected by management activities such as harvesting and natural disturbances such as forest fires. Empirical data on effects of these disturbances on boreal forest albedo are sparse. We conducted ground-based measurements of surface albedo from a series of instrument towers over four years in a replicated chronosequence of mixedwood boreal forest sites differing in stand age (year since disturbance) in both post-harvest and post-fire stands. We investigated the effects of stand age, canopy height, tree species composition, and ground vegetation cover on surface albedo through stand development. Our results indicate that winter and spring albedo values were 63 and 24 % higher, respectively, in post-harvest stands than in post-fire stands. Winter and summer albedos saturated at ~ 50 years of stand age in both post-harvest and post-fire stands. Albedo differences between post-harvest and post-fire stands were most pronounced during winter and spring in young stands (0–19 years post-disturbance). The proportion of deciduous broadleaf species showed a strong positive relationship with seasonal albedo in both post-harvest and post-fire stands. Given that stand composition in mixedwood boreal forests generally shows a gradual replacement of deciduous trees by conifers, our results suggest that successional changes in species composition are likely a key driver of age-related patterns in albedo. Our findings also suggest the efficacy of increasing the proportion of deciduous broadleaf species as a silvicultural option for climate-friendly management of boreal forest.

15

20

25

Keywords

Stand age, species composition, canopy height, ground vegetation cover, harvesting, forest fire, succession, surface albedo, boreal forest.

30

1 Introduction

Surface albedo, the fraction of incoming solar energy reflected from the surface in all directions, is one of the most important biophysical factors affecting both local and global climates. In boreal forest the magnitude of albedo-related forcing on climate is even more important than in other ecosystems because of snow-related feedbacks, low sensible heat flux, and the relative stability of the atmospheric temperature profile (Bright et al., 2015a; Hansen et al., 2005). Even though albedo is increasingly used as an important state variable in climate models (Brown and Caldeira, 2017; Bala et al., 2007; Betts, 2000), forest disturbance effects on net radiative forcing due to local albedo changes and related feedbacks with regional/global mean surface temperature remain highly uncertain (Bright et al., 2015a; Lee et al., 2011). Harvest and fire suppression may differ substantially in their effects on albedo, but empirical data on albedo responses to disturbance type remain particularly sparse.

35



40

Following disturbance events, albedo of boreal forests is expected to change with stand age due to changing surface properties, and forest structure and composition. Age-related stand structural attributes (e.g., tree species composition, leaf area index [LAI], canopy height, and ground vegetation cover) can substantially influence surface albedo of a stand throughout the year. Studies have generally found higher albedos in young stands than in mature stands in the boreal forest (Bright et al., 2015a; Kuusinen et al., 2014; Amiro et al., 2006b), and it has been suggested that albedo stabilizes ~ 25 years after a disturbance event (during the ‘stem exclusion’ phase in the ecological succession trajectory). Early in stand development boreal mixedwood forests are commonly dominated by deciduous broadleaf species (Madoui et al., 2015; Brassard and Chen, 2010; Johnstone et al., 2010), which have higher leaf and canopy reflectance than conifers (Lukeš et al., 2013a; Linacre, 2003), contributing to high summer albedo in young stands (Lukeš et al., 2013b; Betts and Ball, 1997). These deciduous species shed leaves in the winter, which increases canopy openness (lowers LAI) and allows snow albedo to dominate, contributing to the high winter albedo in young stands. Available data suggest that at this stage both LAI and ground vegetation cover usually increase with stand age, depending on site quality and silvicultural practices (Amiro et al., 2006b; Uotila and Kouki, 2005). Low LAI can increase canopy background reflectance both in snow-covered and snow-free conditions, and thus can contribute to the high albedos in young stands (Amiro et al., 2006b). LAI effects on albedo in young stands may be highly modulated by ground vegetation cover in the summer, but probably not much in the winter as ground vegetation is generally leafless or covered with snow (Kuusinen et al., 2015; Lukeš et al., 2013b; Betts and Ball, 1997). In conjunction with other factors, surface albedo tends to decrease with increasing canopy height (Hovi et al., 2016; Linacre, 2003). In the later stages of stand development, albedo is expected to saturate non-linearly as conifers dominate the stand and canopy cover and stand attributes change gradually, but data describing this pattern remain sparse (Amiro et al., 2006b).

Harvesting and fire are the major stand-replacing disturbances in the boreal forest (Brassard et al., 2008). These disturbances may differentially affect surface albedo of post-disturbance stands in complex ways by altering ground surface spectral properties, species composition, and stand structure (Lukeš et al., 2013b; Liu et al., 2005), but field data directly addressing this issue are essentially limited to a single study in Europe (Kuusinen et al., 2016). Structure and composition of post-fire stands are generally more heterogenous than post-harvest stands; for example, post-fire stands are more likely to show a bimodal vertical structure and higher proportion of deciduous broadleaf species during early stand development stages (Brassard and Chen, 2010; Johnstone et al., 2010). Charcoal residues may also strongly reduce albedo in snow-free conditions in the first years following fire disturbances (Amiro et al., 2006b). Both charcoal effects and stand heterogeneity might be expected to reduce surface albedo in post-fire stands relative to post-harvest stands. However, the magnitude of this difference in surface albedo might be less than expected due to the presence of legacy charcoal from historical fires in post-harvest stands (Hart and Luckai, 2013). Immediately after harvesting, albedo of a post-harvest stand can also be reduced because of the presence of coarse woody debris (CWD) and high soil moisture content (Linacre, 2003). In the years following a disturbance event, CWD might be expected to further reduce albedo due to decomposition processes (Brassard and Chen, 2008) and plant colonization (Kumar et al., 2018).

Despite the important role of stand age as a determinant of boreal forest albedo, field measurements are scarce (Kuusinen et al., 2014) and particularly limited for early stand ages that show high variability in surface properties (Bright et al., 2013). This has contributed to poorly constrained estimates of the local albedo changes on net global radiative forcing (Bright et al., 2015a). Data limitations are evident in global climate modeling studies (e.g., Bala et al. 2007, Betts 2000). To estimate



the net change in surface temperature as a function of albedo change from deforestation in boreal forests, these models have used a ‘biome replacement’ approach (replacing boreal forests with grassland or agricultural land cover types) and approximated boreal forests’ albedo as a single value from mature stands (~ 60-year old). Early stand dynamics is reported to determine which mechanism, albedo vs. carbon storage, dominates the net forcing for the boreal forest (Kirschbaum et al., 2011). Such unrealistic simplifications in climate models that do not explicitly consider stand age and successional effects on albedo will likely result in strongly biased estimations of boreal forests’ albedo over the rotation (harvesting/fire) period (Bright et al., 2018).

Given the complex nature and limited understanding of underlying processes (Lukeš et al., 2013b), it is important to conduct field studies to better quantify forest albedo in relation to stand age and disturbance type in boreal forests. Deeper understanding of the local mechanisms that account for variation in albedo will not only enhance global climate models (e.g., by improving the land-surface model: Bright et al., 2018), but also help to design climate-friendly silvicultural practices (Astrup et al., 2018; Matthies and Valsta, 2016; Bright et al., 2015a). In the present study, we set up micrometeorological towers with pyranometers in a replicated chronosequence of post-harvest and post-fire sites to study stand age, disturbance type, and species composition effects on albedo in a mixedwood boreal forest of northwestern Ontario, Canada. We hypothesized: (1) that post-fire stands would show lower albedo values than post-harvest stands as a consequence of stand composition, legacy structures, and fire residues; (2) that all stands would approach albedo values similar to mature stands within ~ 25 years, soon after crown closure; and (3) that stands with higher dominance of deciduous broadleaf species would show higher albedo than conifer-dominated stands, with this effect being most pronounced under snow-covered conditions.

2 Materials and Methods

2.1 Study area

The study was conducted in the boreal forest of the Lake Nipigon region (49.55° N and 89.5° W), Ontario, Canada, approximately 200 km north of Thunder Bay. A series of circular (10-m radius) chronosequence plots were established in the post-harvest (full-tree harvest) and post-fire stands in the study area. Three plots were set up in each of three cutblocks harvested in 1998, 2006, and 2013. Recent (2013) post-fire stands were not present, so we set up three plots only in post-fire stands dating from 1998 and 2006 fire events (Fig. 1). Replicate stands were spatially interspersed to the extent feasible. For each of the 15 plots, albedo and stand attributes (stand age, percentage of deciduous broadleaf species, canopy height, and percentage of ground vegetation cover) were measured from July 2013 to June 2017.

The mesic mixedwood study area is dominated by jack pine (*Pinus banksiana* Lamb.), black spruce (*Picea mariana* (Mill.) BSP), white spruce (*P. glauca* (Moench) Voss), trembling aspen (*Populus tremuloides* Michx.), eastern white cedar (*Thuja occidentalis* L.), balsam fir (*Abies balsamea* (L.) Mill.), and paper birch (*Betula papyrifera* Marsh.) (Chen and Popadiouk, 2002). In study plots over the study period canopy height ranged from 0–7.7 m, ground vegetation cover ranged from 1.8–96.7 %, LAI ranged from 0–2.1, and the proportion of deciduous broadleaf basal area ranged from 10.6–100 %. The study area has an average elevation of 416 m a.s.l. The soil is a moderately deep Brunisol (coarse loamy texture) with 1–15 cm thick organic layer (LFH). The area remains snow covered for 5–6 months with an average snow depth of 9 cm (Environment Canada, 2018; Sims et al., 1997) and the mean annual air temperature of the study plots was –1.1 °C (Halim and Thomas, 2018).



120 2.2 Experimental setup

In the center of each circular plot a pair of upward- and downward-facing pyranometers (Silicon Pyranometer; Onset, Massachusetts, USA; measurement range 0–1280 Wm⁻² over a spectral range of 300–1100 nm, accuracy ± 5 %, resolution 1.25 Wm⁻²) were set up on a mast 3.5 m above the canopy (above the ground for 2013 post-harvest stands) to measure incident and reflected solar radiation every 10 minute. Instrument masts consisted of extendible galvanized steel poles and
125 were set in concrete bases and guyed to mitigate instrument sway. At least once a year pyranometer heads were cleaned and realigned to make sure they were normal to the ground. Average daily albedo was calculated as the ratio of daily total incident and reflected radiation for each plot. The average daily albedo was used to calculate average monthly albedo, which was finally used to calculate mean seasonal albedo for each year in each plot.

130 In addition to albedo, winter (December–February)/spring (March–May) and summer (June–August)/fall (September–November) proportion of deciduous broadleaf area (%), canopy height (m), and ground vegetation cover (%) were measured every year in late October and early July, respectively, in each plot. The proportion of deciduous broadleaf species (%) were determined for trees with diameter at breast height ≥ 5 cm and height > 1 m. Canopy height was determined as the mean height of all trees sampled; the young stands sampled were at stages of development prior to and
135 just after canopy closure, so essentially all trees were “canopy dominants”. The proportion of deciduous broadleaf species of a plot was calculated as the ratio of basal area of the deciduous species to the total basal of area of the plot. In each plot four 1-m² subplots were set up and percent ground vegetation cover was determined visually (Kumar et al., 2018). Stand age was determined as the time (year) since last disturbance (fire/harvesting) for each plot. Fire maps (from the Ontario Ministry of Natural Resources, Canada) and forest management plans (from Resolute Forest Products, Canada) were used to verify type
140 and year of disturbances.

2.3 Sources of secondary data

Since we did not have recent post-fire stands (0–6-year old) in the study area, we used secondary albedo data from studies in post-fire boreal forests with similar stand characteristics in Alaska (USA), Saskatchewan, and Manitoba (Canada) (Fig.
145 1). We also used secondary albedo data for old stands (> 70 years) from these sites along with primary data to develop regression models for both post-fire and post-harvest stands. Here we assumed that at this stage of stand development, there is negligible difference in stand attributes (e.g., species composition, height, LAI) between post-fire and post-harvest stands (Moussaoui et al., 2016). We did not use satellite-based albedo data as secondary sources as they tend to diverge from field measurements depending on a number of factors including stand age, latitude, and cloud cover effects (Halim and Thomas,
150 2017; Bright et al., 2015b).

Data for Alaskan sites were retrieved from Chambers and Chapin (2002) and Liu et al. (2005), for Saskatchewan sites from Amiro et al. (2006a), and for Manitoban sites from Amiro et al., (2006b) by digitizing data points from relevant figures using the WebPlotDigitizer software (Rohatgi, 2018). These stands were dominated by jack pine and black spruce with
155 some intermixing of trembling aspen. Pyranometers were located in Alaskan sites at 7.8–8 m, in Saskatchewan sites at 18–20 m, and in Manitoban sites at 6 m heights. There was no detailed information on how proportions of broadleaf deciduous species were calculated for these sites; however, we assumed they were basal-area based. Detailed description of the study areas and methods can be found in the respective articles.



160 2.4 Measurements of ground surface reflectance

To examine effects of disturbance type on ground surface reflectance, three soil samples (top 10 cm including LFH layer, surface area 78.5 cm²) from each plot were collected in fall 2017 to measure the ground surface reflectance. Samples were all collected within a two-day precipitation-free period, and were brought to the lab in airtight packaging without disturbing the top surface. A spectrometer (SD 2000; Ocean Optics, Florida, USA; measurement spectral range 338.7–1001.8 nm) equipped with an integrating sphere was used to measure their top-surface-specular-included (diffused and direct) reflectance values. As there were no recent post-fire stands in the study area, we collected charcoal samples (of twigs, branches, barks, and stems) from the forest floor of a jack pine dominated post-fire (fire occurred in 2011) stand in summer 2015 from near the Musselwhite mine (52.61° N and 90.37° W), Ontario, Canada. Every sample was measured ten times in ten different locations, and each measurement was performed by scanning 10 times (with Boxcar width 5 and 100 millisecond integration time) to get an average reflectance for each location of a sample. Details of the spectrometer and integrating sphere used can be found in the Materials and Methods section of Baltzer and Thomas (2005). Forest floor reflectance values from the Musselwhite stand (4-year old) were compared to soil sample reflectance values from recent (2013) post-harvest stands (4-year old). For older stands (1998 and 2006 post-harvest and post-fire stands), soil sample reflectance data were compared using samples from the main study plots.

175

2.5 Data analysis

Robust t-tests (Wilcox, 2016) were used to compare mean differences in ground surface reflectance (in visible [400–700 nm] and near-infrared [> 700 –100 nm] spectral bands) and seasonal albedo between post-harvest and post-fire stands. Mean seasonal albedo values of post-harvest and post-fire stands were also compared using Analysis of Covariance (ANCOVA) controlling for the effects of stand age as a covariate. Secondary albedo data for 0–6-year-old post-fire stands were only available for winter and summer seasons. Therefore, in the t-tests (and in ANCOVA) for winter and summer albedo, data from 0–19-year old post-harvest and post-fire stands were used. For spring and fall, albedo data from recent (0–6-year-old) post-harvest stands were omitted (since there were no data from post-fire stands for these seasons), and data from 7–19-year-old stands were used to make the comparisons unbiased. Secondary data from old stands (> 70 years) were not used in the t-tests/ANCOVA.

185

Generalized linear models (GLMs) with the log-linked gaussian family (additive-observation-error model with constant variance) were found to be the best fitted to model seasonal albedo as a function of stand attributes (stand age, proportion of deciduous broadleaf species, canopy height, ground vegetation cover, and their interactions) for both post-harvest and post-fire stands. Best models were chosen using an AIC-based stepwise algorithm. Asymptotic chi-square statistics based on deviance were calculated for each best-fit model to test if the model was significantly better than its counterpart null model. In fall months, some stand attributes were only nonlinearly (double exponentially) related to albedo. To avoid model complexity, for each of these fall attributes a separate nonlinear model was fitted, and for other attributes GLMs with identity-linked gaussian family were found to be the most suitable. The Δ AIC for each best-fit model is calculated as its AIC difference with the corresponding null model (AIC of best-fit model – AIC of corresponding null model). Sample-size corrected AIC values were used in all cases.

195

Data were analysed using the R platform (R Core Team, 2018) and graphs were prepared using the ‘ggplot2’ package (Wickham, 2016). Robust t-tests were done by 10,000 bootstrapped samples considering mean as an estimator for group



200 comparison, and implemented by the *pb2gen* function of the WRS2 R-package (Mair and Wilcox, 2018). Adjusted R^2 values for GLMs were calculated using the *rsq* function of the R-package ‘rsq’ (Zhang, 2018).

3 Results

3.1 Seasonal albedo in post-harvest and post-fire stands

205 Albedo differences between post-harvest and post-fire stands varied among seasons. Albedo values in periods of the year with appreciable snow cover were significantly higher in post-harvest stands than in post-fire stands (for winter: 0.56 vs. 0.34, $p < 0.01$; for spring: 0.32 vs. 0.24, $p = 0.11$). Summer albedo values in post-harvest stands were also marginally higher in post-harvest stands ($p = 0.24$), and fall albedos were similar between disturbance types ($p = 0.73$) (Fig. 2). Considering stand age as a covariate, ANCOVA results also indicate higher albedo of post-harvest stands in winter ($p = 0.02$), spring ($p = 0.15$), summer ($p = 0.04$), and similar in fall ($p = 0.77$) compared to post-fire stands. Data also suggest higher variability
210 in albedo in post-harvest stands than in post-fire stands (Fig. 2).

3.2 Ground surface reflectance in post-harvest and post-fire stands

Specular-included reflectance measurements of ground surface samples suggest that differences in ground surface characteristics contribute to overall surface albedo in the study sites. Summer ground surface reflectance was generally higher in old stands (Fig. 3b) than in young stands (Fig. 3a) particularly in the 600–1000 nm range. Young (4-year old)
215 post-harvest stands showed significantly lower mean ground reflectance values (74.3 %, $p < 0.01$) in the visual spectrum (400–700 nm) and higher (32.3 %, $p < 0.01$) in the near-infrared spectrum (> 700–1000 nm) than those of young post-fire stands (Fig. 3a). Older (11- and 19-year old) post-harvest stands however showed higher mean ground reflectance in both visible (31.7%, $p < 0.01$) and near-infrared (4.6%, $p < 0.01$) spectra compared to post-fire stands (Fig. 3b).

3.3 Seasonal albedo in relation to stand attributes in post-harvest and post-fire stands

220

3.3.1 Winter albedo in post-harvest and post-fire stands

Results from the best-fit GLM ($p < 0.01$, adj. $R^2 = 0.97$) for post-harvest stands indicated that stand age, proportion of deciduous broadleaf species, canopy height, and interactions among these variables were significant predictors of winter albedo (Table 1). Stand age was related to winter albedo via an exponential decay model with a horizontal asymptote
225 ($\Delta AIC = -26.7$), and all estimated model parameters were significant (for 0.2 and 0.55: $p < 0.01$; for -0.06 : $p < 0.05$) (Fig. 4a). The proportion of deciduous broadleaf species (Fig. 6a) and canopy height (Fig. 7a) were also related to winter albedo via negative exponential models with horizontal asymptotes ($\Delta AIC = -6.7$ and -103.13 , respectively), and all estimated parameters for both models were significant ($p < 0.01$).

230 For post-fire stands the best-fit GLM ($p < 0.01$, adj. $R^2 = 0.79$) indicated that stand age, proportion of deciduous broadleaf species, and their interactions were significant predictors of winter albedo (Table 1). Stand age was related to winter albedo via an exponential decay model with a horizontal asymptote ($\Delta AIC = -34.6$), and all estimated model parameters were significant ($p < 0.01$) (Fig. 4b). Proportion of deciduous broadleaf species was related to winter albedo via a negative



235 exponential model with horizontal asymptote ($\Delta\text{AIC} = -15.8$), and all estimated model parameters were significant (for –
0.29: $p < 0.01$; for 1.02: $p = 0.16$; for 0.48: $p < 0.05$) (Fig. 6b).

3.3.2 Spring albedo in post-harvest and post-fire stands

240 For post-harvest stands the best-fit GLM ($p < 0.01$, adj. $R^2 = 0.99$) indicated that stand age, proportion of deciduous
broadleaf species, height, and the interaction of stand age and proportion of deciduous broadleaf species were significant
predictors of spring albedo (Table 1). Stand age (Fig. 5a) and canopy height (Fig. 7c) were related to spring albedo via
exponential decay models with horizontal asymptotes ($\Delta\text{AIC} = -15.1$ and -31.2 , respectively). Estimated parameters of
stand age-albedo (for 0.26: $p < 0.01$; for 0.72 and -0.72 : $p < 0.05$) and canopy height-albedo (for 0.16 and 0.33: $p < 0.01$;
for -1.84 : $p = 0.07$) models were likewise significant. The proportion of deciduous broadleaf species was related to spring
albedo via a negative exponential model ($\Delta\text{AIC} = -6.72$), and all estimated parameters were significant ($p < 0.01$) (Fig. 6c).

245 The best-fit GLM ($p < 0.01$, adj. $R^2 = 0.99$) for post-fire stands indicated that stand age and proportion of deciduous
broadleaf species were the only significant predictors of spring albedo (Table 1). Stand age (Fig. 5b) and proportion of
deciduous broadleaf species (Fig. 6d) were related to spring post-fire stand albedo via exponential negative growth models
($\Delta\text{AIC} = -7.0$ and -7.5 , respectively), and all estimated parameters for both models were significant ($p < 0.01$).

250

3.3.3 Summer albedo in post-harvest and post-fire stands

255 The best-fit GLM ($p < 0.01$, adj. $R^2 = 0.96$) for post-harvest stands indicated that stand age, proportion of deciduous
broadleaf species, ground vegetation cover and its interaction with stand age and proportion of deciduous broadleaf species
were significant predictors of summer albedo (Table 1). Stand age was related to summer albedo via a double exponential
model ($\Delta\text{AIC} = -79.4$), and all estimated model parameters were significant ($p < 0.01$) (Fig. 4c). The pattern described by
this function indicates a sharp peak in albedo with a maximum at 10–15 years of stand age. Proportion of deciduous
broadleaf species is related to summer albedo via a 3-parameter sigmoid model ($\Delta\text{AIC} = -54.6$), and all the estimated
parameters were significant ($p < 0.01$) (Fig. 6e). Ground vegetation cover was related to summer albedo via an exponential
model with a Gumbel distribution without a horizontal asymptote ($\Delta\text{AIC} = -25.8$), and all estimated parameters were
260 significant ($p < 0.01$) (Fig. 8e).

For post-fire stands the best-fit GLM ($p < 0.01$, adj. $R^2 = 0.93$) indicated that stand age, proportion of deciduous broadleaf
species, canopy height, and their interactions with stand age were significant predictors of summer albedo (Table 1). Stand
age (Fig. 4d) and canopy height (Fig. 7f) were related to summer post-fire stand albedo via exponential models with
265 Gumbel distributions with horizontal asymptotes ($\Delta\text{AIC} = -55.7$ and -4.0 , respectively). As in the case of post-harvest
stands, peak albedo was found at ~ 10 –15 years of stand age. All estimated parameters of stand age-albedo ($p < 0.01$) and
canopy height-albedo (for 0.23 and 0.64: $p < 0.05$; for 1.89 and 0.09: $p < 0.01$) models were significant. Proportion of
deciduous broadleaf species was related to summer albedo via a negative exponential growth model ($\Delta\text{AIC} = -6.8$), and all
estimated model parameters were significant ($p < 0.01$) (Fig. 6f).

270

3.3.4 Fall albedo in post-harvest and post-fire stands

The best-fit GLM ($p < 0.01$, adj. $R^2 = 0.94$) for post-harvest stands indicated that stand age, canopy height, ground
vegetation cover, and their interactions were significant predictors of fall albedo (Table 1). Proportion of deciduous



275 broadleaf species was also an important predictor that was modelled separately via a double exponential model (and was not added to the GLM to avoid modelling complexities) ($\Delta\text{AIC} = -0.9$); all estimated model parameters were significant (for 28.9 and 45.4: $p < 0.05$; for 67.6: $p < 0.01$) (Fig. 6g). Stand age (Fig. 5c) and ground vegetation cover (Fig. 8g) were related to albedo via exponential decay models with horizontal asymptotes ($\Delta\text{AIC} = -36.8$ and -28.38 , respectively), and all estimated parameters for both models were significant ($p < 0.01$). Canopy height was also related to albedo via a negative exponential model ($\Delta\text{AIC} = -11.2$), and all estimated parameters were significant ($p < 0.01$) (Fig. 7g).

280

To avoid modelling complexities, stand age and proportion of deciduous broadleaf species were fitted individually with fall albedo of post-fire stands (Table 1). Stand age was related to albedo via a double exponential model ($\Delta\text{AIC} = -3.1$), and all estimated model parameters were significant ($p < 0.01$) (Fig. 5d). Proportion of deciduous broadleaf species was related to albedo via a simple exponential model ($\Delta\text{AIC} = -25.4$), and all estimated model parameters were significant ($p < 0.01$) (Fig. 6h).

285

4 Discussion

Our results provide evidence for dramatic effects of disturbance type on albedo of boreal forest systems, with post-harvest stands showing much higher albedo values in winter and spring months than post-fire stands. Stands of both disturbance types also showed strongly age-dependent patterns in albedo; however, analyses suggest that post-disturbance changes are more gradual than anticipated, with dynamics continuing past stand closure, up to ~ 50 years post-disturbance. The proportion of deciduous species also had large effects on stand albedo – generally larger than stand age effects – showing a positive saturating response in all seasons and for both disturbance types.

290

4.1 Albedo in post-harvest and post-fire stands

295 Mean albedo in post-harvest stands was significantly higher than in post-fire stands in winter and spring, marginally higher in summer, and similar in fall (Fig. 2). A similar pattern in albedo differences was also observed when the stand age effects on albedo were statistically controlled. The magnitude of differences in winter and spring values (0.22 and 0.08, or 63% and 34% increases relative to post-fire values) is large – comparable to albedo differences observed between biomes (Stephens et al., 2015). During snow-covered seasons (winter and spring), charcoal residues in post-fire stands are usually covered with snow, and thus stand structure and composition act as dominant drivers of albedo (Lyons et al., 2008; Amiro et al., 2006b; Liu et al., 2005). Deciduous broadleaf species made up 37.8 % of basal area in post-fire stands and 55.4 % in post-harvest stands: the higher percentage of dark conifer leaves is expected to result in lower winter/spring albedos in post-fire stands compared to post-harvest stands (Betts and Ball, 1997). However, immediately after a stand-replacing fire, the presence of black carbon (charcoal and soot) in the snow can reduce early winter albedo and possibly enhance spring snowmelt by absorbing solar radiation (Qian et al., 2009; Conway et al., 1996). During late spring when snow cover is shallow, it is also likely that charred branches and stems protrude through the snow and reduce albedo. From an energy balance perspective, it is important to note that albedo differences in late spring may be less important as turbulent and latent fluxes likely dominate (Conway et al., 1996).

300

305

310 In snow-free seasons (summer and fall) the marginal differences in mean albedo between post-harvest and post-fire stands can partly be attributed to rapid recovery of ground vegetation in post-fire stands (0–5 years old) compared to post-harvest stands (Bartels et al., 2016), and to the vegetation covering dark charcoals in older (> 5 year) post-fire stands (Randerson et



al., 2006). Soon after a fire, the presence of early-successional plants (Johnstone et al., 2010) can increase surface albedo of post-fire stands because of their higher albedo relative to charcoal (Amiro et al., 2006b; Betts and Ball, 1997). This effect is expected to offset the albedo difference between post-harvest and post-fire stands. In the first year following disturbance events we might expect lower snow-free albedo in post-fire stands than post-harvest stands because of high charcoal occurrence on the soil surface (Lyons et al., 2008; Chambers and Chapin, 2002). However, our soil reflectance data indicate that soils from 4-year old post-fire stands unexpectedly showed significantly higher reflectance in the visible spectrum than did post-harvest stands (with the pattern reversed in the NIR spectrum) (Fig. 3a). Similar patterns in spectral response were recently observed in a biochar-amended agricultural soil relative to the control (Zhang et al., 2013). Soils from older post-harvest stands (11- and 19-year old), as expected, showed higher reflectance in the visible and NIR spectra compared to post-fire stands of similar age (Fig. 3b). Most post-harvest stands exhibited patches of charcoal in surface soils, presumably originating from historical fires (personal observations). The importance of “legacy” soil charcoal on surface albedo of harvested stands has not been considered previously to our knowledge. Charcoal reflectance is highly dependent on charring conditions (e.g., temperature, oxygen content) (Hudspith et al., 2015), and may possibly change with weathering; these processes require additional study in the context of albedo and surface energy balance.

Our results from both snow-covered and snow-free seasons strongly suggest that fire residues on the ground cannot explain the observed differences in albedo between post-harvest and post-fire stands. This result is consistent with the generalization that stand structure and composition are the main drivers of surface albedo and energy balance in the boreal forest (Amiro et al., 2006a).

4.2 Albedo convergence with stand age in post-harvest and post-fire stands

Compiled data for winter and summer albedos from post-harvest and post-fire stands indicate that surface albedo saturates at ~ 50 years of stand age in the boreal forest (Figs. 4a–d). This finding does not support our second hypothesis that albedo saturates after ~ 25 years of stand age (during the ‘stem exclusion’ phase). The rationale behind this hypothesis was that a high cover of deciduous species until ~ 25 years contributes to high albedo in both snow-covered (due to leaf shedding) and snow-free (due to high canopy albedo) seasons; after this age, deciduous species are replaced by conifers and albedo tends to saturate (Kuusinen et al., 2014; Amiro et al., 2006b). Studies using remote sensing techniques also suggest that albedo in both post-harvest and post-fire stands saturates at ~ 50 years after harvest/fire (Bright et al., 2015a; Kuusinen et al., 2014; Lyons et al., 2008; McMillan and Goulden, 2008), consistent with our findings. These results thus suggest that gradual changes in species composition through later stages of succession are an important driver of stand albedo. Stand structural features such as canopy height (in winter) and ground vegetation cover (in summer) usually increase with stand age (Bartels et al., 2016) and might additionally contribute to a gradual reduction in albedo (Hovi et al., 2016) after ~ 25 years (Table 1, Figs. 7 and 8).

The shape of best-fit curves for winter albedo vs. stand age (exponential decay) of post-harvest (Fig. 4a) and post-fire (Fig. 4b) stands are similar to other studies (Bright et al., 2015b; Kuusinen et al., 2014; Lyons et al., 2008; McMillan and Goulden, 2008; Amiro et al., 2006b); however our results diverge markedly for summer albedo. Our best-fit curves for summer albedo vs. stand age for both post-harvest and post-fire stands showed pronounced peaks in early albedo described by double exponential functions (Figs. 4c–d), whereas Amiro et al (2006b) described data with a negative linear relationship, and other remote sensing-based studies have used exponential decay curves (e.g., Kuusinen et al., 2014). However, Lyons et al. (2008) and Randerson et al. (2006) found summer albedo of post-fire stands were related to stand age



via a humped-shape curve, and albedo reached peak at ~ 20 years and gradually levelled off at ~ 50 years after fire, which
355 closely corresponds to our findings. We suggest that most prior studies with sparser or more noisy data sets may have
missed this early peak pattern. Immediately after fires and harvesting (because of high soil moisture, decaying CWD, legacy
charcoal etc.) the summer albedo of post-harvest and post-fire stands is expected to show a low albedo (also see section 4.1
and Fig. 3) which sharply increases as dark ground is covered with early successional pioneer species (Lyons et al., 2008;
Randerson et al., 2006; Amiro et al., 2006b; Betts and Ball, 1997). This sharp increase continues until ~ 20 years of stand
360 age but then decreases slowly until ~ 50 years and saturates – consistent with the patterns found in other seasons.

We did not have albedo data from late-seral post-harvest or post-fire stands for spring and fall. In post-harvest (Fig. 5a) and
post-fire (Fig. 5b) stands spring albedos did not show strong patterns with stand age, and the patterns were disturbance-
specific (exponential decrease vs. negative exponential growth, respectively). Results from Kuusinen et al. (2014), Lyons et
365 al. (2008), and Randerson et al. (2006) also suggest that patterns of spring albedo as a function of stand age can be
disturbance-specific. In post-harvest stands, Kuusinen et al. (2014) found that spring albedo was high immediately after
harvest and decreased exponentially until ~ 50 years and then saturated. However, in post-fire stands, Lyons et al. (2008)
and Randerson et al. (2006), found hump-shaped patterns with peak at ~ 10–15 years, and subsequent declines, similar to
the winter albedo pattern. As discussed in Section 4.1, disturbance-specific responses may partially be attributed to the
370 presence of black carbon (charcoal/soot) in snow immediately after fire, which can substantially reduce snow albedo (Qian
et al., 2009). Trends in fall albedo values with stand age in post-harvest (Fig. 5c) and post-fire (Fig. 5d) stands showed
stronger patterns than spring, but similar disturbance-specific responses. Immediately after harvesting fall albedo was high,
and exponentially decreased as stand age increased. Increased fall albedo in recent post-harvest stands may be due to
contributions to senescing leaves and to snow in the late fall (Amiro et al., 2006b; Liu et al., 2005). In contrast, fall albedo
375 immediately after fire was low (possibly because of charcoal or soot residues as discussed above), and increased with stand
age.

Albedo parameterization has been a classical problem in climate simulations (Bright et al., 2015a; Kuusinen et al., 2012; Qu
and Hall, 2007; Betts, 2000). Our findings indicate that there are important qualitative differences in the post-disturbance
380 albedo patterns between seasons in boreal forests. These differences need to be considered in albedo parameterization for
land surface models.

4.3 Deciduous broadleaf species as a key determinant of surface albedo in the post-harvest and post-fire stands

Our results indicate that the proportion of deciduous broadleaf species is a strong predictor of albedo irrespective of
385 disturbance type, and in most cases a better predictor than stand age (Figs. 4–6). Using remote sensing techniques Kuusinen
et al. (2014) also found that stand age alone was not consistently the best predictor of stand albedo in the boreal forest. We
found a similar mean model residual sum of errors for snow-covered seasons and snow-free seasons (Figs. 4–5 vs. Fig. 6),
indicating that the proportion of deciduous broadleaf species are similarly important in both cases. These findings strongly
support our third hypothesis that stands with a higher proportion of deciduous broadleaf species show higher albedo than
390 conifer-dominated stands, but also that this effect is pronounced under both snow-covered and snow-free conditions. Except
for fall post-fire stands, the relationship between albedo and proportion of deciduous broadleaf species approximated by an
exponential saturating curve in which albedo declined rapidly where the proportion of deciduous broadleaf species fell
below 25–50 %. Fall albedo in post-fire stands, on the other hand, was found to be even more sensitive, with a drop in fall



395 albedo at a proportion of deciduous broadleaf species below 80%. We speculate that this sensitivity was related to exposure
of fire residues in early stand development.

Overall our results indicate a strong dependency of seasonal albedo on the proportion of deciduous broadleaf species both in
post-harvest and post-fire stands. This effect provides a strong link between albedo and successional patterns in mixedwood
boreal forests. Prior studies addressing this relationship (e.g., Lyons et al., 2008; Amiro et al., 2006b) have suggested that
400 increasing deciduous tree cover results in increased albedo values from stand initiation to ~ 25 years of stand age; thereafter,
conifers start dominating the canopy, canopy height increases, and albedo decreases gradually until ~ 50 years of stand age
before reaching a steady state. The data presented in the current study provide a somewhat different picture of these trends,
in that patterns show important quantitative differences depending on season and disturbance type. The importance of
405 deciduous broadleaf species in the albedo signal over ~ 50 years of stand development suggests that slow successional
changes in species composition are likely the main driver of the age-related patterns in mixedwood boreal forests albedo.
The dynamics of this pattern is likely to depend on the intensity and frequency of disturbance, edaphic conditions, species
abundance, and climate (Taylor and Chen, 2011). For example, in dry nutrient-poor boreal stands, deciduous broadleaf
species-driven albedo might never occur, as such stands are commonly dominated by jack pine (Taylor and Chen, 2011);
however, in mesic moderate-nutrient-rich stands, deciduous broadleaf species can dominate for ~ 100 years (Cogbill, 1985).
410 Future studies should prioritize robust modelling of boreal succession pathways under different biotic/abiotic conditions to
properly characterize stand albedo.

5 Conclusions

Our findings have important implications for climate-friendly forest management practices. Since the proportion of
deciduous broadleaf species is a strong predictor of seasonal albedo, stand-level albedo can be increased by enhancing
415 proportion of deciduous broadleaf species in a stand. Precisely this approach has recently been suggested as an adaptation
and mitigation strategy to counter negative climate forcings of boreal forest (Astrup et al., 2018), but empirical data from
actual managed stands have been lacking. Historically, forest managers have commonly sought to decrease or eliminate
deciduous species and enhance conifers. However, there is strong evidence that local tree diversity enhances productivity in
boreal forests as in other systems (Paquette and Messier, 2011), and in particular that mixedwood boreal forests including
420 both conifers and deciduous trees show high productivity (MacPherson et al., 2001; Zhang et al., 2012). Management to
increase the proportion of deciduous broadleaf species in managed boreal forests (for example, simply by avoiding chemical
herbicide used to kill deciduous broadleaf species or retaining deciduous broadleaf species seed-trees) could thus be a “win-
win” scenario for enhanced carbon sequestration via primary productivity, and climate mitigation via enhanced albedo.

425 In modeling studies climate forcings in boreal forests have commonly been estimated using a “biome replacement”
approach, typically simulating harvest by replacing mature coniferous forests with grasslands (Bala et al., 2007; Betts,
2000). These models have essentially conceptualized boreal forest albedo as a binary entity rather than a continuum. In a
recent study, Bright et al. (2018) pointed out that overlooking stand structural and compositional properties over the
successional trajectory is likely to substantially bias radiative forcing estimates in the boreal forest. Ground-based estimates
430 such as those presented are essential: at high latitudes when solar zenith angle is high ($> 70^\circ$), satellites such as MODIS
often provide poor-quality albedo data due to spatial heterogeneity of the landscape pixel signature and performance
degradation of atmospheric correction algorithms (Bright et al., 2015b; Wang et al., 2012). Our findings based on field data



are thus essential in improving albedo parameterization in climate models and in refining albedo estimates derived from remote sensing. In addition, our results point to the importance of slow ecological succession as a driver of age-related patterns in albedo, suggesting that future models should explicitly incorporate these ecological processes to better predict long-term trends in climate forcings in boreal forests.

6 Code availability

R (version 3.5.1) codes used in the data analysis can be requested to M.A.H. (abdul.halim@mail.utoronto.ca).

7 Data availability

Data used to produce the graphs are also available via requesting corresponding author (abdul.halim@mail.utoronto.ca).

8 Author contribution

M.A.H., H.Y.H.C., and S.C.T. designed the experiment. M.A.H. analysed data with inputs from S.C.T. and H.Y.H.C. M.A.H. wrote the manuscript with edits and comments from H.Y.H.C. and S.C.T.

9 Competing interests

The authors declare no competing interests.

10 Acknowledgements

Authors acknowledge assistances from Jillian Bieser, Kyle Gaynor, Lutchemee Sujeun, Jack Richard, Jad Murtada, Anna Almero, and Hiro Sato during experiment setup and data collection. We would also like to thank Mark Horsburgh for his helpful comments on an earlier version of this manuscript. This study was funded by the NSERC (Natural Sciences and Engineering Research Council of Canada) Strategic Grant (grant number: STPGP 428641) and the NSERC Discovery Grant (grant number: RGPIN 06209).

11 References

- Amiro, B., Barr, A., Black, T., Iwashita, H., Kljun, N., McCaughey, J., Morgenstern, K., Murayama, S., Nesic, Z. and Orchansky, A.: Carbon, energy and water fluxes at mature and disturbed forest sites, Saskatchewan, Canada, *Agric. For. Meteorol.*, 136(3–4), 237–251, doi:10.1016/j.agrformet.2004.11.012, 2006a.
- Amiro, B. D., Orchansky, A. L., Barr, A. G., Black, T. A., Chambers, S. D., Chapin III, F. S., Goulden, M. L., Litvak, M., Liu, H. P., McCaughey, J. H., McMillan, A. and Randerson, J. T.: The effect of post-fire stand age on the boreal forest energy balance, *Agric. For. Meteorol.*, 140(1), 41–50, doi:10.1016/j.agrformet.2006.02.014, 2006b.
- Astrup, R., Bernier, P. Y., Genet, H., Lutz, D. A. and Bright, R. M.: A sensible climate solution for the boreal forest, *Nat. Clim. Change*, 8(1), 11–12, doi:10.1038/s41558-017-0043-3, 2018.
- Bala, G., Caldeira, K., Wickett, M., Phillips, T. J., Lobell, D. B., Delire, C. and Mirin, A.: Combined climate and carbon-cycle effects of large-scale deforestation, *Proc. Natl. Acad. Sci.*, 104(16), 6550–6555, doi:10.1073/pnas.0608998104, 2007.
- Baltzer, J. L. and Thomas, S. C.: Leaf optical responses to light and soil nutrient availability in temperature deciduous trees, *Am. J. Bot.*, 92(2), 214–223, 2005.



- 465 Bartels, S. F., Chen, H. Y. H., Wulder, M. A. and White, J. C.: Trends in post-disturbance recovery rates of Canada's forests following wildfire and harvest, *For. Ecol. Manag.*, 361, 194–207, doi:10.1016/j.foreco.2015.11.015, 2016.
- Betts, A. K. and Ball, J. H.: Albedo over the boreal forest, *J. Geophys. Res. Atmospheres*, 102(D24), 28901–28909, doi:10.1029/96JD03876, 1997.
- 470 Betts, R. A.: Offset of the potential carbon sink from boreal forestation by decreases in surface albedo, *Nature*, 408(6809), 187–190, doi:10.1038/35041545, 2000.
- Brassard, B. W. and Chen, H. Y. H.: Effects of Forest Type and Disturbance on Diversity of Coarse Woody Debris in Boreal Forest, *Ecosystems*, 11(7), 1078–1090, doi:10.1007/s10021-008-9180-x, 2008.
- Brassard, B. W. and Chen, H. Y. H.: Stand Structure and Composition Dynamics of Boreal Mixedwood Forest: Implications for Forest Management, *Sustainable Forest Management Network.*, 2010.
- 475 Brassard, B. W., Chen, H. Y. H., Wang, J. R. and Duinker, P. N.: Effects of time since stand-replacing fire and overstory composition on live-tree structural diversity in the boreal forest of central Canada, *Can. J. For. Res.*, 38(1), 52–62, doi:10.1139/X07-125, 2008.
- Bright, R. M., Astrup, R. and Strømman, A. H.: Empirical models of monthly and annual albedo in managed boreal forests of interior Norway, *Clim. Change*, 120(1–2), 183–196, doi:10.1007/s10584-013-0789-1, 2013.
- 480 Bright, R. M., Zhao, K., Jackson, R. B. and Cherubini, F.: Quantifying surface albedo and other direct biogeophysical climate forcings of forestry activities, *Glob. Change Biol.*, 21(9), 3246–3266, doi:10.1111/gcb.12951, 2015a.
- Bright, R. M., Myhre, G., Astrup, R., Antón-Fernández, C. and Strømman, A. H.: Radiative forcing bias of simulated surface albedo modifications linked to forest cover changes at northern latitudes, *Biogeosciences*, 12(7), 2195–2205, doi:10.5194/bg-12-2195-2015, 2015b.
- 485 Bright, R. M., Eisner, S., Lund, M. T., Majasalmi, T., Myhre, G. and Astrup, R.: Inferring Surface Albedo Prediction Error Linked to Forest Structure at High Latitudes, *J. Geophys. Res. Atmospheres*, 123(10), 4910–4925, doi:10.1029/2018JD028293, 2018.
- Brown, P. T. and Caldeira, K.: Greater future global warming inferred from Earth's recent energy budget, *Nature*, 552(7683), 45–50, doi:10.1038/nature24672, 2017.
- 490 Chambers, S. D. and Chapin, F. S.: Fire effects on surface-atmosphere energy exchange in Alaskan black spruce ecosystems: Implications for feedbacks to regional climate, *J. Geophys. Res.*, 108(D1), doi:10.1029/2001JD000530, 2002.
- Chen, H. Y. and Popadiouk, R. V.: Dynamics of North American boreal mixedwoods, *Environ. Rev.*, 10(3), 137–166, doi:10.1139/a02-007, 2002.
- 495 Cogbill, C. V.: Dynamics of the boreal forests of the Laurentian Highlands, Canada, *Can. J. For. Res.*, 15(1), 252–261, doi:10.1139/x85-043, 1985.
- Conway, H., Gades, A. and Raymond, C. F.: Albedo of dirty snow during conditions of melt, *Water Resour. Res.*, 32(6), 1713–1718, doi:10.1029/96WR00712, 1996.
- Environment Canada: Historical climate data, [online] Available from: http://climate.weather.gc.ca/historical_data/search_historic_data_e.html (Accessed 11 January 2018), 2018.
- 500 Halim, M. A. and Thomas, S. C.: Surface albedo in relation to disturbance and early stand dynamics in the boreal forest: Implications for climate models., p. abstract #B21F-2025, American Geophysical Union Fall Meeting, 11-15 December 2017, New Orleans, USA., 2017.
- Halim, M. A. and Thomas, S. C.: A proxy-year analysis shows reduced soil temperatures with climate warming in boreal forest, *Sci. Rep.*, 8(1), doi:10.1038/s41598-018-35213-w, 2018.



505 Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S.,
Bell, N., Cairns, B., Canuto, V., Chandler, M., Cheng, Y., Del, G. A., Faluvegi, G., Fleming, E., Friend A., Hall T.,
Jackman C., Kelley M., Kiang N., Koch D., Lean J., Lerner J., Lo K., Menon S., Miller R., Minnis P., Novakov T., Oinas
V., Perlwitz Ja., Perlwitz Ju., Rind D., Romanou A., Shindell D., Stone P., Sun S., Tausnev N., Thresher D., Wielicki B.,
510 Wong T., Yao M. and Zhang, S.: Efficacy of climate forcings, *J. Geophys. Res. Atmospheres*, 110(D18),
doi:10.1029/2005JD005776, 2005.

Hart, S. and Luckai, N.: Charcoal function and management in boreal ecosystems, edited by P. Brando, *J. Appl. Ecol.*,
1197–1206, doi:10.1111/1365-2664.12136, 2013.

Hovi, A., Liang, J., Korhonen, L., Kobayashi, H. and Rautiainen, M.: Quantifying the missing link between forest albedo
and productivity in the boreal zone, *Biogeosciences*, 13(21), 6015–6030, doi:10.5194/bg-13-6015-2016, 2016.

515 Hudspith, V. A., Belcher, C. M., Kelly, R. and Hu, F. S.: Charcoal Reflectance Reveals Early Holocene Boreal Deciduous
Forests Burned at High Intensities, *PLoS ONE*, 10(4), doi:10.1371/journal.pone.0120835, 2015.

Johnstone, J. F., Hollingsworth, T. N., Chapin, F. S. and Mack, M. C.: Changes in fire regime break the legacy lock on
successional trajectories in Alaskan boreal forest, *Glob. Change Biol.*, 16(4), 1281–1295, doi:10.1111/j.1365-
2486.2009.02051.x, 2010.

520 Kirschbaum, M. U. F., Whitehead, D., Dean, S. M., Beets, P. N., Shepherd, J. D. and Ausseil, A.-G. E.: Implications of
albedo changes following afforestation on the benefits of forests as carbon sinks, *Biogeosciences*, 8(12), 3687–3696,
doi:10.5194/bg-8-3687-2011, 2011.

Kumar, P., Chen, H. Y. H., Thomas, S. C. and Shahi, C.: Epixylic vegetation abundance, diversity, and composition vary
with coarse woody debris decay class and substrate species in boreal forest, *Can. J. For. Res.*, 48(4), 399–411,
525 doi:10.1139/cjfr-2017-0283, 2018.

Kuusinen, N., Kolari, P., Levula, J., Porcar-Castell, A., Stenberg, P. and Berninger, F.: Seasonal variation in boreal pine
forest albedo and effects of canopy snow on forest reflectance, *Agric. For. Meteorol.*, 164, 53–60,
doi:10.1016/j.agrformet.2012.05.009, 2012.

530 Kuusinen, N., Tomppo, E., Shuai, Y. and Berninger, F.: Effects of forest age on albedo in boreal forests estimated from
MODIS and Landsat albedo retrievals, *Remote Sens. Environ.*, 145, 145–153, doi:10.1016/j.rse.2014.02.005, 2014.

Kuusinen, N., Stenberg, P., Tomppo, E., Bernier, P. and Berninger, F.: Variation in understory and canopy reflectance
during stand development in Finnish coniferous forests, *Can. J. For. Res.*, 45(8), 1077–1085, doi:10.1139/cjfr-2014-0538,
2015.

535 Kuusinen, N., Stenberg, P., Korhonen, L., Rautiainen, M. and Tomppo, E.: Structural factors driving boreal forest albedo in
Finland, *Remote Sens. Environ.*, 175, 43–51, doi:10.1016/j.rse.2015.12.035, 2016.

Lee, X., Goulden, M. L., Hollinger, D. Y., Barr, A., Black, T. A., Bohrer, G., Bracho, R., Drake, B., Goldstein, A., Gu, L.,
Katul, G., Kolb, T., Law, B. E., Margolis, H., Meyers, T., Monson, R., Munger, W., Oren, R., Paw U, K. T., Richardson, A.
D., Schmid, H. P., Staebler, R., Wofsy, S. and Zhao, L.: Observed increase in local cooling effect of deforestation at higher
latitudes, *Nature*, 479(7373), 384–387, doi:10.1038/nature10588, 2011.

540 Linacre, E.: *Climate Data and Resources: A Reference and Guide*, e-book., Routledge, London., 2003.

Liu, H., Randerson, J. T., Lindfors, J. and Chapin, F. S.: Changes in the surface energy budget after fire in boreal
ecosystems of interior Alaska: An annual perspective, *J. Geophys. Res.*, 110(D13), doi:10.1029/2004JD005158, 2005.

Lukeš, P., Stenberg, P., Rautiainen, M., Möttus, M. and Vanhatalo, K. M.: Optical properties of leaves and needles for
boreal tree species in Europe, *Remote Sens. Lett.*, 4(7), 667–676, doi:10.1080/2150704X.2013.782112, 2013a.

545 Lukeš, P., Stenberg, P. and Rautiainen, M.: Relationship between forest density and albedo in the boreal zone, *Ecol. Model.*,
261–262, 74–79, doi:10.1016/j.ecolmodel.2013.04.009, 2013b.



- Lyons, E. A., Jin, Y. and Randerson, J. T.: Changes in surface albedo after fire in boreal forest ecosystems of interior Alaska assessed using MODIS satellite observations, *J. Geophys. Res. Biogeosciences*, 113(G02012), doi:10.1029/2007JG000606, 2008.
- 550 MacPherson, D. M., Lieffers, V. J. and Blenis, P. V.: Productivity of aspen stands with and without a spruce understory in Alberta's boreal mixedwood forests, *For. Chron.*, 77(2), 351–356, doi:10.5558/tfc77351-2, 2001.
- Madoui, A., Gauthier, S., Leduc, A., Bergeron, Y. and Valeria, O.: Monitoring Forest Recovery Following Wildfire and Harvest in Boreal Forests Using Satellite Imagery, *Forests*, 6(12), 4105–4134, doi:10.3390/f6114105, 2015.
- 555 Mair, P. and Wilcox, R.: WRS2: Wilcox robust estimation and testing, R package. [online] Available from: <https://r-forge.r-project.org/projects/psychor/>, 2018.
- Matthies, B. D. and Valsta, L. T.: Optimal forest species mixture with carbon storage and albedo effect for climate change mitigation, *Ecol. Econ.*, 123, 95–105, doi:10.1016/j.ecolecon.2016.01.004, 2016.
- McMillan, A. M. S. and Goulden, M. L.: Age-dependent variation in the biophysical properties of boreal forests: Biophysical Properties of boreal forests, *Glob. Biogeochem. Cycles*, 22(2), n/a-n/a, doi:10.1029/2007GB003038, 2008.
- 560 Moussaoui, L., Fenton, N., Leduc, A. and Bergeron, Y.: Can Retention Harvest Maintain Natural Structural Complexity? A Comparison of Post-Harvest and Post-Fire Residual Patches in Boreal Forest, *Forests*, 7(12), 243, doi:10.3390/f7100243, 2016.
- Paquette, A. and Messier, C.: The effect of biodiversity on tree productivity: from temperate to boreal forests: The effect of biodiversity on the productivity, *Glob. Ecol. Biogeogr.*, 20(1), 170–180, doi:10.1111/j.1466-8238.2010.00592.x, 2011.
- 565 Qian, Y., Gustafson, W. I., Leung, L. R. and Ghan, S. J.: Effects of soot-induced snow albedo change on snowpack and hydrological cycle in western United States based on Weather Research and Forecasting chemistry and regional climate simulations, *J. Geophys. Res.*, 114(D3), doi:10.1029/2008JD011039, 2009.
- Qu, X. and Hall, A.: What controls the strength of snow-albedo feedback?, *J. Clim.*, 20(15), 3971–3981, doi:10.1175/JCLI4186.1, 2007.
- 570 R Core Team: R: A language and environment for statistical computing., R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: URL <https://www.R-project.org/>, 2018.
- Randerson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G., Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L., Lyons, E., Neff, J. C., Schuur, E. A. G. and Zender, C. S.: The Impact of Boreal Forest Fire on Climate Warming, *Science*, 314(5802), 1130–1132, doi:10.1126/science.1132075, 2006.
- 575 Rohatgi, A.: WebPlotDigitizer, Austin, Texas, USA. [online] Available from: <https://automeris.io/WebPlotDigitizer>, 2018.
- Sims, R. A., Towill, W. D., Baldwin, K. A. and Wickware, G. M.: Field guide to the forest ecosystem classification for northwestern Ontario., 2nd ed., Ontario Ministry of Natural Resources, Northwest Science and Technology, Thunder Bay, Ontario., 1997.
- 580 Stephens, G. L., O'Brien, D., Webster, P. J., Pilewski, P., Kato, S. and Li, J.: The albedo of Earth, *Rev. Geophys.*, 53(1), 141–163, doi:10.1002/2014RG000449, 2015.
- Taylor, A. R. and Chen, H. Y. H.: Multiple successional pathways of boreal forest stands in central Canada, *Ecography*, 34(2), 208–219, doi:10.1111/j.1600-0587.2010.06455.x, 2011.
- 585 Uotila, A. and Kouki, J.: Understorey vegetation in spruce-dominated forests in eastern Finland and Russian Karelia: Successional patterns after anthropogenic and natural disturbances, *For. Ecol. Manag.*, 215(1–3), 113–137, doi:10.1016/j.foreco.2005.05.008, 2005.
- Wang, Z., Schaaf, C. B., Chopping, M. J., Strahler, A. H., Wang, J., Román, M. O., Rocha, A. V., Woodcock, C. E. and Shuai, Y.: Evaluation of Moderate-resolution Imaging Spectroradiometer (MODIS) snow albedo product (MCD43A) over tundra, *Remote Sens. Environ.*, 117, 264–280, doi:10.1016/j.rse.2011.10.002, 2012.



- 590 Wickham, H.: ggplot2: Elegant Graphics for Data Analysis, R, Springer-Verlag, New York. [online] Available from: <http://ggplot2.org>, 2016.
- Wilcox, R. R.: Introduction to robust estimation and hypothesis testing, 4th edition., Elsevier, Waltham, MA., 2016.
- Zhang, D.: rsq: R-Squared and Related Measures, R. [online] Available from: <https://CRAN.R-project.org/package=rsq>, 2018.
- 595 Zhang, Q., Wang, Y., Wu, Y., Wang, X., Du, Z., Liu, X. and Song, J.: Effects of Biochar Amendment on Soil Thermal Conductivity, Reflectance, and Temperature, *Soil Sci. Soc. Am. J.*, 77(5), 1478–1487, doi:10.2136/sssaj2012.0180, 2013.
- Zhang, Y., Chen, H. Y. H. and Reich, P. B.: Forest productivity increases with evenness, species richness and trait variation: a global meta-analysis: Diversity and productivity relationships, *J. Ecol.*, 100(3), 742–749, doi:10.1111/j.1365-2745.2011.01944.x, 2012.



600 12 Figures and Tables

12.1 Figures




Figure 1. Map of the study area. Labels in the rectangular boxes indicate disturbance types (H: harvest and F: fire) and years (98: 1998, 06: 2006, and 13: 2013) for each plot (grey circles). **Inset:** black squares indicate locations of all data sources including the current study area (LN: Lake Nipigon area, Ontario, Canada; AK: Alaska, USA; SK: Saskatchewan, Canada; MB: Manitoba, Canada).

605





Figure 2. Comparison of seasonal albedo (mean \pm SE) in post-harvest and post-fire stands. Winter (no. of observations for post-fire stands, $n_F = 35$; no. of observations for post-harvest stands, $n_H = 48$) and summer ($n_F = 44$, $n_H = 41$) albedo data were from 0–19-year-old stands, and spring ($n_F = 30$, $n_H = 30$) and fall ($n_F = 30$, $n_H = 30$) albedo data were from 7–19-year-old stands. Albedo of 0–6-year-old post-fire stands were from secondary sources. * and *** indicate significant mean albedo differences between post-harvest and post-fire stands with $p = 0.11$ and $p < 0.01$, respectively.

610



615 **Figure 3.** Specular-included ground surface reflectance (400–1000 nm) of post-harvest and post-fire stands. Lines indicate mean
reflectance (number of sample (n) × 10 replicated measurements/sample) in the corresponding wavelengths, and shades indicate SE. **[a]**
ground surface reflectance of young (4-year old) post-harvest stands (n = 9) and a post-fire stand (n = 12). **[b]** ground surface reflectance
of old (11- and 19-year old) post-harvest (n = 18) and post-fire (n = 18) stands.

620

625




Figure 4. Stand age affecting mean seasonal albedo (\pm SE) in boreal forest over 0–150 years of stand development. Mean winter albedo as a function of stand age in **[a]** post-harvest stands ($n = 42$) and **[b]** post-fire stands ($n = 37$). Mean summer albedo as a function of stand age in **[c]** post-harvest stands ($n = 41$) and **[d]** post-fire stands ($n = 31$). Each field-data point is the average seasonal albedo (error bars indicate standard errors) of three plots from each stand-age category over the study period.

630

635





Figure 5. Stand age affecting mean seasonal albedo (\pm SE) in boreal forest in the early seral stage. Mean spring albedo as a function of stand age in **[a]** post-harvest stands ($n = 24$) and **[b]** post-fire stands ($n = 14$). Mean fall albedo as a function of stand age in **[c]** post-harvest stands ($n = 29$) and **[d]** post-fire stands ($n = 22$). Each field-data point is the average seasonal albedo (error bars indicate standard errors) of three plots from each stand-age category over the study period.

645

650



655

Figure 6. Mean seasonal albedo (\pm SE) as a function of deciduous broadleaf species (%) (proportion of deciduous broadleaf species) in the boreal forest. Proportion of deciduous broadleaf species affecting mean winter albedo in [a] post-harvest stands ($n = 16$) and [b] in post-fire stands ($n = 16$), mean spring albedo in [c] post-harvest stands ($n = 8$) and [d] post-fire stands ($n = 6$), mean summer albedo in [e] post-harvest stands ($n = 21$) and [f] post-fire stands ($n = 17$), and mean fall albedo in [g] post-harvest stands ($n = 8$) and [h] post-fire stands ($n = 7$).

660





Figure 7. Mean seasonal albedo (\pm SE) as a function of canopy height (m) in the boreal forest. Canopy height affecting [a] mean winter albedo in post-harvest stands ($n = 30$), [c] mean spring albedo in post-harvest stands ($n = 16$), [f] mean summer albedo in post-fire stands ($n = 24$), and [g] mean fall albedo in post-harvest stands ($n = 20$). In [b, d, e, h] canopy height is not a significant predictor of the corresponding mean seasonal albedo; thus, no model is fitted to the data points.

665



670

Figure 8. Mean seasonal albedo (\pm SE) as a function of ground vegetation cover (%) in the boreal forest. Ground vegetation cover affecting [e] mean summer albedo in post-harvest stands ($n = 21$) and [g] mean fall albedo in post-harvest stands ($n = 18$). In [a-d, f, h] ground vegetation cover is not a significant predictor of the corresponding mean seasonal albedo; thus, no model is fitted to the data points.

675



12.2 Tables

680 **Table 1. Regression model coefficients and fit statistics for albedo as a function of stand attributes in different seasons in the boreal forest.**

Season	Post-harvest stands				Post-fire stands			
	<i>Parameter Estimates</i>		<i>Model fit</i>		<i>Parameter Estimates</i>		<i>Model fit</i>	
	<i>Coefficients</i>	<i>Estimates</i>	ΔAIC	<i>Adj. R²</i>	<i>Coefficients</i>	<i>Estimates</i>	ΔAIC	<i>Adj. R²</i>
Winter	Intercept	<i>1.679</i>			Intercept	<i>-1.164</i>		
	SA	<i>-0.036</i>			SA	<i>-0.004</i>		
	PDBS	<i>-0.021</i>			PDBS	<i>0.009</i>		
	CH	<i>-0.098</i>	-65.5	0.97	SA: PDBS	<i>-0.0004</i>	-2.8	0.79
	SA:CH	<i>0.002</i>						
	PDBS:CH	<i>-0.007</i>						
Spring	Intercept	<i>-7.195</i>			Intercept	<i>-1.747</i>		
	SA	<i>1.298</i>			SA	<i>0.016</i>		
	PDBS	<i>0.116</i>	-495.4	0.99	PDBS	<i>0.002</i>	-18.8	0.92
	CH	<i>-1.264</i>						
	SA: PDBS	<i>-0.024</i>						
Summer	Intercept	<i>-1.377</i>			Intercept	<i>-2.750</i>		
	SA	<i>0.032</i>			SA	<i>-0.011</i>		
	PDBS	<i>-0.003</i>			PDBS	<i>-0.006</i>		
	GVC	<i>-0.01</i>	-24.8	0.96	CH	<i>0.688</i>	-39.0	0.93
	SA: GVC	<i>-0.0004</i>			SA: PDBS	<i>0.003</i>		
	PDBS: GVC	<i>0.0001</i>			SA: CH	<i>-0.004</i>		
Fall	Intercept	<i>0.398</i>			$\frac{4.5}{6.87} e^{(-\frac{SA-13.2}{6.87} - e^{-\frac{SA-67.62}{45.39}})}$		-3.1	0.045 ¹
	SA	<i>0.013</i>			$0.099 e^{0.013 PDBS}$		-25.4	0.008 ¹
	CH	<i>-0.182</i>						
	GVC	<i>-0.007</i>	-6.1	0.94				
	SA:CH	<i>0.007</i>						
	CH: GVC	<i>0.005</i>						
	SA:CH: GVC	<i>-0.0002</i>						
	$\frac{28.86}{45.39} e^{(-\frac{PDBS-67.62}{45.39} - e^{-\frac{PDBS-67.62}{45.39}})}$		-0.9	0.049 ¹				

685 Notes: SA, PDBS, CH, and GVC indicate stand age (year), proportion of deciduous broadleaf species (%), canopy height (m), and ground vegetation cover (%), respectively. Parameter estimates for GLMs in italic and regular fonts indicate statistical significance at 1% and 5% level, respectively. For fall nonlinear regression models, 28.86 and 45.39 coefficients of post-harvest stands are significant at 5% level and the rest is significant at 1% level. ¹ indicates residual standard error of the nonlinear regression model. ΔAIC = AIC of the best-fit model – AIC of the corresponding null model.