

Dear Editor,

Thank you for handling our manuscript, “Revisiting the boron systematics of aragonite and their application to coral calcification” (bg-2018-77). Two reviewers provided constructive comments that highlighted several areas requiring clarification or additional discussion. We have revised the manuscript following these helpful suggestions, which have improved the quality of our manuscript. Below, we respond (**in bold**) to each reviewer comment (plain text), with the modified text highlighted in yellow throughout the manuscript.

We greatly appreciate the time you have devoted to our manuscript as editor, and we look forward to publishing in *Biogeosciences*.

Sincerely,

Thomas DeCarlo, Michael Holcomb, and Malcolm McCulloch

Responses to reviewer comments:

Reviewer 1:

I really enjoyed reading the manuscript. The authors summarized issues on the selection of K_d value (and its formula) and its potential influence on the calculation of full carbonate chemistry in the calcifying medium. The logic is concise, and I strongly recommend a publication of the manuscript.

The followings are my minor comments that may be helpful for the authors to improve the manuscript.

(pp. 2 Line 20–) I think almost nobody use stable carbon and oxygen isotopes as a proxy of carbonate chemistry, so you can delete the related sentences.

We agree with the reviewer that carbon and oxygen isotope ratios are not commonly applied as carbonate system proxies in corals. This phrasing has been revised to indicate that they are not typically applied in this way (page 2, lines 21-24), but they are theoretically sensitive to carbonate chemistry. We prefer to still mention carbon and oxygen isotopes because they are examples of geochemical proxies that are sensitive to the carbonate system, yet are not very useful proxies in corals due to a variety of vital effects.

(pp. 7 Figure 2 and pp. 14 Figure 8) About pH and $[H^+]$. I think $[H^+]$ presented in the Figure 2 is that of solution used in the precipitation experiment. In Figure 8, on the other hand, they are calcifying fluid pH for coral data as well as solution pH for precipitation experiment. I would be better to clarify what each pH stand for in somewhere in the manuscript (in each figure caption?).

We have made clear the distinction between coral calcifying fluid pH (or H^+) and the abiogenic experimental fluid pH, both in the captions and axis labels (changes to axes made in Figure 2 and Figure 8).

(pp. 10 Figure 4) Why do you use Kd value of 0.002 as an example of constant Kd?

The value of 0.002 was selected simply as an example that intersects the abiogenic data near the range of $[\text{CO}_3^{2-}]$ found in corals (added to page 10, lines 15-16). We could choose any other value, which would be a similar line but further from the abiogenic dataset. The main message is that the constant Kd underestimates the sensitivity of $[\text{CO}_3^{2-}]$ to borate/(B/Ca), which is made clear by the best-case example with Kd of 0.002.

(pp. 12 Figure 6) Is there any better way to plot these data? The difference between New Eq. (12) line and Allison (2017) line are very ambiguous.

We have revised Figure 6 in several ways: Firstly, we now plot only $[\text{CO}_3^{2-}]$ (and not DIC) because the purpose of the plot is to demonstrate differences in derived $[\text{CO}_3^{2-}]$ among Kd formulations, and the derived DIC follows the same pattern. This enables the $[\text{CO}_3^{2-}]$ plot to be larger and thus more clearly visualized. Secondly, we made two separate $[\text{CO}_3^{2-}]$ axes, which allows us to focus more closely on the small differences between the Holcomb et al. (2016), McCulloch et al. (2017), and Eq. (12) formulations. Finally, we added error bars to the lines.

(pp. 14 Line 17- pp. 15 Line 2) It is just a question. Is this the reason why you don't show a cross-plot of Ω_{ar} against the other parameters? (such as Ω_{ar} versus pH)

Yes, we prefer to plot only boron-derived $[\text{CO}_3^{2-}]$, rather than saturation state, because boron systematics really only provide information regarding pH and $[\text{CO}_3^{2-}]$, not $[\text{Ca}^{2+}]$.

Reviewer 2:

DeCarlo et al. synthesize the (very recently developed) joint B/Ca- $\delta^{11}\text{B}$ system in aragonite corals as a proxy for coral calcifying fluid chemistry. Coral aragonite $\delta^{11}\text{B}$ has previously been applied as a calcifying fluid pH proxy, while recent studies of synthetic aragonite B/Ca suggest control by $[\text{CO}_3^{2-}]$. If these results apply to corals, then coral aragonite B/Ca may reflect $[\text{CO}_3^{2-}]$ in the calcifying fluid. The ability to reconstruct $[\text{CO}_3^{2-}]$ (from B/Ca) and pH (from $\delta^{11}\text{B}$) allows for solving the carbonate chemistry of coral calcifying fluid, which permits reconstructions of calcifying fluid DIC (among other parameters). This new approach hinges on the veracity of coral B/Ca to $[\text{CO}_3^{2-}]$ reconstructions, which these authors test in detail. They also present a calcifying fluid calculation routine that propagates all uncertainties associated with the above calculations.

This is a nicely written and useful contribution, and I do support its publication, but I think it is missing one key component:

Primary concern/recommendation: Coral B/Ca as a $[\text{CO}_3^{2-}]$ proxy exploded in the last two years, in large part due to the works of these authors. While this contribution cites requisite previous reasoning (Holcomb et al. Chem Geol. 2016 for synthetic aragonite, and McCulloch et al. Nat. Comm. 2017), I do not find that the rationale for this approach has been sufficiently explored in previous publications. As the authors use this manuscript to comprehensively and quantitatively analyze KD formulations, I strongly encourage them to

also take a step back and comprehensively evaluate the B/Ca-[CO₃²⁻] proxy system in corals and its inherent assumptions. Adding this to the quantitative treatment already provided would greatly enhance this contribution's readability and utility.

Guiding questions for this background:

1) What is known about patterns in coral B/Ca? How do features of these patterns (seasonal cycles, etc.) imply a relationship to [CO₃²⁻]cf and/or [DIC]cf? It seems previously published B/Ca data are already compiled in Figure 8, so this won't require much work.

We added some discussion on patterns of B/Ca in coral skeletons (e.g. acknowledging previous reports of seasonality on page 11, lines 12-14). However, it is difficult to interpret B/Ca alone because it is not directly related to [CO₃²⁻], but rather depends also on borate concentration (i.e. B/Ca depends on both pH and [CO₃²⁻]), which we now highlight on page 13, lines 5-8. Additionally, we state directly that our focus for this study is on the combined application of B/Ca and δ¹¹B (page 3, lines 16-17).

2) What is known about coral [DIC]cf, both naturally and in controlled experiments (e.g., Cai et al., 2016; Comeau et al., 2017)? What are the limitations to direct measurements? (Schoepf et al. 2017 gave a nice overview of this, but I would appreciate seeing that reasoning here)

We added an extended discussion of calcifying fluid DIC (page 13, lines 18-30). Here, we discuss that there is a substantial, and currently unresolved, difference between DIC derived from boron systematics (DIC_{cf} > seawater) and from microsensors (DIC_{cf} < seawater). Additionally, we describe the implications for understanding coral calcification, and acknowledge that there is some independent supporting evidence for the high DIC_{cf} scenario (page 13, lines 24-28).

3) Two previous studies of paired foraminifera B/Ca and δ¹¹B concluded that joint reconstructions of [CO₃²⁻] and pH could not be used to reconstruct full ocean carbonate chemistry because the relative uncertainties in reconstructing Alk and DIC were larger than the entire range of these parameters in the modern ocean (Yu et al., EPSL 2010; Rae et al., EPSL 2011). What is different in corals that make this application feasible? I think it probably relates to the much bigger ranges of [CO₃²⁻] and/or [DIC] in coral calcifying fluids vs. seawater, but I'd like to hear that from the authors. In general, the coral joint B/Ca and δ¹¹B approach needs to be presented within the context of previous (unsuccessful) open ocean efforts.

We added a discussion of the difficulty in applying boron systematics to reconstruct seawater chemistry (page 13, lines 13-17). Like the foraminifera studies mentioned by the reviewer, efforts to reconstruct ocean carbonate chemistry with corals are not very successful because the changes within the calcifying fluid often far exceed natural variability of seawater. Thus, while boron systematics is a useful tool for understanding coral calcification and its sensitivity to changes in reef environments, it may not be generally applicable for deriving ocean chemistry.

Specific comments:

Page 3, L1 (relevant for Section 2): For most boron proxy applications, inorganic carbonate precipitation experiments do not reflect biogenic carbonates as well as our community would

like (see, e.g., Allen and Hönisch, 2011; 2012; Uchikawa et al. 2015, 2017, review in Rae and Foster, 2016; Rae 2018 book chapters). Please defend why applying a KD derived from synthetic aragonite B/Ca is appropriate for coral aragonite in light of issues observed in other boron applications. This discussion could fit well in Section 7 (p. 13).

In general, it is difficult to validate the application of Kd derived in abiogenic experiments to coral skeletons because independent data of coral calcifying fluid chemistry are scarce. Microsensor and fluorescent dye measurements of calcifying fluid pH are broadly similar to boron isotope-derived pH, but the one study of calcifying fluid DIC derived from microsensors differs from boron systematics results. However, boron systematics are broadly similar with constraints from U/Ca and Raman spectroscopy, which we have added to the manuscript (page 13, lines 24-28).

Page 5, L31: What might compositional effects on B/Ca partitioning look like? This is a critical point for two reasons: 1) If compositional effects do exist, then B/Ca partitioning is not effectively described by KD, and instead requires additional parameters related to varying solution chemistry than only $[\text{CO}_3^{2-}]$ and $[\text{Ca}^{2+}]$. 2) If compositional effects do exist, then application of B/Ca- $[\text{CO}_3^{2-}]$ approach to coral calcifying fluid would carry additional uncertainty because the calcifying fluid composition is not unaltered seawater (because of ion pumps such as Ca-ATPase) Note: I feel that the authors nicely dealt with comparing the B/Ca data from Mavromatis and Holcomb nicely throughout the manuscript, and their approach of using both datasets to define KD in terms of CO_3^{2-} (Equation 12) implies that compositional effects do not matter. But I think it is important for them to note that compositional effects could undermine the application of the B/Ca- $[\text{CO}_3^{2-}]$ approach to non-seawater media (which includes the calcifying fluid).

The reviewer makes a good point here. It is important to note that the Holcomb et al. (2016) experiments include a range of seawater chemical manipulations, including [Mg], [Ca], and [Sr] exceeding changes typically thought to occur within the calcifying fluid, without clear effects on Kd. Thus, we do not think there are strong sensitivities of Kd to trace element variations. Yet it is possible that there are subtle effects, which are not apparent in Holcomb et al. (2016) because the fluids are all broadly similar to seawater, but do become apparent in Mavromatis et al. (2015) since the fluid chemistry departs substantially from seawater for many elements. We have added this discussion to page 6, lines 1-11.

Page 12, L14-19 and Figure 8: Suggest you change the order of figures, starting from the measured parameters ($\delta^{11}\text{B}$ and B/Ca, in a), then each converted to their independent parameters (pH and CO_3^{2-}), and finally plots vs. DIC, which requires both parameters. It is tough to say whether the correlation between DIC and CO_3^{2-} is "interesting" or even surprising, because the calculation of DIC depends on pH and CO_3^{2-} . Because pH and DIC do not correlate well, changes in DIC are probably principally driven by changes in $[\text{CO}_3^{2-}]$ (and hence coral B/Ca). This could be worth exploring with a sensitivity test.

We agree with this suggestion, and we have revised the order of panels in Figure 8 as suggested. In terms of deriving DIC, yes it appears to depend most strongly on $[\text{CO}_3^{2-}]$. However, in terms of modification within the calcifying fluid, it may be that CO_2 diffusion drives DIC changes, which in turn affect $[\text{CO}_3^{2-}]$.

Page 14, L16-17: In section 2, the authors state that Holcomb et al. (2016) only performed two experiments at each offset temperature, and that this was insufficient to quantify temperature effects on precipitation rate. Are the data also too limited to find a temperature dependence on B/Ca partitioning?

We revised the statement regarding the quantification of temperature effects on precipitation rate (page 5, lines 12-15). The Holcomb et al. (2016) data are generally consistent with Burton and Walter (1987) in that precipitation rate increases with temperature, and the data are sufficient to demonstrate this. However, Burton and Walter (1987) show that the order of the reaction changes with temperature, which requires a full calibration dataset (i.e. more than 2 experiments) for each temperature. Thus, we now describe that we do see a temperature dependence of reaction rate, but that we cannot go as far as Burton and Walter (1987) in quantifying changes in the reaction order. In addition, Holcomb et al. (2016) already reported that there was no apparent temperature effect on B/Ca partitioning between 20 and 40 °C (page 16, lines 6-7).

Figure comments.

Please label panels a) through d) (or however many panels) in each figure (some are missing). I would recommend increasing the font size of these labels; they are difficult to see.

We added panel labels to all figures (Figure 7 has been revised).

Figure 6. I only see three line types on here (solid-McCulloch, gray dash-Allison, and then a dot dash that may be both the Holcomb and Equation 12 lines?) If the Holcomb and Equation 12 lines fall on top of each other, please say so in the text and figure caption. Additionally, while the authors MATLAB routine calculates a propagated uncertainty on derived $[\text{CO}_3^{2-}]_{\text{cf}}$ and $[\text{DIC}]_{\text{cf}}$, no uncertainties are plotted. Please illustrate this uncertainty on Figure 6. How does the propagated uncertainty affect the conclusion about applicability of McCulloch, Holcomb, and Equation 12 lines? Are they truly any different from each other (tested statistically)?

As described above in response to comments from Reviewer 1, we revised Figure 6 to more clearly show the separate lines and we included error bars.

Figure 7. Panel labeling. Also, do not use $\Delta[\text{CO}_3^{2-}]$ in titles, as this is a well-used carbonate chemistry term. Suggest changing titles to “[CO_3^{2-}]_{cf} difference” or “[CO_3^{2-}]_{cf} M17 – [CO_3^{2-}]_{cf} H16”. Please specify that [CO_3^{2-}] is [CO_3^{2-}]_{cf} on figures and in caption. Finally, the color schemes are a bit tough to follow. In b) through d), white is good, right?

We changed the panel titles as suggested (Figure 7). We prefer to keep the color scheme as it is a common (and we believe effective) way to visualize anomalies because it is easy to see where the two formulas are consistent (white) or one higher than the other (red or blue).

Figure 8. Tough figure to read, recommend brighter symbol colors and making the ~ gray shading for the Holcomb et al. data lighter.

We made the symbol colors clearer and the gray shading lighter in Figure 8.

Reviews and syntheses: Revisiting the boron systematics of aragonite and their application to coral calcification

Thomas M. DeCarlo^{1,2}, Michael Holcomb^{1,2}, and Malcolm T. McCulloch^{1,2}

¹Oceans Institute and School of Earth Sciences, The University of Western Australia, 35 Stirling Hwy, Crawley 6009, Australia

²ARC Centre of Excellence for Coral Reef Studies, The University of Western Australia, 35 Stirling Hwy, Crawley 6009, Australia

Correspondence to: Thomas M. DeCarlo (thomas.decarlo@uwa.edu.au)

Abstract. The isotopic and elemental systematics of boron in aragonitic coral skeletons have recently been developed as a proxy for the carbonate chemistry of the coral extracellular calcifying fluid. With knowledge of the boron isotopic fractionation in seawater and the B/Ca partition coefficient (K_D) between aragonite and seawater, measurements of coral skeleton $\delta^{11}\text{B}$ and B/Ca can potentially constrain the full carbonate system. Two sets of abiogenic aragonite precipitation experiments designed to quantify K_D have recently made possible the application of this proxy system. However, while different K_D formulations have been proposed, there has not yet been a comprehensive analysis that considers both experimental datasets and explores the implications for interpreting coral skeletons. Here, we evaluate four potential K_D formulations: three previously presented in the literature and one newly developed. We assess how well each formulation reconstructs the known fluid carbonate chemistry from the abiogenic experiments, and we evaluate the implications for deriving the carbonate chemistry of coral calcifying fluid. Three of the K_D formulations performed similarly when applied to abiogenic aragonites precipitated from seawater and to coral skeletons. Critically, we find that some uncertainty remains in understanding the mechanism of boron elemental partitioning between aragonite and seawater, and addressing this question should be a target of additional abiogenic precipitation experiments. Despite this, boron systematics can already be applied to quantify the coral calcifying fluid carbonate system, although uncertainties associated with the proxy system should be carefully considered for each application. Finally, we present a user-friendly computer code that calculates coral calcifying fluid carbonate chemistry, including propagation of uncertainties, given inputs of boron systematics measured in coral skeleton.

1 Introduction

Quantifying the carbonate chemistry of the fluid from which corals accrete their skeletons is essential for understanding the mechanisms of skeletal growth and the sensitivity of skeletal composition to environmental variability. It is generally thought that corals precipitate aragonite (CaCO_3) crystals within an extracellular fluid-filled space between the living polyp and the skeleton (Barnes, 1970). Evidence from skeletal geochemistry and fluorescent dye experiments suggests that while seawater is the initial source of the calcifying fluid (McConnaughey, 1989; Adkins et al., 2003; Cohen and McConnaughey, 2003; Gagnon et al., 2012; Tambutté et al., 2012), the carbonate chemistry of the calcifying fluid is subject to substantial modifications (*i.e.*

pH and dissolved inorganic carbon, or DIC) that enhance the rapid nucleation and growth of aragonite crystals (Al-Horani et al., 2003; Venn et al., 2011). Because the isolation and small size of the calcifying fluid makes it difficult to sample directly, a variety of techniques have been employed to characterize its composition. These include microelectrodes inserted into tissue incisions or through the mouth (Al-Horani et al., 2003; Ries, 2011; Cai et al., 2016), pH-sensitive dyes (Venn et al., 2011, 5 2013; Holcomb et al., 2014; Comeau et al., 2017), Raman spectroscopy (DeCarlo et al., 2017), and a variety of skeletal-based geochemical proxies (Rollion-Bard et al., 2010, 2011; Inoue et al., 2011; Trotter et al., 2011; McCulloch et al., 2012b; Allison et al., 2014; Holcomb et al., 2014; DeCarlo et al., 2015). Although microelectrodes and pH-sensitive dyes are arguably the most direct methods, their utilities are limited by difficulties of applying them to corals living in their natural environment or developing seasonally-resolved time series. Geochemical proxies, although indirect, can be readily applied to the skeletons of 10 corals living in both laboratory and natural environments, and to skeletons accreted years or even centuries ago.

In recent years, boron systematics (including $\delta^{11}\text{B}$ and B/Ca) have become one of the most commonly applied proxies for the carbonate chemistry of coral calcifying fluid (cf) (Hönisch et al., 2004; Trotter et al., 2011; McCulloch et al., 2012b, a, 2017; Allison et al., 2014; DeCarlo et al., 2016; Stewart et al., 2016; Comeau et al., 2017; Wu et al., 2017; D'Olivo and McCulloch, 2017; Kubota et al., 2017; Ross et al., 2017; Schoepf et al., 2017). The sensitivity of boron isotopes to seawater pH arises from 15 the borate versus boric acid speciation being pH-dependent and the isotopic fractionation between these species being constant (Klochko et al., 2006). Since the $\delta^{11}\text{B}$ composition of aragonite precipitating from seawater reflects that of the borate species (Klochko et al., 2006; Trotter et al., 2011; Noireaux et al., 2015), the $\delta^{11}\text{B}$ composition of the skeletal carbonate records the pH of the calcifying fluid. Furthermore, the B/Ca ratio depends inversely on the concentration of carbonate ion ($[\text{CO}_3^{2-}]$) since borate substitutes for carbonate ion in the aragonite lattice (Holcomb et al., 2016).

20 Use of combined boron isotopic ($\delta^{11}\text{B}$) and elemental (B/Ca) systematics has several advantages relative to other geochemical proxies. For example, while stable carbon and oxygen isotopes are sensitive to carbonate chemistry, they are complicated by kinetic effects, strong sensitivities to the photosynthetic activity of coral symbionts, and variable compositions in seawater, which together have precluded their utility as acceptable carbonate system proxies (Adkins et al., 2003; Cohen and McConnaughey, 2003; Schoepf et al., 2014). The U/Ca ratio of aragonite is also sensitive to $[\text{CO}_3^{2-}]$, but the amount of U in coral 25 skeleton relative to its concentration in seawater suggests that $[\text{U}]_{cf}$ is depleted substantially, complicating its utility as a direct $[\text{CO}_3^{2-}]_{cf}$ proxy (DeCarlo et al., 2015). Conversely, the B/Ca and $\delta^{11}\text{B}$ compositions of seawater are homogeneous (Foster et al., 2010; Lee et al., 2010) and likely not modified substantially by photosynthetic activity (Hönisch et al., 2004). Further, incorporation into the skeleton is less important for B/Ca than U/Ca because the partition coefficient between B and $[\text{CO}_3^{2-}]$ is at least 2 orders of magnitude smaller than that of $\text{U}/\text{CO}_3^{2-}$ (DeCarlo et al., 2015; Mavromatis et al., 2015; Holcomb et al., 30 2016), meaning that $[\text{B}]_{cf}$ is depleted much less than $[\text{U}]_{cf}$ as skeletal aragonite precipitates. While a low partition coefficient causes Rayleigh fractionation for elements in a closed system (e.g. coral $[\text{Mg}]/[\text{Ca}]_{cf}$) (Gaetani and Cohen, 2006), $[\text{CO}_3^{2-}]_{cf}$ is elevated relative to seawater and is modified by CO_2 diffusion and pH up-regulation (i.e. it is not in a closed system) (Adkins et al., 2003; Cai et al., 2016), meaning that $[\text{B}]/[\text{CO}_3^{2-}]_{cf}$ is likely not changed substantially due to skeletal aragonite precipitation. Therefore, boron-based proxies are thought to be largely dependent on carbonate chemistry alone (Trotter et al.,

2011; McCulloch et al., 2017). Finally, the combination of two carbonate system proxies (pH and $[\text{CO}_3^{2-}]$) derived from boron systematics allows for computation of the full carbonate system (Zeebe and Wolf-Gladrow, 2001).

Abiogenic laboratory experiments provide the underlying quantitative foundation necessary to apply these proxies to aragonitic coral skeletons. Klochko et al. (2006) determined the fractionation factor (α_{B3-B4}) between boric acid and borate in seawater, which allows $\delta^{11}\text{B}$ of carbonates to be used as a pH proxy when combined with knowledge of $\text{p}K_B$ (Dickson, 1990) and seawater $\delta^{11}\text{B}$ (Foster et al., 2010). Although there is potential for B isotopic fractionation between aragonite and seawater (Balan et al., 2018), the veracity of the $\delta^{11}\text{B}$ proxy has been largely confirmed by comparison with direct in-situ measurements using either pH micro-electrodes or confocal microscopy of pH-sensitive dyes in the calcifying fluid (Ries, 2011; Venn et al., 2011; Holcomb et al., 2014; Cai et al., 2016). Additionally, results from two sets of abiogenic precipitation experiments can be used to constrain the partitioning of B/Ca between fluid and aragonite (Mavromatis et al., 2015; Holcomb et al., 2016). Thus, while all the information theoretically required to constrain the full seawater carbonate system from boron systematics is now available, a variety of different approaches have been presented, especially regarding the interpretation of B/Ca partitioning (Mavromatis et al., 2015; Holcomb et al., 2016; Allison, 2017; McCulloch et al., 2017). Here, we assess the abiogenic partitioning data (Mavromatis et al., 2015; Holcomb et al., 2016), and the subsequent fitting of those data (Allison, 2017; McCulloch et al., 2017). We consider which mechanisms of B incorporation and sensitivities of B/Ca partitioning are plausible, and the implications for interpreting coral skeletons. Our focus is on the combined application of $\delta^{11}\text{B}$ and B/Ca because it is only when the two are used in tandem that it is possible to calculate the full calcifying fluid carbonate system. Finally, we present a user-friendly computer code to calculate coral calcifying fluid carbonate chemistry from measurements of $\delta^{11}\text{B}$ and B/Ca. The code also propagates known uncertainties for deriving calcifying fluid $[\text{CO}_3^{2-}]_{cf}$ and DIC_{cf} , and allows for evaluating the effects of using different constants and partition coefficient formulations.

2 Partitioning of B/Ca between aragonite and seawater

The main discrepancy among various applications of boron systematics to coral skeletons relates to the partition coefficient of boron between aragonite and seawater. Given the variety of possible exchange reactions and partition coefficients that have been proposed (Allen and Hönisch, 2012; Mavromatis et al., 2015; Holcomb et al., 2016; Allison, 2017; McCulloch et al., 2017), we begin with a brief review of how partition coefficients are derived. In general, the substitution of minor elements into a solid is described by an exchange reaction such that:



For example, the substitution of Sr^{2+} for Ca^{2+} in aragonite follows (Gaetani and Cohen, 2006):



Element distribution described by this exchange is quantified through a partition coefficient, expressed as the concentration ratio of products over reactants:

$$K_D^{Sr/Ca} = \frac{[Sr]^{aragonite}[Ca]^{fluid}}{[Sr]^{fluid}[Ca]^{aragonite}} \quad (3)$$

Equation (3) is typically rearranged as:

$$5 \quad K_D^{Sr/Ca} = \left(\frac{[Sr]^{aragonite}}{[Sr]^{fluid}} \right) \left(\frac{[Ca]^{fluid}}{[Ca]^{aragonite}} \right) = \left(\frac{[Sr]^{aragonite}}{[Sr]^{fluid}} \right) \left(\frac{[Ca]^{aragonite}}{[Ca]^{fluid}} \right)^{-1} = \frac{Sr/Ca^{aragonite}}{Sr/Ca^{fluid}} \quad (4)$$

The case of Sr^{2+} substituting for Ca^{2+} is straightforward in that the exchange reaction (Eq. 2) is charge-balanced. Boron is more complicated because it is commonly thought that the singly charged $B(OH)_4^-$ is incorporated into aragonite in place of the doubly charged CO_3^{2-} (Mavromatis et al., 2015; Noireaux et al., 2015). There are at least two possible exchange reactions for $B(OH)_4^-$ to substitute for CO_3^{2-} that maintain charge balance:



following Holcomb et al. (2016), or:



following McCulloch et al. (2017). The K_D for Eq. (5) is:

$$K_D^{B/Ca} = \frac{[B(OH)_4^-/[CO_3^{2-}]^{0.5}]^{aragonite}}{[B(OH)_4^-/[CO_3^{2-}]^{0.5}]^{fluid}} = \frac{[B/Ca]^{aragonite}}{[B(OH)_4^-/[CO_3^{2-}]^{0.5}]^{fluid}} \quad (7)$$

15 and for Eq. (6) is:


$$K_D^{B/Ca} = \frac{[B(OH)_4^-/CO_3^{2-}]^{aragonite}}{[B(OH)_4^-/CO_3^{2-}]^{fluid}} = \frac{[B/Ca]^{aragonite}}{[B(OH)_4^-/CO_3^{2-}]^{fluid}} \quad (8)$$

where $[CO_3^{2-}]^{aragonite}$ is assumed equal to $[Ca^{2+}]^{aragonite}$, and Eq. (7) and Eq. (8) differ by whether or not the square root of CO_3^{2-} is used. Since Eq. (6) includes H^+ in the products, this reaction implies that the K_D may be pH-dependent (McCulloch et al., 2017). Incorporation of B into aragonite may also involve adsorption of $B(OH)_4^-$ onto crystal surfaces, incorporation at
20 defect sites, or local charge balance by Na^+ (Balan et al., 2018).

Conversely, Allison et al. (2014) and Allison (2017) considered exchange reactions in which borate substitutes for bicarbonate (HCO_3^-), with the partition coefficient:

$$K_D^{B/Ca} = \frac{[B/Ca]^{aragonite}}{[B(OH)_4^-/HCO_3^-]^{fluid}} \quad (9)$$

This approach resolves the issue of charge balance and would account for a CO_3^{2-} reacting with H^+ , thus removing the pH
25 dependence expected from Eq. (8). However, Eq. (9) implies that aragonite forms via the reaction:



rather than:



Whether aragonite precipitates via Eq. (10) or Eq. (11) is testable because the rate of the net forward reaction should depend on the concentrations of the reactants. Burton and Walter (1987) demonstrated that the rate of aragonite precipitation increases as a function of Ω_{Ar} (where $\Omega_{Ar} = [\text{Ca}^{2+}][\text{CO}_3^{2-}]/K_{sp}$) and temperature, although they did not explicitly consider the relationship between $[\text{HCO}_3^-]$ and precipitation rate. Holcomb et al. (2016) reported bulk precipitation rates for aragonites precipitated from seawater with various $[\text{CO}_3^{2-}]$ and $[\text{HCO}_3^-]$, with independence between these two variables achieved by manipulating pH and DIC. While the bulk precipitation rates were not normalized to surface area as in Burton and Walter (1987), the experimental vessels used by Holcomb et al. (2016) were of consistent dimensions and material. Thus, the bulk precipitation rate data of Holcomb et al. (2016) should be comparable among their experiments, allowing us to evaluate between the reactions of Eq. (10) and Eq. (11). The aragonite precipitation rates reported by Holcomb et al. (2016) at 25 °C are significantly correlated with both $[\text{CO}_3^{2-}]$ ($r^2 = 0.56$, $p < 0.01$) and Ω_{Ar} ($r^2 = 0.62$, $p < 0.01$) (Figure 1a,b). Experiments conducted at 20 °C, 33 °C, and 40 °C are consistent with this trend (Figure 1a,b), and with previous observations that precipitation rate increases with temperature (Burton and Walter, 1987), although we do not attempt to quantify temperature effects on the order of the reaction (as done by Burton and Walter, 1987) since only two experiments were conducted at each temperature other than 25 °C. Conversely, there are no significant correlations between aragonite precipitation rate at 25 °C and either $[\text{HCO}_3^-]$ ($r^2 = 0.00$, $p = 0.95$) or $[\text{Ca}^{2+}][\text{HCO}_3^-]$ ($r^2 = 0.01$, $p = 0.54$) as would be expected based on Eq. (10). Other possibilities include precipitation reactions involving both CO_3^{2-} and HCO_3^- , or total DIC (Allison et al., 2014; Allison, 2017). However, there are no significant correlations between precipitation rate and either $[\text{CO}_3^{2-}] + [\text{HCO}_3^-]$ ($r^2 = 0.01$, $p = 0.59$) or DIC ($r^2 = 0.01$, $p = 0.59$) (Figure 1e,f). Together, these data lead us to conclude that aragonite precipitates from seawater via Eq. (11). Therefore, since B/Ca partition coefficients expressed with $[\text{HCO}_3^-]$ do not have a chemical reaction basis, we do not consider them further. Rather, we consider only the B/Ca partition coefficients that are based on borate substituting for CO_3^{2-} (Eqs. 7-8).

3 Fitting the experimental B/Ca partitioning data

The second source of discrepancies between various applications of boron systematics to coral skeletons is the dependence of the K_D on fluid chemistry. Holcomb et al. (2016) fit the K_D as either a function of $[\text{CO}_3^{2-}]$ or Ω_{Ar} , McCulloch et al. (2017) refit the Holcomb et al. (2016) data as a function of $[\text{H}^+]$, and Allison (2017) fit data from both Mavromatis et al. (2015) and Holcomb et al. (2016) as a function of Ω_{Ar} .

At the outset, it is important to recognize that there are two key differences between the abiogenic experiments of Mavromatis et al. (2015) and Holcomb et al. (2016). Firstly, Mavromatis et al. (2015) precipitated aragonite from NaCl solutions, whereas Holcomb et al. (2016) used filtered seawater. Secondly, $[\text{CO}_3^{2-}]$ and Ω_{Ar} are lower in the experiments of Mavromatis et al. (2015) relative to Holcomb et al. (2016). Potentially as result of one or both of these differences, Mavromatis et al. (2015) found much lower K_D values than Holcomb et al. (2016). Here, we consider four possible K_D dependencies based on these two experimental datasets (Figure 2).




Figure 1. Aragonite precipitation rates as functions of fluid chemistry based on data from Holcomb et al. (2016). Each point represents a separate abiogenic aragonite precipitation experiment conducted at 20 °C (blue), 25 °C (black), 33 °C (green), and 40 °C (red). Bulk aragonite precipitation rates (R) are plotted against mean fluid $[\text{CO}_3^{2-}]$ (a), Ω_{Ar} (b), $[\text{HCO}_3^-]$ (c), $[\text{Ca}^{2+}][\text{HCO}_3^-]$ (d), $[\text{CO}_3^{2-}] + [\text{HCO}_3^-]$ (e), and DIC (f). Solid lines show regression fits at each temperature (note that there are only two experiments at each temperature other than 25 °C, and thus lines fit for these temperatures should be interpreted with caution).

The first two formulations assume that there are substantial compositional effects on B/Ca partitioning, and thus the offsets in K_D between Mavromatis et al. (2015) and Holcomb et al. (2016) arise due to the use of NaCl versus seawater solutions, respectively (Figure 2a,b). If this is correct, the Holcomb et al. (2016) data are more appropriate for application to corals based on evidence that they precipitate their skeletons from seawater-based solutions (McConnaughey, 1989; Cohen and

5 McConnaughey, 2003; Gagnon et al., 2012; Tambutté et al., 2012). Holcomb et al. (2016) precipitated their aragonites from seawater solutions modified with $[\text{Ca}^{2+}]$ and $[\text{Mg}^{2+}]$ ranging between 6 and 20 mmol kg^{-1} , and 48 and 98 mmol kg^{-1} , respectively, without any apparent effects on the B/Ca K_D . While this suggests that the K_D is not highly sensitive to seawater elemental chemistry, it is still possible that there are subtle compositional effects that have little influence on K_D in modified seawater, but become apparent in the NaCl solutions used by Mavromatis et al. (2015). Assuming some compositional effects

10 do exist, we are left with the two plausible K_D expressions (Eqs. 7-8), and the previously presented dependencies on either $[\text{CO}_3^{2-}]$ (Holcomb et al., 2016) or $[\text{H}^+]$ McCulloch et al. (2017).

Alternatively, it is possible that there are negligible effects from using NaCl or seawater solutions and, therefore, the data from both Mavromatis et al. (2015) and Holcomb et al. (2016) should be fit by a single, continuous function. There are again two plausible formulations: K_D increases as a function of $[\text{CO}_3^{2-}]$ or Ω_{Ar} (Figure 2c,d). Allison (2017) proposed a linear fit between K_D and Ω_{Ar} that includes both the Mavromatis et al. (2015) and Holcomb et al. (2016) data. From a practical standpoint, however, this latter approach is problematic in that it requires an independent proxy for $[\text{Ca}^{2+}]$ (see section 8) and the linear fit effectively precludes its use for deriving coral calcifying fluid chemistry (see section 4). In an attempt to avoid these issues, we introduce a logarithmic relationship between K_D and $[\text{CO}_3^{2-}]$, which fits both the Mavromatis et al. (2015) and Holcomb et al. (2016) data (Figure 2c):

$$K_D^{\text{B/Ca}} = 0.00077(\pm 0.00007) * \ln([\text{CO}_3^{2-}]) - 0.0028(\pm 0.0004) \quad (12)$$

where parentheses indicate 95% confidence, K_D is defined by Eq. (8), and $[\text{CO}_3^{2-}]$ is in units of $\mu\text{mol kg}^{-1}$. Mechanistically, the increase in K_D with $[\text{CO}_3^{2-}]$ or Ω_{Ar} (or precipitation rate) is consistent with the surface entrapment model proposed by Watson (2004). In this model, minor element impurities, such as B, are incorporated in the near-surface layer of a growing crystal. Slower growing crystals allow these impurities to diffuse out of the near-surface region into the fluid, whereas faster growing crystals bury the near-surface impurities into the bulk crystal. The sensitivity of K_D to $[\text{CO}_3^{2-}]$ or Ω_{Ar} is also consistent with a surface kinetic model (DePaolo, 2011), in which trace element partitioning depends on the net rate of precipitation relative to dissolution. Thus, both the surface entrapment and kinetic models offer potential explanations as to why the low- Ω_{Ar} experiments of Mavromatis et al. (2015) produced lower K_D than the higher- Ω_{Ar} experiments of Holcomb et al. (2016).

4 Back-application of partition coefficient formulations to abiogenic datasets

We conducted a simple test to evaluate the utility of the four K_D dependencies considered above. For each K_D formulation, we used the reported aragonite B/Ca, fluid $[\text{B}(\text{OH})_4^-]$, and pH data of Mavromatis et al. (2015) and Holcomb et al. (2016) to calculate the fluid $[\text{CO}_3^{2-}]$, and then we compared the predicted $[\text{CO}_3^{2-}]$ to the concentrations measured during the experiments (Figure 3) (see also Ross et al. (2017) for a similar analysis). The basis for this approach is to assess how well the experimental fluid $[\text{CO}_3^{2-}]$ can be reconstructed using boron systematics alone. When boron systematics are applied to coral skeletons, $[\text{CO}_3^{2-}]$ is predicted from only B/Ca and $\delta^{11}\text{B}$. However, since $\delta^{11}\text{B}$ was not reported by Holcomb et al. (2016), we instead use the measured pH for the McCulloch et al. (2017) K_D formulation. Additionally, since [B] was manipulated in some experiments, we use reported fluid $[\text{B}(\text{OH})_4^-]$ instead of calculating it from pH as is done in applications to corals (Allison et al., 2014; McCulloch et al., 2017). Nevertheless, since pH (and thus seawater $[\text{B}(\text{OH})_4^-]$) are readily calculated from $\delta^{11}\text{B}$, our approach is suitable for evaluating the utility of each K_D formulation for reconstructing $[\text{CO}_3^{2-}]$ with B/Ca.

Since three of the K_D formulations (Holcomb et al. (2016), Allison (2017), and our new Eq. 12) themselves depend on $[\text{CO}_3^{2-}]$, we solved for $[\text{CO}_3^{2-}]$ as follows. An initial guess of $[\text{CO}_3^{2-}]$ was used to calculate an initial K_D , and this K_D was used to solve for $[\text{CO}_3^{2-}]$ by rearranging Eq. (7) to:

$$[\text{CO}_3^{2-}] = \left(K_D^{\text{B/Ca}} \frac{[\text{B}(\text{OH})_4^-]_{\text{fluid}}}{[\text{B/Ca}]_{\text{aragonite}}} \right)^2 \quad (13)$$




Figure 2. B/Ca K_D formulations. Abiogenic B/Ca partitioning data from Mavromatis et al. (2015) (red circles) and Holcomb et al. (2016) (blue triangles) fit as functions of fluid chemistry: $[\text{CO}_3^{2-}]$ (a,c) (Holcomb et al., 2016), $[\text{H}^+]$ (b) (McCulloch et al., 2017), and Ω_{Ar} (d) (Allison, 2017). Note that K_D in (a) is defined with Eq. (7) and in (b-d) is defined with Eq. (8). We use only the Mavromatis et al. (2015) with $[\text{B}] < 1000 \mu\text{mol kg}^{-1}$ due to the apparent effect of $[\text{B}]$ on K_D (Holcomb et al., 2016).

and Eq. (8) to:

$$[\text{CO}_3^{2-}] = K_D^{\text{B/Ca}} \frac{[\text{B(OH)}_4^-]_{\text{fluid}}}{[\text{B/Ca}]_{\text{aragonite}}} \quad (14)$$

where Eq. (14) is used for Allison (2017) and our new Eq. (12), and Eq. (13) is used for Holcomb et al. (2016). We then calculated the residual between the calculated (Eqs. 13-14) and initially estimated $[\text{CO}_3^{2-}]$. Finally, we iteratively adjusted the initial $[\text{CO}_3^{2-}]$ estimate for each data point until it equaled the $[\text{CO}_3^{2-}]$ derived from Eqs. (13-14).

Both the Holcomb et al. (2016) fit (their equation 7) and the McCulloch et al. (2017) refit perform similarly, effectively reconstructing the fluid $[\text{CO}_3^{2-}]$ of the Holcomb et al. (2016) experimental data (root mean square error, RMSE = 151 and 163 $\mu\text{mol kg}^{-1}$, respectively), but performing poorly for the Mavromatis et al. (2015) data (RMSE = 1370 and 1385 $\mu\text{mol kg}^{-1}$, respectively) (Figure 3a,b). This is not surprising because these K_D dependencies are offset from the Mavromatis et al. (2015) data (Figure 2a,b). Our new logarithmic equation performs well for both datasets (RMSE = 42 and 204 $\mu\text{mol kg}^{-1}$ for Mavromatis et al. (2015) and Holcomb et al. (2016), respectively). The Allison (2017) formulation (assuming $[\text{Ca}^{2+}]$ of 10 mmol kg^{-1}) performs well for the Mavromatis et al. (2015) data (RMSE = 51 $\mu\text{mol kg}^{-1}$), but creates a trend opposite




Figure 3. Reconstructing experimental fluid $[\text{CO}_3^{2-}]$ using the K_D formulations presented in Figure 2. Symbols are the same as Figure 2. In panel (d), negative $[\text{CO}_3^{2-}]$ solutions have been excluded (see Appendix). Calculations using the Allison (2017) K_D formulation have been performed with both assuming seawater $[\text{Ca}^{2+}]$ (blue) and using the $[\text{Ca}^{2+}]$ reported from the experiments (cyan).

that expected for the Holcomb et al. (2016) data (RMSE = $1375 \mu\text{mol kg}^{-1}$) (Figure 3d). Using the reported $[\text{Ca}^{2+}]$ and K_{sp} from the Holcomb et al. (2016) experiments in the Allison (2017) formulation improves the results slightly and generates more positive solutions, but the RMSE is still $950 \mu\text{mol kg}^{-1}$.

An alternative way to understand these patterns is to investigate the relationship between $[\text{CO}_3^{2-}]$ and the ratio of fluid $[\text{B}(\text{OH})_4^-]$ to solid B/Ca (Figure 4). Following Eqs. (13-14), $[\text{CO}_3^{2-}]$ should be positively related to $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}$, and this behavior is clearly evident in the abiogenic aragonites of Holcomb et al. (2016) (blue triangles in Figure 4). The K_D formulations of Holcomb et al. (2016), McCulloch et al. (2017), and our new Eq. (12) all closely track the abiogenic data, especially for $[\text{CO}_3^{2-}] < 2000 \mu\text{mol kg}^{-1}$. Conversely, the Allison (2017) fit (assuming $[\text{Ca}^{2+}]$ of 10 mmol kg^{-1}) produces the opposite trend and is invalid or negative below a $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}$ of $\sim 0.44 \text{ mol kg}^{-1}$ (see Appendix for derivation of an analytical solution).

The behavior of the K_D formulations can be understood by inspecting the residuals between initial $[\text{CO}_3^{2-}]$ estimates and those derived from Eqs. (13-14) (Figure 5). The Holcomb et al. (2016) K_D formulation generates unique $[\text{CO}_3^{2-}]$ solutions (*i.e.* where the residual equals zero) that increase with $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}$ (Figure 5a), which is the ideal behavior. Our new Eq. (12) also produces increasing $[\text{CO}_3^{2-}]$ solutions with increasing $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}$ (Figure 5b), however, a major issue of this formulation is




Figure 4. Experimental fluid $[\text{CO}_3^{2-}]$ as a function of $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$. The K_D formulations of Holcomb et al. (2016) (dotted black line), McCulloch et al. (2017) (black crosses), and Eq. (12) (dashed black line) all capture the trend of increasing $[\text{CO}_3^{2-}]$ with increasing $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ that is apparent in the abigenic data (blue triangles). A constant K_D (solid grey line) underestimates the slope between $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ and $[\text{CO}_3^{2-}]$. The pink shaded region shows the range of $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ derived for *Porites* corals by McCulloch et al. (2017).

that there may be two $[\text{CO}_3^{2-}]$ solutions for each $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$. Finally, although the Allison (2017) K_D formulation produces unique $[\text{CO}_3^{2-}]$ solutions, they increase with decreasing $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ (Figure 5c), opposite to that expected (Figure 4).

The reason for the poor behavior of the Allison (2017) formulation is the linear fit between K_D and Ω_{Ar} with an intercept near the origin. When using this formulation to predict $[\text{CO}_3^{2-}]$ from boron systematics alone, we must assume $[\text{Ca}^{2+}]$ is approximately equal to seawater ($\sim 10 \text{ mmol kg}^{-1}$), meaning that Ω_{Ar} is directly related to $[\text{CO}_3^{2-}]$. Since the intercept in the Allison (2017) K_D formulation is close to the origin, any change in $[\text{CO}_3^{2-}]$ results in an almost proportional change in K_D . It can be seen why this is problematic by inspecting how $[\text{CO}_3^{2-}]$ is derived from Eq. (14). The $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ is derived from pH (or $\delta^{11}\text{B}$) and measured B/Ca, so this ratio remains constant while we find the appropriate K_D that minimizes the residual $[\text{CO}_3^{2-}]$, as in Figure 5. Therefore, Eq. (14) is effectively reduced to $[\text{CO}_3^{2-}]$ being a function of K_D multiplied by a constant. However, since K_D changes almost directly proportional to $[\text{CO}_3^{2-}]$ according to Allison (2017), it is difficult to find a $[\text{CO}_3^{2-}]$ that explains different $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$. Although Allison (2017) recognized the difficulty of explaining the range of B/Ca observed in corals (see their Figure 8g), the implication of applying this K_D formulation to predict $[\text{CO}_3^{2-}]$ was not discussed. Our analysis suggests that this K_D formulation is poorly suited for accurately reconstructing fluid $[\text{CO}_3^{2-}]$ from boron systematics (Figure 3d, Figure 4).

Another approach presented by Allison (2017) is to use a constant K_D . We selected a K_D value of 0.02 as an example that fits the abigenic data near the low- $[\text{CO}_3^{2-}]$ range of the data (Figure 4). While a constant K_D performs better than the linear fit to Ω_{Ar} , it underestimates the slope of the relationship between $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ and $[\text{CO}_3^{2-}]$ (Figure 4). This is not surprising




Figure 5. Predicting $[\text{CO}_3^{2-}]$ from the K_D formulations which themselves depend on $[\text{CO}_3^{2-}]$: Holcomb et al. (2016) (a), Eq. (12) (b), and Allison (2017) (c). Each panel shows the residual between a guess of $[\text{CO}_3^{2-}]$ used to calculate K_D and that calculated from Eqs. (13-14), plotted against the $[\text{CO}_3^{2-}]$ guess. The final $[\text{CO}_3^{2-}]$ is derived by finding where the residual is minimized for a particular $\frac{[\text{B}(\text{OH})_4^-]_{\text{fluid}}}{[\text{B}/\text{Ca}]_{\text{aragonite}}}$ (three of which are plotted as examples in red, blue, and black).

because the abiogenic data clearly show the K_D does not remain constant as $[\text{CO}_3^{2-}]$ changes (Figure 2). Since using a constant K_D will underestimate variability in $[\text{CO}_3^{2-}]_{cf}$ when applied to corals, we do not recommend this approach.

5 Application to deriving coral calcifying fluid carbonate chemistry

The ability of boron systematics to predict two independent carbonate chemistry parameters allows for calculation of the full carbonate system. This has prompted several recent applications deriving the carbonate chemistry of coral calcifying fluids (Allison et al., 2014; Comeau et al., 2017; D’Olivo and McCulloch, 2017; Kubota et al., 2017; McCulloch et al., 2017; Ross et al., 2017; Schoepf et al., 2017). Here, we investigate the differences in derived coral calcifying fluid $[\text{CO}_3^{2-}]$ that arise from the choice of K_D formulation. We use the paired $\delta^{11}\text{B}$ and B/Ca data of the “Davies 2” coral from McCulloch et al. (2017) as an example.

- Derived $[\text{CO}_3^{2-}]_{cf}$ shows similar seasonality when using the K_D formulations of Holcomb et al. (2016), McCulloch et al. (2017), or our new Eq. (12) (Figure 6). Regardless of which of these three K_D formulations are used, $[\text{CO}_3^{2-}]_{cf}$ is highest in summer and lowest in winter over a multi-year time series. This is consistent with other reports of B/Ca seasonality in coral skeletons (Sinclair, 2005; D’Olivo and McCulloch, 2017; McCulloch et al., 2017; Ross et al., 2017), and with an independent approach based on Rayleigh modelling of minor elements in coral skeleton (Gaetani and Cohen, 2006; Gaetani et al., 2011). The primary difference among the derived values is that the K_D formulations from Holcomb et al. (2016) and our Eq. (12) produce seasonal cycles with ~50% greater amplitude relative to the McCulloch et al. (2017) K_D formulation. The absolute values of derived $[\text{CO}_3^{2-}]_{cf}$ is approximately equal for all three formulations at the summertime maxima, but are lower during winter when using the K_D formulations from Holcomb et al. (2016) or our Eq. (12), relative to McCulloch et al. (2017). Conversely, using the Allison (2017) K_D formulation produces the opposite seasonal pattern with amplitude several times greater than the other K_D formulations. This large discrepancy is not surprising given the behavior of the Allison (2017) K_D formulation when retrospectively applied to the fluid composition of abiogenic aragonites (Figure 3).




Figure 6. Application of the four K_D formulations for the “Davies 2” *Porites* coral data from McCulloch et al. (2017). Derived $[\text{CO}_3^{2-}]_{cf}$ is plotted over multiple years using the K_D formulations of Holcomb et al. (2016) (black), McCulloch et al. (2017) (red), Eq. (12) (blue), and Allison (2017) (dashed grey line). Shading represents 1 standard deviation of the systematic errors due to uncertainty in each K_D formulation. Note that (1) the Holcomb et al. (2016) and the Eq. (12) lines plot nearly on top of each other, and (2) $[\text{CO}_3^{2-}]_{cf}$ derived from the Allison (2017) K_D formulation corresponds to the right y-axis.

6 A computer code for applying boron systematics to coral skeletons

We present here a user-friendly computer code for deriving $[\text{CO}_3^{2-}]_{cf}$ and DIC_{cf} from boron systematics (supplemental files). The function is provided in both MATLAB and R formats, and it calculates $[\text{CO}_3^{2-}]_{cf}$ and DIC_{cf} given inputs of $\delta^{11}\text{B}$, B/Ca, temperature, salinity, and water depth. It allows easy toggling between what we consider the three plausible K_D formulations (Holcomb et al. (2016), McCulloch et al. (2017), and our new Eq. 12). Furthermore, the code permits a choice of $[\text{B}]_{sw}$ functions since Allison et al. (2014) and Allison (2017) used the relation between salinity and $[\text{B}]_{sw}$ from Uppstrom (1974), whereas D’Olivio and McCulloch (2017) and McCulloch et al. (2017) used that of Lee et al. (2010). The carbonate dissociation constants can also be toggled between Dickson and Millero (1987) and Lueker et al. (2000). The code follows the calculations of CO2SYS (Lewis et al., 1998) for converting between pH scales and accounting for pressure effects on equilibrium constants, and uses the $\delta^{11}\text{B}_{sw}$ of Foster et al. (2010) and the α_{B3-B4} of Klochko et al. (2006).

Perhaps most importantly, the code propagates known uncertainties into the derivation of $[\text{CO}_3^{2-}]_{cf}$ and DIC_{cf} . These uncertainties are estimated using a Monte Carlo scheme, in which random errors (assuming Gaussian distributions) are added to parameters while repeating the calculations many times. The non-systematic uncertainty of derived values depends on the measurement precisions of $\delta^{11}\text{B}$, B/Ca, temperature, and salinity. These will depend on the instruments and protocols used, and for $\delta^{11}\text{B}$ and B/Ca should be estimated by each laboratory, for example by repeated measurements of an external consistency standard. The systemic errors of derived values depend on the uncertainties of the various K_D formulations, uncertainties associated with $\delta^{11}\text{B}_{sw}$ (Foster et al., 2010), $[\text{B}]_{sw}$ (Lee et al., 2010), α_{B3-B4} (Klochko et al., 2006), and pK_B (Dickson, 1990); and if known, any uncertainties in the accuracy of $\delta^{11}\text{B}$, B/Ca, temperature, and salinity measurements.

7 Relationships among coral calcifying fluid carbonate chemistry parameters

With our code, the parameter space of $[\text{CO}_3^{2-}]_{cf}$ derived from $\delta^{11}\text{B}$ and B/Ca, and the differences among K_D formulations, can be readily visualized (Figure 7). This enables future applications of boron systematics to coral skeletons to consider how the choice of K_D formulation affects the particular question being investigated. We also apply the code to calculate carbonate system parameters using published $\delta^{11}\text{B}$ and B/Ca datasets (Figure 8). Coral $\delta^{11}\text{B}$ is tightly related to pH_{cf} , varying only slightly with changes in seawater temperature and salinity (Figure 8b). Likewise, B/Ca is primarily a function of $[\text{CO}_3^{2-}]_{cf}$, but also depends in part on borate concentration, and hence on pH_{cf} (Figure 8c). For this reason, it is difficult to directly interpret coral B/Ca, and instead we recommend pairing $\delta^{11}\text{B}$ and B/Ca to calculate the full calcifying fluid carbonate chemistry.

Interestingly, this analysis shows that coral calcifying fluid $[\text{CO}_3^{2-}]_{cf}$ and DIC are consistently positively correlated across studies (Figure 8f), whereas the sign of correlations between pH and both $[\text{CO}_3^{2-}]_{cf}$ and DIC varies (Figure 8d-e). Assuming $[\text{CO}_3^{2-}]_{cf}$ is the carbonate system parameter most important for aragonite precipitation, these patterns may suggest that elevating DIC_{cf} is critical to the coral calcification process, although up-regulating pH is still important for shifting the carbonate system to favor CO_3^{2-} over HCO_3^- . In addition, the large changes in pH, DIC, and $[\text{CO}_3^{2-}]$ that occur within the calcifying fluid relative to natural variability in the open ocean likely precludes the utility of boron systematics for reconstructing seawater carbonate chemistry, reinforcing previous conclusions made for both corals and foraminifera (Yu et al., 2010; Rae et al., 2011; Rollion-Bard et al., 2011). Rather, the boron systematics of coral skeletons are primarily useful for investigating calcifying fluid dynamics and understanding coral biomineralization.

In contrast to boron systematics, which consistently show elevated DIC_{cf} , micro-electrode measurements of $[\text{CO}_3^{2-}]_{cf}$ and pH_{cf} imply that DIC_{cf} is depleted (Cai et al., 2016), relative to that of seawater (typically $< 2000 \mu\text{mol kg}^{-1}$). The reason for this discrepancy is not yet clear, and resolving it should be a high priority because whether DIC_{cf} is greater than or less than seawater implies different calcification strategies. Reducing DIC_{cf} may be an efficient strategy to increase pH_{cf} because the reduced buffering capacity means that less energy is required to elevate pH via proton pumping (Cai et al., 2016). Alternatively, increasing DIC_{cf} means that a higher Ω_{Ar} is achieved for a given pH_{cf} . Deciphering between these possibilities has key implications for whether calcification is limited by DIC_{cf} (or CO_2 diffusion into the calcifying fluid), or by Ω_{Ar} . Two independent approaches to quantifying calcifying fluid carbonate chemistry are consistent with the high DIC_{cf} scenario. First, coral U/Ca ratios imply that DIC_{cf} is between 2600 and 6100 $\mu\text{mol kg}^{-1}$ (DeCarlo et al., 2015), which is a similar range to that derived from boron systematics (Figure 8). Second, boron systematics-derived $[\text{CO}_3^{2-}]$ is consistent with a combination of Raman spectroscopy-derived Ω_{Ar} and trace element ratios (Mg/Ca and Sr/Ca) (DeCarlo et al., 2017). Nevertheless, since low DIC_{cf} has been derived from micro-electrodes in several species (Cai et al., 2016), studies combining multiple approaches (*i.e.* geochemistry and micro-electrodes) on the same specimens will be essential for resolving the DIC_{cf} discrepancy.




Figure 7. Application of our computer code to visualizing the parameter space of $[\text{CO}_3^{2-}]$ (in $\mu\text{mol kg}^{-1}$) derived from B/Ca and $\delta^{11}\text{B}$ at 25 °C and salinity 35. The upper left panel shows absolute $[\text{CO}_3^{2-}]$ derived with the K_D of Holcomb et al. (2016) (“H16”), whereas the other panels show the differences in $[\text{CO}_3^{2-}]$ between the K_D formulations of H16, McCulloch et al. (2017) (“M17”), and our new Eq. (12). The black dots show coral data from the literature (see Figure 8 legend below). Note that the actual $[\text{CO}_3^{2-}]$ derived for the coral data will also depend on variations of the *in situ* temperature and salinity, which are not accounted for in the plots.

8 Which K_D formulation to use?

Despite the availability of abiogenic B/Ca partitioning data from two experiments (Mavromatis et al., 2015; Holcomb et al., 2016), and several attempts to fit the data (Holcomb et al., 2016; Allison, 2017; McCulloch et al., 2017), it is important to recognize that uncertainties still remain, in particular an understanding of the controlling factors, and thus the appropriate K_D formulation to apply. From a mechanistic viewpoint, the key fundamental question that remains is whether the abiogenic data of Mavromatis et al. (2015) and Holcomb et al. (2016) are directly comparable and thus should be fit with a continuous function (*e.g.* Eq. 12), or if they are incomparable because Mavromatis et al. (2015) used NaCl solutions and Holcomb et al. (2016) used seawater. If they are comparable, then our new Eq. (12) or a similar fit to both datasets is the most appropriate K_D formulation. Calcite precipitation studies provide some support for the hypothesis that crystal growth rate or Ω_{Ar} influences B/Ca partitioning (Ruiz-Agudo et al., 2012; Uchikawa et al., 2015, 2017), but it is not yet known if these results can be extended to aragonite precipitation from seawater. Alternatively, if the solution chemistry makes the two experiments incomparable, the





Figure 8. Correlations among coral calcifying fluid carbonate system parameters based on published boron systematics datasets: (a) B/Ca and $\delta^{11}\text{B}$, (b) pH_{cf} and $\delta^{11}\text{B}$, (c) $[\text{CO}_3^{2-}]_{cf}$ and B/Ca, (d) pH and $[\text{CO}_3^{2-}]_{cf}$, (e) pH_{cf} and DIC_{cf} , and (f) $[\text{CO}_3^{2-}]_{cf}$ and DIC_{cf} . Colors show different studies, and lines are plotted for significant ($p < 0.05$) correlations using all the data within each study. Grey area shows the convex hull of the parameter space covered in the abiogenic experiments of Holcomb et al. (2016). Calculations are performed using the Holcomb et al. (2016) K_D formulation.

Holcomb et al. (2016) K_D data are most likely the more suitable choice for corals because the experiments were conducted with seawater at comparable Ω_{Ar} to that of coral calcifying fluids (DeCarlo et al., 2017), and they can be fit as either a function of $[\text{CO}_3^{2-}]$ or $[\text{H}^+]$. However, it is important to recognize that the parameter space of CO_2 system parameters covered in the Holcomb et al. (2016) experiments includes some, but not all, of the published coral data (Figure 8). Further, since we are

5 unable to conclusively distinguish whether the two abiogenic datasets are directly comparable, all three K_D formulations may be considered equally valid until proven otherwise. Additional abiogenic experiments aimed at this question will clearly be useful in refining the boron systematics proxies.

From a practical standpoint, the K_D formulations of Holcomb et al. (2016) and McCulloch et al. (2017) may be the most appropriate. Both produce unique solutions of $[\text{CO}_3^{2-}]$ that increase with $\frac{[\text{B}(\text{OH})_4^-]_{fluid}}{[\text{B}/\text{Ca}]_{aragonite}}$, and they effectively reconstruct fluid

10 $[\text{CO}_3^{2-}]$ using the abiogenic aragonites precipitated from seawater. While our Eq. (12) produces $[\text{CO}_3^{2-}]_{cf}$ estimates that are

nearly identical under most $\delta^{11}\text{B}$ and B/Ca combinations to those derived using the Holcomb et al. (2016) K_D formulation (Figure 7), Eq. (12) can have non-unique solutions, which could complicate interpretations of $[\text{CO}_3^{2-}]_{cf}$ in some cases.

A final consideration is that two of the K_D formulations (Holcomb et al. (2016) and our new Eq. 12) are fit to $[\text{CO}_3^{2-}]$. Fitting Eq. (12) to a wider range of $[\text{CO}_3^{2-}]$ helps to account for the different solution chemistries and associated growth rates of the two abiogenic precipitation studies (Mavromatis et al., 2015; Holcomb et al., 2016), but Ω_{Ar} or crystal growth rate may be the true controlling factor (Watson, 2004; van der Weijden and van der Weijden, 2014). However, Holcomb et al. (2016) did not find a temperature dependence of B/Ca partitioning, as would be expected if precipitation rate influenced K_D . While growth rate is likely related to $[\text{CO}_3^{2-}]$ (Burton and Walter, 1987), the two could decouple with changes in temperature or if coral calcifying fluid $[\text{Ca}^{2+}]_{cf}$ departs from seawater levels. Recent evidence combining Raman spectroscopy with boron systematics suggests $[\text{Ca}^{2+}]_{cf}$ is within ~20% of seawater (DeCarlo et al., 2017), but this has yet to be tested on a range of coral species and locations. Thus, future abiogenic experiments designed to test under what conditions $[\text{CO}_3^{2-}]$ or crystal growth rates control B/Ca partitioning, as well as development of proxies for $[\text{Ca}^{2+}]_{cf}$, may improve the accuracy of deriving calcifying fluid carbonate chemistry from boron systematics.

9 Conclusions

Recent abiogenic aragonite precipitation experiments have made possible the application of boron systematics to quantifying the full carbonate system of coral calcifying fluid. However, a number of approaches to doing so have been utilized (Allison et al., 2014; Allison, 2017; D’Olivo and McCulloch, 2017; McCulloch et al., 2017), without a comprehensive analysis of which K_D formulations are plausible (*i.e.* can reproduce the experimental fluid chemistry) or the implications for interpreting coral skeletons. We evaluated four potential B/Ca K_D formulations involving $\text{B}(\text{OH})_4^-$ substituting for CO_3^{2-} in the aragonite lattice. Our analysis suggests that there are at least three plausible K_D formulations (Holcomb et al. (2016), McCulloch et al. (2017), and our new Eq. 12) that can be used to determine the K_D and its dependence on fluid chemistry. Despite the differences in plausible approaches, we show that all three produce similar patterns in derived coral calcifying fluid carbonate chemistry. Nevertheless, subtle differences in derived carbonate chemistry remain among the approaches, and addressing these differences should be the target of future abiogenic aragonite precipitation experiments. Finally, we present a code that computes coral calcifying fluid carbonate chemistry from boron systematics, and allows for comparison among different K_D formulations.

Code availability. Codes are available in the Supplement

Appendix A

In the main text, we used numerical solutions to predict $[\text{CO}_3^{2-}]$ based on Eq. (14). Here, we show an analytical solution to Eq. (14) for the Allison (2017) K_D formulation. Allison (2017) fit K_D to Ω_{Ar} with a linear regression in the form:

$$K_D^{\text{B/Ca}} = a\Omega + b \quad (\text{A1})$$

5 where

$$\Omega = \frac{[\text{CO}_3^{2-}][\text{Ca}^{2+}]}{K_{sp}} \quad (\text{A2})$$

with concentrations in units of mol kg^{-1} and where K_{sp} is the solubility product. Inserting Eq. (A2) into Eq. (A1):

$$K_D^{\text{B/Ca}} = a \frac{[\text{CO}_3^{2-}][\text{Ca}^{2+}]}{K_{sp}} + b \quad (\text{A3})$$

and then inserting Eq. (A3) into Eq. (14) of the main text:

$$10 \quad [\text{CO}_3^{2-}] = \left(a \frac{[\text{CO}_3^{2-}][\text{Ca}^{2+}]}{K_{sp}} + b \right) \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} \quad (\text{A4})$$

where $[\text{CO}_3^{2-}]$ is in units of mol kg^{-1} . We must now solve Eq. (A4) for $[\text{CO}_3^{2-}]$. First, expand the right side of the equation:

$$[\text{CO}_3^{2-}] = a \frac{[\text{CO}_3^{2-}][\text{Ca}^{2+}]}{K_{sp}} \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} + b \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} \quad (\text{A5})$$

Multiply both sides by K_{sp} :

$$[\text{CO}_3^{2-}]K_{sp} = a[\text{CO}_3^{2-}][\text{Ca}^{2+}] \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} + b \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} K_{sp} \quad (\text{A6})$$

15 Collect all the $[\text{CO}_3^{2-}]$ terms on the left side of the equation:

$$[\text{CO}_3^{2-}]K_{sp} - a[\text{CO}_3^{2-}][\text{Ca}^{2+}] \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} = b \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} K_{sp} \quad (\text{A7})$$

Factor out $[\text{CO}_3^{2-}]$:

$$[\text{CO}_3^{2-}] \left(K_{sp} - a[\text{Ca}^{2+}] \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} \right) = b \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} K_{sp} \quad (\text{A8})$$

Solve for $[\text{CO}_3^{2-}]$:

$$20 \quad [\text{CO}_3^{2-}] = \frac{b \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} K_{sp}}{K_{sp} - a[\text{Ca}^{2+}] \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}} \quad (\text{A9})$$

In seawater at 25 °C and salinity 34, $[\text{Ca}^{2+}]$ is approximately 0.01 mol kg^{-1} and K_{sp} is 6.54×10^{-7} (Riley and Tongudai, 1967; Lewis et al., 1998). According to Allison (2017), a is 1.48×10^{-4} and b is -1.30×10^{-4} . Inserting these values in Eq. (A9):

$$[\text{CO}_3^{2-}] = \frac{(-1.30 \times 10^{-4}) \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}} (6.54 \times 10^{-7})}{(6.54 \times 10^{-7}) - (1.48 \times 10^{-4})(0.01) \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}} = \frac{-8.50 \times 10^{-11} \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}}{6.54 \times 10^{-7} - 1.48 \times 10^{-6} \frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B/Ca}]^{aragonite}}} \quad (\text{A10})$$

The denominator equals zero (*i.e.* the solution is undefined) when $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}} = \frac{6.54 \times 10^{-7}}{1.48 \times 10^{-6}} = 0.44$. If $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}} < 0.44$, then the denominator is positive, and since the numerator is always negative, this means that the predicted $[\text{CO}_3^{2-}]$ will be negative. Predicted $[\text{CO}_3^{2-}]$ will be highest when the denominator is a small negative number, which occurs when $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ is slightly greater than 0.44. As $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ increases $\gg 0.44$, the absolute value of the denominator increases more than that of the numerator because the coefficient attached to $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ is raised to the -6 power in the denominator and to the -11 power in the numerator. The implication is that predicted $[\text{CO}_3^{2-}]$ will decrease as $\frac{[\text{B}(\text{OH})_4^-]^{fluid}}{[\text{B}/\text{Ca}]^{aragonite}}$ increases beyond 0.44. This is the same conclusion reached in the main text, and is the opposite trend to that observed in the abiogenic aragonites (Figure 4).

Competing interests. The authors declare no competing interests.

Acknowledgements. The authors thank Glenn Gaetani for valuable comments. This study was funded by an ARC Laureate Fellowship (FL120100049) awarded to M.T.M., and the ARC Centre of Excellence for Coral Reef Studies (CE140100020).

References

- Adkins, J. F., Boyle, E. A., Curry, W. B., and Lutringer, A.: Stable isotopes in deep-sea corals and a new mechanism for "vital effects", *Geochimica et cosmochimica acta*, 67, 1129–1143, 2003.
- Al-Horani, F. A., Al-Moghrabi, S. M., and De Beer, D.: The mechanism of calcification and its relation to photosynthesis and respiration in the scleractinian coral *Galaxea fascicularis*, *Marine Biology*, 142, 419–426, <https://doi.org/10.1007/s00227-002-0981-8>, 2003.
- Allen, K. A. and Hönisch, B.: The planktic foraminiferal B/Ca proxy for seawater carbonate chemistry: A critical evaluation, *Earth and Planetary Science Letters*, 345, 203–211, 2012.
- Allison, N.: Reconstructing coral calcification fluid dissolved inorganic carbon chemistry from skeletal boron: An exploration of potential controls on coral aragonite B/Ca, *Heliyon*, 3, e00387, <https://doi.org/10.1016/j.heliyon.2017.e00387>, 2017.
- Allison, N., Cohen, I., Finch, A. A., Erez, J., and Tudhope, A. W.: Corals concentrate dissolved inorganic carbon to facilitate calcification., *Nature communications*, 5, 5741, <https://doi.org/10.1038/ncomms6741>, 2014.
- Balan, E., Noireaux, J., Mavromatis, V., Saldi, G. D., Montouillout, V., Blanchard, M., Pietrucci, F., Gervais, C., Rustad, J. R., Schott, J., and Gaillardet, J.: Theoretical isotopic fractionation between structural boron in carbonates and aqueous boric acid and borate ion, *Geochimica et Cosmochimica Acta*, 222, 117–129, <https://doi.org/10.1016/j.gca.2017.10.017>, 2018.
- Barnes, D. J.: Coral skeletons: an explanation of their growth and structure, *Science*, 170, 1305–1308, <https://doi.org/10.1126/science.170.3964.1305>, 1970.
- Burton, E. A. and Walter, L. M.: Relative precipitation rates of aragonite and Mg calcite from seawater: Temperature or carbonate ion control?, *Geology*, 15, 111–114, 1987.
- Cai, W.-J., Ma, Y., Hopkinson, B. M., Grottole, A. G., Warner, M. E., Ding, Q., Hu, X., Yuan, X., Schoepf, V., Xu, H., Han, C., Melman, T. F., Hoadley, K. D., Pettay, D. T., Matsui, Y., Baumann, J. H., Levas, S., Ying, Y., and Wang, Y.: Microelectrode characterization of coral daytime interior pH and carbonate chemistry., *Nature communications*, 7, 11144, <https://doi.org/10.1038/ncomms11144>, 2016.
- Cohen, A. L. and McConnaughey, T. A.: Geochemical Perspectives on Coral Mineralization, *Rev. Mineral. Geochem.*, 54, 151–187, <https://doi.org/10.2113/0540151>, 2003.
- Comeau, S., Tambutté, E., Carpenter, R. C., Edmunds, P. J., Evensen, N. R., Allemand, D., Ferrier-Pagès, C., Tambutté, S., and Venn, A. A.: Coral calcifying fluid pH is modulated by seawater carbonate chemistry not solely seawater pH, *Proceedings of the Royal Society of London B: Biological Sciences*, 284, 2017.
- DeCarlo, T. M., Gaetani, G. A., Holcomb, M., and Cohen, A. L.: Experimental determination of factors controlling U/Ca of aragonite precipitated from seawater: implications for interpreting coral skeleton, *Geochimica et cosmochimica acta*, 162, 151–165, <https://doi.org/doi:10.1016/j.gca.2015.04.016>, 2015.
- DeCarlo, T. M., Gaetani, G. A., Cohen, A. L., Foster, G. L., Alpert, A. E., and Stewart, J.: Coral Sr-U Thermometry, *Paleoceanography*, 31, 626–638, <https://doi.org/10.1002/2015PA002908>, 2016.
- DeCarlo, T. M., D’Olivo, J. P., Foster, T., Holcomb, M., Becker, T., and McCulloch, M. T.: Coral calcifying fluid aragonite saturation states derived from Raman spectroscopy, *Biogeosciences*, 14, 5253–5269, <https://doi.org/10.5194/bg-14-5253-2017>, 2017.
- DePaolo, D. J.: Surface kinetic model for isotopic and trace element fractionation during precipitation of calcite from aqueous solutions, *Geochimica et Cosmochimica Acta*, 75, 1039–1056, <https://doi.org/10.1016/J.GCA.2010.11.020>, 2011.
- Dickson, A. G.: Standard potential of the reaction: $\text{AgCl (s)} + \frac{1}{2}\text{H}_2 \text{(g)} = \text{Ag (s)} + \text{HCl (aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K, *The Journal of Chemical Thermodynamics*, 22, 113–127, 1990.

- Dickson, A. G. and Millero, F. J.: A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media, *Deep Sea Research Part A. Oceanographic Research Papers*, 34, 1733–1743, [https://doi.org/10.1016/0198-0149\(87\)90021-5](https://doi.org/10.1016/0198-0149(87)90021-5), 1987.
- D’Olivo, J. P. and McCulloch, M. T.: Response of coral calcification and calcifying fluid composition to thermally induced bleaching stress, *Scientific Reports*, 7, 2207, 2017.
- 5 Foster, G. L., Pogge von Strandmann, P. A. E., and Rae, J. W. B.: Boron and magnesium isotopic composition of seawater, *Geochemistry, Geophysics, Geosystems*, 11, Q08015, <https://doi.org/10.1029/2010GC003201>, 2010.
- Gaetani, G. A. and Cohen, A. L.: Element partitioning during precipitation of aragonite from seawater: A framework for understanding paleoproxies, *Geochimica et cosmochimica acta*, 70, 4617–4634, <https://doi.org/10.1016/j.gca.2006.07.008>, 2006.
- Gaetani, G. A., Cohen, A. L., Wang, Z., and Crusius, J.: Rayleigh-Based, Multi-Element Coral Thermometry: a Biomineralization Approach
10 to Developing Climate Proxies, *Geochimica et cosmochimica acta*, 75, 1920–1932, <https://doi.org/10.1016/j.gca.2011.01.010>, 2011.
- Gagnon, A. C., Adkins, J. F., and Erez, J.: Seawater transport during coral biomineralization, *Earth and Planetary Science Letters*, 329, 150–161, 2012.
- Holcomb, M., Venn, A. A., Tambutté, E., Tambutté, S., Allemand, D., Trotter, J., and McCulloch, M.: Coral calcifying fluid pH dictates response to ocean acidification, *Scientific Reports*, 4, 2014.
- 15 Holcomb, M., DeCarlo, T., Gaetani, G., and McCulloch, M.: Factors affecting B/Ca ratios in synthetic aragonite, *Chemical Geology*, 437, 67–76, <https://doi.org/10.1016/j.chemgeo.2016.05.007>, 2016.
- Hönisch, B., Hemming, N. G., Grottole, A. G., Amat, A., Hanson, G. N., and Bijma, J.: Assessing scleractinian corals as recorders for paleo-pH: Empirical calibration and vital effects, *Geochimica et cosmochimica acta*, 68, 3675–3685, 2004.
- Inoue, M., Suwa, R., Suzuki, A., Sakai, K., and Kawahata, H.: Effects of seawater pH on growth and skeletal U/Ca ratios of *Acropora digitifera* coral polyps, *Geophysical Research Letters*, 38, L12809, 2011.
- 20 Klochko, K., Kaufman, A. J., Yao, W., Byrne, R. H., and Tossell, J. A.: Experimental measurement of boron isotope fractionation in seawater, *Earth and Planetary Science Letters*, 248, 276–285, <https://doi.org/10.1016/j.epsl.2006.05.034>, 2006.
- Kubota, K., Yokoyama, Y., Ishikawa, T., Suzuki, A., and Ishii, M.: Rapid decline in pH of coral calcification fluid due to incorporation of anthropogenic CO₂, *Scientific Reports*, 7, 7694, <https://doi.org/10.1038/s41598-017-07680-0>, 2017.
- 25 Lee, K., Kim, T.-W., Byrne, R., Millero, F., Feely, R., and Liu, Y.-M.: The universal ratio of boron to chlorinity for the North Pacific and North Atlantic oceans, *Geochimica et Cosmochimica Acta*, 74, 1801–1811, <https://doi.org/10.1016/J.GCA.2009.12.027>, 2010.
- Lewis, E., Wallace, D., and Allison, L. J.: Program developed for CO₂ system calculations, Tech. rep., 1998.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO₂ calculated from dissolved inorganic carbon, *Marine Chemistry*, 70, 105–119, 2000.
- 30 Mavromatis, V., Montouillout, V., Noireaux, J., Gaillardet, J., and Schott, J.: Characterization of boron incorporation and speciation in calcite and aragonite from co-precipitation experiments under controlled pH, temperature and precipitation rate, *Geochimica et Cosmochimica Acta*, 150, 299–313, <https://doi.org/10.1016/j.gca.2014.10.024>, 2015.
- McConnaughey, T.: ¹³C and ¹⁸O isotopic disequilibrium in biological carbonates: I. Patterns, *Geochimica et Cosmochimica Acta*, 53, 151–162, [https://doi.org/10.1016/0016-7037\(89\)90282-2](https://doi.org/10.1016/0016-7037(89)90282-2), 1989.
- 35 McCulloch, M., Trotter, J., Montagna, P., Falter, J., Dunbar, R., Freiwald, A., Försterra, G., López Correa, M., Maier, C., and Rüggeberg, A.: Resilience of cold-water scleractinian corals to ocean acidification: Boron isotopic systematics of pH and saturation state up-regulation, *Geochimica et cosmochimica acta*, 87, 21–34, 2012a.

- McCulloch, M. T., Falter, J., Trotter, J., and Montagna, P.: Coral resilience to ocean acidification and global warming through pH up-regulation, *Nature Climate Change*, 2, 623–627, 2012b.
- McCulloch, M. T., D’Olivo Cordero, J. P., Falter, J., Holcomb, M., and Trotter, J. A.: Coral calcification in a changing World: the interactive dynamics of pH and DIC up-regulation, *Nature Communications*, 8, 15 686, 2017.
- 5 Noireaux, J., Mavromatis, V., Gaillardet, J., Schott, J., Montouillout, V., Louvat, P., Rollion-Bard, C., and Neuville, D.: Crystallographic control on the boron isotope paleo-pH proxy, *Earth and Planetary Science Letters*, 430, 398–407, <https://doi.org/10.1016/j.epsl.2015.07.063>, 2015.
- Rae, J. W., Foster, G. L., Schmidt, D. N., and Elliott, T.: Boron isotopes and B/Ca in benthic foraminifera: Proxies for the deep ocean carbonate system, *Earth and Planetary Science Letters*, 302, 403–413, <https://doi.org/10.1016/J.EPSL.2010.12.034>, <https://www.sciencedirect.com/science/article/pii/S0012821X1000796X>, 2011.
- 10 Ries, J. B.: A physicochemical framework for interpreting the biological calcification response to CO₂-induced ocean acidification, *Geochimica et cosmochimica acta*, 75, 4053–4064, 2011.
- Riley, J. P. and Tongudai, M.: The major cation/chlorinity ratios in sea water, *Chemical Geology*, 2, 263–269, 1967.
- Rollion-Bard, C., Blamart, D., Cuif, J. P., and Dauphin, Y.: In situ measurements of oxygen isotopic composition in deep-sea coral, *Lophelia*
15 *pertusa*: Re-examination of the current geochemical models of biomineralization, *Geochimica et cosmochimica acta*, 74, 1338–1349, 2010.
- Rollion-Bard, C., Blamart, D., Trebosc, J., Tricot, G., Mussi, A., and Cuif, J. P.: Boron isotopes as pH proxy: A new look at boron speciation in deep-sea corals using 11B MAS NMR and EELS, *Geochimica et cosmochimica acta*, 75, 1003–1012, 2011.
- Ross, C. L., Falter, J. L., and McCulloch, M. T.: Active modulation of the calcifying fluid carbonate chemistry ($\delta^{11}\text{B}$, B/Ca) and seasonally
20 invariant coral calcification at sub-tropical limits, *Scientific Reports*, 7, 13 830, <https://doi.org/10.1038/s41598-017-14066-9>, <http://www.nature.com/articles/s41598-017-14066-9>, 2017.
- Ruiz-Agudo, E., Putnis, C., Kowacz, M., Ortega-Huertas, M., and Putnis, A.: Boron incorporation into calcite during growth: Implications for the use of boron in carbonates as a pH proxy, *Earth and Planetary Science Letters*, 345-348, 9–17, <https://doi.org/10.1016/j.epsl.2012.06.032>, 2012.
- 25 Schoepf, V., Levas, S. J., Rodrigues, L. J., McBride, M. O., Aschaffenburg, M. D., Matsui, Y., Warner, M. E., Hughes, A. D., and Grottoli, A. G.: Kinetic and metabolic isotope effects in coral skeletal carbon isotopes: A re-evaluation using experimental coral bleaching as a case study, *Geochimica et Cosmochimica Acta*, 146, 164–178, <https://doi.org/10.1016/j.gca.2014.09.033>, 2014.
- Schoepf, V., Jury, C. P., Toonen, R. J., and McCulloch, M. T.: Coral calcification mechanisms facilitate adaptive responses to ocean acidification., *Proceedings. Biological sciences*, 284, 20172 117, <https://doi.org/10.1098/rspb.2017.2117>, 2017.
- 30 Sinclair, D. J.: Correlated trace element ‘vital effects’ in tropical corals: a new geochemical tool for probing biomineralization, *Geochimica et cosmochimica acta*, 69, 3265–3284, 2005.
- Stewart, J. A., Anagnostou, E., and Foster, G. L.: An improved boron isotope pH proxy calibration for the deep-sea coral *Desmophyllum dianthus* through sub-sampling of fibrous aragonite, *Chemical Geology*, 447, 148–160, <https://doi.org/10.1016/j.chemgeo.2016.10.029>, 2016.
- 35 Tambutté, E., Tambutté, S., Segonds, N., Zoccola, D., Venn, A., Erez, J., and Allemand, D.: Calcein labelling and electrophysiology: insights on coral tissue permeability and calcification, *Proceedings of the Royal Society B: Biological Sciences*, 279, 19–27, <https://doi.org/10.1098/rspb.2011.0733>, 2012.

- Trotter, J., Montagna, P., McCulloch, M., Silenzi, S., Reynaud, S., Mortimer, G., Martin, S., Ferrier-Pagès, C., Gattuso, J. P., and Rodolfo-Metalpa, R.: Quantifying the pH 'vital effect' in the temperate zooxanthellate coral *Cladocora caespitosa*: Validation of the boron seawater pH proxy, *Earth and Planetary Science Letters*, 303, 163–173, 2011.
- Uchikawa, J., Penman, D. E., Zachos, J. C., and Zeebe, R. E.: Experimental evidence for kinetic effects on B/Ca in synthetic calcite: Implications for potential B(OH)₄⁻ and B(OH)₃ incorporation, *Geochimica et Cosmochimica Acta*, 150, 171–191, <https://doi.org/10.1016/j.gca.2014.11.022>, 2015.
- Uchikawa, J., Harper, D. T., Penman, D. E., Zachos, J. C., and Zeebe, R. E.: Influence of solution chemistry on the boron content in inorganic calcite grown in artificial seawater, *Geochimica et Cosmochimica Acta*, 218, 291–307, <https://doi.org/10.1016/j.gca.2017.09.016>, 2017.
- Uppstrom, L.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean, *Deep Sea Research and Oceanographic Abstracts*, 1974.
- van der Weijden, C. and van der Weijden, R.: Calcite growth: Rate dependence on saturation, on ratios of dissolved calcium and (bi)carbonate and on their complexes, *Journal of Crystal Growth*, 394, 137–144, <https://doi.org/10.1016/J.JCRYSGRO.2014.02.042>, 2014.
- Venn, A., Tambutte, E., Holcomb, M., Allemand, D., and Tambutte, S.: Live tissue imaging shows reef corals elevate pH under their calcifying tissue relative to seawater, *PLoS One*, 6, e20013, <https://doi.org/10.1371/journal.pone.0020013>, 2011.
- Venn, A. A., Tambutté, E., Holcomb, M., Laurent, J., Allemand, D., and Tambutté, S.: Impact of seawater acidification on pH at the tissue-skeleton interface and calcification in reef corals., *Proceedings of the National Academy of Sciences of the United States of America*, 110, 1634–9, <https://doi.org/10.1073/pnas.1216153110>, 2013.
- Watson, E. B.: A conceptual model for near-surface kinetic controls on the trace-element and stable isotope composition of abiogenic calcite crystals, *Geochimica et cosmochimica acta*, 68, 1473–1488, 2004.
- Wu, H. C., Dissard, D., Le Cornec, F., Thil, F., Tribollet, A., Moya, A., and Douville, E.: Primary Life Stage Boron Isotope and Trace Elements Incorporation in Aposymbiotic *Acropora millepora* Coral under Ocean Acidification and Warming, *Frontiers in Marine Science*, 4, 129, <https://doi.org/10.3389/fmars.2017.00129>, 2017.
- Yu, J., Foster, G. L., Elderfield, H., Broecker, W. S., and Clark, E.: An evaluation of benthic foraminiferal B/Ca and $\delta^{11}\text{B}$ for deep ocean carbonate ion and pH reconstructions, *Earth and Planetary Science Letters*, 293, 114–120, <https://doi.org/10.1016/J.EPSL.2010.02.029>, <https://www.sciencedirect.com/science/article/pii/S0012821X10001317>, 2010.
- Zeebe, R. E. and Wolf-Gladrow, D. A.: *CO₂ in seawater: equilibrium, kinetics, isotopes*, vol. 65, Elsevier Science Limited, 2001.