

Validation of carbon isotope fractionation in algal lipids as a PCO_2 proxy using a natural CO_2 seep (Shikine Island, Japan)

Caitlyn R Witkowski^{1*}, Sylvain Agostini², Ben P Harvey², Marcel TJ van der Meer¹, Jaap S Sinninghe Damsté^{1,3}, Stefan Schouten^{1,3}

5 ¹Department of Marine Microbiology and Biogeochemistry, Royal Netherlands Institute for Sea Research, Den Burg (Texel), 1790AB, and Utrecht University, The Netherlands

²Shimoda Marine Research Center, University of Tsukuba, Shimoda, 415-0025, Japan

³Department of Geosciences, Utrecht University, Utrecht, 3508 TA, The Netherlands

*Present address: School of Earth Sciences, University of Bristol, Bristol, UK

10 Correspondence to: Caitlyn R Witkowski (caitlyn.witkowski@bristol.ac.uk)

Abstract. Carbon dioxide concentrations in the atmosphere play an integral role in many earth system dynamics, including its influence on global temperature. The past can provide insights into these dynamics, but unfortunately reconstructing long-term trends of atmospheric carbon dioxide (expressed in partial pressure; PCO_2) remains a challenge in paleoclimatology. One promising approach for reconstructing past PCO_2 utilizes the isotopic fractionation associated with CO_2 -fixation during photosynthesis into organic matter (ε_p). Previous studies have focused primarily on testing estimates of ε_p derived from the $\delta^{13}C$ of species-specific alkenone compounds in laboratory cultures and mesocosm experiments. Here, we analyze ε_p derived from the $\delta^{13}C$ of more general algal biomarkers, i.e. compounds derived from a multitude of species, from sites near a CO_2 seep off the coast of Shikine Island (Japan), a natural environment with CO_2 concentrations ranging from ambient (ca. 310 μ atm) to elevated (ca. 770 μ atm) PCO_2 . We observed strong, consistent $\delta^{13}C$ shifts in several algal biomarkers from a variety of sample matrices over the steep CO_2 gradient. Of the three general algal biomarkers explored here, namely loliolide, phytol, and cholesterol, ε_p positively correlates with PCO_2 , in agreement with ε_p theory and previous culture studies. PCO_2 reconstructed from the ε_p of general algal biomarkers show the same trends throughout, as well as the correct control values, but with lower absolute reconstructed values than the measured values at the elevated PCO_2 sites. Our results show that naturally occurring CO_2 seeps may provide useful testing grounds for PCO_2 proxies and that general algal biomarkers show promise for reconstructing past PCO_2 .

1 Introduction

The current increase in the atmospheric concentration of carbon dioxide (expressed in partial pressure, PCO_2) plays a leading role in climate change (Forster et al., 2007). PCO_2 is significantly higher now than it has been in the past 800 ka (Lüthi et al., 2008) and although long-term changes in PCO_2 are not uncommon over millions of years (Foster et al., 2017), this current spike in PCO_2 has occurred within only the past two centuries (IPCC, 2013). Uncertainties remain on the exact magnitude to which PCO_2 influences climate, as well as the exact response of the environment to these climate changes. Long-term PCO_2

trends help us better understand the context for these changes and are reconstructed via indirect means, i.e. environmental proxies. Two proxies can span timescales over 100 Ma (Foster et al., 2017), e.g. the terrestrial proxies paleosols and leaf stomata. The paleosol proxy has large uncertainties due to the difficulties in constraining soil respiration (Breecker et al., 2010; Cotton and Sheldon, 2012) due to carbon isotopic fractionation during microbial decomposition (Bowen and Beerling, 2004), carbonate diagenesis (Quast et al., 2006), and other local and regional influences on carbon cycles in these terrestrial settings. Although the leaf stomata proxy is often better constrained than paleosols, some experiments do not show the expected trends (Ellsworth et al., 2011; Ward et al., 2013; DaMatta et al., 2016), suggesting that factors other than PCO_2 , e.g. ecological systems, species, and development stage, also impact the leaf stomata proxy (Xu et al., 2016). The development of new proxies for PCO_2 may help us better constrain past long-term trends, particularly marine-based proxies which tend to have more homogenized signals but are currently relatively limited in time.

A proxy that has been explored with mixed success over the past several decades is the stable carbon isotopic fractionation associated with photosynthetic inorganic carbon fixations (ε_p), which has been shown to positively correlate with PCO_2 (Bidigare et al., 1997; Jasper and Hayes, 1990; Zhang et al., 2013). ε_p occurs as the CO_2 -fixing enzyme in photoautotrophs, Rubisco (ribulose 1,5-biphosphate carboxylase oxygenase), favors ^{12}C which consequently results in photosynthates isotopically more depleted in ^{13}C than the original carbon source. A greater abundance of CO_2 increases Rubisco-based isotopic discrimination, resulting in an even lower $^{13}C/^{12}C$ ratio ($\delta^{13}C$) in photoautotroph biomass (Farquhar et al., 1989; Farquhar et al., 1982; Francois et al., 1993; Popp et al., 1989). When this phototrophic biomass is preserved in the geologic record, the $\delta^{13}C$ of sedimentary organic matter can be used to reconstruct PCO_2 (Hayes et al., 1999). The largely mixed contributions and diagenetic processes on bulk organic matter can, however, mask this signal (Hayes, 1993; Hayes et al., 1999). Thus, isotope analysis of specific biomarker lipids is preferred in order to better define the source of the $\delta^{13}C$ signal (Jasper and Hayes, 1990; Pagani, 2002).

The most studied biomarkers for calculating ε_p are alkenones, i.e. long-chain unsaturated methyl and ethyl ketones produced by select Haptophytes (Volkman et al., 1980; de Leeuw et al., 1980). The stable carbon isotopic fractionation of alkenones has been studied using cultures and mesocosms with controlled environments (Laws et al., 1995; Benthien et al., 2007), but conditions do not always mimic natural environments and the natural variation in carbonate chemistry that occurs on a daily to seasonal time scales. Furthermore, these experiments are also time-consuming given that they must have delicately balanced water chemistry including $CO_{2[aq]}$ concentrations, pH, and alkalinity, as well as nutrients such as nitrate and phosphate (Popp et al., 1998; Laws et al., 1995; Bidigare et al., 1997), along with the additional challenge of maintaining a constant $\delta^{13}C$ of the $CO_{2[aq]}$ while photoautotrophs continually enrich the growth water as they fix CO_2 . Water column studies (Bidigare et al., 1997) and surface sediments (Pagani, 2002) have been applied but rarely reach elevated PCO_2 levels such as those encountered in the past.

Here we use an alternative approach by analyzing algal lipids near natural CO_2 seep systems. In tectonically active zones, volcanically induced seeps consistently bubble high PCO_2 concentrations into the surrounding water, substantially increasing the local CO_2 concentrations in the water and providing an environment referent to past and future high- CO_2 worlds. CO_2

seeps were previously overlooked for biological studies due to the typically high sulfide (H_2S) concentrations associated with volcanic degassing that make these environments largely uninhabitable (Dando et al., 1999). However, certain CO_2 seep systems have been found to have low H_2S concentrations making them suitable for ocean acidification experiments (Hall-Spencer et al., 2008), prompting further research in e.g. the Mediterranean (Boatta et al., 2013), in Japan (Agostini et al., 2015),
5 Papua-New-Guinea (Fabricius et al., 2011), and New Zealand (Brinkman and Smith, 2015). These sites may provide an ideal testing ground for the impact of isotopic fractionation on algal lipids where environmental conditions are at naturally balanced levels with the exception of the large gradient of CO_2 concentrations.

In our study, we explore the relationship between ε_p and $\text{CO}_{2[\text{aq}]}$ across several pre-established sites with different (temporally consistent) levels of PCO_2 at the warm-temperate CO_2 seep at Mikama Bay off the shore of Shikine Island, Japan. We test this
10 relationship using general algal biomarkers, i.e. compounds derived from a multitude of species and which have rarely been used for ε_p -based PCO_2 reconstructions despite their potential utility (Witkowski et al., 2018; Popp et al., 1989; Freeman and Hayes, 1992).

2 Materials and Methods

2.1 Sample site

15 The site is briefly described here. For further details, we refer to Agostini et al. (2018). Mikama Bay is located on the south-southwest corner of Shikine Island off the Izu Peninsula, Japan bay (34.320865 N, 139.204868 E) with several venting locations in the north of the bay (Fig. 1). The gas emitted from the seep contains 98% CO_2 and the bay has a spatially and temporally constant total alkalinity averaging at $2265 \pm 10 \mu\text{mol kg}^{-1}$. Samples were collected from three preestablished PCO_2 sites (Agostini et al., 2015), “Control PCO_2 ” site in an adjacent bay outside the influence of the CO_2 seep ($\text{PCO}_2 309 \pm 46 \mu\text{atm}$),
20 a “Mid PCO_2 ” site ($\text{PCO}_2 \text{ ca. } 460 \pm 40 \mu\text{atm}$), and a “High PCO_2 ” site ($\text{PCO}_2 769 \pm 225$) (Fig. 1). PCO_2 estimates are based on the carbonate chemistry parameters (pH_{NBS} , temperature, salinity, and total alkalinity) of water in the bay and calculated using the program CO2sys (Agostini et al., 2018; Harvey et al., 2018). Temperature (annual range ca. 14 to 28°C) and salinity (ca. 34‰) are relatively homogenous throughout the bay and do not differ between the elevated PCO_2 sites and control PCO_2 sites (Agostini et al. 2018). Currents and wind were measured in October 2014 and April 2015 (Agostini et al., 2015). October
25 2014 measurements showed moderate turbulent winds (ranging from 0.6-11.5 m s^{-1} , average 4.5 m s^{-1}) associated with current velocities (ranging from 0 to 1.6 knots, average 0.4 knots) at 5 m in the surface water, whereas April 2015 measurements showed moderate north-northeast winds (1.5-8.6 m s^{-1} , average 5.1 m s^{-1}) associated with low current velocities (0-0.2 knots, average 0.04 knots). Monthly surveys in the bays over the past five years show that these sites have similar annual mean values for temperature, salinity, and currents. Weather station data shows that the severity of seasonal extreme weather event (e.g.
30 typhoons) varies on an annual basis (Japan Meteorological Agency, <https://www.jma.go.jp/en/typh/>).

2.2 Materials

Samples were collected in June and September of 2016 (indicated in Fig. 1). All samples were collected in at least triplicate for each site (“Control PCO₂”, “Mid PCO₂” and “High PCO₂” site). Additional control sites (at ca. 1.8 km and 2.4 km away from the CO₂ seep) around the island were taken to ensure the fidelity of the Control site closest to the seep. June sampling included surface waters for dissolved inorganic carbon (DIC) measurements, surface sediments, and benthic diatoms attached to surface sediment through extracellular polymeric substance production. In September, macroalgae and plankton tows were collected, in addition to surfaced water DIC and surface sediments, taken in triplicate at each site on four separate days.

Water for the δ¹³C of DIC analysis was collected by overfilling glass vials with sea surface water and adding mercury chloride (0.5%) before closing with a septa cap and sealing with electrical tape. Surface sediments were collected by divers using geochemical sample bags. Macroalgae and benthic diatoms were scraped off submerged rocks at each respective site. A 25 μm mesh plankton net (“small plankton net”, Rigo, Saitama, Japan) was towed 50 m three times per site and filtered using a portable hand aspirator on the boat over a single 0.7 μm GFF (combusted prior to sampling for 4 h at 450°C).. All samples were immediately frozen; once back in the lab, these were freeze-dried and kept in a refrigerator.

2.3 Methods

Each seawater sample was measured for the δ¹³C of DIC in duplicate on a Thermo Scientific Gas Bench II coupled to a Thermo Scientific Delta V mass spectrometer. Prior to running samples, 12 ml vials were prepared with 100 μL of 85% H₃PO₄ and flushed with He. 500 μL of seawater was added and left to react for at least 1 h prior to measuring the headspace. Standards were run at the start, end, and every six runs of a sequence. Standards were prepared with 0.3 mg of Na₂CO₃ and 0.4 mg of Ca₂CO₃ (all calibrated against NBS-19) flushed with He, injected with 500 μL of 85% H₃PO₄, and reacted for 1 h. The headspace was measured and average values and standard deviation errors reported are based on six measurements for June (three at the High PCO₂ site and three at the Control) and thirty-six measurements for September (three each at the High PCO₂ site, Mid PCO₂ site, and Control collected on four separate days).

Freeze-dried sediments, benthic diatoms, and macroalgae were homogenized using mortar and pestle and extracted using a Dionex 250 accelerated solvent extractor at 100°C, 7.6x10⁶ Pa using dichloromethane (DCM): MeOH (9:1 v/v). GFFs containing plankton net material were cut into 1 mm x 1 mm squares and extracted using ultrasonication (5x) with 2 ml dichloromethane (DCM): MeOH (9:1 v/v). All total lipid extracts (TLEs) were hydrolyzed by refluxing the TLE with 1N of KOH in MeOH for one hour and neutralized to pH 5 using 2 N of HCl in MeOH. Bi-distilled water (2 ml) and DCM (2 ml) were added (5x) to the hydrolyzed centrifuge tubes and the organic matter in the DCM layers were pooled and dried over Na₂SO₄. The resulting hydrolyzed TLEs were eluted over an alumina packed column and separated into apolar (hexane: DCM, 9:1 v/v), ketone (DCM), and polar (DCM: MeOH, 1:1 v/v) fractions. Polar fractions were silylated using pyridine: N,O-Bis(trimethylsilyl)trifluoroacetamide (BSTFA) (1:1 v/v) and heated at 60°C for 1 h prior to analyses on the gas

chromatography-flame ionization detector (GC-FID), gas chromatography-mass spectrometry (GC-MS), and gas chromatography isotope-ratio mass spectrometry (GC-IRMS).

Silylated polar fractions were analyzed by GC-FID for quantification. Based on the quantities, fractions were diluted with ethyl acetate and ca. 1 ug of polar fraction was injected on-column for GC-MS to identify compounds and for GC-IRMS to measure the isotopic composition of specific compounds. Each instrument is equipped with the same CP-Sil 5 column (25 m x 0.32 mm; df 0.12 μ m) and He is used as carrier gas. GC oven was programmed from 70°C to 130°C at 20°C/min and then to 320°C at 4°C/min which was held for 10 min. All three instruments use the same in-house mixture of *n*-alkanes and fatty acids to check chromatography performance at the start of each day (GC-standard). For compound specific stable carbon isotope analysis using GC-IRMS, additional standards with known isotopic values (-32.7 and -27.0‰) of per deuterated (99.1%) *n*-alkanes (C_{20} and C_{24} , respectively), were co-injected with the GC-standard. Samples were also co-injected with the same GC-IRMS standards to monitor instrument performance. Every day, the Isolink II combustion reactor of the GC-IRMS was oxidized for at least 10 min, backflushed with He for 10 min, and purged for 5 min; a shorter version of this sequence is conducted in post-sample seed oxidation which includes 2 min oxidation, 2 min He backflush, and 2 min purge conditioning line. Longer oxidations were run weekly. Each derivatized compound was corrected for the $\delta^{13}\text{C}$ of the BSTFA used in silylation (-32.2‰).

3 Results

Samples from the different matrices were collected at several Control PCO_2 sites (309 ± 46 , at a “Mid PCO_2 ” site (ca. 100 m from the venting area; $460 \pm 40 \mu\text{atm}$), and near the venting area (“High PCO_2 ” site; $769 \pm 225 \mu\text{atm}$) during June and September 2016 (Fig. 1), which included June-collected surface waters (for DIC), surface sediments, and benthic diatoms, and September-collected surface waters (for DIC), surface sediments, plankton net tows, and macroalgae.

The $\delta^{13}\text{C}$ of DIC demonstrated minimal change over the gradient of CO_2 and minimal change between the two seasons (Fig. S1). The June $\delta^{13}\text{C}$ of DIC was $0.2 \pm 0.2 \text{‰}$ ($\pm \text{SD}$; N=3) at the Control site and $0.5 \pm 0.04 \text{‰}$ (N=3) at the High PCO_2 site. The September $\delta^{13}\text{C}$ of DIC was $-0.4 \pm 0.2 \text{‰}$ (N=8) at the Control site, $-0.1 \pm 0.1 \text{‰}$ (N=8) at the Mid PCO_2 site, and $0.2 \pm 0.4 \text{‰}$ (N=8) at the High PCO_2 site.

The polar fractions of the extracts of the surface sediments, plankton, macroalgae, and benthic diatoms showed a similar suite of compounds, observed across all sites and during both seasons. The most prominent compounds were loliolide, phytol, C_{14} - C_{16} alkanols, and sterols such as cholesta-5,22E-dien-3 β -ol, cholesterol, 23-methylcholesta-5,22dienol, campesterol, stigmasterol, and β -sitosterol (e.g. Fig. 2). Terrestrial biomarkers, such as long chain alcohols and triterpenoids were not detected. Loliolide, phytol, and cholesterol were targeted for stable carbon isotope analysis as the most abundant general algal biomarkers and with relatively good separation in the GC. The biological sources of these compounds will be discussed in Section 4.1.

Among the sample matrices, the $\delta^{13}\text{C}$ of loliolide ranges from -19.8 to -22.0 ‰ at the Control sites, from -20.5 to -22.9 ‰ at the Mid PCO_2 site, and from -23.1 to -29.0 ‰ at the High PCO_2 site (Fig. 3A; Table S1). The $\delta^{13}\text{C}$ of loliolide from June surface sediments shows the strongest change from the Control site to the High PCO_2 site (-21.2 to -29.0 ‰), followed by the $\delta^{13}\text{C}$ of loliolide from September macroalgae (-21.3 to -25.7 ‰). A lesser $\delta^{13}\text{C}$ shift is observed in the September surface sediment-derived loliolide (-19.8 to -23.1 ‰). The $\delta^{13}\text{C}$ of the benthic diatom-derived loliolide (-20.2 to 23.6 ‰) and the plankton tow-derived loliolide show the smallest shifts from the Control to High PCO_2 site (-22.0 to -23.6 ‰).

Similar to the results of the $\delta^{13}\text{C}$ of loliolide, the $\delta^{13}\text{C}$ of phytol also consistently shows higher $\delta^{13}\text{C}$ values in the Control sites and lower $\delta^{13}\text{C}$ values in the elevated PCO_2 sites among all samples types collected in both seasons (Fig. 3B; Table S1). For the whole sample set, the $\delta^{13}\text{C}$ of phytol ranges from -18.9 to -22.6 ‰ at the Control site, from -19.4 to -22.4 ‰ at the Mid PCO_2 site, and from -22.6 to -27.8 ‰ at the High PCO_2 site (Fig 3B), similar ranges as observed for loliolide. A similar shift in $\delta^{13}\text{C}$ values of phytol is observed with increasing PCO_2 in the June surface sediments (-22.6 to -27.8 ‰), the June benthic diatoms (-18.9 to -24.4 ‰), and the September macroalgae (-21.5 to -26.9 ‰). Smaller changes in the $\delta^{13}\text{C}$ of phytol are observed for September plankton (-21.7 to -24.4 ‰) and September sediment (-20.5 to -22.6 ‰).

The $\delta^{13}\text{C}$ of cholesterol likewise shows a similar trend to the other two biomarkers but with a smaller shift in the $\delta^{13}\text{C}$ values from the Control PCO_2 sites to the elevated PCO_2 sites. Among the different sample matrices, the $\delta^{13}\text{C}$ of cholesterol ranges from -21.2 ‰ to -25.1 ‰ at the Control site, -22.1 to -23.4 ‰ at the Mid PCO_2 site and -23.1 to -27.4 ‰ at the High PCO_2 site (Fig. 3C; Table S1). The strongest change in the $\delta^{13}\text{C}$ of cholesterol with increase PCO_2 occurs in the June surface sediments from -22.6 ‰ in the Control to -27.8 ‰ at the High PCO_2 site. The June benthic diatoms also have a large isotopic shift in the $\delta^{13}\text{C}$ of cholesterol (-21.2 to -25.8 ‰), as does the September macroalgae (-22.7 to -25.8 ‰). The September surface sediments (-22.2 to -23.1 ‰) and plankton tow-derived cholesterol (-25.1 to -26.7 ‰), however, have a smaller shift from the control to the elevated PCO_2 sites.

4 Discussion

4.1 The $\delta^{13}\text{C}$ differences in biomarkers among matrices and seasons

All three biomarkers, phytol, loliolide and cholesterol, show a negative shift in $\delta^{13}\text{C}$ values with increasing PCO_2 in each matrix and each season (Fig. 3), agreeing with the theory that higher PCO_2 conditions result in lower $\delta^{13}\text{C}$ values in biomass (Farquhar et al., 1982). However, despite all having algal sources, the absolute isotope values vary for 1) each compound, 2) each matrix, and 3) both seasons, which we will now discuss.

First, the absolute values of $\delta^{13}\text{C}$ values vary among the three compounds. This may be expected given the different biosynthetic pathways leading to formation of each compound (Schouten et al., 1998), as well as the different contributors to each compound. Loliolide, considered a diatom biomarker in paleoreconstructions (e.g. Castañeda et al., 2009), is a diagenetic product of fucoxanthin (Repeta, 1989; Klok et al., 1984), a xanthophyll which contributes to approximately 10% of all

carotenoids found in nature (Liaaen-Jensen, 1978). Phytol, considered a photoautotroph biomarker in paleoreconstructions (Hayes et al., 1990), is the side-chain of the vital and omnipresent pigment chlorophyll *a* that directly transfers sunlight energy into the photosynthetic pathway in nearly all photosynthetic organisms. Sterols, considered a general eukaryotic biomarker in paleoreconstructions, are the eukaryotic tetracyclic triterpenoid lipids used for critical regulatory roles of cellular functions 5 e.g. maintaining membrane fluidity (Nes et al., 1993). Although sterols are virtually restricted to eukaryotes, some exceptions have been found in bacteria (Wei et al., 2016). Here we only examine cholesterol, which is universally absent in prokaryotes and composes of up to 20-40% of eukaryotic plasma membranes (Mouritsen and Zuckermann, 2004). Phytol and cholesterol may also have terrestrial sources given that they are derived from all photoautotrophs and all eukaryotes, respectively. However, these samples were taken off the coast of a small island in open ocean and the absence of characteristic terrestrial 10 biomarkers indicates that terrestrial contributions can be considered to be minimal. The close resemblance of the isotopic composition among all three compounds, including the primarily diatom-limited compound loliolide, suggests that these compounds share relatively similar source organisms. Cholesterol shows a lessened isotopic shift than the other two compounds from the ambient to elevated PCO_2 sites. Although we cannot fully exclude that this is due to terrestrial input, it is more likely due to the mobile eukaryotic zooplankton in the water column which also contribute to the cholesterol signal.

15 Within the same biomarker and same season, some differences among matrices were observed. This difference may be due to the mobility of the matrix, as well as the algal assemblages. The plankton tow which captured free-floating surface water algae from that specific growth season is more readily transported by wind than the surface sediment, which likely reflects the culmination of multiple growth seasons throughout the water column. This is seen, for example, in the $\delta^{13}C$ of cholesterol collected in September from the same Control site where surface sediments are -22.2 ‰ and plankton tows are -25.1 ‰, where 20 the latter has possibly been transported from sites with elevated PCO_2 levels. Similar differences among matrices are also observed in phytol and loliolide. The hypothesis of transportation affecting the isotopic signal in certain matrices is supported by the results from the macroalgae. The macroalgae, in contrast to the algae collected by plankton tows, were unaffected by transportation due to being fixed to the nearby rocks at each site. Thus, the isotopic composition of compounds of the macroalgae was similar to that of the long-accumulated surface sediments, e.g. -22.7 ‰ for the $\delta^{13}C$ of cholesterol at the 25 September Control site.

Finally, there is a difference in the $\delta^{13}C$ values for biomarkers between seasons. The June-collected surface sediments and algae yielded a larger difference in $\delta^{13}C$ values along the CO_2 gradient than the September-collected surface sediments and algae. This seasonal difference may be due to extreme weather conditions experienced between the two sampling campaigns. Although typhoons are common in this region, in the weeks preceding the fieldwork in September, Shikine Island experienced 30 an unusually high quantity of storms. The storms were also of unusual strength for this region of the Pacific, including Typhoons Mindulle and Kompasu, the severe tropic storms Omais and Chanthu, and the long-lived, erratic Lionrock typhoon. This atypical abundance and severity of storms observably ripped corals out of the rocks around Shikine Island and thus likely resuspended and transported some sediment around the bay. This would explain the reduced $\delta^{13}C$ difference between the Control and High PCO_2 site in the surface sediments collected in September, as well as the readily transportable algae collected

by the plankton tow, and would explain why the rock-affixed macroalgae, also collected in September, maintained a strong $\delta^{13}\text{C}$ change across the transect.

4.2 The ε_p among general algal biomarkers

To further validate the impact of PCO_2 , we calculated the isotopic fractionation of algal biomass based on the $\delta^{13}\text{C}$ of the three biomarkers. Here we focus on surface sediments as they are a close analogue to the geological sediment records. Although the macroalgae and benthic diatoms also show strong isotopic fractionation, they represent a limited number of species and a single growth season. Furthermore, we calculated the ε_p from the June-collected surface sediments, which appear to be the least affected by typhoon activity and represent fractionation over multiple seasons.

To calculate ε_p in the June-collected surface sediments, we correct the $\delta^{13}\text{C}$ of the organic matter (δ_p) for the $\delta^{13}\text{C}$ of the inorganic carbon source for the producers of these compounds (δ_d) in Eq. (1):

$$\varepsilon_p = 1000 \cdot [(\delta_d + 1000) / (\delta_p + 1000) - 1], \quad (1)$$

δ_p is calculated by correcting the $\delta^{13}\text{C}$ for each individual biomarker for the offset with photosynthetic biomass caused by isotopic fractionation during biosynthesis. The isotopic offset between phytol and biomass is $3.5 \pm 1.3\text{‰}$ based on the average of twenty-three species compiled in Witkowski et al. (2018) and the isotopic offset between sterols and biomass is $4.5 \pm 3.0\text{‰}$ based on the average of eight algal species (Schouten et al., 1998). The isotopic offset for loliolide from biomass, however, has not been determined. Because isoprenoids are formed from the same biosynthetic pathway, we here average the offset of the other two isoprenoids here (4.0‰) to estimate a value for the difference between loliolide and biomass.

δ_d is calculated by correcting the measured $\delta^{13}\text{C}$ of DIC for temperature (Mook, 1974) and pH (Madigan et al., 1989), which considers the relative contribution of different inorganic carbon species to the measured DIC. Based on the equations of Mook et al. (1974), we correct for the temperature-dependent carbon isotopic fractionation of dissolved CO_2 with respect to HCO_3^- using the annual mean sea surface temperature for Shikine Island of 20.4°C (Agostini et al., 2018). Based on the equations of Madigan et al. (1989), we corrected for the $\delta^{13}\text{C}$ of HCO_3^- and $\delta^{13}\text{C}$ of $\text{CO}_{2[\text{aq}]}$ mass balance calculation that accounts for the relative abundance of these inorganic carbon species based on pH (Lewis and Wallace, 1998) at the High PCO_2 site (7.81 pH_T) and Mid PCO_2 site (7.99 pH_T) relative to the ambient Control (8.14 pH_T). The corrected δ_d values yield -10.1‰ at the Control site, -10.0‰ at the Mid PCO_2 site, and -9.5‰ at the High PCO_2 site (Table S2).

ε_p values consistently yield much higher values at the elevated PCO_2 sites than the ambient Control sites for all three biomarkers, which share similar trends and absolute values (Fig. 4; Table S3). ε_p derived from loliolide averages $7.2 \pm 1.6\text{‰}$ at the Control, $9.2 \pm 1.6\text{‰}$ at the Mid PCO_2 site, and $15.9 \pm 1.6\text{‰}$ at the High PCO_2 site, ε_p derived from phytol at $8.6 \pm 0.4\text{‰}$, $8.6 \pm 0.9\text{‰}$, and $14.9 \pm 1.0\text{‰}$, respectively, and ε_p derived from cholesterol at $7.6 \pm 3.0\text{‰}$, $9.2 \pm 3.1\text{‰}$, to $13.7 \pm 3.1\text{‰}$, respectively, where errors represent the standard deviation of the triplicate samples taken at each site. These results show that CO_2 has a profound impact on ε_p as it is the only variable with a large gradient in the bay. Given that maximum fractionation for algae species is ca. 25 to 28 ‰ in laboratory cultures (Goericke and Fry, 1994), the CO_2 seep values suggests strong, but does not approach maximum fractionation (ε_f) at the high CO_2 site. This may be due to presence of carbon concentrating

mechanism in phytoplankton which utilize ^{13}C -enriched bicarbonate or possible due to the presence of Rubisco types with different ε_f values than previously assumed (Thomas et al., 2018).

4.3 PCO_2 reconstructed from general algal biomarkers

We estimate PCO_2 from the ε_p values, a relationship first derived for higher plants (Farquhar et al., 1989; Farquhar et al., 1982)

5 and later adapted for algae (Jasper et al., 1994; Rau et al., 1996) in Eq. (2):

$$\text{PCO}_2 = [b / (\varepsilon_f - \varepsilon_p)] / K_0 , \quad (2)$$

where ε_f reflects the maximum Rubisco-based isotopic fractionation, b reflects species carbon demand per supply such as growth rate and cell-size (Jasper et al., 1994), and K_0 reflects a constant to convert $\text{CO}_{2\text{[aq]}}$ to PCO_2 based on temperature and salinity (Weiss, 1974). ε_f for algal species range from 25 to 28 ‰ in laboratory cultures (Goericke and Fry, 1994); we use an

10 average 26.5 ‰ with an uncertainty of 1.5 ‰ uniformly distributed for these general algal biomarkers (Witkowski et al., 2018).

The b value is difficult to estimate as it is a catchall for factors other than PCO_2 that affect fractionation and is particularly difficult to estimate for general algal biomarkers because they are derived from a multitude of species. Previous studies using phytol's diagenetic product phytane as a PCO_2 proxy (Bice et al., 2006; Sinninghe Damsté et al., 2008; van Bentum et al., 2012) have used a mean value of 170 ‰ $\text{kg } \mu\text{M}^{-1}$, similar to the mean of alkenone-producers. This is supported by a compilation 15 of the $\delta^{13}\text{C}$ values of modern surface sediment organic matter mean average of $168 \pm 43 \text{ ‰ kg } \mu\text{M}^{-1}$ (Witkowski et al., 2018) and a single study on phytol in the equatorial Pacific Ocean (Bidigare et al., 1997). We apply this average, rounded to $170 \pm 50 \text{ ‰ kg } \mu\text{M}^{-1}$ to all three general algal biomarkers.

The resulting reconstructed PCO_2 estimations show the expected values in the Control sites and much higher values in the elevated CO_2 sites among all three biomarkers (Fig. 5; Table S3). Loliolide shows the biggest shift, from $239 +50/-49 \mu\text{atm}$ at

20 the Control, $266 +57/-54 \mu\text{atm}$ at Mid PCO_2 site, and $437 +113/96 \mu\text{atm}$ at the High PCO_2 site. Phytol has similar but a slightly smaller shift in PCO_2 estimates to loliolide, with estimations of $264 +55/-54 \mu\text{atm}$, $291 +56/-53 \mu\text{atm}$, and $444 +98/-87 \mu\text{atm}$ at the Control, Mid PCO_2 site, and High PCO_2 site, respectively. Cholesterol shifts similarly to the other two biomarkers with $244 +67/-54 \mu\text{atm}$, $266 +77/-61 \mu\text{atm}$, and $358 +136/-90 \mu\text{atm}$, respectively. These reconstructed values closely match each other and trend in the same direction as the actual values.

25 The reconstructed PCO_2 values derived from the $\delta^{13}\text{C}$ of general algal biomarkers closely match the actual measured PCO_2 values of the Control (Fig. 5), i.e. $309 \pm 46 \mu\text{atm}$ (Agostini et al., 2018; Harvey et al., 2018), when considering the uncertainty in the reconstructed estimations. However, the proxies underestimate the absolute values measured at the elevated PCO_2 sites (Fig. 5; Table S3), i.e. $460 \pm 40 \mu\text{atm}$ at the Mid PCO_2 site and $769 \pm 225 \mu\text{atm}$ at the High PCO_2 site (Agostini et al., 2018; Harvey et al., 2018). There are several possible explanations to why there is an underestimation. As discussed above, carbonate 30 concentration mechanisms may be operating in a large number of phytoplankton, such that they become relatively enriched in ^{13}C and thus lead to lower reconstructed PCO_2 values (Stoll et al., 2019; Badger et al., 2019). There is also a large uncertainty in the b value applied, which may be much lower than the value assumed here. However, if so, then PCO_2 values reconstructed for past times may be much higher, leading to considerable discrepancies with other PCO_2 proxies (c.f. Witkowski et al.,

2018). A simple explanation for this underestimation may be some site limitations. The high variability of PCO_2 at these sites could have impacted the reconstructed values, as these algae could have been exposed to much different, and perhaps lower, levels than those observed during the times that PCO_2 values were measured. Furthermore, there is a strong possibility of allochthonous marine input of sediment at the Mid and High PCO_2 site, i.e. input from sediment outside of the bay area. This
5 allochthonous input seems likely given the intense weather conditions that occur annually in this small bay in which lateral transport of sediment could bring algal material grown in ambient PCO_2 conditions into the bay and dampen the overall PCO_2 signal picked up in the biomarkers. Future research conducted at another CO_2 seep settings with different weather and current conditions could illuminate this.

5 Conclusion

10 We analyzed the $\delta^{13}C$ of general algal biomarkers in surface sediments, plankton, benthic diatoms, and macroalgae collected in a transect from a CO_2 vent during two seasons. The strong $\delta^{13}C$ change between the Control and elevated PCO_2 sites suggest that the increased CO_2 concentrations in the seawater does indeed influence fractionation of photoautotrophic biomass and validates previous PCO_2 reconstructions which have considered utilizing general algal biomarkers for this purpose. Reconstructions correctly estimate control values, though reconstructions at the elevated PCO_2 sites show underestimations of
15 the actual PCO_2 , possibly due to the allochthonous input from nearby marine sediments deposited under normal PCO_2 levels caused by the intense annual typhoon activity in this region. Our results show that CO_2 seeps may offer testing grounds for exploring new PCO_2 proxies under natural conditions at high PCO_2 levels such as those encountered in the geological past.

Acknowledgements

We thank Jason M. Hall-Spencer, Marco Milazzo and Yasutaka Tsuchiya for their help in sample-collection. We also thank
20 Jort Ossebaar and Ronald van Bommel at the NIOZ for technical support. **Funding:** This study received funding from the Netherlands Earth System Science Center (NESSC) through a gravitation grant (024.002.001) to JSSD and SS from the Dutch Ministry for Education, Culture and Science. **Author contributions:** CRW, SS, and JSSD designed the study. SA, BPH, and CRW collected field samples. CRW analyzed samples and wrote the manuscript. CRW, MvdM, JSSD, and SS interpreted the data. **Competing interests:** The authors declare that they have no competing interests. **Data and materials availability:** All
25 data are present in the paper and/or the Supplementary Materials.

References

Agostini, S., Wada, S., Kon, K., Omori, A., Kohtsuka, H., Fujimura, H., Tsuchiya, Y., Sato, T., Shinagawa, H., Yamada, Y., and Inaba, K.: Geochemistry of two shallow CO_2 seeps in Shikine Island (Japan) and their potential for ocean acidification research, *Regional Studies in Marine Science*, 2, 45-53, 2015.

Agostini, S., Harvey, B. P., Wada, S., Kon, K., Milazzo, M., Inaba, K., and Hall-Spencer, J. M.: Ocean acidification drives community shifts towards simplified non-calcified habitats in a subtropical–temperate transition zone *Sci Rep-Uk*, 8:11354 DOI:10.1038/s41598-018-29251-7, 2018.

5 Badger, M. P. S., Chalk, T. B., Foster, G. L., Bown, P. R., Gibbs, S. J., Sexton, P. F., Schmidt, D. N., Pälike, H., Mackensen, A., and Pancost, R. D.: Insensitivity of alkenone carbon isotopes to atmospheric CO₂ at low to moderate CO₂ levels, *Clim. Past*, 15, 539-554, 10.5194/cp-15-539-2019, 2019.

Benthien, A., Zondervan, I., Engel, A., Hefta, J., Terbruggen, A., and Riebesell, U.: Carbon isotopic fractionation during a mesocosm bloom experiment dominated by *Emiliania huxleyi*: Effects of CO₂ concentration and primary production, *Geochim Cosmochim Ac*, 71, 1528-1541, 10.1016/j.gca.2006.12.015, 2007.

10 Bice, K. L., Birgel, D., Meyers, P. A., Dahl, K. A., Hinrichs, K. U., and Norris, R. D.: A multiple proxy and model study of Cretaceous upper ocean temperatures and atmospheric CO₂ concentrations, *Paleoceanography*, 21, 2006.

Bidigare, R. R., Fluegge, A., Freeman, K. H., Hanson, K. L., Hayes, J. M., Hollander, D., Jasper, J. P., King, L. L., Laws, E. A., Milder, J., Millero, F. J., Pancost, R., Popp, B. N., Steinberg, P. A., and Wakeham, S. G.: Consistent fractionation of C-13 in nature and in the laboratory: Growth-rate effects in some haptophyte algae, *Global Biogeochem Cy*, 11, 279-292, 1997.

15 Boatta, F., D'Alessandro, W., Gagliano, A. L., Liotta, M., Milazzo, M., Rodolfo-Metalpa, R., Hall-Spencer, J. M., and Parella, F.: Geochemical survey of Levante Bay, Vulcano Island (Italy), a natural laboratory for the study of ocean acidification, *Marine Pollution Bulletin*, 73, 485-494, 2013.

Bowen, G. J., and Beerling, D. J.: An integrated model for soil organic carbon and CO₂: Implications for paleosol carbonate pCO(2) paleobarometry, *Global Biogeochem Cy*, 18, Artn Gb1026 10.1029/2003gb002117, 2004.

20 Breecker, D. O., Sharp, Z. D., and McFadden, L. D.: Atmospheric CO₂ concentrations during ancient greenhouse climates were similar to those predicted for AD 2100, *P Natl Acad Sci USA*, 107, 576-580, 2010.

Brinkman, T. J., and Smith, A. M.: Effect of climate change on crustose coralline algae at a temperate vent site, White Island, New Zealand, *Marine and Freshwater Research*, 66, 360-370, 2015.

25 Castañeda, I. S., Werne, J. P., and Johnson, T. C.: Influence of climate change on algal community structure and primary productivity of Lake Malawi (East Africa) from the Last Glacial Maximum to present, *Limnology and Oceanography*, 54, 2431-2447, 2009.

Cotton, J. M., and Sheldon, N. D.: New constraints on using paleosols to reconstruct atmospheric PCO₂, *Geol Soc Am Bull*, 124, 1411-1423, 2012.

30 Dando, P. R., Stuben, D., and Varnavas, S. P.: Hydrothermalism in the Mediterranean Sea, *Progress in Oceanography*, 44, 333-367, 1999.

de Leeuw, J. W., Meer, F. W. V. D., Rijpstra, W. I. C., and Schenck, P. A.: On the occurrence and structural identification of long chain unsaturated ketones and hydrocarbons in sediments, in: *Advances in Organic Geochemistry*, edited by: Douglas, A. E., and Maxwell, J. R., Pergamon, Oxford, 211-217, 1980.

35 Fabricius, K. E., Langdon, C., Uthicke, S., Humphrey, C., Noonan, S., De'ath, G., Okazaki, R., Muehllehner, N., Glas, M. S., and Lough, J. M.: Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations, *Nat Clim Change*, 1, 165-169, 2011.

Farquhar, G. D., Oleary, M. H., and Berry, J. A.: On the Relationship between Carbon Isotope Discrimination and the Inter-Cellular Carbon-Dioxide Concentration in Leaves, *Australian Journal of Plant Physiology*, 9, 121-137, 1982.

40 Farquhar, G. D., Ehleringer, J. R., and Hubick, K. T.: Carbon Isotope Discrimination and Photosynthesis, *Annual Review of Plant Physiology and Plant Molecular Biology*, 40, 503-537, 1989.

Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Contribution of Working Group I to the Fourth Assessment

Report of the Intergovernmental Panel on Climate Change, in: Climate Change 2007: The Physical Science Basis., edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, Cambridge, United Kingdom, 131-217, 2007.

5 Foster, G. L., Royer, D. L., and Lunt, D. J.: Future climate forcing potentially without precedent in the last 420 million years, Nature Communications, 8, 2017.

Francois, R., Altabet, M. A., Goericke, R., McCorkle, D. C., Brunet, C., and Poisson, A.: Changes in the Delta-C-13 of Surface-Water Particulate Organic-Matter across the Subtropical Convergence in the SW Indian-Ocean, Global Biogeochem Cy, 7, 627-644, 1993.

10 Freeman, K. H., and Hayes, J. M.: Fractionation of Carbon Isotopes by Phytoplankton and Estimates of Ancient CO₂ Levels, Global Biogeochem Cy, 6, 185-198, 1992.

Goericke, R., and Fry, B.: Variations of Marine Plankton Delta-C-13 with Latitude, Temperature, and Dissolved CO₂ in the World Ocean, Global Biogeochem Cy, 8, 85-90, 1994.

Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D., and Buia, M. C.: Volcanic carbon dioxide vents show ecosystem effects of ocean acidification, Nature, 454, 96-99, 2008.

15 Harvey, B. P., Agostini, S., Wada, S., Inaba, K., and Hall-Spencer, J. M.: Dissolution: The Achilles' Heel of the Triton Shell in an Acidifying Ocean, Front Mar Sci, 5, 2018.

Hayes, J. M., Freeman, K. H., Popp, B. N., and Hoham, C. H.: Compound-Specific Isotopic Analyses - a Novel Tool for Reconstruction of Ancient Biogeochemical Processes, Organic Geochemistry, 16, 1115-1128, 1990.

20 Hayes, J. M.: Factors Controlling C-13 Contents of Sedimentary Organic-Compounds - Principles and Evidence, Marine Geology, 113, 111-125, 1993.

Hayes, J. M., Strauss, H., and Kaufman, A. J.: The abundance of C-13 in marine organic matter and isotopic fractionation in the global biogeochemical cycle of carbon during the past 800 Ma, Chemical Geology, 161, 103-125, 1999.

IPCC: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge, 1535, 2013.

25 Jasper, J. P., and Hayes, J. M.: A Carbon Isotope Record of CO₂ Levels during the Late Quaternary, Nature, 347, 462-464, 1990.

Jasper, J. P., Hayes, J. M., Mix, A. C., and Prahl, F. G.: Photosynthetic Fractionation of C-13 and Concentrations of Dissolved CO₂ in the Central Equatorial Pacific during the Last 255,000 Years, Paleoceanography, 9, 781-798, 1994.

30 Klok, J., Cox, H. C., Deleeuw, J. W., and Schenck, P. A.: Loliolides and Dihydroactinidiolide in a Recent Marine Sediment Probably Indicate a Major Transformation Pathway of Carotenoids, Tetrahedron Letters, 25, 5577-5580, 1984.

Laws, E. A., Popp, B. N., Bidigare, R. R., Kennicutt, M. C., and Macko, S. A.: Dependence of Phytoplankton Carbon Isotopic Composition on Growth-Rate and [CO₂](Aq) - Theoretical Considerations and Experimental Results, Geochim Cosmochim Ac, 59, 1131-1138, 1995.

35 Lewis, E., and Wallace, D.: Program developed for CO₂ system calculations, Oak Ridge National Laboratory, Oak Ridge, Tennessee, 1998.

Liaaen-Jensen, S.: Marine carotenoids, in: Marine natural products, edited by: Scheuer, P. J., Academic Press, 1978.

Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J. M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000-800,000 years before present, Nature, 453, 379-382, 2008.

- Madigan, M. T., Takigiku, R., Lee, R. G., Gest, H., and Hayes, J. M.: Carbon Isotope Fractionation by Thermophilic Phototrophic Sulfur Bacteria - Evidence for Autotrophic Growth in Natural-Populations, *Applied and Environmental Microbiology*, 55, 639-644, 1989.
- Mook, W. G. B., J.C.; Staverman, W.H.: Carbon isotope fractionation between dissolved bicarbonate and gaseous carbon dioxide, *Earth and Planetary Science Letters*, 22, 169-176, 10.1016/0012-821X(74)90078-8, 1974.
- Mouritsen, O. G., and Zuckermann, M. J.: What's so special about cholesterol?, *Lipids*, 39, 1101-1113, 2004.
- Nes, W. D., Janssen, G. G., Crumley, F. G., Kalinowska, M., and Akihisa, T.: The Structural Requirements of Sterols for Membrane-Function in *Saccharomyces-Cerevisiae*, *Archives of Biochemistry and Biophysics*, 300, 724-733, 1993.
- Pagani, M.: The alkenone- CO_2 proxy and ancient atmospheric carbon dioxide, *Philosophical Transactions of the Royal Society of London Series a-Mathematical Physical and Engineering Sciences*, 360, 609-632, 2002.
- Popp, B. N., Takigiku, R., Hayes, J. M., Louda, J. W., and Baker, E. W.: The Post-Paleozoic Chronology and Mechanism of C-13 Depletion in Primary Marine Organic-Matter, *American Journal of Science*, 289, 436-454, 1989.
- Popp, B. N., Laws, E. A., Bidigare, R. R., Dore, J. E., Hanson, K. L., and Wakeham, S. G.: Effect of phytoplankton cell geometry on carbon isotopic fractionation, *Geochim Cosmochim Ac*, 62, 69-77, 1998.
- Quast, A., Hoefs, J., and Paul, J.: Pedogenic carbonates as a proxy for palaeo- CO_2 in the Palaeozoic atmosphere, *Palaeogeogr Palaeocl*, 242, 110-125, 10.1016/j.palaeo.2006.05.017, 2006.
- Rau, G. H., Riebesell, U., and WolfGladrow, D.: A model of photosynthetic C-13 fractionation by marine phytoplankton based on diffusive molecular CO_2 uptake, *Mar Ecol Prog Ser*, 133, 275-285, 1996.
- Repeta, D. J.: Carotenoid Diagenesis in Recent Marine-Sediments .2. Degradation of Fucoxanthin to Loliolide, *Geochim Cosmochim Ac*, 53, 699-707, 1989.
- Schouten, S., Breteler, W. C. M. K., Blokker, P., Schogt, N., Rijpstra, W. I. C., Grice, K., Baas, M., and Sinninghe Damsté, J. S.: Biosynthetic effects on the stable carbon isotopic compositions of algal lipids: Implications for deciphering the carbon isotopic biomarker record, *Geochim Cosmochim Ac*, 62, 1397-1406, 1998.
- Sinninghe Damsté, J. S., Kuypers, M. M. M., Pancost, R. D., and Schouten, S.: The carbon isotopic response of algae, (cyano)bacteria, archaea and higher plants to the late Cenomanian perturbation of the global carbon cycle: Insights from biomarkers in black shales from the Cape Verde Basin (DSDP Site 367), *Org Geochem*, 39, 1703-1718, 2008.
- Stoll, H. M., Guitian, J., Hernandez-Almeida, I., Mejia, L. M., Phelps, S., Polissar, P., Rosenthal, Y., Zhang, H. R., and Ziveri, P.: Upregulation of phytoplankton carbon concentrating mechanisms during low CO_2 glacial periods and implications for the phytoplankton p CO_2 proxy, *Quaternary Sci Rev*, 208, 1-20, 10.1016/j.quascirev.2019.01.012, 2019.
- van Bentum, E. C., Reichart, G. J., and Sinnenhe Damsté, J. S.: Organic matter provenance, palaeoproductivity and bottom water anoxia during the Cenomanian/Turonian oceanic anoxic event in the Newfoundland Basin (northern proto North Atlantic Ocean), *Org Geochem*, 50, 11-18, 2012.
- Volkman, J. K., Eglinton, G., Corner, E. D. S., and Forsberg, T. E. V.: Long-Chain Alkenes and Alkenones in the Marine Coccolithophorid *Emiliania-Huxleyi*, *Phytochemistry*, 19, 2619-2622, 1980.
- Wei, J. H., Yin, X. C., and Welander, P. V.: Sterol Synthesis in Diverse Bacteria, *Frontiers in Microbiology*, 7, 2016.
- Weiss, R. F.: Carbon dioxide in water and seawater: The solubility of a non-deal gas, *Marine Chemistry*, 2, 203-215, 10.1016/0304-4203(74)90015-2, 1974.
- Witkowski, C. R., Weijers, J. W. H., Blais, B., Schouten, S., and Sinnenhe Damsté, J. S.: Molecular fossils from phytoplankton reveal secular PCO₂ trend over the Phanerozoic, *Science Advances*, 4, eaat4556, 10.1126/sciadv.aat4556, 2018.
- Zhang, Y. G., Pagani, M., Liu, Z. H., Bohaty, S. M., and DeConto, R.: A 40-million-year history of atmospheric CO₂, *Philosophical Transactions of the Royal Society a-Mathematical Physical and Engineering Sciences*, 371, 2013.

Figure Captions

Figure 1: Map of PCO₂ in the study region at Shikine Island (Japan). Top panel shows geographical context. Lower panel shows the bay on Shikine Island, with spatial variability in PCO₂ (Agostini et al., 2018), computed using the nearest neighbor algorithm in ArcGIS 10.2 software (<http://www.esri.com/software/arcgis/>).

Figure 2: GC-FID trace of silylated polar fraction. June sediment collected at the A) Control site and B) CO₂ vent, showing saturated fatty alcohols (asterisk) and sterols (square), and the representative compounds found among all sample matrices, seasons, and CO₂ concentrations: loliolide, phytol, and cholesterol.

10

Figure 3: The δ¹³C of general algal biomarkers in sediments. A) Loliolide, B) phytol, and C) cholesterol from the Control, Mid, and High PCO₂ sites during June and September from different sample matrices, including surface sediment (square), benthic diatoms (diamond), plankton tow (triangle), and macroalgae (circle).

15 **Figure 4: The ε_p of general algal biomarkers in sediments.** Loliolide (triangle), phytol (circle), and cholesterol (square) from the Control, Mid and High PCO₂ sites during June sediment collection.

Figure 5: Reconstructed PCO₂ from general algal biomarkers. PCO₂ reconstructed from the δ¹³C of loliolide (triangle), phytol (circle), and cholesterol (square) in June-collected sediments versus the actual PCO₂ measured at each location (Agostini et al., 2018; Harvey et al. 2018).

Supplementary material

Table S1. The $\delta^{13}\text{C}$ of biomarkers from different matrices. The $\delta^{13}\text{C}$ of general biomarkers loliolide, phytol, and cholesterol measured during two seasons from sea surface sediments, diatom mats, plankton net tows, and macroalgae.

5

Table S2. The $\delta^{13}\text{C}$ of CO₂. All parameters used to calculate the $\delta^{13}\text{C}$ of CO₂ (represented in the equations as δ_d), including corrections for sea surface temperature and pH.

Table S3. The $\delta^{13}\text{C}$ of algal biomarkers and all parameters used to estimate PCO₂. All parameters used to calculate PCO₂

10 from the $\delta^{13}\text{C}$ of general algal biomarkers: loliolide, phytol, and cholesterol.